

INTERNATIONAL UNION OF
PURE AND APPLIED CHEMISTRY
AND
INTERNATIONAL UNION OF BIOCHEMISTRY

**NOMENCLATURE
OF
CAROTENOIDS
(RULES APPROVED 1974)**

*Issued by the
IUPAC Commission on the Nomenclature of Organic Chemistry
and the
IUPAC-IUB Commission on Biochemical Nomenclature*

LONDON
BUTTERWORTHS

NOMENCLATURE OF CAROTENOIDS

(RULES APPROVED 1974)

*Issued by the
IUPAC Commission on Nomenclature of Organic Chemistry
and the
IUPAC-IUB Commission on Biochemical Nomenclature*

INTRODUCTION

Definitive Rules for the Nomenclature of Carotenoids were approved at the IUPAC 14th Conference (London, 1947; *Comptes Rendus*, pp 142–143) and the 16th Conference (New York, 1951; *Comptes Rendus*, p 110). Many recent developments in the carotenoid field have resulted in situations that are not covered adequately by the existing rules; since these rules were adopted, the number of known naturally occurring carotenoids has increased from some 80 to about 300. In 1965, the IUPAC Commission on Nomenclature of Organic Chemistry (CNOC) and the IUPAC-IUB Commission on Biochemical Nomenclature (CBN) approved the formation of a subcommittee to revise the carotenoid rules. The subcommittee consisted of Professor W. Klyne (representative of CBN; and Chairman), Professor S. Veibel (representative of CNOC), Professor C. Bodea, Professor T. W. Goodwin, Dr F. Haxo, Dr O. Isler, Dr S. Liaaen-Jensen, Dr W. Powell, Dr W. Quackenbush, Professor N. A. Sørensen, and Professor B. C. L. Weedon.

The subcommittee met in Trondheim in 1966 and considered a number of approaches to the naming of carotenoids. Subsequently the set of proposals selected by the Trondheim subcommittee was refined by an *ad hoc* 'working party' (S. Liaaen-Jensen, B. C. L. Weedon, W. Klyne and L. C. Cross) to give a set of tentative rules for the nomenclature of this group of naturally occurring chemical compounds that was in accord with both the general traditions of biochemical usage and principles of organic chemical nomenclature¹.

These rules are now issued in revised form¹ jointly by the IUPAC Commission on Nomenclature of Organic Chemistry² and the IUPAC-IUB

¹ Tentative Nomenclature Appendix No. 19 (February 1972) to IUPAC *Information Bulletin*. The revised rules shall be known as the IUPAC-IUB 1974 Recommendations.

² Those who have served on the Commission on the Nomenclature of Organic Chemistry for varying periods during 1967–1973 are the following (present members are shown by an asterisk): P. E. Verkade (Chairman to 1971), N. Lozac'h* (Chairman from 1971), K. Bláha*, L. C. Cross*, G. M. Dyson, H. Grünewald*, S. P. Klesney*, W. Klyne, K. L. Loening*, H. S. Nutting, J. Rigaudy*, Associate members: R. S. Cahn, K. Hirayama*, S. Veibel*. Observer: K. A. Jensen.

NOMENCLATURE OF CAROTENOIDS


Commission on Biochemical Nomenclature³. Comments should be sent to one or other of the Commission Chairmen or Secretaries⁴.

Contents

	Page
1. Definition of class of compound	408
2. The stem name	409
3. Specific names; end-group designations	409
4. Numbering of carotenoid hydrocarbons	410
5. Nor carotenoids and seco carotenoids	411
6. Changes in hydrogenation level	412
7. Oxygenated derivatives	413
8. Numbering of oxygenated derivatives	414
9. <i>retro</i> Nomenclature	415
10. Apo nomenclature	416
11. Higher carotenoids	417
12. Stereochemistry	417
13. Trivial names	418
Appendix: Naturally occurring carotenoids having trivial names (compiled by Drs O. Straub and H. Gutmann of F. Hoffmann-La Roche & Co. Limited, Basle)	

Rule Carotenoid 1. Class of compound

Carotenoids are a class of hydrocarbons (carotenes) and their oxygenated derivatives (xanthophylls) consisting of eight isoprenoid units joined in such a manner that the arrangement of isoprenoid units is reversed at the centre of the molecule so that the two central methyl groups are in a 1,6-positional relationship and the remaining nonterminal methyl groups are in a 1,5-positional relationship. All carotenoids may be formally derived from the acyclic $C_{40}H_{56}$ structure (I below), having a long central chain of conjugated double bonds, by (i) hydrogenation, (ii) dehydrogenation, (iii) cyclization, or (iv) oxidation, or any combination of these processes.




³ Those who have served on the Commission on Biochemical Nomenclature for varying periods during 1967-1973 are the following (present members are shown by an asterisk): O. Hoffmann-Ostenhof (Chairman), A. E. Braunstein*, W. E. Cohn*, J. S. Fruton, B. L. Horecker*, P. Karlson*, B. Keil*, W. Klyne*, C. Liébecq*, E. C. Slater, E. C. Webb*, W. J. Whelan, Observer: S. Veibel*.

⁴ Chairman (CNC): Professor N. Lozac'h, Ecole nationale supérieure de Chimie, 5 Avenue d'Edimbourg, 14032 Caen Cedex, France; Secretary: S. P. Klesney, Central Report Index, 566 Building, The Dow Chemical Company, Midland, Michigan 48640, U.S.A. Chairman (CBN): Professor O. Hoffmann-Ostenhof, Institut für allgemeine Biochemie der Universität Wien, Währingerstrasse 38, A1090 Vienna, Austria; Secretary: Dr. Waldo E. Cohn, Oak Ridge National Laboratory, Post Office Box Y, Oak Ridge, Tennessee 37830, U.S.A.

NOMENCLATURE OF CAROTENOIDS

This class also includes certain compounds that arise from certain rearrangements of the carbon skeleton (I), or by the (formal) removal of part of this structure.


For convenience carotenoid formulae are often written in a shorthand form as



(General formula ; broken lines indicate formal division into isoprenoid units).

Rule Carotenoid 2. The stem name

All specific names are based on the stem name 'carotene', which corresponds to the structure and numbering shown in II, where the broken lines at the two terminations are intended only to represent two 'double-bond equivalents'. Individual compounds may have C₉ acyclic end groups with two double bonds at positions 1,2 and 5,6 (e.g., III) or cyclic end groups (such as IV, V, VI, VII, VIII and IX).











Rule Carotenoid 3. Specific names; End-group designations

3.1. The name of a specific carotenoid hydrocarbon is constructed by adding two Greek letters as prefixes to the stem name 'carotene' (defined in Rule Carotenoid 2), these prefixes being characteristic of the two C₉ end groups.

3.2. The prefixes are

Type	Prefix	Formula	Structure
Acyclic	ψ	C ₉ H ₁₅	III
Cyclohexene	β, ε	C ₉ H ₁₅	IV, V
Methylenecyclohexane	γ	C ₉ H ₁₅	VI
Cyclopentane	κ	C ₉ H ₁₇	VII
Aryl	φ, χ	C ₉ H ₁₁	VIII, IX

and correspond to the following end-group modifications





NOMENCLATURE OF CAROTENOIDS

Note: The choice of locants 16 and 17 for the two methyl groups at C-1 is considered in connection with stereochemistry in Rule Carotenoid 12.4.

3.3. The Greek-letter prefixes are cited in alphabetical order; the first is separated from the second by a comma, and the second is connected to the stem name by a hyphen.

Note: The Greek-letter alphabetical order is β (beta), γ (gamma), ε (epsilon), κ (kappa), ϕ (phi), χ (chi), ψ (psi).

Examples:



Notes: (i) The Greek-letter prefixes are derived: β and ε from the symmetrical carotenoids with the trivial names 'beta-carotene' and 'epsilon-carotene', κ from the symmetrical capsorubin, γ and ψ from γ - and ψ -ionone, ϕ for phenyl, and χ , the next Greek letter after ϕ . (ii) α and δ , from the trivial names α - and δ -carotene, are not used.

3.4. When, in a modified or degraded carotenoid, an end group can be derived from more than one specific end group, the end group chosen as the basis of the name is that occurring earliest in the alphabetical order (see Rule Carotenoid 3.3).

Example:



is a derivative of the β end group, not the ε end group.


Rule Carotenoid 4. Numbering of carotenoid hydrocarbons

The basic system of numbering is that shown in structure II of Rule Carotenoid 2; the end groups are numbered as indicated in Rule Carotenoid 3. If the two end groups are dissimilar, lower (unprimed) numbers are given to that end of the molecule which is associated with the Greek-letter prefix cited first in the name. (All unprimed locants are cited before primed locants⁵.)


⁵ IUPAC Nomenclature of Organic Chemistry (*Pure Appl. Chem.* **11**, Nos. 1 and 2 (1965); Sections A, B and C published by Butterworths, London, 1971) indicates that, in ranking locants for priority, a primed numeral ranks immediately after the same numeral unprimed. In general organic nomenclature the common practice is therefore to cite locants in the order $x, x', (x + n), (x + n')$. The established sequence in the carotenoid field is, however, to cite all unprimed numerals before any primed numerals, and this practice is followed in these rules.

NOMENCLATURE OF CAROTENOIDS

Examples :



ϵ,ϵ -Carotene



ϵ,γ -Carotene


Note : It is recommended that formulae be drawn so that unprimed numbers are on the left-hand side.

Rule Carotenoid 5. Nor carotenoids and seco carotenoids


5.1. Nor Carotenoids

Elimination of a CH_3 , CH_2 , or CH group from a carotenoid is indicated by the prefix "nor", which in all cases is preceded by the locant of the carbon atom that has been eliminated. The prefix is nondetachable. When alternatives are possible, the locant attached to nor is the lowest possible⁶. The basic numbering of the carotenoid is retained in the nor carotenoid.

Examples :




2,2'-Dinor- β,β -carotene



(12,13,20 formally 'removed' at this point)

12,13,20-Trinor- β,β -carotene



16,17,16',17'-Tetranor- ϵ,ϵ -carotene


⁶ The contrast with steroid usage [IUPAC-IUB 1971 Definitive Rules for Steroid Nomenclature, *Pure Appl. Chem.* 31, Nos. 1-2 (1972)] Rule 2S-6, where the prefix 'nor' is associated with the highest permissible number, is to be noted.

NOMENCLATURE OF CAROTENOIDS

5.2. Seco Carotenoids

Fission of the bond between two adjacent carbon atoms (other than carbon atoms 1 and 6 of a cyclic end group) with addition of one or more hydrogen atoms at each terminal group thus created, is indicated by the prefix "seco," the original carotenoid numbering being retained.

Example :



Rule Carotenoid 6. Changes in hydrogenation level




Carotenoid hydrocarbons, including carotenoid acetylenes and allenes, that differ in hydrogenation level from the corresponding carotenoid hydrocarbon defined by Rule Carotenoid 3, are named from the latter by use of the prefixes "hydro" and "dehydro" together with the locants specifying the carbon atoms at which hydrogen atoms have been added or removed.

These prefixes are nondetachable⁷ and immediately precede the Greek-letter prefixes denoting the end groups, and, if both occur in one name, are cited in the order : dehydro *before* hydro (multipliers do not affect the order).

Example : tetradehydronaphthalene

Note : Since, to maintain valency requirements, hydrogen atoms are, formally, always added or removed in pairs these prefixes will always be used with an even-number multiplier, e.g. tetrahydro, didehydro.

Examples :



⁷ IUPAC Nomenclature of Organic Chemistry (reference in footnote 5), Rule C-16.1. allows the prefix 'hydro' to be detachable or nondetachable and the former has become the established usage in general organic chemistry. However, the common practice in carotenoid names is now to use this prefix as nondetachable, a practice that is followed in this set of rules.


NOMENCLATURE OF CAROTENOIDS

Rule Carotenoid 7. Oxygenated derivatives


7.1. Oxygenated (and other) derivatives of carotenoid hydrocarbons are named by use of suffixes and prefixes according to the rules of general organic chemical nomenclature⁸.

Of the oxygen-containing characteristic groups present, that occurring earliest in the sequence carboxylic acid, ester of carotenoid acid, aldehyde, ketone, alcohol, ester of carotenoid alcohol, is chosen as principal group (following IUPAC Rule C-10.3) and is cited by use of a suffix; all other groups are cited as prefixes.

Examples :




3-Hydroxy- β , ϵ -caroten-3'-one



3-Hydroxy-3'-oxo- β , ϵ -caroten-16-oic acid

7.2. A nonbridging ether group is named by use of the appropriate alkoxy or aryloxy prefix (Rule C-211.2).




1-Ethoxy-1,2,7',8'-tetrahydro- ψ , ψ -carotene


7.3. Oxygen bridges are indicated by use of the prefix 'epoxy'; this prefix is preceded by the locants of the two carbon atoms that form the bridge-heads of the oxygen bridge.

Note: The prefix 'epoxy' denotes replacement, by an oxygen bridge, of a hydrogen atom at each of two carbon atoms already otherwise connected to one another. An epoxide, notionally formed by adding an oxygen atom to a double bond, is therefore an epoxydihydro derivative of the original compound.

Examples :



5,6:5',6'-Diepoxy-5,6,5',6'-tetrahydro- β , β -carotene-3,3'-diol




5,8:5',8'-Diepoxy-5,8,5',8'-tetrahydro- β , β -carotene

⁸ IUPAC Rules, Nomenclature of Organic Chemistry (reference in footnote 5), Subsections C-2 to C-4.


NOMENCLATURE OF CAROTENOIDS

7.4. Compounds that may be formally derived from a carotenoid hydrocarbon by the addition of the elements of water (H, OH) or of methanol (H, OCH₃) to a double bond are named as 'hydroxydihydro' or 'methoxydihydro' derivatives.

Examples :



3,7'-Dihydroxy-7',8'-dihydro- β, ϵ -caroten-3'-one




7,7'-Dimethoxy-7,8,7',8'-tetrahydro- ϵ, ϵ -carotene-3,3'-dione


Rule Carotenoid 8. Numbering of oxygenated derivatives

8.1. If the two C₉ end groups of the parent carotenoid hydrocarbon are dissimilar, their oxygenated derivatives are numbered according to Rule Carotenoid 4, i.e. the end group designated by the Greek letter occurring earlier in the Greek alphabet receives unprimed locants.

Examples :



2,4-Dihydroxy- β, ϵ -caroten-3'-one



2,2',3'-Trimethoxy- β, ϵ -carotene

8.2. (a) If the two C₉ end groups of the parent carotenoid hydrocarbon are identical, then the lowest* locant possible is assigned to the principal group, cited as suffix.


(b) If more than one of the group chosen to be cited as suffix is present, the numbering is determined by the principle of lowest locants** applied to the suffixes.

(c) If no group qualifies to be cited as suffix, then the numbering is determined by the principle of lowest locants* for all groups cited as prefixes.


NOMENCLATURE OF CAROTENOIDS

Examples :

(a)




2'-Methoxy- ϵ,ϵ -caroten-3-one (not 2-Methoxy- ϵ,ϵ -caroten-3'-one)



β,β -Caroten-19,11-olide


(The position of the lactone group is indicated following the principles of Rule C-472.2)

(b)



2'-Methoxy- ϵ,ϵ -carotene-2,3'-dione [not 2-Methoxy- ϵ,ϵ -carotene-3,2'-dione ; see para (b)]

(c)



3,2'-Dimethoxy-1,2,5',6'-tetrahydro- ψ,ψ -carotene [not 2,3'-Dimethoxy-1',2',5,6-tetrahydro- ψ,ψ -carotene, 1,2,3,2', 5', 6' having priority over 2,5,6,1',2',3' ; see para (c)]

Notes : * In the carotenoid series *all* unprimed numbers are cited *before* primed numbers and the former are therefore considered as 'lower than' primed numbers, e.g. 2,6,6,1',2',6', etc. ** When series of locants containing the same number of terms are compared term by term, that series is 'lowest' that contains the lowest number on the occasion of the first difference [IUPAC Rule C-13.11(e), footnote].

Rule Carotenoid 9. *retro* Nomenclature

9.1. The prefix '*retro*' (printed in *italics*) and a pair of locants are used to indicate a shift, by one position, of *all* single and double bonds of the conjugated polyene system delineated by the pair of locants.


9.2. The pair of locants precedes the prefix *retro*. The first locant is that of the carbon atom that has lost a proton, the second that of the carbon atom that has gained one.

9.3. The prefix and its accompanying locants are placed immediately before, and hyphenated to, the Greek-letter prefixes of the name defined according to Rule Carotenoid 3.


9.4. The prefix and its associated locants are not detachable from the names defined according to Rule Carotenoid 3.

NOMENCLATURE OF CAROTENOIDS


Examples :



3-Hydroxy-6',7-retro- β , ϵ -caroten-3'-one



6',7-retro- ϵ , κ -Caroten-3-one



8',11-retro- β , ψ -Caroten-11-one

Rule Carotenoid 10. Apo nomenclature

It is often necessary to designate derivatives in which the carbon skeleton has been shortened by the formal removal of fragments from one or both ends of a carotenoid.

10.1. The unitalicized prefix 'apo', preceded by a locant, is used to indicate that all of the molecule beyond the carbon atom corresponding to that locant has been replaced by hydrogen atoms. A side-chain methyl group is not considered to be 'beyond' the carbon atom to which it is attached.


10.2. The prefix and its locant immediately precede the specific name (Rule Carotenoid 3) unless the locant associated with the prefix 'apo' is greater than 5, in which case there is no need to give a Greek-letter end-group designation for that end of the molecule.

10.3. For purposes of numbering, etc., an end that has been shortened by 5 or less skeletal carbon atoms is considered a ψ (acyclic) end group.


10.4. The prefix diapo, preceded by two locants, is used to indicate removal of fragments from both ends of the molecule.

10.5. If, in a diapo compound, the two ends of the carotenoid skeleton have been shortened unequally, the lower locant associated with the prefix 'diapo' is unprimed.

Examples :



2'-Apo- β , ψ -caroten-2'-al





Methyl hydrogen 6,6'-diapocarotene-6,6'-dioate (trans-Bixin)

NOMENCLATURE OF CAROTENOIDS

Rule Carotenoid 11. Higher carotenoids


The higher carotenoids are a class of hydrocarbons and their oxygenated derivatives consisting of *more than* eight isoprenoid units joined in a manner similar to that of the C₄₀ carotenoids. They are named as mono- or disubstituted C₄₀ carotenoids. The numbering of the normal carotenoid is retained. Examples:



Rule Carotenoid 12. Stereochemistry


12.1. Absolute Configuration at Chiral Centers. The absolute configuration at chiral centers is designated by use of the RS convention⁹, the symbols being placed, with the corresponding locants, before the carotenoid name.

Example:



(3S,5R,3'S,5'R)-3,3'-Dihydroxy- ϵ, ϵ -carotene-6,6'-dione (Capsorubin)

12.2. Absolute Configuration of Allenic Compounds. The absolute configuration around allene groups will be similarly designated, when known.



(3'S,5'R,6'R)-3'-Acetoxy-5,6-epoxy-3,5'-dihydroxy-6,7'-didehydro-5,6,7,8,5',6'-hexahydro- β, β -caroten-8-one (Fucoxanthin)


12.3. Geometrical Configuration around Double Bonds. The stem name 'carotene' implies *trans* configuration about all double bonds unless the contrary is indicated. Following the designation of absolute configuration

⁹ For a discussion on the RS convention and the use of thickened lines, broken lines, wavy lines, and wedges in displayed formulae, see IUPAC Tentative Rules for the Nomenclature of Organic Chemistry, Section E, Fundamental Stereochemistry, IUPAC *Information Bulletin* No. 35, p 68.

NOMENCLATURE OF CAROTENOIDS

(if any), geometrical configuration is indicated by citing the double bond or bonds with a *cis* configuration.

Example:



Natural phytoene = 15-*cis*-7,8,11,12,7',8',11',12'-octahydro- β,β -carotene


At trisubstituted double bonds, the term *cis* refers to the relative position of the two substituents forming parts of the main chain of carbon atoms.

Example: natural bixin = methyl hydrogen 9'-*cis*-6,6'-diapocarotene-6,6'-dioate.


In the absence of definite information on geometrical configuration, *cis* isomers may be distinguished by prefixes such as neo A, neo U, etc. (cf. Zechmeister, "Cis-Trans Isomeric Carotenoids, Vitamins A and Aryl-polyenes," Springer-Verlag, Vienna, 1962).

The stereochemical prefixes *E* and *Z* (see footnote to Rule Carotenoid 12.1 and *J. Amer. Chem. Soc.* 90, 509 (1968)) may be used, especially when the prefixes *cis* and *trans* might lead to ambiguity.

12.4. Numbering of *gem*-Dimethyl Groups at C-1. In an end group of β , γ , κ or ϵ type, the two methyl groups attached to C-1 are distinguished as follows: when the potential chirality is as shown in formula A, i.e. with the polyene chain (R) to the right of C-1, the methyl groups *below* and *above* the plane of the paper are numbered 16 and 17, respectively; if, with the polyene chain (R) to the left, the end group is as shown in B¹, then these designations are unaltered; if the end group is as shown in B², then they are reversed.



In an acyclic end group, the methyl group that is *trans* to the main skeletal chain is numbered 16 and the methyl group that is *cis* is numbered 17, as shown in C.



Rule Carotenoid 13. Trivial names

The preceding rules are designed to define precisely the structure of a given carotenoid by its name. Use of the semisystematic names derived from these rules will greatly assist communication between scientists and enable work to be more readily retrieved from the literature.

NOMENCLATURE OF CAROTENOIDS

The appendix contains a list of some trivial names currently in use for naturally occurring carotenoids; it also gives their semisystematic names and structures. These trivial names will be of value in natural-product and biochemical work, but their use in systematic organic work should be restricted. If trivial names are used in a paper, the semisystematic names should always be clearly indicated.


While the need to coin a new trivial name must occasionally arise (e.g. because the structure of the compound is unknown), the list in the appendix should not be unnecessarily enlarged. Simple derivatives of known carotenoids should not be given new trivial names; they may, however, be named by modification of existing trivial names (c.f. Steroid Rule 2S-4.2, Note)¹⁰, but the full semisystematic name according to these Rules should be given for each compound at the first mention in each paper.

¹⁰ For reference to the Definitive Rules for Steroid Nomenclature, see footnote 6.




APPENDIX

Some Naturally Occurring Carotenoids Having Trivial Names. (The examples in this appendix are based on an exhaustive list of naturally occurring carotenoids prepared by Dr O. Straub, which forms Chapter XII of a book "Carotenoids," O. Isler, Ed., Birkhäuser, Basel and Stuttgart, 1971. The two commissions are greatly indebted to Dr Isler, Dr Straub, and their colleagues of F. Hoffmann-La Roche Co. Ltd., Basel, for providing this list of compounds. The list includes a few compounds whose natural occurrence or structure is doubtful, but whose systematic names illustrate particular points of nomenclature.)

Hydrocarbons


3',4'-Didehydro- β,ψ -carotene

Torulene


7',8'-Dihydro- α,ψ -carotene α -Zeacarotene15-cis-7,8,11,12,7',8'-Hexahydro- ψ,ψ -carotenePhytofluene
Hexahydrolycopene7,8,11,12,15,7',8',11',12',15'-Decahydro- ψ,ψ -carotene

Lycopersene


Alcohols

(3*R*)-5'-*cis*- β,ψ -caroten-3-ol


Gazaniaxanthin

1',2'-Dihydro- ϕ,ψ -caroten-1'-ol


OH-Chlorobactene

1,2-Dihydro- ψ,ψ -caroten-1-ol


Rhodopin

 ψ,ψ -Caroten-16-ol


Lycoxanthin

3',4'-Didehydro-1',2'-dihydro- β,ψ -carotene-3,1'-diol

Saproxanthin


13-*cis*-1,2-Dihydro- ψ,ψ -carotene-1,20-diolRhodopinol
Warmingol

$(3R,3'R)-7,8,7',8'-Tetrahydro-\beta,\beta\text{-carotene-3,3'-diol}$




Alloxanthin
Cynthiaxanthin
Pectenoxanthin
Cryptomonaxanthin

$\beta,\beta\text{-Carotene-3,4,3',4'-tetrol}$



Crustaxanthin


$\beta,\epsilon\text{-Carotene-3,19,3'-triol}$



Loroxanthin


Glycosides

$1'-(\beta\text{-D-Glucopyranosyloxy})-3',4'-didehydro-1',2'-dihydro-\beta,\psi\text{-carotene-2'-ol}$



Phleixanthophyll


$2,2'\text{-Bis}(\beta\text{-L-rhamnopyranosyloxy})-3,4,3',4'\text{-tetrahydro-1,2,1',2'-tetrahydro-\psi,\psi\text{-carotene-1,1'-diol}$



Oscillaxanthin


Ethers

$1\text{-Methoxy-3,4-didehydro-1,2,7',8'-tetrahydro-\psi,\psi\text{-carotene}}$



Spheroidene


$1\text{-Methoxy-3',4'-didehydro-1,2,1',2'-tetrahydro-\psi,\psi\text{-caroten-1-ol}}$



Rhodovibrin


Epoxides

5',6-Epoxy-7',8'-didehydro-
5,6-dihydro- β,β -carotene-
3,3'-diol




Diadinoxanthin

5',6'-Epoxy-6,7-didehydro-
5,6,5',6'-tetrahydro- β,β -caro-
tene-3,5,19,3'-tetrol




Vaucheraxanthin

5,8-Epoxy-5,8-dihydro- β,β -caro-
tene-3,3'-diol




Mutatoxanthin
Citroxanthin
Zeaxanthin furanoxide

5',8'-Epoxy-6,7-didehydro-
5,6,5',8'-tetrahydro- β,β -
carotene-3,5,3'-triol




Neochrome
Foliachrome
Trollichrome

5,6:5',8'-Diepoxy-5,6,5',8'-tet-
rahydro- β,β -carotene-3,3'-diol









Luteoxanthin


3',4'-Didehydro- β,ψ -caroten-
16'-al




Aldehydes

Torularhodinaldehyde


1-cis-1-Hydroxy-1,2-dihydro- β,ψ -caroten-20-al**Rhodopinal**
Warmingone*3',4'-Didehydro- β,ψ -caroten-16'-oic acid***Torularhodin***Methyl 3',4'-didehydro- β,ψ -caroten-16'-oate***Torularhodin methyl ester***β,ε-Caroten-4-one***Phoenicopterone***β,β-Carotene-4,4'-dione***Canthaxanthin**
Aphanicin
Chlorellaxanthin*3-Hydroxy- β,ψ -caroten-4'-one***Rubixanthone**

1'-Methoxy-1',2'-dihydro- χ,ψ -caroten-4'-one


Okenone

1'-Hydroxy-1-methoxy-3,4-didehydro-1,2,-1',2',7',8'-hexahydro- ψ,ψ -caroten-2-one


Hydroxyspheroidenone

1,1'-Dimethoxy-3,4,3',4'-tetrahydro-1,2,1',2'-tetrahydro- ψ,ψ -carotene-2,2'-dione


2,2'-Diketospirilloxanthin

3,19,3'-Trihydroxy-7,8-dihydro- β,ϵ -caroten-8-one


Siphonaxanthin

3,1'-Dihydroxy-3',4'-didehydro-1',2'-dihydro- β,ψ -caroten-4-one

Flexixanthin


(3'S,5'R)-3'-Hydroxy- β,κ -caroten-6'-one

Cryptocapsin

(3R,3'S,5'R)-3,3'-Dihydroxy- β,κ -caroten-6'-one


Capsanthin

(3*S*,5*R*,3'*S*,5'*R*)-3,3'-Dihydroxy- κ,κ -carotene-6,6'-dione




Capsorubin

3,3'-Dihydroxy-7',8'-didehydro- β,β -carotene-4-one




Pectenolone

3-Hydroxy- β,β -carotene-4,4'-dione



Adonirubin
Phenicoxanthin
3-OH-Canthaxanthin


3-Hydroxy-2,3-didehydro- β,β -carotene-4,4'-dione



Phoeniconone
Dehydroadonirubin


Esters of Alcohols

(3*R*,3'*R*)-3,3'-Bispalmitoyloxy- β,β -carotene or (3*R*,3'*R*)- β,β -carotene-3,3'-diol dipalmitate




Zeaxanthin
dipalmitate
Physalien

3,3'-Bispalmitoyloxy-2,3,2',3'-tetrahydro- β,β -carotene-4,4'-dione or 3,3'-dihydroxy-2,3,2',3'-tetrahydro- β,β -carotene-4,4'-dione dipalmitate




Astacein

3,3'-Dihydroxy-19-lauroyloxy-7,8-dihydro- β,ϵ -caroten-8-one or 3,19,3'-trihydroxy-7,8-dihydro- β,ϵ -caroten-8-one 19-laurate




Siphonein

3'-Acetoxy-3,5,5'-trihydroxy-6',7'-didehydro-5,8,5',6'-tetrahydro- β,β -caroten-8-one



Isofucoxanthin


3'-Acetoxy-5,6-epoxy-3,5'-dihydroxy-6',7'-didehydro-5,6,7,8,5',6'-hexahydro- β,β -caroten-8-one



Fucoxanthin


Apo Carotenoids

3',4'-Didehydro-2'-apo- β -caroten-2'-al




β -Apo-2'-carotenal

5,6-Dihydroxy-5,6-dihydro-10'-apo- β -caroten-10'-al












Azafrinaldehyde


6'-Apo- ψ -caroten-6'-al






Apo-2'-lycopenal
Apo-6'-lycopenal

Methyl 6'-apo- ψ -caroten-6'-oate**Methyl apo-6'-lycopenoate****5'6'-Dihydro-5'-apo-18'-nor- β -caroten-6'-one or 6'-methyl-6'-apo- β -caroten-6'-one****Citranaxanthin****7',8'-Dihydro-7'-apo- β -caroten-8'-one or 8'-methyl-8'-apo- β -caroten-8'-one****Sintaxanthin****3,5-Dihydroxy-6,7-dihydro-5,6,7',8'-tetrahydro-7'-apo- β -caroten-8'-one or 3,5-dihydroxy-8'-methyl-6,7-dihydro-5,6-dihydro-8'-apo- β -caroten-8'-one****Paracentrone****3-Hydroxy-7,8-dihydro-7',8'-dihydro-7'-apo- β -carotene-4,8'-dione or 3-hydroxy-8'-methyl-7,8-dihydro-8'-apo- β -carotene-4,8'-dione****Hopkinsxanthin****6'-Methyl hydrogen 9'-cis-6,6'-diapocarotene-6,6'-dioate****Bixin****8'-Oxo-8,8'-diapo caroten-8-oic acid****Crocetinsemialdehyde****8,8'-Diapocarotene-8,8'-dioic acid****Crocetin****Digentiobiosyl 8,8'-diapo caroten-8,8'-dioate****Crocin**



Nor and Seco Carotenoids

3,3'-Bisacyloxy-2,2'-dinor- β,β -carotene-4,4'-dione


Actinioerythrin

5,6-Seco- β,β -carotene-5,6-dioneSemi- β -carotenone5,6-Seco- β,ϵ -carotene-5,6-dioneSemi- α -carotenone5,6:5',6'-Diseco- β,β -carotene-5,6,5',6'-tetrone


β-Carotenone

3'-Hydroxy-5,6-seco- β,β -carotene-5,6-dioneTriphasiaxanthin
3-Hydroxysemi- β -carotenone3'-Acetoxy-5,6-epoxy-3,5'-dihydroxy-6',7'-didehydro-5,6,5',6'-tetrahydro-12',13',20'-trinor- β,β -carotene-19,11-olide


Peridinin

5,6-Epoxy-3,3'-dihydroxy-7',8'-didehydro-5,6-dihydro-12',13',20'-trinor- β,β -carotene-19,11-olide


Pyrrhoxanthinol

retro Carotenoids and *retro* Apo Carotenoids4',5'-Didehydro-4,5'-*retro*- β,β -carotene-3,3'-diol


Eschscholtzxanthin

3'-Hydroxy-4',5'-didehydro-4,5'-*retro*- β,β -carotene-3-one


Eschscholtzxanthone

4',5'-Didehydro-4,5'-*retro*- β,β -carotene-3,3'-dione

Rhodoxanthin


7,8-Dihydro-8,7'-*retro*- β,β -carotene or 7,7'-dihydro- β,β -carotene

No trivial name


3-Hydroxy-5'-methyl-4,5'-*retro*-5'-apo- β -caroten-5'-one or 3-hydroxy-4,5'-*retro*-5'-apo- β -caroten-5'-one

Tangeraxanthin

Higher Carotenoids


2-(4-Hydroxy-3-methyl-2-butenyl)-7',8',11',12'-tetrahydro- ϵ,ψ -carotene

Nonaprenoxanthin

2,2'-Bis(4-hydroxy-3-methyl-2-butenyl)- ϵ,ϵ -carotene


Decaprenoxanthin

2-[4-Hydroxy-3-(hydroxymethyl)-2-butenyl]-2'-(3-methyl-2-butenyl)- β,β -carotene




C.p. 450

2'-(4-Hydroxy-3-methyl-2-butenyl)-2-(3-methyl-2-butenyl)-3',4'-didehydro-1',2'-dihydro- β,ψ -caroten-1'-ol



C.p. 473

2,2'-Bis(3-hydroxy-3-methylbutyl)-3,4,3',4'-tetradehydro-1,2,1',2'-tetrahydro- ψ,ψ -carotene-1,1'-diol



Bacterioruberin