




The simplest natural product prepared from the telomer 2 is Matsutake alcohol 4, a fragrant compound contained in a Japanese mushroom (6). The synthesis was achieved by highly selective reduction of one of the terminal double bonds at C₇ without attacking the other one at C₁. Hydroalumination catalyzed by a titanium compound was used for the selective reduction (7). Formation of alkoxyaluminum compound 5 by the treatment of 1,7-octadien-3-ol 3 with one mole of dibutylaluminum hydride protects the hydroxy group and also gives steric hindrance to the neighbouring double bond. Addition of LiAlH₄ in the presence of a catalytic amount of Cp₂TiCl₂ caused the selective hydroalumination of the terminal olefin. The hydrolysis afforded 1-octen-3-ol (Matsutake alcohol 4) in a high yield with 90% selectivity.




The alcohol 3 is a good starting material for the synthesis of lipoic acid 10 without changing carbon numbers (8). Hydroboration of the two double bonds produced the triol 6. The 1,3-diol system was protected by acetal formation 7 and the terminal free alcohol was oxidized to the carboxylic acid 8. The conversion of the 1,3-diol to 1,3-dithiol is a known reaction. The oxidation of the dithiol 9 catalyzed by FeCl₃ afforded lipoic acid 10.




We have explored a method for macrolide synthesis based on intramolecular carbon-carbon bond formation using butadiene telomers as starting materials. The acetate 1 has a suitable functionality for the synthesis of diploidalide B 14 (9). At first the acetate was converted to tetrahydropyranyl ether, and the terminal olefin was oxidized to the methyl ketone 11 with PdCl₂/CuCl/O₂ and then reduced to alcohol. The bromoacetate 12 was formed by the reaction of the alcohol with bromoacetyl bromide. The tetrahydropyranyl ether was removed and the alcohol was oxidized with CrO₃ to the aldehyde 13 in 60% yield (10). The final step is the intramolecular Reformatsky reaction to form diploidalide B 14 in 45% yield. The cyclization was promoted by Zn in the presence of AlEt₂Cl as an activator (11). In this reaction, aluminum protects the hydroxy group formed by the reaction from dehydration.




The ring chain of the ten-membered lactones can be conveniently prepared from the acetate 1 (12). Again the terminal double bond was oxidized to the methyl ketone 15 and then reduced to the alcohol. The allylic acetate was converted, after hydrolysis, to the allylic chloride 16 by the treatment with $\text{CCl}_4/\text{PPh}_3$. The ester 17 was prepared by the reaction of phenylthioacetyl chloride. The intramolecular alkylation of the anion generated by $\text{KN}(\text{SiMe}_3)_2$ as a base afforded the unsaturated lactone 18 in 71% yield. The saturated and unsaturated lactones 19 and 20 were prepared from 18.



The ring chain of lasiodiplodin 24 can be conveniently derived from the allylic chloride 16 (13). The ester 22 was prepared from 16 and 21 and the intramolecular alkylation of the anion generated by $\text{KN}(\text{SiMe}_3)_2$ afforded the unsaturated lactone 23 in 40% yield. The reduction of the double bond and removal of the phenylthio group were achieved by the treatment with Raney nickel to give the methyl derivative of lasiodiplodin 24.




Zearalenone is another type of orsellinic acid type macrolide and its ring chain was prepared easily from the acetate 2 by the following sequence of reaction (13). The allylic alcohol 25 was oxidized to the enone 26, which underwent Michael reaction with malonate to give the diester 27 in 70% yield. One of the ester was removed by heating in HMPA and the ketone was protected to give 28. The ester was reduced to the alcohol and converted to tosylate. The terminal olefin was oxidized with $\text{PdCl}_2/\text{CuCl}/\text{O}_2$ to give the methyl ketone 29. The tosylate was converted to the iodide and the ketone was reduced to the alcohol 30, which was esterified with 21 to give the ester 31. The cyclization was carried out as before to give the lactone 32 in 50% yield. Oxidative removal of the phenylthio group and deketalization produced dimethyl ether of zearalenone 33.




Elimination reactions of allylic acetates and phenyl ethers


We found that allylic acetates and allylic phenyl ethers can be converted to conjugated dienes by the treatment with a catalyst system of Pd(OAc)₂/PPh₃ at 100° with expulsion of acetic acid and phenol (14).



When these allylic compounds were treated with alkylammonium formate in the presence of Pd(OAc)₂/PPh₃ as the catalyst, reductive elimination of acetic acid and phenol took place to give olefins (15). Interestingly, the olefins produced from both isomeric 1- and 3-oxyallylic compounds, were predominantly terminal olefins.




The above shown catalytic reaction was applied to the synthesis of 12-acetoxy-1,3-dodecadiene 39, a pheromone of *Diparopsis castanea* (16). Starting from 8-phenoxy-1,6-octadiene 34, easily prepared by the reaction of butadiene and phenol, a four carbon unit was added. The alcohol 35a was formed by selective hydroboration of the terminal double bond and tosylated. The tosylate was converted to iodide 35b with sodium iodide. The iodide 35b was coupled in 80% yield with the Grignard reagent 36 prepared from THF using CuI/bipyr as a catalyst system. Then the phenyl ether 37 was converted to the diene 38 with Pd(OAc)₂/PPh₃ in 71% yield. The acetylation completes the synthesis of the pheromone 39.




Telomers of nitroalkanes


Active hydrogens of nitroalkanes are displaced in a stepwise-manner with the octadienyl chain by the catalysis of $\text{Pd}(\text{OAc})_2/\text{PPh}_3$ at room temperature to give long chain unsaturated nitro compounds (17).




The telomer of nitroethane **40** was used for the synthesis of Recifeiolide **44**, a naturally occurring 12-membered lactone (12, 18). The usefulness of the compound **40** is apparent by the comparison of **40** with **44**. The internal double bond of **40** is at the right position with the *trans* configuration, and the nitro group can be converted to ketone easily. The nitro group was converted to the ketone by Nef reaction ($\text{MeONa}/\text{TiCl}_4$) and protected by ketal formation. The terminal double bond was selectively hydroaluminated with LiAlH_4 using TiCl_4 and quenched with iodine to give the iodide. The protected ketone was liberated and reduced to the alcohol **41**. The phenylthioacetate **42** was prepared by the treatment of the alcohol **41** with phenylthioacetyl chloride. The intramolecular alkylation of **42** using $\text{KN}(\text{SiMe}_3)_2$ afforded the lactone **43** in 75% yield. The phenylthio group was removed with deactivated Raney nickel to give Recifeiolide **44** in 80% yield.



9-Nitro-1,6,11,16-heptadecatetraene 45 is one of the telomerization product of butadiene and nitromethane. The telomer 45 has a linear 17 carbon chain and the nitro group at the center of the carbon chain. Civetonedicarboxylic acid 46, a precursor of civetone, was prepared by the following scheme without changing the carbon numbers.




The nitro group was converted to ketone by the Nef reaction ($\text{MeONa}/\text{H}_2\text{SO}_4$) and protected as a ketal. The terminal double bonds were selectively hydroborated with 9-BBN and the internal double bonds were hydrogenated. Then the terminal alcohols were oxidized to carboxylic acids to afford civetone-dicarboxylic acid 46. Civetonedicarboxylate was cyclized by acyloin condensation to give the acyloin 47 in 65% yield. The ketol was oxidized to α -diketone and then converted to dihydrazone 48, which was oxidized to the acetylene 49 with CuCl in pyridine. The hydrogenation catalyzed by Lindler catalyst afforded *cis*-civetone 50 (19).




3,8-Nonadienoate


Carbonylation of butadiene in alcohol catalyzed by $\text{Pd}(\text{OAc})_2/\text{PPh}_3$ affords 3,8-nonadienoate 51 in a high yield (20).



Synthesis of 2-decenedioic acid 53 (royal jelly acid) was carried out by the carbonylation of 51 and the double bond migration (21). $\text{Co}_2(\text{CO})_8/\text{pyridine}$ was used as a catalyst, and the linear diester 52 was obtained as a main product. The hydrolysis and double bond migration were achieved by the treatment of the diester with strong alcoholic KOH to give royal jelly acid 53 as a crystalline compound.



The terminal double bond of **51** was oxidized with $\text{PdCl}_2/\text{CuCl}/\text{O}_2$ to the methyl ketone **54** and the double bond was hydrogenated. Hydrolysis of the ester afforded 8-oxononanoic acid **55**, which was subjected to Kolbe electrolysis. By this way, 1,15-hexadecanedione **56** was obtained in a high yield (22). Cyclization of this dione and hydrogenation produced muscone **57**.



Synthesis of Brevicommin from **51** has been reported (23).

REFERENCES

1. J. Smidt, W. Hafner, R. Jira, R. Sieber, J. Sedlmeier, and A. Sabel, *Angew. Chem. Intern. Ed.* **1**, 80-88 (1962).
2. W. H. Clement and C. M. Selwitz, *J. Org. Chem.* **29**, 241-243 (1964); W. G. Lloyd and B. J. Juberoff, *J. Org. Chem.* **34**, 3949-3952 (1969).
3. J. Tsuji, I. Shimizu, and K. Yamamoto, *Tetrahedron Lett.* 2975-2976 (1976).
4. J. Tsuji, *Accounts Chem. Res.* **6**, 8-15 (1973).
5. S. Takahashi, T. Shibano, and N. Hagihara, *Tetrahedron Lett.* 2451-2453 (1967).
6. J. Tsuji and T. Mandai, *Chem. Lett.* 975-976 (1977).
7. F. Sato, S. Sato, and M. Sato, *J. Organometal. Chem.* **122**, C25-27 (1976); **131**, C26-28 (1977).
8. J. Tsuji, H. Yasuda, and T. Mandai, *J. Org. Chem.*, in press.
9. T. Ishida and K. Wada, *Chem. Commun.* 209-210 (1975).
10. J. Tsuji and T. Mandai, *Tetrahedron Lett.* 1817-1820 (1978).
11. K. Maruoka, S. Hashimoto, Y. Kitagawa, H. Yamamoto, and H. Nozaki, *J. Am. Chem. Soc.* **99**, 7705-7709 (1977).
12. T. Takahashi, K. Kasuga, S. Hashiguchi, and J. Tsuji, in press.
13. T. Takahashi, K. Kasuga, and J. Tsuji, in preparation.
14. J. Tsuji, T. Yamakawa, M. Kaito, and T. Mandai, *Tetrahedron Lett.* 2075-2078 (1978).
15. J. Tsuji, T. Yamakawa, and T. Mandai, in preparation.
16. T. Mandai, H. Yasuda, M. Kaito, J. Tsuji, R. Yamaoka, and H. Fukami, *Tetrahedron*, in press.
17. T. Mitsuyasu and J. Tsuji, *Tetrahedron* **30**, 831-834 (1974).
18. J. Tsuji, T. Yamakawa, and T. Mandai, *Tetrahedron Lett.* 565-568 (1978).
19. J. Tsuji and T. Mandai, *Tetrahedron Lett.* 3285-3286 (1977).
20. J. Tsuji, Y. Mori, and M. Hara, *Tetrahedron* **28**, 3721-3725 (1972).
21. J. Tsuji and H. Yasuda, *J. Organometal. Chem.* **131**, 133-135 (1977).
22. J. Tsuji, M. Kaito, Y. Yamada, and T. Mandai, *Bull. Chem. Soc. Jpn.* **51**, 1915-1916 (1978).
23. N. T. Byron, R. Grigg, and B. Kongathip, *Chem. Comm.* 216-217 (1976).