

Contents

Preface — V

Preface to the third edition — VII

Preface to the second edition — IX

About the authors — XI

1 Crystal density — 1

- 1.1 Group additivity method — 2
 - 1.1.1 The method of atomic contributions — 7
 - 1.1.2 Benzene-derived energetic compounds using atomic volumes — 7
 - 1.1.3 The method of group additivity for estimating densities of energetic ionic liquids and salts — 9
- 1.2 Quantum mechanical approach — 11
 - 1.2.1 Quantum mechanical approach for neutral energetic compounds — 11
 - 1.2.2 Energetic ionic liquids and salts as room-temperature energetic materials — 13
- 1.3 Empirical methods for the calculation of the crystal density of different classes of energetic materials — 14
 - 1.3.1 Nitroaromatic energetic compounds — 14
 - 1.3.2 Acyclic and cyclic nitramines, nitrate esters, and nitroaliphatic compounds — 16
 - 1.3.3 Improved method for the prediction of the crystal densities of nitroaliphatics, nitrate esters, and nitramines — 18
 - 1.3.4 Reliable correlation for the prediction of the crystal densities of polynitroarenes and polynitroheteroarenes — 19
 - 1.3.5 The extended correlation for the prediction of the crystal density of energetic compounds — 21
 - 1.3.6 Energetic azido compounds — 23
 - 1.3.7 High-nitrogen-content organic compounds — 24
- 1.4 Empirical methods for the assessment of the crystal density of hazardous ionic molecular energetic materials using the molecular structures — 31
 - 1.4.1 Two general empirical methods — 32
 - 1.4.2 The effects of various substituents on the density of tetrazolium nitrate salts — 33
 - 1.4.3 Predicting the density of tetrazole-*N*-oxide salts — 34
 - 1.4.4 High-nitrogen-containing salts and ionic liquids (HNCSILs) — 35
- 1.5 Machine learning for predicting properties of energetic materials — 39

1.5.1	Machine learning for predicting density in energetic materials —	39
1.5.2	Limitations of ML in predicting energetic compound densities —	41
1.6	Summary —	42
2	Heat of formation —	44
2.1	Gas-phase heats of formation of energetic compounds —	44
2.1.1	The evolution of computational thermochemistry from semiempirical to composite quantum methods —	45
2.1.2	ML advances in predicting formation enthalpies outperform traditional quantum methods —	46
2.2	Condensed phase heats of formation of energetic compounds —	48
2.2.1	Thermochemical prediction of condensed phase heats of formation in energetic materials —	49
2.2.2	Empirical approaches or QSPR methods on the basis of structural parameters —	51
2.3	Energetic compounds with high nitrogen contents —	63
2.3.1	Using the molecular structure —	64
2.3.2	Gas-phase information —	64
2.4	The condensed phase heat of formation of energetic ionic liquids and salts —	66
2.4.1	Complex approach —	66
2.4.2	QSPR methods —	67
2.5	Summary —	73
3	Melting point —	74
3.1	Group additivity, QSPR, ML, and quantum mechanical methods —	75
3.2	Simple empirical methods on the basis of molecular structure —	78
3.2.1	Nitroaromatic compounds —	79
3.2.2	Polynitroarene and polynitroheteroarene compounds —	80
3.2.3	Nitramines, nitrate esters, nitrate salts, and nitroaliphatics —	82
3.2.4	Nonaromatic energetic compounds —	82
3.2.5	Improved method for predicting the melting points of energetic compounds —	84
3.2.6	Organic molecules containing hazardous peroxide groups —	87
3.2.7	Organic azides —	89
3.2.8	General method for the prediction of melting points of energetic compounds including organic peroxides, organic azides, organic nitrates, polynitroarenes, polynitroheteroarenes, acyclic and cyclic nitramines, nitrate esters, and nitroaliphatic compounds —	90
3.2.9	Cyclic saturated and unsaturated hydrocarbons —	99
3.3	Melting points of ionic liquids —	103
3.3.1	Group additivity approach —	104

3.3.2	QSPR approaches based on complex descriptors —	109
3.3.3	Simple approach based on the structure of cations and anions —	111
3.4	Summary —	114
4	Enthalpy and entropy of fusion —	116
4.1	Different approaches for the prediction of the enthalpy of fusion —	116
4.1.1	Group additivity method for prediction of the enthalpy of fusion —	117
4.1.2	Nitroaromatic carbocyclic energetic compounds —	119
4.1.3	Nitroaromatic energetic compounds —	120
4.1.4	Nonaromatic energetic compounds containing nitramine, nitrate, and nitro functional groups —	121
4.1.5	Improved method for the reliable prediction of the enthalpy of fusion of energetic compounds —	122
4.1.6	A reliable method to predict the enthalpy of fusion of energetic materials —	125
4.1.7	A generalized predictive model for enthalpy of fusion in diverse hydrocarbons —	127
4.2	Different methods to predict the entropy of fusion —	134
4.3	A group contribution approach incorporating cation-anion interactions and structural effects for ionic liquids —	138
4.4	Summary —	141
5	Heat of sublimation —	142
5.1	Group additivity method for prediction of the heat of sublimation —	143
5.2	Quantum mechanical and complex approaches for predicting the heat of sublimation —	151
5.3	The use of structural parameters —	152
5.3.1	Nitroaromatic compounds —	152
5.3.2	Nitramines —	153
5.3.3	Nitroaromatics, nitramines, nitroaliphatics, and nitrate esters —	154
5.3.4	General method for polynitroarenes, polynitroheteroarenes, acyclic and cyclic nitramines, nitrate esters, nitroaliphatics, cyclic and acyclic peroxides, as well as nitrogen-rich compounds —	155
5.4	Summary —	158
6	Impact sensitivity —	159
6.1	Complex computational methods —	160
6.2	Advances in modeling impact sensitivity: phonon-vibration coupling and multi-phonon interactions in energetic materials —	166
6.3	Simple methods on the basis of molecular structure for neutral energetic compounds —	168
6.3.1	Oxygen balance correlations —	169

6.3.2	Elemental composition and molecular moieties —	169
6.4	Impact sensitivity of quaternary ammonium-based energetic ionic liquids or salts —	181
6.5	Quantitative impact sensitivity prediction: multiplicative incremental theory for energetic materials —	183
6.5.1	Aromatic systems exhibit distinct behavior —	191
6.5.2	Understanding symmetry in fused ring systems —	193
6.5.3	Identifying trigger bonds in symmetric structures —	193
6.5.4	Handling complex cases: when structures defy simple analysis —	194
6.5.5	Breaking down the approach —	194
6.5.6	When the rules don't apply: unexplained cases in impact sensitivity —	196
6.5.7	The puzzling case of compounds (I)–(IV) —	196
6.5.8	Nitrogen's subtle influence: compounds (II) vs. (V) —	196
6.5.9	The azo-linked surprise: compounds (VI)–(VII) —	198
6.5.10	The curious case of salt (VIII) —	198
6.5.11	Looking ahead —	198
6.6	Summary —	198
7	Electric spark sensitivity —	200
7.1	Measurement of electric spark sensitivity —	200
7.2	Different methods for predicting electric spark sensitivity —	201
7.3	Simple methods for predicting electrostatic spark sensitivity based on the RDAD instrument —	202
7.3.1	Polynitroaromatic compounds —	202
7.3.2	Cyclic and acyclic nitramines —	203
7.3.3	General correlation for polynitroaromatics as well as cyclic and acyclic nitramines —	204
7.4	Simple prediction of electrostatic spark sensitivity based on the new ESZ KTTV instrument —	205
7.4.1	Polynitroarenes based on ESZ KTTV —	205
7.4.2	Nitramines based on ESZ KTTV —	207
7.4.3	Quaternary ammonium-based energetic ionic liquids or salts based on ESZ KTTV —	209
7.5	Electrostatic discharge (ESD) sensitivity of nitrogen-rich heterocyclic energetic compounds (NRHECs) —	211
7.5.1	Influence of functional groups on ESD sensitivity —	211
7.5.2	Predictive modeling via MLR —	212
7.5.3	Practical implications —	213
7.6	Some aspects of predictive methods —	214
7.7	Summary —	214

8 Shock sensitivity — 216

- 8.1 Small-scale gap test — 217
- 8.2 Large-scale gap test — 219
- 8.3 Critical diameter of solid pure and composite high explosives — 220
- 8.4 Summary — 223

9 Friction sensitivity — 224

- 9.1 Friction sensitivity of nitramines — 226
- 9.2 Friction sensitivity of quaternary ammonium-based energetic ionic liquids — 227
- 9.3 Summary — 229

10 Heat sensitivity — 230

- 10.1 Thermal kinetics correlations — 230
- 10.1.1 Nitroparaffins — 231
- 10.1.2 Nitramines — 232
- 10.1.3 Polynitroarenes — 232
- 10.1.4 Organic energetic compounds — 234
- 10.2 Heat of decomposition and temperature of thermal decomposition — 235
- 10.2.1 Heat of decomposition of nitroaromatics — 238
- 10.2.2 Heats of decomposition of organic peroxides — 239
- 10.2.3 Onset temperature of polynitroarenes and organic peroxides as well as maximum loss temperature of organic azides — 240
- 10.3 Deflagration temperature — 244
- 10.3.1 F_{nonadd}^+ — 245
- 10.3.2 F_{nonadd}^- — 245
- 10.3.3 Energetic compounds containing both F_{nonadd}^+ and F_{nonadd}^- — 245
- 10.4 Thermal stability of selected classes of energetic ionic liquids and salts — 246
- 10.4.1 Predicting activation energy of thermolysis of some selected ionic liquids — 246
- 10.4.2 Decomposition temperature of imidazolium-based energetic ionic liquids or salts — 248
- 10.5 Decomposition temperature of azole-based energetic compounds — 252
- 10.6 Predictive model for thermal decomposition onset in heterocyclic aromatic compounds and salts, incorporating elemental composition and structural features salts — 255
- 10.7 Summary — 263

11	Relationships between different sensitivities — 265
11.1	Relationship between impact sensitivity of energetic compounds and activation energies of thermal decomposition — 265
11.1.1	Nitroaromatics — 265
11.1.2	Nitramines — 266
11.2	Relationship between electric spark sensitivity and impact sensitivity of nitroaromatics — 267
11.3	A general correlation between electric spark sensitivity and impact sensitivity of nitroaromatics and nitramines — 267
11.4	Relationship between electric spark sensitivity and activation energy of the thermal decomposition of nitramines — 269
11.5	Correlation of the electrostatic sensitivity and activation energies for the thermal decomposition of nitroaromatics — 270
11.6	Relationship between the activation energy of thermolysis and friction sensitivity of cyclic and acyclic nitramines — 271
11.7	Relationship between shock sensitivity of nitramine energetic compounds based on small-scale gap test and their electric spark sensitivity — 272
11.8	Summary — 274
12	Estimation of the properties of metal-containing energetic complexes and energetic metal-organic frameworks (MOFs) — 275
12.1	A molecular weight-dependent model for predicting the density of EMOFs — 276
12.2	Predictive model for condensed-phase heat of formation in EMOFs — 280
12.3	A reliable predictive model for the onset temperature of energetic complexes — 283
12.4	Summary — 289
13	Estimating the properties of energetic polymers — 290
13.1	Predicting solubility of energetic polymers: computational models, group contribution methods, and experimental validation — 291
13.2	Characterization and prediction of intrinsic viscosity in energetic polymer solutions — 294
13.3	Glass transition in energetic polymers: experimental characterization, predictive modeling, and performance implications — 296
13.4	Energetic polymers with engineered refractive index for multifunctional applications — 299
13.5	Summary — 302

14 Prediction of the properties of energetic materials — 304

- 14.1 Computational prediction of energetic materials: methods, challenges, and applications — **304**
- 14.2 Summary — **315**

Problems — 316**Answers to Problems — 331****List of symbols — 337**

- A Glossary of compound names and heats of formation for pure as well as composite explosives — 347**
- B Calculation of the gas-phase standard enthalpies of formation — 351**
- C Glossary of compound names, as well as the measured and calculated values of the condensed-phase heats of formation for some energetic ionic liquids and salts — 359**
- D Common ligand abbreviations in energetic metal-organic frameworks (EMOFs) — 363**

Bibliography — 365**Index — 403**

