

REACTIONS OF 1,2,4-TRIAZINE AZIDES WITH α -KETO AND α -ESTER PHOSPHORUS YLIDES. SYNTHESIS OF SOME NEW 1-TRIAZINO-1,2,3-TRIAZOLES

YEHA O. EL-KHOSHNEH


National Research Centre, Dokki, Cairo, Egypt.

Abstract : 3-Azidotriazines 1 and 2 react with acetylmethylenetriphenylphosphorane 3a to give 1-triazino-5-substituted-1,2,3-triazoles 6 and 7, respectively. The regiochemistry of the reaction was resolved by comparison of the triazole product 6 with an authentic sample, prepared by the Dimroth reaction. On the other hand, azides 1 and 2 react with ethoxycarbonylmethylenetriphenylphosphorane 3b to afford the corresponding iminophosphoranes 12 and 13. Possible reaction mechanisms are proposed for the formation of the new products.


Introduction

1,2,4-Triazines possess a wide spectrum of biological activities whereas their numerous derivatives are in clinical use (1-6). A slight variation in nuclear substitution causes a marked difference in the biological activities and therefore it has been considered worthwhile to synthesize new heterocyclic compounds incorporating 1,2,4-triazine moiety.

As an extension of our work in the 1,2,4-triazine field (7-9), the present investigation was undertaken to examine the reaction of 3-azido-phenanthro[9,10-e]-1,2,4-triazine 1 and 3-azido-5,6-diphenyl-1,2,4-triazine 2 with Wittig reagents 3. The condensation of azidotriazine 1 with the active methylene compound 4 in the presence of a base was also examined.




Phosphorus ylides 3a,b are commonly represented in three resonance forms (A, B and C) and can, a priori, react with azides (10-15) either as carbanions (3A) or as unsaturated systems $\text{C}=\text{P}$ (3B) as well as $\text{C}=\text{C}$ (3C).



Results and Discussion

3-Azidotriazine **1** reacts with acetylmethylenetriphenylphosphorane **3a** in toluene solution at reflux temperature for 10 h to give a brown crystalline product formulated as **6** (Scheme 1). Triphenylphosphine oxide (TPPO) was also isolated from the reaction medium. The triazole structure **6** was deduced from the following: (a) Correct elemental analysis and molecular weight determination (MS) correspond to $C_{18}H_{12}N_6$; (b) The IR spectrum of **6**, in KBr, reveals the presence of the triazole absorption band at 1180 cm^{-1} (16) (c) The ^1H -NMR spectrum of **6** showed signals at δ 2.60 (d, 3H, CH_3 , $J_{\text{HH}}=1\text{ Hz}$) and at 7.8 (q, 1H, $J_{\text{HH}}=1\text{ Hz}$). The aromatic protons appeared as a multiplet in the range 7.33-7.72 ppm (8H).


A mechanism which accounts for the reaction of azidotriazine **1** with acetylmethylenetriphenylphosphorane **3a** is depicted in Scheme 1. The reaction can be visualized as 1,3-dipolar cycloaddition of the azide **1** onto the $\text{C}=\text{C}$ bond of the ylide **3a** (form C) followed by the loss of triphenylphosphine oxide (TPPO) from the cyclic intermediate **5** to give N-1 substituted 5-methyl-1,2,3-triazole (11) (Scheme 1).



Scheme 1

Azidotriazine **2**, similarly, reacts with acetylmethylenetriphenylphosphorane **3a** to give N-1 substituted 5-methyl-1,2,3-triazole **7** in 85% yield (Scheme 1), based on correct elemental, spectroscopic data and by analogy with **6** (cf. Experimental).

Further structural proof for the triazole product **6** was obtained through an independent synthesis by reacting the β -keto ester : ethyl acetoacetate **4**, with azidotriazine **1** in ethanol in the presence of sodium ethoxide, followed by decarboxylation of the formed triazole-4-carboxylic acid **9** (Scheme 2) (cf. Experimental). This reaction, Dimroth reaction (11,17) is known to afford a triazole with the R substituent ($-\text{CH}_3$) in the 5-position. The identity of the triazole obtained from **3a** with that from **4** adequately demonstrates the regiochemistry of the reaction (**1, 2** + **3a**) whereby an exclusive formation of 1,5- instead of 1,4-**8**-disubstituted triazoles (cf. Scheme 1).




Scheme 2

Conversely, the reaction of azidotriazine **1** with ethoxycarbonylmethylenetriphenylphosphorane **3b** in dry toluene at 50 °C affords the iminophosphorane **12** and ethyl diazoacetate **11** (Scheme 3). The iminophosphorane structure **12** is assigned for the following reasons : (a) Correct elemental analysis and molecular weight determination (MS) correspond to $\text{C}_{33}\text{H}_{23}\text{N}_4\text{P}$, (b) Its ^{31}P -NMR spectrum has one signal at $\delta = 18.5$ ppm (vs. 85% H_3PO_4) which clearly indicates an iminophosphorane structure (18), (c) The IR spectrum of **12** reveals the presence of an absorption band at 1355.7 cm^{-1} characteristic for N=P group absorption (19) and the absence of N_3 group absorption which is recorded in the starting material at 2135.3 cm^{-1} (9), (d) The ^1H -NMR spectrum (CDCl_3) showed only a multiplet in the range $\delta\ 7.25\text{--}7.98$ ppm (Ar-H).


The iminophosphorane **13** was likewise obtained from the reaction of **2** and **3b**. Assignment of **13** was based on analytical, spectroscopic data and by analogy with **12**. The ^{31}P -NMR measurement of **13** exhibited one signal at $\delta\text{p} = 17.6$ ppm (vs. 85% H_3PO_4).

A possible explanation for the course of the reaction of azidotriazine **1** with ethoxycarbonyltriphenylphosphorane **3b** is shown in Scheme 3. The reaction can be regarded to proceed by a concerted 1,3-dipolar cycloaddition of the azide **1** or **2** onto the C=P bond of the ylide **3b**, (form B) in the first step of the reaction, followed by collapse of the cyclic intermediate **10**, to afford ethyl diazoacetate **11** and the iminophosphoranes **12** or **13** (12-14). The success of the reaction depends upon the phosphinimine formed being unreactive toward ethyl diazoacetate.



Scheme 3

Moreover, compound **12** was further synthesized by unequivocal route, namely by heating of the phosphazene **14** (9), initially produced by reacting azidotriazine **1** with triphenylphosphine (Staudinger reaction) to give rise the corresponding iminophosphorane **12** accompanied with loss of nitrogen (20) (Scheme 4) (*cf.* Experimental).



Scheme 4

Experimental

All melting points are uncorrected. Ethoxycarbonylmethylene- (21) and acetylmethylene-triphenylphosphorane (22) were prepared according to established procedure. IR spectra were recorded on a Perkin-Elmer Spectrophotometer model 197 (Grating) using KBr disk. ^1H -NMR spectra were recorded on a Jeol-270 MHz Spectrometer, using TMS as an internal reference. ^{31}P -NMR were carried out on a Varian CFT-20 Spectrometer (vs. external 85% H_3PO_4). Mass spectra were performed at 70 eV on MS-50 Kratos (A.E.I.) Spectrometer provided with a data system. The appropriate precautions in handling moisture-sensitive compounds were observed. Solvents were dried by standard techniques.

Reaction of ylide 3a with azidotriazine 1 and 2. the ylide **3a** (22) (0.01 mol) was allowed to react with 3-azidotriazine **1** or **2** (9) (0.01 mol) in 50 ml of refluxing toluene. After the reaction was completed (TLC, 10 h) the solvent was evaporated in vacuo and the residual substance was crystallized from the suitable solvent to give triazoles **6** and **7**. The mother liquors from the crystallization were concentrated and the residue (2.5 g, 90%) crystallized from hexane and proved to be triphenylphosphine oxide (mp, mixed mp and comparative IR spectra) (23).

Compound **6**, brown needles (2.18 g, 70%), m.p. 194 °C (chloroform). Anal. Calcd. for $\text{C}_{18}\text{H}_{12}\text{N}_6$ (312.336): C, 69.22; H, 3.87; N, 26.91. Found: C, 69.16; H, 3.82; N, 26.84. Mol. wt. (MS) = 312. IR (KBr) cm^{-1} : 1180 (triazole). ^1H -NMR (CDCl_3): δ 2.60 (d, 3H, CH_3 , $J_{\text{HH}}=1$ Hz), 7.80 (q, 1H, $J_{\text{HH}}=1$ Hz), 7.33-7.72 (m, 8H, Ar-H).

Compound **7**, yellow needles (2.67 g, 85%), m.p. 147 °C (chloroform). Anal. Calcd. for $\text{C}_{18}\text{H}_{14}\text{N}_6$ (314.352): C, 68.78; H, 4.49; N, 26.73. Found: C, 68.75; H, 4.44; N, 26.70. Mol. wt. (MS) = 314. IR (KBr) cm^{-1} : 1189.86 (triazole). ^1H -NMR (CDCl_3): δ 2.55 (d, 3H, CH_3 , $J_{\text{HH}}=1$ Hz), 7.1 (q, 1H, $J_{\text{HH}}=1$ Hz), 7.25-7.75 (m, 10H, Ar-H).

Synthesis of triazole 6 by the Dimroth method. Equimolecular amounts (0.01 mol) of 3-azidotriazine **1** (9), ethyl acetoacetate **4** and sodium ethoxide were refluxed in 75 ml ethanol for 2 h. The product was

dissolved in the minimum of hot water and heated under reflux for further 1 h. The reaction mixture was poured into cold water, acidified with conc. HCl. 1-Triazino-5-methyl-1,2,3-triazole-4-carboxylic acid **9** separated (2.56 g, 72%) and was crystallized from water and dried, m.p. 212 °C. Anal. Calcd. for $C_{19}H_{12}N_6O_2$ (356.347) : C, 64.04; H, 3.40; N, 23.58. Found : C, 64.00; H, 3.32; N, 23.56. IR (KBr) cm^{-1} : 1695 (C=O). The triazole-4-carboxylic acid **9** was converted quantitatively into triazole **6**, (mp, mixed mps and comparative IR spectra) when heated above its m.p. until gas evolution ceases.

*Reaction of ylide **3b** with azidotriazines **1** and **2**.* A mixture of 3-azidotriazine **1** or **2** (**9**) (0.01 mol) and ylide **3b**, (**21**) (0.01 mol) was heated in dry toluene (50 ml) at 50 °C. After the reaction was completed (TLC, 50-60 h), the volatile materials were evaporated in vacuo and the residual substance was treated with petroleum ether (b.r. 60-80 °C) in order to precipitate the iminophosphoranes **12** or **13**. The mother liquor was concentrated to give ethyl diazoacetate **11**, as light yellow oil, 0.8 g (70%), which was identified by comparison its IR(**24**) and 1H -NMR (**13**) spectra.

Compound **12**, yellow substance (3.9 g, 78%), m.p. 178 °C (chloroform-light petroleum). Anal. Calcd. for $C_{33}H_{23}N_4P$ (506.55) : C, 78.25; H, 4.58; N, 11.06; P, 6.11. Found : C, 78.19; H, 4.53; N, 11.02; P, 6.07. Mol. wt. (MS) = 506. IR (KBr) cm^{-1} : 1355.7 (N=P), 1027.87 (P-C, phenyl). ^{31}P -NMR ($CDCl_3$) : δ = 18.5 ppm.

Compound **13**, yellow substance (4.4 g, 88%), m.p. 125 °C (chloroform-light petroleum). Anal. Calcd. for $C_{33}H_{25}N_4P$ (508.57) : C, 77.94; H, 4.95; N, 11.02; P, 6.09. Found : C, 77.90; H, 4.89; N, 10.97; P, 6.00. Mol. wt. (MS) = 508. IR (KBr) cm^{-1} : 1369.21 (N=P), 1025.94 (P-C, phenyl). ^{31}P -NMR ($CDCl_3$) : δ = 17.6 ppm.

*Action of heat on **14**. Adduct **14** (**9**).* (0.5 g) was heated in a cold Finger sublimation at 150 °C (bath temperature) under reduced pressure (2mm/Hg) for 10 min. The substance that sublimed was collected, crystallized from chloroform-pet. ether (60-80 °C) to give a yellow substance proved to be iminophosphorane **12** in 85% yield (mp, mixed mps and comparative IR spectra).

References

1. L.M. Mironovich, V.N. Promonenkov, L.I. Minaev, P.S. Balesta, Z.A. Bukashkina and E.P. Fedorov, *Fiziol. Akt. Veshchestva*, **23**, 64 (1991); C.A. **118**, 96159 (1993).
2. A. Mong, J. Palop, C. Ramirez, M. Font and E. Fernandez-Alvarez, *Eur. J. Med. Chem.*, **26**, 179 (1990); C.A. **115**, 136057 (1991).
3. K.H. Büchel, *Chemistry of Pesticides*, Wiley-Interscience, New York (1971).
4. S.V. Galushko, I.P. Shishkina and I.V. Alekseeva, *J. Chromatogr.*, **547**, 161 (1991).
5. H. Uhr, A. Widdig, D. Berg and G. Haenssler, *Eur. Pat.*, **4**, 002, 997 (1990); C.A. **115**, 183375 (1991).
6. C. Fest and K.J. Schmidt, *The Chemistry of Organophosphorus Pesticides*, Springer-New York, 1973, pp. 109.
7. Y.O. El-Khoshnieh, Y.A. Ibrahim and W.M. Abdou, *Phosphorus, Sulfur and Silicon*, **101**, 67 (1995).

8. Y.O. El-Khoshnieh, I.T. Hennawy and W.M. Abdou, *Heterocyclic Communication*, 1, 2-3. 167 (1995).
9. Y.O. El-Khoshnieh, *Phosphorus, Sulfur and Silicon*, 1998, in press.
10. G.R Harvey, *J. Org. Chem.*, 31, 1587 (1966).
11. P. Ykman, G. L'abbe' and G. Smets, *Tetrahedron*, 27, 845 (1971).
12. G. L'abbe' and H.J. Bestman, *Tetrahedron Lett.*, 63 (1969).
13. G. L'abbe', P. Ykman and G. Smets, *Tetrahedron*, 25, 5421 (1969).
14. P. Ykman, G. L'abbe' and G. Smets, *Tetrahedron*, 29, 195 (1973).
15. L.S. Boulos and N. Khir Eldin, *Tetrahedron*, 49, 3871 (1993).
16. (a) J.H. Boyer, "Heterocyclic Compounds", Vol. 7, R.C. Elderfield, Ed., John Wiley and Sons, Inc., New York, N.Y., 1957, p. 422; (b) L.W. Hartzel and F.R. Benson, *J. Am. Chem. Soc.*, 76, 667 (1954); (c) C.S. Rondestvedt, Jr. and P.K. Chang, *ibid.* 77, 6532 (1955).
17. O. Dimroth, *Chem. Ber.*, 35, 1029 (1902); *ibid.* 35, 4041 (1902).
18. M.M. Grutchield, O.H. Dungan, J.H. Letcher, V. Mark and J.R. Van Wazer, "Topics in Phosphorus Chemistry", Interscience Publishers, 1967, Vol. 5.
19. L.C. Thomas, "The Identification of Functional Groups in Organo-phosphorus Compounds", Academic Press Inc., London, 1974.
20. E. Bergmann and H.A. Wolff, *Chem. Ber.*, 63, 1176 (1930).
21. Th. Kappe, E. Lender and E. Ziegler, *Monatsh. Chem.*, 99, 2157 (1968).
22. F. Ramirez and S. Dershowitz, *J. Org. Chem.*, 22, 41 (1957).
23. A. Michaelis and L. Gleichman, *Chem. Ber.*, 15, 801 (1882).
24. P. Yates, B.L. Shapiro, N. Yoda and J. Fugger, *J. Am. Chem. Soc.*, 79, 5756 (1957).

Received on September 25, 1998