

HETEROCYCLIC COMMUNICATIONS

An International Journal in Heterocyclic Chemistry

Published bimonthly by Freund Publishing House Ltd.,
Suite 500, Chesham House, 150 Regent Street, London W1R 5FA, England

Notes for the preparation of Camera-Ready Manuscripts

Heterocyclic Communications seeks to publish preliminary communications and full length research papers on all phases of heterocyclic chemistry including inorganic ring systems. All papers to the journal will be reproduced from author's typescripts (camera-ready manuscript) by a reduction of 25%.

Guidelines for Authors

1. Preparation and Mailing of Manuscripts

Author(s) should mail manuscript to respective editor in triplicate. All manuscript should be made on a computer and printed out by a laser printer preferably at 600 dpi or higher resolution. Author(s) should send the editor a floppy disc (3 1/2" or 5 1/4") in DOS format (1.4MB for 3 1/2" and 1.25MB for 5 1/4") which contains a combined manuscript file including computer drawn structural formulas, figures, tables etc. For the all type setting, Times Roman 10 Pt. font or corresponding font set should be used. Preferably, file format of the manuscript is Adobe PageMaker 6.0/5.0 or MS Word 97 versions for MS Windows. Structures may be drawn by programs like Chem Draw, Chem Win, Chem Window/Coral Draw, Chem Text. Apple Macintosh and other word processor users should convert the file format to those readable by these softwares. Some times RTF (Rich Text Format) and PDF (Portable Document Format) are effective. Details of the program(s) used should be also provided. If structures/diagrams have been prepared using program(s) other than those stated, then structures/diagrams should be converted into windows meta file format. Details of the program(s) used to draw structures/diagrams should be also provided. Manuscript should end with even numbered page (viz 2,4,6,...). Normally manuscript not exceeding 6 pages (excluding graphical abstract) will be accepted. Editor can accept even manuscript exceeding 6 pages, if they feel that it is appropriate to do so.

All manuscripts except from North America, European countries and Japan should be mailed to Dr. R.R. Gupta, Editor-in-Chief, Heterocyclic Communications, 10A, Vasundhara Colony, Tonk Road, Jaipur-302018, India. Authors from Europe should mail manuscripts to Prof. Jacques Barbe, Editor for European Countries, Faculte de Pharmacie, 27, Bd. Jean-Moulin, 13385 Marseille Cedex 5, France. Authors from North America should mail manuscripts to Prof. David W. Boykin, Department of Chemistry, Georgia State University, University Plaza, Atlanta, Georgia 30303-3083, USA. Authors from Japan should mail manuscripts to Prof. S. Eguchi, Department of Molecular Design and Engineering, Graduate School of Engineering, Nagoya University, Furo-cho, Chikusa-ku, Nagoya 464-8603, Japan.

2. Organization of the Manuscript

2.1 Title : It should be clear, concise, and informative. It should be typed with words all in bold face capitals, single spaced, centered on the width of the first page, 2 cm below the top of typing area.

2.2 Author(s) : After 4 single spaces from the title, authors names and addresses should be given on separate lines, typed single spaced, centered on the width of the first page. The name of the author to whom correspondence should be directed should be marked with an asterisk as superscript.

2.3. Abstract : After 4 single spaces from authors address, a short abstract (50-100 words) describing results, methodology employed, conclusions etc., should be placed. It should be labelled as the abstract with the word "Abstract" in bold face and should be typed single spaced. Total typing area on the first page should be 17×22 cm. Typing area for other pages should be 17×24 cm. Each manuscript should be accompanied by a Graphical Abstract, which should contain the title in bold face single spaced, authors names and addresses. It should summarise concisely the contents of the communications in pictorial form designed to capture the attention of a wide readership and to facilitate compilation of database. Chemical Structures designed to illustrate the theme of research work presented should be included. It should be prepared on a separate sheet of paper sized 19×7 cm. An example of graphical presentation is shown as :

Graphical Abstract

Space	Leave blank		1 cm
	10 cm		
For			
Abstract			

2.4 Text : Text should be divided into sections like **Introduction, Experimental**, including spectra screening, instruments specifications, **Results, Discussions, Conclusions, Acknowledgements** (if applicable) and **References**. However, the experimental section should be excluded in preliminary Communications as far as possible. Between the end of one section and the beginning of another, there should be a gap of three single spaces. Text should be typed in one and a half spacing throughout.

2.5 References : These should be numbered consecutively throughout the text as they appear and should be superscripted. All compound number in text should not be bold e.g., phenothiazines **7** can be prepared... Abbreviations for journals should be those used in Chemical Abstracts. Abbreviations for other terms should also be taken from Chemical Abstracts /ACS journal. Each reference should be typed single spaced. However, there should be one empty line between two consecutive references. Some examples for journal and books references are given:

- 1 R.R. Gupta, V. Saraswat, A. Gupta, M. Jain and V. Gupta, J. Heterocycl. Chem. **29**, 1703 (1992)
- 2 R.R. Gupta, Diamagnetic Susceptibility, Springer Verlag, Berlin, 1986
- 3 R.R. Gupta (Ed.), Physical Methods in Heterocyclic Chemistry, John Wiley, New York, 1984
- 4 R.R. Gupta and M. Kumar, Synthesis, Properties and Reactions of Phenothiazines, in R.R. Gupta (Ed.), Phenothiazines and 1,4-Benzothiazines-Chemical and Biomedical Aspects, Elsevier, Amsterdam, 1988, pp. 1

2.6 Illustrations

Both line drawings and photographs should be used sparingly. All illustrations must be numbered in sequence using arabic numbers, and each illustration must be referred to in the text. Each illustration must be provided with a caption which is complete enough for the figure to be appreciated without referring to the text. Line drawings must be submitted as the perfectly drawn original or as a glossy photograph print. Photocopies, Multiliths, Verifax or Xerox copies are not acceptable substitutes. If you are using desktop publishing software with high quality graphics capability, which is recommended, it is desirable that the graphics be sized and printed along with the text. A laser printer should be used. Photographs must be supplied as glossy prints in which any lettering must be part of the photographs. No lettering should be applied to the surface of the photograph. Micrographs and similar material in which linear dimensions are important should have a scale of length.

2.7 Tables

Tabular presentation of data is an economic way of condensing many items. Refer to tables by using Latin numerals in text and to head the table itself. Provide each table with a caption which will allow it to be read without reference to the text. The table caption should be typed to the width of the table itself. Exceptionally large tables may be placed landscape on the page with the top of table at the left-hand margin.

2.8 Page Numbers

When your typescript is complete and in its final form, number each page sequentially from 1, using a light blue pencil, in the top right-hand corner of each sheet. Final page numbers will be inserted by the editor or the publisher.

2.9 Layout

The typing area should be 17×24 cm for all pages except the first. On the first page typing should start 2 cm below top of the typing area, e.g., its typing area should be 17×22 cm. After abstract text should be typed in one and a half spacing throughout on all pages. Photographic reduction will be by 25%.

2.10 Displayed Equations and Formulas

Leave a blank line above and below all display. Break equations, where possible, only at an equals sign (=) or equivalent (>, <, >., <.). Do not repeat equations sign at the end of one line and the beginning of the next, but carry it over to the beginning of the following line.

3 Cost of Reprints

Orders for purchase of reprints should be placed to the publisher after receiving the acceptance of paper. Cost of reprints is given in the following table.

Number of Reprints	Number of Pages					
	1-2pp	3-4pp	5-8pp	9-12pp	13-16pp	Each additional unit (1-4pp)
100	\$ 83	122	159	253	341	111
200	\$ 109	150	259	380	516	136
300	\$ 128	245	380	569	759	172

Orders for quantities over 300 copies are subject to special quotations.

Index of Vol. 5 (Issue 1-6), 1999

Heterocyclic Communications, Vol. 5, No. 1, 1999

Vol. 5, Issue 1 (1999)

	1	Contributors to this Issue.
	2	Graphical Abstracts.
L. Streckowski, M.A. Ismail and H.H. Zoorob	9	A general method for acylation of 1,3-dialkyl-substituted barbituric and 2-thiobarbituric acids.
A.K. Singh and J.W. Lown	11	Synthesis of Hoechst 33258 analogues designed to target human tumor helicases.
K. Matsumoto, A. Shigeta, A. Okada, T. Kakara and M. Toda	19	Binding properties of some new mono-, bis-, and tris-crown ethers for silver ions.
P. Sonnet, H. Miel, J. Guillon, P. Dallemagne and S. Rault	25	Synthesis of new phenylpyrrolizinones derivatives, via a Mannich reaction.
N. Kizilcan, A. Akar, N. Talinli and S. Yaslak	27	Phosphorous esters of 2,13-dihydro-7a,14c-dihydronaphtho[1',2':4,5]furo-[3,2-d]furan.
T. Xu, J. Su, K. Chen and H. Tian	31	Synthesis of bis- and tris-1,8-naphthalimides bridged by N-N bond.
K. Wurst and M.R. Buchmeiser	37	Chiral chelating systems: Unusual formation and X-ray-structure of RAC-3A(R),4(S),5,9B(S)-tetrahydro-4-(pyrid-2-yl)-3H-cyclopenta[C]quinoline.
T. Eszenyi, T. Timar, P. Sebők and J. Jekő	45	On the synthesis of benzodipyrans.
M. Vlassa, C. Afloreaei, N. Dulamita, P. Brouant and J. Barbe	51	Application of phase transfer catalysis in the acridine series VII (1). Synthesis of 9-cyanoacridine derivatives.
Y. Ikemi, A. Okada, H. Katsura, S. Otani and K. Matsumoto	53	Inverse electron demand Diels-Alder reactions of 3,6-diaryl-1,2,4,5-tetrazines with cyclooctyne.
E. Abele, R. Abele, I. Sleiksa and E. Lukevics	59	A simple phase transfer catalyzed synthesis of benzyl and hetaryl methyl silacycloalkyl ethers from acetates.
A.R. Subramanian, S.A.V. Raghavan, R.J. Babu, C.N.V.H.B. Gupta, N. Sridhar, A. Veeranjanyulu, P. Parimoo and P. Srinivas	63	Synthesis of new piperazinyl indolyl propanones and preliminary CNS pharmacological evaluation in mice.
V.F. Traven, A.Y. Tolmachev, N.Y. Podhaluzina, D.S. Kanevskii and N. P. Solovieva	69	New ways of lactone ring shortening and cyclopropanation in coumarin derivatives.
Y.O. El-Khoshnief	77	Reactions of 1,2,4-triazine azides with α -keto and α -ester phosphorus ylides. Synthesis of some new 1-triazino-1,2,3-triazoles.
Ö. Sezer, K. Dabak, O. Anaç and A. Akar	83	Synthesis of triazolylbenzoxazole derivatives.
M. Balogh, A. Gerstmans and I. Hermecz	89	Solid acid catalyzed reaction of amins with methyl 3-aminocrotonate.
P.S. Verma, R. Gupta, N. Sharma, M.Y. Hamadi, V. Gupta and R.R. Gupta	93	Single step synthesis of 3-methyl-5/7-substituted 4H-1,4-benzothiazines.
	97	Notes for the preparation of manuscript.

Heterocyclic Communications, Vol. 5, No. 2, 1999

Vol. 5, Issue 2 (1999)

	99	Contributors to this Issue.
	100	Graphical Abstracts.
L. Streckowski, M.A. Ismail and H.H. Zoorob	107	Novel synthesis transformation of 5-(ω -chloroalkanoyl)-1,3-dimethyl-barbituric acids.
F.A. Chowdhury, H. Nishino and K. Kurusawa	111	Manganese(III)-based oxidative radical cyclization of terminal alkadienes with <i>N,N'</i> -bis(3-oxobutanoyl)- α,ω -alkanediamines. Formation of macro-diamides.
P. Norris	113	Chiral tetrahydrofuran synthesis from D-ribose diphenyl dithioacetal.
J. Quiroga, B. Insuasty, A. Hormaza, P. Cabildo, R.M. Claramunt and J. Elguero	115	Synthesis, molecular structure and tautomerism of 1(2 <i>H</i>)-dihydropyrazolo-[3,4- <i>b</i>]pyridin-6-ones.
R. Demadrille, C. Moustrou, A. Samat and R. Guglielmetti	123	Palladium-catalyzed amination of photochromic triflate-substituted 3 <i>H</i> -naphtho[1,2- <i>b</i>]pyrans.
A. Perjéssy, P. Meyer, W.-D. Rudolf, D. Loos, A. Kolbe and L. Schaller	127	Transmission of substituent effect in ethyl 2-aryloxy- and ethyl 2-aryl-carbamoyl-4,5-dimethyl-1,2,3,6-tetrahydropyridazine-1-carboxylates : An infrared and theoretical study.
M. Noguchi, S. Nishimura, H. Fujii and A. Kakehi	133	A simple oxime-nitrone isomerization and intramolecular cycloaddition reaction of 2-(alk-2-enyl)oxy and 2-[<i>N</i> -(alk-2-enyl)benzylamino]-cyclohexanone oximes.
C. Foces-Foces, A.L. Llamas-Saiz and J. Elguero	137	The structure of <i>N</i> -acetylazoles (azolines) : A semiempirical (AMI) computational study.
I.A. Silberg, I. Silaghi-Dumitrescu, C. Cristea, P. Tordo and D. Gígmes	147	Molecular orbital calculations and physical properties of 1,4-benzothiazino-[2,3- <i>b</i>]phenothiazine and its substituted derivatives.
A. Lévai	151	Fused heterocycles 9. synthesis of tricyclic pyrazolines by the reaction of <i>E</i> -3-arylidene flavanones with phenylhydrazine.
J. Styskala and J. Slouka	157	Cyclization reactions of hydrazones XXV : Synthesis and study of reactivity of some derivatives of [1]benzothieno[2,3- <i>e</i>]1,2,4-triazine.
T. Okano, H. Yamada and S. Eguchi	163	Addition-elimination reaction of highly electrophilic difluoroolefin, 5-difluoromethylene-3-pyrrolin-2-ones.
M.A. Raslan, M.A. Khalil, S.M. Sayed and A.M. Farag	167	Facile synthesis of novel polysubstituted thiophene and benzo [g] thiazo-[3,2- <i>b</i>]pyridine.
Y. Dürüst, H.A. Döndas and N. Ozkoru	173	Synthesis of some bicyclo and spiro isoxazolines.
V.S. Rao, S.V.S.A.K. Gupta and B.S. Reddy	179	Ortho amino cyano aromatic compounds as precursors for the synthesis of some novel heterocyclic compounds.
V.F. Traven, A.Y. Tolmachev, N.A. Podhaluzina and D.S. Kanevskii	183	Cholesteryl esters of furocoumarin and coumarin carboxylic acids.
K. Li, L.-N. He, X.-H. Qing, Y.-P. Luo and M.-W. Ding	189	A facile synthesis of 1-alkyl-1,3,2-diazaphospholidin-4-thione-2-sulfide via Lawesson's reagent.
	193	Notes for the preparation of manuscript.

Heterocyclic Communications, Vol. 5, No. 3, 1999

Vol. 5, Issue 3 (1999)

	195	Contributors to this Issue.
	196	Graphical Abstracts.
Z. Wang, R. Neidlein and C. Krieger	203	Reactions of 3,3-diamino-2-cyanoacrylates with substituted ureas and dichloromethylenedialkyliminium chlorides.
E. Fourmaintraux, P. Depreux and I. Lesieur	213	Synthesis of new phthalazinyl compounds as potential inhibitors of aldose reductase and sorbitol dehydrogenase.
B.F. Bonini, G. Mazzanti and P. Zani	217	Chemistry of silyl thioketones : Studies concerning the regiochemistry of cycloaddition with substituted 1,3-dienes.
A.P. Aguiar and W.B. Kover	227	Chemo- and regioselective synthesis of new phosphorated 4,5-dihydro-isoxazoles from different monoterpenes.
P. Bilek and J. Slouka	231	Cyclocondensation reactions of heterocyclic carbonyl compounds VI: Cyclocondensation of the isomeric 1-(2-aminophenyl)-6-azauracil-5-carboxylic acid N-methyl derivatives.
S. Alibert-Franco, C. Santelli-Rouvier, J. Barbe, B. Pradines, C. Houdoin and D. Parzy	235	9,10-(3',4'-Pyrrolidino)-9,10-dihydroanthracene and structurally related compounds as synergistic antimalarial drugs.
M.V. Vasylyev, Y.V. Bilokin, S.M. Branytska S.M. Kovalenko and V.P. Chernykh	241	A facile method for synthesis of heterocycles containing tetrahydrobenzo-[4,5]thieno[2,3-d]pyrimidine and coumarin moieties.
A.H. Atta	243	Reactions of 1-(2-benzothiazolyl)-4-(dicyanomethylene)-3-methyl-2-pyrazolin-5-one towards amines.
C. Afloroaei, M. Vlassa, A. Becze, P. Brouant and J. Barbe	249	Microwave action on 2-(arylamino)-nicotinic acid derivatives.
F.D. Irimie, C. Afloroaei, M. Tosa and C. Paizs	253	Bioreduction with bakers' yeast of π -deficient heterocyclic aldehydes.
L. Cabanal-Duvillard and J.-F. Berrien	257	A simple access to key pyridine building blocks.
R. Gupta, V. Gupta, N. Sharma, M.Y. Hamadi, P.S. Verma and R.R. Gupta	263	Synthesis of 1-methyl-3,7-disubstituted phenothiazines.
Z.A. Hozien, A.A.O. Mohamed	269	Synthesis of spiroheterocycles related to spiro[fluren-9,2'(1',3'-oxathiolan)]-5'-one and spiro[anthracen-9(10)H,2'-(1',3'-oxathiolan)]-5'-one.
S. Deprets and G. Kirsch	275	Synthesis of new heterocyclic analogues of isocoumestane.
I.E. Bylov, Y.V. Bilokin and S.M. Kovalenko	281	Specific features of reactions of 2-aminobenzotrifluoride and anthranilates with ethyl cyanoacetate-Expedition routes to 3-substituted 4-amino- and 4-hydroxyquinolin-2(1H)-ones.
B.B. Reddy, A. Balaiah, V. Padmavathi and A. Padmaja	285	Selena and thiadiazole fused polycyclic polythia compounds-Part-III.
	291	Notes for the preparation of manuscript.

Heterocyclic Communications, Vol. 5, No. 4, 1999

Vol. 5, Issue 4 (1999)

	293	Contributors to this Issue.
	294	Graphical Abstracts.
A. Kumar, C.E. Stephens and D.W. Boykin	301	Palladium catalyzed cross-coupling reactions for the synthesis of 2,5-disubstituted furans.
M. Welch and R.S. Phillips	305	Improved synthesis of [3,2- <i>b</i>]- and [3,2- <i>b</i>]- fused selenolo- and thienopyrroles, and of furo[3,2- <i>b</i>]pyrrole.
L.H. Klemm, T.J.R. Weakley and M. Yoon	311	X-Ray crystallographic studies of four monosubstituted thienopyridines. Comparison of experimental data with calculated or measured values for the parent compounds.
K. Ito, T. Ohta, Y. Ohba and T. Sone	319	Syntheses of chiral tetrahomodiazacalix[4]arenes incorporating amino acid residues. Chiral induction of cyclophane moiety by the chirality of amino acid residue.
T. Kurz and D. Geffken	325	Cyclization of thiocyanato malonates with hydroxylamines to 5-carboxy-alkyl-4-oxo-thiazoli(di)nes.
R. Barret, A. Belaissaoui, Z.M. Du, T. Mulamba, J.Y. Laronze and J. Levy	329	Oxidation of indolo-quinoline structures.
R. Miranda, R. Osnaya, I. Oviedo, A. Ciprian, T. Cruz and M. Martinez	331	Facile route to amino phthalimides and isothiocyanate analogues; Novel reagents to prepare fluorescent protein conjugates.
E. Yanase and S.-i. Nakatsuka	339	Synthesis of monomethyl and dimethyl derivatives of epicatechin gallate (ECg) and their photo-sensitivity.
S. Danoun, G. Baziard-Mouysset, J.-L. Stigliani, M. Ané-Margail, M. Payard, J.-M. Léger, X. Canron, H. Vial, P.M. Loiseau and C. Bories	343	Synthesis and photozoocidal activity of new 1,4-naphthoquinones.
J. Styskala and J. Slouka	349	Cyclization reactions of hydrazones XXVI: Synthesis of aryl-2,3-dihydro-naphto[1'.2':4,5]furo[2,3- <i>e</i>]1,2,4-triazin-3-ones and their use for the preparation of 1-aryl-5-(2-hydroxy-1-naphthyl)-6-azauracils.
G.-F. Yang and H.-Z. Yang	355	Synthesis and herbicidal activity of novel α -(1,2,4-triazolo[1,5- <i>a</i>]-pyrimidine-2-oxyl)phosphonate derivatives.
A. Lévai	359	Oxazepines and thiazepines 38, Synthesis of 2,4-diaryl-2,3-dihydro-1,5-benzothiazepines by the reaction of 2-hydroxychalcones with 2-amino-thiophenol.
H. Wamhoff, J. Muhr, M. Horn, P. Sohár and A. Csámpai	365	Subsequent reactions of 7-ethoxypyrimido[4,5- <i>d</i>]pyrimidines: Nucleophilic exchange, pyrimido[4,5:4',5']pyrimido[1,2- <i>a</i>]quinazoline, benzo[f]-pyrimido[4,5:4',5']pyrimido[1,2- <i>d</i>][1,3,4]triazepine.
D. Graham and G. McAnally	377	Synthesis of aminobenzotriazoles.
V.F. Traven, LL Saharuk, D.V. Kravtchenko and LG. Makarov	379	The first pyrrolofurocoumarins.
S.B. Desai, P.B. Desai and K.R. Desai	385	Synthesis of new heterocyclic thiazolidinone and azetidinone compounds and their anticancer activity.
	391	Notes for the preparation of manuscript.

Heterocyclic Communications, Vol. 5, No. 5, 1999

Vol. 5, Issue 5 (1999)

	393	Contributors to this issue.
	394	Graphical Abstracts.
D.J. Koza and W.B. Euler	399	Convenient syntheses of 2,2'-biindole.
L. Lamarque, A. Meou and P. Brun	403	Manganese (III)-promoted addition of monomethyl malonate on dihydropyran in the presence of molecular oxygen.
L.M.F. Santos, A.J. Alves, A.R. de Faria, D.J. Brondani, A.C.C. Freitas, J.G. de Lima and A.C.L. Leite	407	A new series of 1-phenylpyrazole peptidomimetics.
A. Gaz, F. Ammadi, S. Boukhris, A. Souizi and G. Coudert	413	One-pot synthesis of 1,4-benzoxazin-2-ones and 1,4-benzoxazines from epoxides.
A. Levai	419	An efficient procedure for the preparation of 4-thioflavones by the reaction of flavones with Lawesson's reagent.
M.A. Metwally and M. Abdel-Mogib	423	The use of thiochroman-4-one in the synthesis of some benzothiopyranopyrimidines and bis-thiochrom-2-ene-4-one of pharmaceutical interest.
M. Al-Talib, S. Bataineh and H. Tashtoush	429	Reaction of 1,3- and 1,4-bis(1,3,4-oxadiazole-5(3H)-thione-2-yl) benzenes with aryl aldehydes.
N.R. Mohamed	433	Synthesis of α -thiocarbamoyl phosphoranes and their utility for formation of some pyrazole and thiophene derivatives.
A. Levai, C. Nemes and T. Patonay	441	Synthesis of new (Z)-3-arylidenechromanones by the photoisomerization of (E)-3-arylidenechromanones.
M.A. Metwally, M. Hammouda, S.S. El-Morsy and M.M. El-Hussini	445	The use of 4-hydroxymethyl-1-phenyl-2-pyrazolin-5-one in the synthesis of new heterocycles of pharmaceutical interest.
V. Padmavathi, K. Sharmila, A. Padmaja and D.B. Reddy	451	An efficient synthesis of 6,8-diaryl carbazoles via Fischer indole cyclizations.
C.A. Câmara, E.T. Silva, E.J. Barreiro and C.A.M. Fraga	457	Synthesis of new isosteric heterocyclic derivatives: Pyrazolo[3,4-b]thieno[3,2-e]pyridine, pyrazolo[3,4-b]pyrrolo[3,2-e]pyridine and furo[2,3-b]-pyrazolo[4,3-e]pyridine.
M.A. Khalil, M.A. Raslan, K.M. Dawood and S.M. Sayed	463	Synthesis of 1,2,4-triazole, 1,2,4-triazolo[3,4-b][1,3,4]thiadiazole and 1,2,4-triazolo[3,4-b][1,3,4]thiadiazine derivatives of 3-[5-(benzothiazol-2-yl)thieno[2,3-d]pyrimidine-4-one] acetic acid hydrazide.
H.A. Hamid, A. Mousaad, E.S. Ramadan and S.S.H.E. Ashry	473	Synthesis of fused and nonfused heterocycles from 5- and 8-substituted 3-hydrazino-1,2,4-triazino[5,6-b] indole.
C. Kavitha and K.J.R. Prasad	481	Tetracyclic compounds from indolo[2,3-b]cycloheptan-1-ones. Synthesis of isoxazolo[4'3':6,7]cyclohepta[b]indoles.
	489	Notes for the preparation of manuscript.

Heterocyclic Communications, Vol. 5, No. 6, 1999

Vol. 5, Issue 6 (1999)

	491	Contributors to this issue.
	492	Graphical Abstracts.
G. Jia, H. Iida and J.W. Lown	497	Design and synthesis of 1,2,9,9a-tetrahydrocyclopropa[c]benz[e]indole-4-one (CBI) dimers.
J. Garner and A. McCluskey	503	Regiocontrolled amination of dichloropyrimidines in LiClO ₄ – Et ₂ O solutions.
H. Iida, Y. Misumi, K. Matsumoto and J.W. Lown	509	S _N Ar Reactions of methyl and ethyl 2-nitro-5-fluorobenzoates in the synthesis of pyrro[2,1-c][1,4]benzodiazepine precursors.
E. Gavrielatos, J. Markopoulos and O. Iglessi-Markopoulou	515	Synthesis and nmr spectroscopic studies of novel N-acetyl-3-aminoalkyl tetramic acids.
R.F. Vieira, D.J. Brondani, A.R. Faria, R.M. Srivastava and A.C.L. Leite	521	Synthesis of new 1,2,4-oxadiazoles-derived dipeptidomimetics, a potential class of antiinflammatory drugs.
L. Strekowski and M.A. Ismail	525	Sodium borohydride mediated benzylation of 1,3-dimethylbarbituric acid.
D.J. Aitken, L. Besson, K. Partogyan-Halim and H.-P. Husson	529	Stereochemical effects in the efficient cyclisation of <i>N</i> -(1-cyano-3-methoxybut-3-enyl)-4-phenyloxazolidine.
S.T. Selvi and P.S. Mohan	533	Synthesis of furo- thieno- and pyrrolo- [3,2-a]acridones.
S.A. El-Kafrawy and E.B. Pedersen	539	Multiple lid stabilization of the DNA three-way junction. Insertion of <i>N</i> ³ -(1-pyrenylmethyl)thymidine.
L. Cristea, M.M. Popovici, M.T. Mendel, L. Silaghi-Dumitrescu and E. Kozma	543	1-(Acridin-9'-yl)-pyrazolin-3 and 5-ones. A new class of heterocycles with potential biological activity.
R. Gawinecki, E. Kolehmainen, B. Osmialowski, P. Palkovič and M. Nissinen	549	Synthesis and nmr spectra of 2-methyl-2-quinolin-2-yl-propiophenones.
A.M.E. Massry	555	Synthesis and antimicrobial activity of condensed and uncondensed quinoxalines.
R. Chandra and N.N. Ghosh	565	A. simple and economical method for the synthesis of intermediates of cefazolin.
A. Krauze, R. Vitoliņa, V. Garalene, H.-J. Jänsch and G. Duburs.	569	Synthesis of 4-substituted pyridin-2(1H)-ones, pyridine-2(1H)-thiones, related derivatives as analogues of cardiotonic drug of milrinone.
A.I. Khodair, M.M.A. Abbasi, E.-S.L. Ibrahim, A.H. Soliman and E.-S.H. El-Ashry	577	Synthesis of substituted quinolines and heterocyclo[x,y-c]quinolines by the nucleophilic substitution and rearrangements of 4-chloro-2-methyl-3-nitroquinolines.
	585	Notes for the preparation of manuscript.
	587	Index of Vol. 5, 1999