


ACRIDINE FORMATION BY THE REACTION OF 2-NONAFLUOROBUTYLANILINE WITH 2-MESITYLMAGNESIUM BROMIDE

Lucjan Strekowski* and Hyeran Lee
Department of Chemistry, Georgia State University, Atlanta, Georgia 30303, USA

Abstract: The title reaction conducted in THF at 23 °C and under reflux furnishes 1,3-dimethyl-9-perfluoropropylacridine (**8**) and 1,3-dimethyl-9-[(*E*)-1-perfluoropropenyl]acridine (**5**), respectively.

Recently, we have reported synthesis of perfluoroalkyltriarylmethanes (molecular propellers) such as **4** by the reaction of 2-perfluoroalkylanilines with phenylmagnesium bromide.¹ The mechanistic pathway, shown for **4**, involves elimination of fluoride from anion derived from **1** followed by addition of PhMgBr to the resultant intermediate product **2** to generate an adduct **3**. Compound **4** is produced in a similar elimination-addition reaction with **3**.

In an attempt to synthesize a triarylmethane that would be more sterically congested than **4**, the substrate **1** (3 mmol) was allowed to react with 2-mesitylmagnesium bromide (10 mmol) in anhydrous THF (50 mL). The reactions were conducted at 23 °C and under reflux conditions. For the 23 °C reaction the GC-MS analysis showed the absence of **1** after 24 h, and under the reflux conditions the reaction was completed after 6 h. A single, albeit different, major product was observed in each case. To our surprise, however, none of the mass spectra, including those for a number of minor products, could be correlated with the expected analog of **4**.



Following standard workup¹ and chromatography on silica gel (hexanes/ether, 4:1) the product of the reaction at 23 °C was identified as 1,3-dimethyl-9-perfluoropropylacridine [**8**, yield 30%, mp 90-93 °C (from hexanes)]. A similar chromatographic separation of the mixture obtained under reflux conditions gave 1,3-dimethyl-9[(*E*)-1-perfluoropropenyl]acridine [**5**, yield 65%, mp 85-87 °C]. Compounds **5** and **8** gave satisfactory results of elemental analysis and were characterized² by MS, ¹H NMR and ¹⁹F NMR. In particular, their ¹H NMR spectra are similar as expected for a similarly substituted acridine system. In the ¹⁹F NMR spectrum of **5** the large coupling constant of 140 Hz between fluorine atoms F1 and F2 of the perfluoropropenyl substituent is indicative of *E*-stereochemistry.³

It is suggested that the formation of either acridine **5** or **8** involves the intermediary of the same dihydroacridine derivative **7** that is generated by intramolecular S_N2' cyclization of **3** (R = Me) or electrocyclization of a mesityl-substituted analog of **2** (not shown) derived from **3**. This unusual intramolecular reaction can be explained in terms of greatly increased steric congestion in **3** (R = Me) in comparison to that of **3** (R = H). Aromatization of **7** to give **8** requires a formal elimination of MeMgBr. This process may take place within aggregated higher-order structures of **7** that are expected to be present at low temperatures. On the other hand, an increase in temperature causes a decrease in aggregation of organometallic compounds. This phenomenon may be responsible for the elimination of fluoride ion and a C4a-methyl group from **7** by a nucleophilic attack of an external nucleophile at the C4a-methyl group to generate **6**. The intermediate product **6** is the suggested direct precursor to **5**. Additional studies of these novel transformations by using a variety of aryl Grignard reagents are in progress.

Acknowledgment. We thank the donors of the Petroleum Research Fund, administered by the American Chemical Society, for partial support of this research.

References and Notes

1. L. Strekowski, H. Lee, S.-Y. Lin, A. Czarny and D. Van Derveer, *J. Org. Chem.* **65**, 7703 (2000)
2. **5**: ¹H NMR (400 MHz, CDCl₃) δ 2.57 (s, 3-Me), 2.85 (s, 1-Me), 7.33 (s, 2-H), 7.65 (t, J=8 Hz, 6-H or 7-H), 7.82 (t, J=8 Hz, 6-H or 7-H), 7.96 (s, 4-H), 8.03 (d, J=8 Hz, 8-H), 8.26 (d, J=8 Hz, 5-H); MS m/z 253 (40, M⁺-CF₃-Me), 268 (80, M⁺-CF₃), 322 (35, M⁺-Me), 337 (100, M⁺). **8**: ¹H NMR (400 MHz, CDCl₃) δ 2.53 (s, 3-Me), 2.71 (narrow m, due to H-F couplings, 1-Me), 7.35 (s, 2-H), 7.57 (t, J=8 Hz, 6-H or 7-H), 7.75 (t, J=8 Hz, 6-H or 7-H), 7.85 (s, 4-H), 8.21 (d, J=8 Hz, 8-H), 8.34 (d, J=8 Hz, 5-H); MS m/z 256 (100, M⁺-C₂F₅), 356 (10, M⁺-F), 375 (75, M⁺)
3. R.S. Macomber, *A Complete Introduction to Modern NMR*, Wiley-Interscience, New York, 1998, p 141

Received on October 7, 2001