

OXAZEPINES AND THIAZEPINES 39*
SYNTHESIS OF TETRACYCLIC 1,5-BENZOTHAZEPINES BY THE REACTION
OF (Z)-3-ARYLIDENE-1-THIOFLAVANONES WITH 2-AMINOTHIOPHENOL

Albert Lévai

Department of Organic Chemistry, University of Debrecen, Egyetem tér 1,
H-4010 Debrecen, Hungary

Dedicated to Prof. Dr. András Messmer on the occasion of his 80th birthday

Abstract: Tetracyclic benzothiazepines **3a-j** have been synthesized by the reaction of 2-aminothiophenol (**1**) and (Z)-3-arylidene-1-thioflavanones **2a-j** in hot toluene by using trifluoroacetic acid catalyst. Structures of all new compounds have been elucidated by IR, ¹H and ¹³C NMR spectroscopic measurements.


Introduction

Owing to their diverse bioactivities (1-10), 1,5-benzothiazepines are important compounds in the drug research. For this reason, numerous representatives have been published in the literature and different methods were introduced for their synthesis. A thoroughly investigated procedure is the preparation of 2,3-dihydro-1,5-benzothiazepines by the reaction of α,β -enones with 2-aminothiophenol (11-20). Their syntheses have also been summarized and discussed in several reviews (21-24). However, only few examples have hitherto been published for the synthesis of tetracyclic 1,5-benzothiazepines (25-30). This fact prompted us to study the synthesis of the tetracyclic 1,5-benzothiazepines in detail and herein we report on the preparation of such compounds by the reaction of (Z)-3-arylidene-1-thioflavanones with 2-aminothiophenol.

Results and Discussion

The 3-arylidene-1-thioflavanones belong to the exocyclic α,β -unsaturated ketones and their (Z)-isomers, where the carbonyl and aryl moieties are on the opposite sides of the carbon-carbon double bond, were synthesized by acid- (31-33) and base-catalyzed (34) condensation of 1-thioflavanones and aromatic aldehydes. The diastereomeric (E)-isomers were obtained by the photoisomerization of the appropriate (Z)-isomers (35,36). Formerly, we have investigated not only the synthesis, but also some chemical transformations of the 3-arylidene-1-thioflavanones. These studies comprised their rearrangement into 3-benzyl-1-thioflavones (37), 1,3-dipolar cycloaddition with diazomethane (38-40) and their oxidative transformations (41-43). As a continuation, now we have performed their reaction with 2-aminothiophenol to afford tetracyclic 1,5-benzothiazepines.

Scheme 1

**2a, 3a:** Ar = C₆H₅**2b, 3b:** Ar = 2-Me-C₆H₄**2c, 3c:** Ar = 3-Me-C₆H₄**2d, 3d:** Ar = 4-Me-C₆H₄**2e, 3e:** Ar = 4-iPr-C₆H₄**2f, 3f:** Ar = 4-MeO-C₆H₄**2g, 3g:** Ar = 4-F-C₆H₄**2h, 3h:** Ar = 2-Cl-C₆H₄**2i, 3i:** Ar = 2-thienyl**2j, 3j:** Ar = 2-naphthyl

2-Aminothiophenol **1** was allowed to react with (*Z*)-3-arylidene-1-thioflavanones **2a-j** in hot toluene by using trifluoroacetic acid as catalyst and tetracyclic 1,5-benzothiazepines **3a-j** were obtained in high (76-93%) yields (Scheme 1). The appropriate 1,5-benzothiazepine derivative was the sole isolable product in each case.

Structures of all new compounds **3a-j** have been elucidated by elemental analyses, IR, ¹H- and ¹³C-NMR spectroscopic measurements. Absence of C=O and NH₂ bands in the IR spectra reveals that not only the Michael addition of the thiol group to the β-carbon atoms of the α,β-unsaturated ketones but also the ring closure took place under these reaction conditions providing 1,5-benzothiazepines **3a-j** which was corroborated by the presence of a characteristic C=N band at ca. 1600 cm⁻¹. In the ¹H-NMR spectra a doublet signal between 3.60 and 3.83 ppm belongs to the 6-H hydrogen, while a doublet-doublet signal of the 6a-H proton is found between 3.57 and 3.90 ppm. The *J*_{6-H,6a-H} = 1.7-2.5 Hz coupling constant values refer to a *gauche*-orientation of these two neighbouring hydrogen atoms. A doublet signal of the 7-H proton is between 5.14 and 5.85 ppm and the ca. 12 Hz *J*_{7-H,6a-H} coupling constant values reveal an *antiperiplanar*-arrangement of these two protons. ¹³C-NMR spectra of compounds **3a-j** corroborate the structures deduced from the IR and ¹H-NMR spectra by the assignment of the aliphatic and the imine carbon atoms (*cf.* Experimental). All these spectroscopic measurements unequivocally prove that 1,5-benzothiazepines **3a-j** were formed

as stereohomogeneous products and no other diastereomers were obtained even as minor components.

In summary, it can be concluded that this simple and convenient procedure can be advantageously utilized for a completeness stereoselective preparation of these tetracyclic 1,5-benzothiazepines with three centres of chirality.

Experimental

Melting points were determined with a Koffler hot-stage apparatus and are uncorrected. ^1H and ^{13}C NMR spectra were recorded on a Varian Gemini 200 spectrometer at 200/50 MHz in CDCl_3 (internal standard TMS, $\delta = 0.0$ ppm) at room temperature. The IR spectra (KBr discs) were measured with a Perkin-Elmer 16 PC instrument. Elemental analyses were performed in-house with a Carlo Erba 1106 EA instrument. TLC was performed on Kieselgel 60 F₂₅₄ (Merck) layer using hexane:acetone (7:3 v/v) as eluent. Starting materials **2a-j** were synthesized by the piperidine-catalyzed reaction 1-thioflavanone and aromatic aldehydes as described earlier by us (34).

General procedure for the synthesis of compounds **3a-j**

A mixture of 2-aminothiophenol (**1**, 6.0 mmol), (*Z*)-3-arylidene-1-thioflavanone (**2a-j**, 5.0 mmol), trifluoroacetic acid (0.5 ml) and toluene (30 ml) was refluxed for 3 h. The solvent was evaporated under reduced pressure and the residue was crystallized from methanol to afford tetracyclic benzothiazepines **3a-j**.

6a,7-Dihydro-6,7-diphenyl-6H-[1]benzothiopyrano[3,4-c][1,5]benzothiazepine (3a): Yield 89%, m.p. 174-175 °C; IR: $\nu_{\text{C=N}}$ 1596 cm^{-1} ; ^1H -NMR (δ): 3.62 (1H, d, $J = 2.3$ Hz, 6-H), 3.78 (1H, dd, $J = 12.0, 2.4$ Hz, 6a-H), 5.16 (1H, d, $J = 12.0$ Hz, 7-H), 6.97-8.62 (18 arom. H, m); ^{13}C -NMR (δ): 44.6 (C-6a), 47.1 (C-6), 61.5 (C-7), 164.1 (C-14).

Anal. Calcd. for $\text{C}_{28}\text{H}_{21}\text{NS}_2$: C, 77.22; H, 4.86; N, 3.21. Found: C, 77.28; H, 4.84; N, 3.23.

6a,7-Dihydro-7-(2-methylphenyl)-6-phenyl-6H-[1]benzothiopyrano[3,4-c][1,5]benzothiazepine (3b): Yield 88%, m.p. 192-193 °C; IR: $\nu_{\text{C=N}}$ 1598 cm^{-1} ; ^1H -NMR (δ): 2.58 (3H, s, CH_3), 3.78 (1H, d, $J = 2.3$ Hz, 6-H), 3.80 (1H, dd, $J = 11.8, 2.5$ Hz, 6a-H), 5.54 (1H, d, $J = 12.0$ Hz, 7-H), 7.02-8.61 (17 arom. H, m); ^{13}C -NMR (δ): 20.1 (CH_3), 44.8 (C-6a), 46.7 (C-6), 55.8 (C-7), 164.0 (C-14).

Anal. Calcd. for $\text{C}_{29}\text{H}_{23}\text{NS}_2$: C, 77.48; H, 5.16; N, 3.11. Found: C, 77.53; H, 5.14; N, 3.09.

6a,7-Dihydro-7-(3-methylphenyl)-6-phenyl-6H-[1]benzothiopyrano[3,4-c][1,5]benzothiazepine (3c): Yield 84%, m.p. 165-166 °C; IR: $\nu_{\text{C=N}}$ 1600 cm^{-1} ; ^1H -NMR (δ): 2.34 (3H, s, CH_3), 3.68 (1H, d, $J = 2.2$ Hz, 6-H), 3.80 (1H, dd, $J = 11.8, 2.6$ Hz, 6a-H), 5.55 (1H, d, $J = 12.1$ Hz, 7-H), 6.94-8.61 (17 arom. H, m); ^{13}C -NMR (δ): 21.3 (CH_3), 44.7 (C-6a), 46.9 (C-6), 61.5 (C-7), 164.2 (C-14).

Anal. Calcd. for $\text{C}_{29}\text{H}_{23}\text{NS}_2$: C, 77.48; H, 5.16; N, 3.11. Found: C, 77.44; H, 5.17; N, 3.13.

6a,7-Dihydro-7-(4-methylphenyl)-6-phenyl-6H-[1]benzothiopyrano[3,4-c][1,5]benzothiazepine (3d): Yield 93%, m.p. 126-127 °C; IR: $\nu_{\text{C=N}}$ 1604 cm^{-1} ; ^1H -NMR (δ): 2.37 (3H, s,

CH₃), 3.68 (1H, d, J = 2.3 Hz, 6-H), 3.78 (1H, dd, J = 12.0, 2.4 Hz, 6a-H), 5.17 (1H, d, J = 11.9 Hz, 7-H), 7.02-8.64 (17 arom. H, m); ¹³C-NMR (δ): 21.0 (CH₃), 44.8 (C-6a), 47.1 (C-6), 61.4 (C-7), 164.2 (C-14).

Anal. Calcd. for C₂₉H₂₃NS₂: C, 77.48; H, 5.16; N, 3.11. Found: C, 77.52; H, 5.18; N, 3.12.

6a,7-Dihydro-7-(4-isopropylphenyl)-6-phenyl-6H-[1]benzothiopyrano[3,4-c][1,5]benzothiazepine (3e): Yield 85%, m.p. 160-161 °C; IR: νC=N 1602 cm⁻¹; ¹H-NMR (δ): (6H, d, CH(CH₃)₂), 2.90 (1H, m, CH(CH₃)₂), 3.66 (1H, d, J = 2.5 Hz, 6-H), 3.78 (1H, dd, J = 11.9, 2.4 Hz, 6a-H), 5.17 (1H, d, J = 11.9 Hz, 7-H), 7.02-8.61 (17 arom. H, m); ¹³C-NMR (δ): 23.6 (2 CH₃), 33.6 (CH(CH₃)₂), 44.8 (C-6a), 47.2 (C-6), 61.4 (C-7), 164.2 (C-14).

Anal. Calcd. for C₃₁H₂₇NS₂: C, 77.96; H, 5.70; N, 2.93. Found: C, 77.91; H, 5.72; N, 2.94.

6a,7-Dihydro-7-(4-methoxyphenyl)-6-phenyl-6H-[1]benzothiopyrano[3,4-c][1,5]benzothiazepine (3f): Yield 81%, m.p. 177-178 °C; IR: νC=N 1604 cm⁻¹; ¹H-NMR (δ): 3.66 (1H, d, J = 2.5 Hz, 6-H), 3.78 (1H, dd, J = 12.1, 2.8 Hz, 6a-H), 3.83 (3H, s, OCH₃), 5.14 (1H, d, J = 11.8 Hz, 7-H), 6.83-8.62 (17 arom. H, m); ¹³C-NMR (δ): 44.8 (C-6a), 47.3 (C-6), 55.2 (OCH₃), 61.1 (C-7), 164.3 (C-14).

Anal. Calcd. for C₂₉H₂₃NOS₂: C, 74.82; H, 4.98; N, 3.01. Found: C, 74.78; H, 4.96; N, 3.02.

6a,7-Dihydro-7-(4-fluorophenyl)-6-phenyl-6H-[1]benzothiopyrano[3,4-c][1,5]benzothiazepine (3g): Yield 84%, m.p. 169-170 °C; IR: νC=N 1600 cm⁻¹; ¹H-NMR (δ): 3.60 (1H, d, J = 2.5 Hz, 6-H), 3.71 (1H, dd, J = 12.0, 2.4 Hz, 6a-H), 5.15 (1H, d, J = 12.0 Hz, 7-H), 6.97-8.60 (17 arom. H, m); ¹³C-NMR (δ): 44.7 (C-6a), 47.2 (C-6), 60.7 (C-7), 163.9 (C-14).

Anal. Calcd. for C₂₈H₂₀FNS₂: C, 74.16; H, 4.45; N, 3.09. Found: C, 74.19; H, 4.47; N, 3.07.

6a,7-Dihydro-7-(2-chlorophenyl)-6-phenyl-6H-[1]benzothiopyrano[3,4-c][1,5]benzothiazepine (3h): Yield 80%, m.p. 169-170 °C; IR: νC=N 1598 cm⁻¹; ¹H-NMR (δ): 3.70 (1H, d, J = 1.7 Hz, 6-H), 3.86 (1H, dd, J = 12.0 Hz, 2.2 Hz, 6a-H), 5.85 (1H, d, J = 12.1 Hz, 7-H), 6.98-8.63 (17 arom. H, m); ¹³C-NMR (δ): 44.7 (C-6a), 46.8 (C-6), 55.6 (C-7), 163.8 (C-14).

Anal. Calcd. for C₂₈H₂₀ClNS₂: C, 71.56; H, 4.29; N, 2.98. Found: C, 71.59; H, 4.31; N, 2.97.

6,7-Dihydro-6-phenyl-7-(2-thienyl)-6H-[1]benzothiopyrano[3,4-c][1,5]benzothiazepine (3i): Yield 76%, m.p. 166-167 °C; IR: νC=N 1598 cm⁻¹; ¹H-NMR (δ): 3.57 (1H, dd, J = 11.8, 2.4 Hz, 6a-H), 3.83 (1H, d, J = 2.3 Hz, 6-H), 5.49 (1H, d, J = 12.0 Hz, 7-H), 6.90-8.57 (16 arom. H, m); ¹³C-NMR (δ): 44.9 (C-6a), 48.7 (C-6), 57.1 (C-7), 163.6 (C-14).

Anal. Calcd. for C₂₆H₁₉NS₃: C, 70.74; H, 4.34; N, 3.17. Found: C, 70.77; H, 4.36; N, 3.18.

6a,7-Dihydro-7-(2-naphthyl)-6-phenyl-6H-[1]benzothiopyrano[3,4-c][1,5]benzothiazepine (3j): Yield 78%, m.p. 177-178 °C; IR: νC=N 1600 cm⁻¹; ¹H-NMR (δ): 3.64 (1H, d, J = 2.4 Hz, 6-H), 3.90 (1H, dd, J = 12.0, 2.5 Hz, 6a-H), 5.35 (1H, d, J = 12.1 Hz, 7-H), 6.98-8.64 (20 arom. H, m); ¹³C-NMR (δ): 45.3 (C-6a), 47.0 (C-6), 62.3 (C-7), 164.6 (C-14).

Anal. Calcd. for C₃₂H₂₃NS₂: C, 79.17; H, 4.77; N, 2.88. Found: C, 79.14; H, 4.75; N, 2.87.

Acknowledgements

The present study was sponsored by the Hungarian National Research Foundation (Grant No. OTKA-T034123) for which our gratitude is expressed. Technical assistance of Mrs. M. Nagy is highly appreciated.

References

*For Part 38, see A. Lévai, *Heterocycl. Commun.* **5**, 359 (1999)

- (1) M. Sato, T. Nagao, I. Yamaguchi, H. Nakiyama and A. Kiyomoto, *Arzneim-Forsch*, **21**, 1338 (1971)
- (2) K. Yamada, T. Shimamura and H. Nakiyama, *Japan J. Pharmacol.*, **23**, 321 (1973)
- (3) T. Nagao, M. Sato, H. Nakiyama and A. Kiyomoto, *Chem. Pharm. Bull.*, **21**, 92 (1973)
- (4) J. Slade, J. L. Stanton, D. Ben-David and G. Mazzenga, *J. Med. Chem.*, **28**, 1517 (1986)
- (5) H. Yamamoto and H. Asai, *Chem. Pharm. Bull.*, **34**, 3844 (1986)
- (6) M. J. Kendall and J. Okopski, *J. Clin. Hospital Pharm.*, **11**, 159 (1986)
- (7) Y. Inada, K. Itoh, K. Kamiya, H. Sugihara and K. Nishikawa, *Japan J. Pharmacol.*, **47**, 135 (1988)
- (8) S. Murata, K. Kikkawa, H. Yabana and T. Nagao, *Arzneim.-Forsch.*, **38**, 521 (1988)
- (9) K. Kikkawa, S. Murata and T. Nagao, *Arzneim.-Forsch.*, **38**, 526 (1988)
- (10) H. Narita, M. Gaino, T. Suzuki, H. Kurosawa, H. Inoue and T. Nagao, *Chem. Pharm. Bull.*, **38**, 407 (1990)
- (11) W. Ried and W. Marx, *Chem. Ber.*, **90**, 2683 (1957)
- (12) W. D. Stephens and L. Field, *J. Org. Chem.*, **24**, 1576 (1959)
- (13) A. Lévai and R. Bognár, *Acta Chim. Acad. Sci. Hung.*, **88**, 293 (1976)
- (14) A. Lévai and R. Bognár, *Acta Chim. Acad. Sci. Hung.*, **92**, 415 (1977)
- (15) A. Lévai, *Pharmazie*, **34**, 439 (1979)
- (16) A. Lévai, R. Bognár and J. Kajtár, *Acta Chim. Acad. Sci. Hung.*, **103**, 27 (1980)
- (17) A. Lévai, *Pharmazie*, **36**, 449 (1981)
- (18) K. P. Jadhav and D. B. Ingle, *Indian J. Chem.*, **22B**, 180 (1983)
- (19) A. K. Gupta, V. K. Singh and U. C. Pant, *Indian J. Chem.*, **22B**, 1057 (1983)
- (20) U. C. Pant, B. S. Gaur and M. Chugh, *Indian J. Chem.*, **27B**, 752 (1988)
- (21) A. Lévai, *Trends Heterocycl. Chem.*, **4**, 51 (1995)
- (22) A. Chimirri, R. Gitto, S. Grasso, A. M. Monforte and M. Zappala, *Adv. Heterocycl. Chem.*, **63**, 61 (1995)

- (23) A. Lévai, *Pharmazie*, **54**, 719 (1999)
- (24) A. Lévai, *J. Heterocycl. Chem.*, **37**, 199 (2000)
- (25) O. Hideg-Hankovszky and K. Hideg, *Acta Chim. Acad. Sci. Hung.*, **68**, 403 (1971)
- (26) A. Lévai, G. Tóth and Á. Szöllősy, *Stud. Org. Chem. (Amsterdam)*, **11**, 41 (1982)
- (27) G. Tóth, Á. Szöllősy, A. Lévai and H. Duddeck, *Org. Magn. Reson.*, **20**, 133 (1982)
- (28) S. Pant, B. C. Joshi and U. C. Pant, *Indian J. Chem.*, **33B**, 869 (1993)
- (29) A. Lévai, *Sci. Pharm.*, **64**, 523 (1996)
- (30) A. Lévai, *Heterocycl. Commun.*, **3**, 211 (1997)
- (31) K. Auwers and F. Arndt, *Ber. Dtsch. Chem. Ges.*, **42**, 2706 (1909)
- (32) F. Arndt, W. Fleming, E. Scholz and V. Löwensohn, *Ber. Dtsch. Chem. Ges.*, **56**, 1269 (1923)
- (33) J. J. Biernes and W. I. O'Sullivan, *Proc. Roy. Irish Acad.*, **77B**, 331 (1977)
- (34) A. Lévai, Á. Szöllősy and G. Tóth, *Acta Chim. Hung.*, **128**, 359 (1991)
- (35) G. Tóth, Á. Szöllősy, A. Lévai, G. Oszbach, W. Dietrich and H. Kühne, *Magn. Reson. Chem.*, **29**, 801 (1991)
- (36) G. Tóth, J. Halász, A. Lévai, C. Nemes and T. Patonay, *J. Chem. Soc., Perkin Trans. 2*, 547 (1996)
- (37) A. Lévai and Z. Szabó, *Bull. Soc. Chim. Fr.*, **128**, 976 (1991)
- (38) G. Tóth, A. Lévai, Á. Szöllősy and H. Duddeck, *Tetrahedron*, **49**, 863 (1993)
- (39) A. Lévai, A. M. S. Silva, T. Patonay and J. A. S. Cavaleiro, *J. Heterocycl. Chem.*, **36**, 1215 (1999)
- (40) A. Lévai, *Org. Prep. Proced. Int.*, in press
- (41) W. Adam, D. Golsch, L. Hadjarapoglou, A. Lévai, C. Nemes and T. Patonay, *Tetrahedron*, **50**, 13113 (1994)
- (42) A. Lévai, G. Tóth, J. Kovács, T. Patonay and E. B. Vass, *Monatsh. Chem.*, **132**, 707 (2001)
- (43) G. Tóth, J. Kovács, A. Lévai, E. Kleinpeter and A. Koch, *Magn. Reson. Chem.*, **39**, 251 (2001)