


A SHORT METHOD FOR THE SYNTHESIS OF 4,6-DIMETHOXY-1-AZAAURONES

Martin Ata Lawson, Anne-Marie Mariotte and Ahcène Boumendjel*
Département de Pharmacochimie Moléculaire, UMR-CNRS 5063. Faculté de Pharmacie de
Grenoble, 38706 La Tronche – France

Abstract: Aurones (Benzylidenebenzofuran-3(2*H*)-one) are naturally occurring compounds, which are isomers of the well known flavones. Like flavones, ring A dioxygenated aurones especially on 4 and 6 positions are frequently found in nature and this substitution pattern is surely responsible for aurones bioactivity. In the present communication, we report the synthesis of ring A-oxygenated 1-azaurones. These molecules adopt conformations very close to aurones and the presence of the nitrogen can allow derivitization for SAR studies and also making ammonium salts, suitable for the molecule hydrosolubility. The synthesis of 4,6-dimethoxy-1-azaurones was achieved by condensation of a benzaldehyde derivative with 1-acetyl-4,6-dimethoxy-2,3-dihydro-1*H*-indole-3-one. The latter was prepared in three steps starting from 3,5-dimethoxyaniline and chloroacetonitrile. If the benzaldehyde derivative is appropriately chosen, the method can allow the preparation of a large number of azaurones.

Introduction


Aurones are secondary metabolites belonging to the flavonoid family of naturally occurring compounds (1). In the plant arena, these 2-phenylidenebenzofuranone which are structurally isomeric of flavones contribute to plants color. The biological activities of flavones are well investigated. In particular, ring A methoxylated or hydroxylated flavones are of current interest (2-4). Compared to flavones, aurones still poorly investigated as therapeutics (5-7). Our interest in the biological properties of ring-A methoxylated and/or hydroxylated flavonoids prompted us to investigate the synthesis of 4,6-oxygenated aurones and their 1-azaanalogs (azaurones) (8). The latter are structurally similar to the naturally occurring aurones and in addition, they also have an available nitrogen for further extension and for branching different substituents that may give rise to beneficial interactions with the biological active site. Therefore, the search for an approach toward 4,6-dimethoxy-1-azaurones was undertaken in the framework of a project related to the synthesis of biologically active flavonoid analogs (9-11).

**Figure 1.**

The literature survey indicated that only simple azaaurones (unsubstituted or monosubstituted on the A-ring) have been investigated (12). Owing to the important role played by the simultaneous presence of hydroxyl or methoxy groups on 5 and 7 positions, we report herein a simple method for preparation of 4,6-dimethoxyazaaurones.

Results and Discussion:


Azaaurones were prepared from 1-acetyl-4,6-dimethoxy-2,3-dihydro-1*H*-indole **2** by base-catalyzed condensation with a benzaldehyde derivative. Oxindole **2** was prepared starting from 3,5-dimethoxyaniline and chloroacetonitrile (Scheme 1).

**Scheme 1.** Synthesis of 4,6-dimethoxy-1-azaurones

Ketone **1** was synthesized by a specific ortho chloroacetylation of 3,5-dimethoxyaniline, according to the method reported by Sugawara (13). Cyclisation of **1** with K_2CO_3 in acetone followed by acetylation with acetic acid anhydride gave 1-acetyl-4,6-dimethoxy-2,3-dihydro-1*H*-indole-3-one **2** in 50% yield. It should be mentioned that analogs of **2** (mono or unsubstituted on ring-A) have been used as starting blocks for the synthesis of biologically active compounds (14-17). The key intermediate **2** can be easily prepared in multi-grams scale without yield lowering. The final step is the condensation of **2** with a benzaldehyde derivative which gives directly the desired 4,6-

dimethoxyazaurone. The structure of azaurones **3** were determined by NMR and mass spectrometry. The configuration of the double bond is exclusively *Z* as can be assigned on the basis of the chemical shift of the olefinic proton (δ 6.7 ppm) and by comparing with literature (18).

Table 1. Structures and physicochemical data of synthesized 4,6-dimethoxy-1-azaurones **3**.



R	Yield (%) ^{a,b}	mp (°C)	¹ H-NMR (CDCl ₃) δ
4'-H	75	152 - 154	7.5 (dd, 2H, $J_1 = 7.7$ Hz, $J_2 = 1.4$ Hz); 7.4 (dd, 1H, $J_1 = 7.7$ Hz, $J_2 = 1.4$ Hz, H-4'); 7.3 (dd, 2H, $J_1 = 7.7$ Hz, $J_2 = 1.4$ Hz); 6.8 (sl, 1H, NH); 6.7 (s, 1H); 6.1 (d, 1H, $J = 1.9$ Hz); 5.9 (d, 1H, $J = 1.9$ Hz), 3.9 (s, 3H); 3.8 (s, 3H).
4'-Et	70	215 - 217	7.45 (dd, 2H, $J_1 = 8.2$ Hz, $J_2 = 0.4$ Hz); 7.25 (dd, 2H, $J_1 = 8.2$ Hz, $J_2 = 0.4$ Hz); 7.1 (sl, 1H); 6.7 (s, 1H); 6.1 (d, 1H, $J = 1.8$ Hz); 5.9 (d, 1H, $J = 1.8$ Hz), 3.9 (s, 3H); 3.8 (s, 3H); 2.7 (q, 2H, $J = 6$ Hz); 1.3 (t, $J = 6$ Hz).
4'-SMe	60	228 - 230	7.4 (dd, 2H, $J_1 = 8.5$ Hz, $J_2 = 0.4$ Hz); 7.3 (dd, 2H, $J_1 = 8.5$ Hz, $J_2 = 0.4$ Hz); 6.8 (sl, 1H); 6.7 (s, 1H); 6.1 (d, 1H, $J = 1.6$ Hz, H-5); 5.9 (d, 1H, $J = 1.6$ Hz), 3.9 (s, 3H); 3.8 (s, 3H); 2.5 (s, 3H).
4'-Cl	80	146 - 148	7.4 (dd, 2H, $J_1 = 8.5$ Hz, $J_2 = 0.4$ Hz); 7.3 (dd, 2H, $J_1 = 8.5$ Hz, $J_2 = 0.4$ Hz); 6.6 (s, 1H); 6.1 (d, 1H, $J = 1.6$ Hz); 5.85 (d, 1H, $J = 1.6$ Hz), 3.85 (s, 3H); 3.8 (s, 3H).
2',4',6'-triOMe	60	109 - 111	7.6 (sl, 1H); 6.95 (s); 6.7 (s, 1H); 6.55 (s, 1H); 6.0 (d, 1H, $J = 1.8$ Hz); 5.85 (d, 1H, $J = 1.8$ Hz), 3.9 (5s, 15H).
Ph-R = 2-naphthyl	63	254 - 256	7.9 (ddd, 2H, $J_1 = 8.7$ Hz, $J_2 = 1.5$ Hz, $J_3 = 0.4$ Hz); 7.8 (dd, 2H, $J_1 = 8.7$ Hz, $J_2 = 1.5$ Hz); 7.6 (dd, 1H, $J_1 = 8.7$ Hz, $J_2 = 0.4$ Hz); 7.4 (dd, 2H, $J_1 = 8.7$ Hz, $J_2 = 0.4$ Hz); 7.15 (sl, 1H); 6.85 (s, 1H); 6.1 (d, 1H, $J = 1.5$ Hz); 5.9 (d, 1H, $J = 1.5$ Hz), 3.9 (s, 3H); 3.8 (s, 3H).

^aReported yields are those of purified compounds. ^bMass spectrometry performed on intermediates and final compounds gave satisfactory analysis.

In conclusion, if appropriately substituted benzaldehydes are selected, our synthetic approach constitutes a simple and high yield procedure for the synthesis of divers 4,6-dimethoxy-1-

azaaurones which can be used as analogs of aurones, especially in studying aurones structure-activity relationship.

References

- (1) J.R. Harborne (Ed.), *The Flavonoids: Advances in Research Since 1986*, Chapman and Hall, London, 1994.
- (2) G. Conseil, H. Baubichon-Cortay, G. Dayan G, J.M. Jault, D. Barron and A. Di Pietro, *Proc. Natl. Acad. Sci. USA*. **95**, 9831 (1998).
- (3) F. Sicheri, I. Moarefi, J. Kuriyan, *Nature* **385**, 602 (1997).
- (4) F.W. De Azevedo, H.J. Mueller-Dieckmann, U. Schulze-Gahmen, P.J. Worland, E. Sausville and S.H. Kim, *Proc. Natl. Acad. Sci. USA*. **93**, 2735 (1996).
- (5) M. Aufmkolk, J. Koerhle, R.D. Hesch and V. Cody, *J. Biol. Chem.* **261**, 11623 (1986).
- (6) O. Kayser, A.F. Kiderlen, U. Folkens and H. Kolodziej, *Planta Medica* **65**, 316 (1999) and references therein.
- (7) C. J. W. Brooks and D.G. Watson, *Nat. Prod. Rep.* 427 (1985).
- (8) C. Beney, A.M. Mariotte and A. Boumendjel, *Heterocycles* **55**, 967 (2001).
- (9) F. Bois, C. Beney, A. Boumendjel, A.M. Mariotte, G. Conseil and A. Di Pietro *A. J. Med. Chem.* **41**, 4161 (1998).
- (10) F. Bois, A. Boumendjel, A.M. Mariotte, G. Conseil and A. Di Pietro, *Bioorg. Med. Chem.* **7**, 2691 (1999).
- (11) A. Boumendjel, F. Bois, C. Beney, A.M. Mariotte, G. Conseil and A. Di Pietro, *Bioorg. Med. Chem. Lett.* **11**, 75 (2001).
- (12) A.R. Katritzky, Q. L. Li and Q. Fan, *J. Heterocycl. Chem.* **25**, 1287 (1998).
- (13) T. Sugawara, M. Adachi, K. Sasakura and A. Kitagawa, *J. Org. Chem.* **44**, 578 (1979).
- (14) D.E. Bierer, D.M. Fort, C.D. Mendez, J. Luo, P. A. Imbach, L. G. Dubenko, S.D. Jolad, R.E. Gerber, J. Litvak, Q. Lu, P. Zhang, M. J. Reed, N. Waldeck, R.C. Bruening, B.K. Noamesi, R.F. Hector, T.J. Carlson and S.R. King *J. Med. Chem.* **41**, 894 (1998).
- (15) J.Y. Mérour, P. Gadonneix, B. Malapel-Andrieu and E. Desarbre, *Tetrahedron* **57**, 1995 (2001).
- (16) J.Y. Mérour, J.Y. Coadou, F. Tatibouet. *Synthesis* 1053 (1982).
- (17) J.Y. Mérour and A. Mérour. *Synthesis* 767 (1994).
- (18) K. Thakkar, M. Cushman. *J. Org. Chem.* **60**, 6499 (1995).

Received on December 16, 2002