

FACILE SYNTHESIS OF DIMERIC HEPTAMETHINE CYANINE DYES CONTAINING A LINKER AT THE MESO POSITIONS

Mariusz Mojzych,[#] Aldona Raszkievicz and Lucjan Strekowski*

Department of Chemistry, Georgia State University, Atlanta, Georgia 30302-4098, USA

[#]On leave of absence from Department of Chemistry, Academy of Podlasie, 08-110 Siedlce, Poland

e-mail: lucjan@gsu.edu

Abstract: The synthesized dyes contain 1-butyl-3,3-dimethyl-3*H*-indolium or benzo[*e*] analogs as the terminal subunits and diverse chains linking the central meso positions of the individual chromophores. The dimeric dyes show absorption in the red (> 600 nm) or near-infrared (> 700 nm) regions.


Introduction

Red and near-infrared dyes have found broad analytical applications in enzymatic assays,¹ immunoassays,² drug displacement studies,³ DNA sequencing,⁴ determination of solvent hydrophobicity⁵ and pH studies,⁶ among other things. Currently there is a substantial interest in hydrophobic dimeric dyes in which two chromophoric subunits are linked by a conformationally flexible chain. Under low concentration conditions in aqueous solution, these hydrophobic bichromophoric molecules tend to exist in an intramolecular clam-shell conformation with the two chromophores in close proximity to each other. In general, the stacking interactions are less important in solvents of low polarity. Some dimeric dyes undergo dissociation of the intramolecular foldamer before binding of the open form to proteins⁷⁻¹⁰ and nucleic acids.¹¹ Binding of the open form of the dimeric dye to a biopolymer usually results in a bathochromic shift in absorption and a greatly increased quantum yield of fluorescence. Several bichromophoric dyes bind metal cations and the resulting complexes show different spectral properties in comparison to non-complexed dyes.¹²⁻¹⁴ It has also been found that some bichromophoric cyanine dyes exhibit strongly enhanced fluorescence upon interaction with hydrophobic fats of fingerprints, which results in a clear fluorescence image of the fingerprint.^{8,15} The bichromophoric dyes reported to date are dimeric cyanines and analogs that contain a chain linking the nitrogen atoms of the terminal heterocyclic systems of the two chromophores.

Results and discussion

Synthesis of new dimeric cyanine dyes **5** – **10** (Scheme 1) with a linker connecting the two cyanine components at their central meso position is described in this report. In compounds **5** and **6** the linker is a relatively inflexible 1,4-phenylene unit that joins two cyanines through amino, oxy or thio groups. Compounds **7** – **10** contain a flexible oligomethylene chain that is α,ω -disubstituted with amino or thio functions. Dimers **5**, **7**, and **9** are derivatives of 3,3-dimethyl-3*H*-indolium heptamethine cyanines while dimers **6**, **8**, and **10** are their benzo[*e*] analogs. The amino-substituted dyes show absorption in the red region of the visible spectrum and the oxy- and thio-substituted dyes absorb in the near-infrared region of the electromagnetic spectrum.


Scheme 1



The key intermediate products are meso-chloro-substituted cyanines **3** and **4**. These dyes were obtained by condensation of indolium salts **1** with Vilsmeier-Hack reagent **2** by using a standard procedure. The desired dimeric dyes **5** – **10** were synthesized by treatment of chloro-substituted cyanine **3** or **4** with a corresponding bifunctional nucleophilic reagent, as shown in Scheme 1. All final products were purified by silica gel chromatography followed by crystallization. Although the yields of the analytically pure dyes thus obtained ranged only from 10% to 15%, the procedure is experimentally simple and the yields

are highly reproducible. An attempt to improve the efficiency of the preparation of the dimeric dyes is illustrated in Scheme 2. In this modified synthesis the chloro-substituted dye **3** was allowed to react with an excess of 1,4-phenylenediamine or 1,4-hydroquinone to give the respective mono-substituted dye **11a** or **11b**. The reaction of these intermediate products with dye **3** provided dimeric dyes **5a** and **5b** that were identical with the respective products obtained by using the one-pot procedure discussed above. Although successful, the two-pot procedure does not result in an improved yield.

Scheme 2



Experimental

General: Indolium salts **1**⁹ and Vilsmeier-Haack reagent **2**¹⁶ were synthesized as previously described. All dimeric dyes were purified by silica gel chromatography eluting with a mixture of dichloromethane and methanol (30:1). For the subsequent crystallization, dye was dissolved in a minimum amount of ethyl acetate and the solution was treated with an equal volume of ether. Melting points for **6**, **8**, and **10** are provided below. The remaining dyes undergo decomposition before melting. All products were characterized by mass spectrometry (MALDI) and ¹H NMR spectroscopy and gave satisfactory results of elemental analysis (C, ± 0.4; H ± 0.4; N ± 0.4). Their composition, as obtained from microanalysis, is provided below. Visible and near-infrared absorption spectra were obtained in methanol.

Synthesis of 3 and 4: A mixture of the corresponding salt **1** (4 mmol), Vilsmeier-Haack reagent **2** (0.72 g, 2 mmol) and anhydrous sodium acetate (0.66 g, 8 mmol) in ethanol (100 mL) was heated under reflux for 5 h, then cooled and concentrated under reduced pressure. The residue was extracted with dichloromethane, the solution was concentrated, and the solid residue was crystallized from methanol/ether (1:5). Compounds **3** and **4** gave spectral data virtually identical with those published previously.⁹

Synthesis of 5 – 10: A mixture of dye **3** or **4** (1.0 mmol), 1,4-phenylenediamine, 1,4-hydroquinone, 1,4-phenylenedithiol, an α,ω -alkanediamine or α,ω -alkanedithiol (0.5 mmol), and anhydrous potassium carbonate (0.17 g, 1.0 mmol) in anhydrous acetonitrile (50 mL) was heated under reflux under a nitrogen atmosphere until a TLC analysis (silica gel, dichloromethane/methanol, 20:1) showed the absence of the dye substrate (several hours). Heating was continued for an additional 2 h. Concentration under a reduced pressure was followed by purification of the dimeric dye as described above. The yields of analytically pure products **5** – **10** were between 10% and 15%. Compound **5a**: C₈₂H₁₀₂I₂N₆, VIS λ_{max} = 656 nm. Compound **5b**: C₈₂H₁₀₀I₂N₄O₂, NIR λ_{max} = 699 nm. Compound **5c**: C₈₂H₁₀₀I₂N₄S₂•H₂O, NIR λ_{max} = 722 nm. Compound **6a**: C₉₈H₁₁₀I₂N₆, NIR λ_{max} = 675 nm. Compound **6b**: C₉₈H₁₀₈I₂N₄O₂•H₂O, mp 194 – 195 °C, NIR λ_{max} = 783 nm. Compound **6c**: C₉₈H₁₀₈I₂N₄S₂•2H₂O, mp 178 – 179 °C, NIR λ_{max} = 815 nm. Compound **7a**: C₈₀H₁₀₆I₂N₆•0.5H₂O, VIS λ_{max} = 619 nm. Compound **7b**: C₈₄H₁₁₄I₂N₆, C₉₈H₁₀₈I₂N₄O₂•H₂O, VIS λ_{max} = 622 nm. Compound **7c**: C₈₈H₁₂₂I₂N₆•0.5H₂O, VIS λ_{max} = 622 nm. Compound **8a**: C₉₆H₁₁₄I₂N₆•2H₂O, mp 168 – 169 °C, VIS λ_{max} = 653 nm. Compound **8b**:

$C_{100}H_{122}I_2N_6 \cdot 2H_2O$, mp 189 – 190 °C, VIS λ_{max} = 656 nm. Compound **8c**: $C_{104}H_{130}I_2N_6 \cdot 3H_2O$, mp 175 – 176 °C, VIS λ_{max} = 656 nm. Compound **9a**: $C_{80}H_{104}I_2N_4S_2 \cdot H_2O$, NIR λ_{max} = 777 nm. Compound **9b**: $C_{84}H_{112}I_2N_4S_2 \cdot 2.5H_2O$, NIR λ_{max} = 779 nm. Compound **10a**: $C_{96}H_{112}I_2N_4S_2 \cdot 2H_2O$, mp 182 – 183 °C, NIR λ_{max} = 800 nm. Compound **10b**: $C_{100}H_{120}I_2N_4S_2 \cdot 3H_2O$, mp 170 – 171 °C, NIR λ_{max} = 808 nm.

Synthesis of 11: A mixture of dye **3** (0.35 g, 0.5 mmol), 1,4-phenylenediamine (0.11 g, 1.0 mmol) or 1,4-hydroquinone (0.11 g, 1.0 mmol), and anhydrous potassium carbonate (0.086 g, 0.5 mmol) in anhydrous acetonitrile (10 mL) was heated under reflux under a nitrogen atmosphere until a TLC analysis (silica gel, dichloromethane/methanol, 20:1) showed the absence of **3** (several hours). Concentration of the mixture under a reduced pressure followed by chromatography and crystallization as described above gave **11a** or **11b**. Compound **11a**: yield 40%, $C_{44}H_{55}I_2N_4$, VIS λ_{max} = 648 nm. Compound **11b**: yield 43%, $C_{44}H_{53}I_2N_2O_2$, NIR λ_{max} = 760 nm.

Synthesis of 5 from 11: An equimolar mixture of **11a** or **11b** and **3** was allowed to react by using the procedure for **5** described above.

References

1. K.B. Male and J.H. Luong, *Appl. Biochem. Biotechnol.* **61**, 267 (1996)
2. M.I. Daneshvar, J.M. Peralta, G.A. Casay, N. Narayanan, L. Evans, G. Patonay and L. Strekowski, *J. Immunol. Methods* **226**, 119 (1999)
3. J. Sowell, J.C. Mason, L. Strekowski and G. Patonay, *Electrophoresis* **22**, 2512 (2001)
4. E. Waddell, Y. Wang, W. Stryjewski, S. McWhorter, A.C. Henry, D. Evans, R.L. McCarley and S.A. Soper, *Anal. Chem.* **72**, 5907 (2000)
5. C.Q. Zhu, H. Zheng, D.H. Li, S.H. Li and J.G. Xu, *Chin. J. Chem.* **22**, 1319 (2004)
6. L. Tarazi, N. Narayanan and G. Patonay, *Microchem. J.* **64**, 247 (2000)
7. J. Kim, A. Watson, M. Henary and G. Patonay, *Near-Infrared Cyanine Dye-Protein Interactions*, in R.R. Gupta (Ser. Ed.), L. Strekowski (Vol. Ed.), *Topics Heterocycl. Chem.*, Vol. 14, Polymethine Heterocyclic Dyes, Springer, Berlin, 2008, pp. 11 – 29
8. G. Patonay, L. Strekowski, J.S. Kim and M. Henary, *NIR News* **18**, 7 (2007)
9. G. Patonay, J.S. Kim, R. Kodagahally and L. Strekowski, *Appl. Spectrosc.* **59**, 682 (2005)
10. J.S. Kim, R. Kodagahally, L. Strekowski and G. Patonay, *Talanta* **67**, 947 (2005)
11. B.A. Armitage, *Cyanine Dye-Nucleic Acid Interactions*, in R.R. Gupta (Ser. Ed.), L. Strekowski (Vol. Ed.), *Topics Heterocycl. Chem.*, Vol. 14, Polymethine Heterocyclic Dyes, Springer, Berlin, 2008, pp. 31 – 39
12. S. Yagi, A. Nakamura, D. Watanabe and H. Nakazumi, *Dyes Pigments* **80**, 98 (2009)
13. S. Yagi, Y. Hyodo, M. Hirose, H. Nakazumi, Y. Sakurai and A. Ajayaghosh, *Org. Lett.* **9**, 1999 (2007)
14. E. Arunkumar, A. Ajayaghosh and J. Daub, *J. Am. Chem. Soc.* **127**, 3156 (2005)
15. G. Patonay, B. Eckenrode, J.J. Krutak, J. Salon and L. Strekowski, *Latent Invisible Trace Evidence: Chemical Detection Strategies*, in R.D. Blackledge (Ed.), *Forensic Analysis on the Cutting Edge*, Wiley – Interscience, 2007, pp. 115 – 140
16. S.M. Makin, L.I. Boiko and O.A. Shavrygina, *Zh. Org. Khim.* **13**, 1189 (1977).

Received on 24 February 2009.