

Graphical Abstract


Heterocycl. Commun.4(2009) 239-244

Alternative Products In One-Pot Reaction Of Benzylidene-Malononitrile, Thiocarbamoylacetamide And Alkyl Halides

A.Krauze, M. Vīlums, L.Sīle, G.Duburs

Latvian Institute of Organic Synthesis, Riga, Aizkraukles 21, LV-1006, Latvia

6-Oxo-5-(1,3-thiazol-2-yl)-1,4,5,6-tetrahydropyridine-3-carbonitriles **17**, corresponding 6-oxo-5-(1,3-thiazol-2-yl)-5,6-dihydropyridine-3-carbonitriles **18** and intermediates - 2-(1,3-thiazol-2-yl)acetamides **15** and 4,4-dicyano-2-(1,3-thiazol-2-yl)-butyramides **16** have been obtained as alternative products to 3-hydroxy-2,3-dihydro-7H-thiazolo[3,2-a]pyridines **5** carrying out one-pot condensation of 2-thiocarbamoylacetamide **2**, base **3**, halomethyl ketones **4** and benzylidenemalononitrile **1**.




Heterocycl. Commun.4(2009) 245-250

One-pot synthesis of 3-(2'-Benzoxazole)-2H-1-Benzopyran -2-One Derivatives with Benzoic Acid Catalysis

Liang Han, Shi-Hai Zhou, Jianhong Jia, Yujin Li, Shengwei Jian, Jianrong Gao

State Key Laboratory Breeding Base of Green Chemistry-Synthesis Technology, Zhejiang University of Technology, Hangzhou 310014 China



Title compounds **4** were synthesized in one pot with benzoic acid catalysis from salicylaldehydes, ethyl cyanoacetate and o-aminophenols.

Heterocycl. Commun.4(2009) 251-261

Syntheses Of New Heterocycles Derived From 2-Phenyl-3,1-Benzoxazin-4-One And Their Antibacterial And Antifungal Activity

Freddy H. Havaladar* and Abhay R. Patil

Nadkarny-Sacasa Research Laboratory, Department of Chemistry,
St. Xavier's College, Mumbai - 400 001, India.


A series of novel substituted N-(6-substituted-4H-benzo[e][1,3]oxazin-3-yl)- 2-[4-(4-oxo-2-phenyl-4H-quinazolin-3-yl)-phenoxy]-acetamides (**9a-b**) and N-(6-substituted-2-thione-4H-benzo[e][1,3]oxazin-3-yl)-2-[4-(4-oxo-2-phenyl-4H-quinazolin-3-yl)-phenoxy]-acetamides (**10a-b**) were designed and synthesized. The structures of the newly synthesized compounds have been confirmed by IR, ¹H NMR and mass spectra. The compounds have also been screened for their biological activity.

Synthesis and spectral investigations of covalently linked phthalocyanine-C₆₀ dyad via flexible carbon linker

Daisuke Sukeguchi, Hideyuki Yoshiyama, Surya Prakash Singh, Norio Shibata,* Shuichi Nakamura, Toru Takeshi,* Yasuhiko Hayashi, and Tetsuo Soga

Department of Frontier Materials, Graduate School of Engineering, Nagoya Institute of Technology, Gokiso, Showa-ku, Nagoya, 466 8555, Japan.

Phthalocyanine-C₆₀ dyad **1** via a flexible saturated carbon linker was synthesized. The UV/Vis and fluorescence spectra studies of **1** revealed that intramolecular, not intermolecular, photoinduced electronic communication was observed in the solution state. We also examined UV/Vis and fluorescence spectra of a mixture of C₆₀ and conventional tetra *tert*-Bu phthalocyanine **2** for comparison



A Novel Phospha Sugar Analogue: Synthesis And Evaluation Of 2,3-Dibromo-3-Methyl-1-Phenylphospholane 1-Oxide As A New Class Of Potential Anti-Proliferative Materials For Leukemia Cells

Michio Fujie^{a,b,*}, Satoki Nakamura^c, Kazuhide Asai^a, Taishi Niimi^a, Junko Yamashita^d, Keita Kiyofuji^d, Kiyoshi Shibata^b, Masako Suzuki^b, Reiji Aoshima^b, Tetsumei Urano^{b,e}, and Mitsuji Yamashita^{d,*}

^a Department of Materials Science, Graduate School of Science and Engineering, Shizuoka University, Hamamatsu 432-8561, Japan.


^b Research Equipment Center, Hamamatsu University School of Medicine, Hamamatsu 431-3192, Japan.

^c Department of Internal Medicine III, Hamamatsu University School of Medicine, Hamamatsu 431-3192, Japan.

^d Department of Nano Materials, Graduate School of Science and Technology, Shizuoka University, Hamamatsu 432-8011, Japan.

^e Department of Physiology II, Hamamatsu University School of Medicine, Hamamatsu 431-3192, Japan.

Abstract: A novel dideoxydibromophospha sugar analogue, 2,3-dibromo-3-methyl-1-phenylphospholane 1-oxide (**1**), was prepared from 1-phenyl-3-methyl-2-phospholene 1-oxide (**2**) and evaluated by MTT *in vitro* methods for leukemia cells. The evaluation of dibromide **1** (mixture of diastereomers) as well as the separated diastereomers of dibromide **1a** to **1d** revealed clearly that the synthesized phospha sugar analogue has competent potentials and excellent anti-tumor activities for the human leukemia cells of K562 and U937 in selective and specific manner. The cell cycle analyses by flow cytometry for K562 cells clearly demonstrated that the mechanism of the anti-proliferative effect on the human tumor cells is apoptosis induced by the phospha sugar.




Synthesis and Reactions of Some 1*H*-Pyrazole-3 Carboxylic Acid Chloride

İlhan Özer İlhan*, Sevgi Zühal, Zülbiye Önal, Emin Sarıpınar

Department of Chemistry, Erciyes University, 38039, Kayseri, Turkey

The pyrazole-carboxylic acid chloride 2 was obtained from the reaction of 4-Benzoyl-1-(2,4-dinitrophenyl)-5-phenyl-1*H*-pyrazole-3-carboxylic acid 1 and thionyl chloride. 1*H*-Pyrazole-3 carboxylic acid chlorides 2 can easily be converted into corresponding 1*H*-pyrazole-3-carboxylic acid amide derivatives 4 and 1*H*-pyrazole-3-carboxamide derivatives 6 from the reaction with various aliphatic and aromatic amines. The structures of these new synthesized compounds were determined from the IR, ¹H and ¹³C NMR spectroscopic data and elemental analysis.




Fungicidal Activities of 1,4-Bis[(3-aryl)-s-triazolo-[3,4-*b*]-[1,3,4]thiadiazole-6-yl]naphthalenes

De-Jiang Li

College of Chemistry and Life Science, China Three Gorges University, Yichang 443002, P. R. China

1,4-Bis[(3-aryl)-s-triazolo[3,4-*b*]-[1,3,4]thiadiazole-6-yl]naphthalenes 2 were synthesized in high yields by reaction of 3-aryl 4-amino-5-mercapto-1,2,4-triazole 1 with 1,4-naphthalic acid in the presence of POCl₃ and tetrabutylammonium iodide as catalyst. The structures have been established on the basis of elemental analysis and spectral data. The preliminary antibacterial tests showed that 2b, 2c and 2d exhibited good fungicidal activities against *Cerospora beticola* sacc.




A facile synthesis of 3-[4'-(2',6'-dimethyl-3',5'-dicarbethoxy-1',4'-dihydropyridyl)]chromenes

S. Satyanarayana Reddy, Kamalesh. K. Sharma and G. L. David Krupadanam*

Department of Chemistry, Osmania University, Hyderabad-500007, A.P., India.

e-mail: davidkrupa@hotmail.com, Ph: 91-40-27007360


Abstract: 2H-3-Chromenecarbaldehydes (**3a-h**) on reaction with ethyl 3-aminocrotonate (**4**) in the presence of acetic acid gave 3-[4'-(2',6'-dimethyl-3',5'-dicarbethoxy-1',4'-dihydropyridyl)]chromenes (**5a-h**) in good yields.

**Synthesis Of 6- Alkoxy carbonylmethylsulfanyl-5-Cyano-2-Methyl-4-Phenyl-1,4-Dihydropyridine-3-Carboxylates**

A.Krauze, L.Sile, L.Chernova, Z. Andzans, G.Duburs

Latvian Institute of Organic Synthesis, Riga, Aizkraukles 21, LV-1006, Latvia

6-Alkoxy carbonylmethylsulfanyl-5-cyano-2-methyl-4-phenyl-1,4-dihydropyridine-3-carboxylates **3** which are more lipophilic derivatives of the biologically active 6-methylsulfanyl-1,4-DHP-3-carboxylates **2** have been prepared by alkylation of 1,4-dihydropyridine-6-thiolates **1** with alkyl bromoacetates. Reactivity of **2** with KOH/H₂O was investigated.



a) R = Me; b) R = Et; c) R = i-Pr; d) R = t-Bu

Synthesis and antimicrobial activities of 5-(2', 5'-disubstituted 1H-indol-3'-yl)-5H-thiazolo[4,3-b]1,3,4-oxadiazole, -1,3,4-thiadiazole, -1,2,4-triazole and their derivatives

Saundane. A.R*, Yarlakatti Manjunatha and Prabhaker Walmik.

Department of Postgraduate Studies and Research in Chemistry

Gulbarga University, Gulbarga-585 106

Karnataka, India.

5-(2', 5'-disubstituted 1H-indol-3'-yl)-5H-thiazolo[4,3-b]1,2,4-triazoles (5) -1,3,4-thiadiazoles (6) and -1,3,4-oxadiazoles (7) and their analogues were synthesized from 4.

