

# Synthesis and Spectral studies of Nitrosourea derivatives of 3-Methyl- 5/7- Substituted -2- (3,4-dichloro) benzoyl-4H-1,4-Benzothiazines as Bifunctional Anticancer Agents.

Rajni Gupta\* and Vandana Gupta

*Department of Chemistry, University of Rajasthan, Jaipur-302004, India*

*Email: rajni187@yahoo.co.in*

**Abstract:** The synthesis of 3-methyl-5/7- substituted-4- (N-propyl-N-nitrosoamido)- 2- (3,4-dichloro benzoyl) -4H-1,4-Benzothiazines by the isocyanation and successive nitrosation of 3-methyl -5/7- substituted- 2- (3,4-dichlorobenzoyl) -4H-1,4-Benzothiazines has been reported. The synthesized compounds have been characterized by their elemental analyses and spectral characteristics.

## Introduction:

Analogous to phenothiazines, benzothiazines possess a wide spectrum of biological activities<sup>1</sup>. Their several derivatives are in clinical use<sup>2-7</sup>. They exhibit significant anticancer activities, which are assigned due to their interaction with DNA by complexation.

Nitrosourea derivatives constitute an important class of anticancer agents and its several derivatives like MNNG, CNU, MNU, GANU, and CDL-7 etc. are clinically significant. They interact with DNA via alkylation<sup>8-9</sup>. However their clinical use is limited because of cumulative and delayed side effects exerted by these compounds. Bone marrow toxicity being dose limiting, therefore it is worthwhile to develop a new series of nitrosoureas with minimum toxicity and side effects. 4H-1, 4- Benzothiazines are much less toxic and therefore it is anticipated that their nitrosourea derivatives will be potent anticancer agents with minimum toxicity, side effects etc.

In 3-methyl-5/7- substituted-4- (N-propyl-N-nitrosoamido)- 2- (3,4-dichloro benzoyl) -4H-1,4-Benzothiazines heterocyclic nitrogen with a side chain at 4-position constitutes N-nitrosourea linkage and possess both 1,4-benzothiazines nucleus and a nitrosourea moiety. They would show two fold interaction with DNA via complexation<sup>10</sup> as well as alkylation and will constitute bifunctional anticancer agents.



## Experimental:

Melting points of the synthesized compounds were determined on an electric melting point apparatus and are uncorrected. IR spectra were recorded in KBr on SHIMADZU 8400S FT IR spectrophotometer. The <sup>1</sup>H NMR and <sup>13</sup>C NMR spectra were recorded on a model Bruker-DRX-300 NMR spectrometer at 300 MHz and 75 MHz respectively using CDCl<sub>3</sub> as a solvent and TMS as an internal standard. The Mass spectrum of the representative compound was recorded on JEOL-SX-102/DA-6000 mass spectrometer.

### (i) Preparation of 3-methyl -5/7- substituted- 2- (3,4-dichlorobenzoyl) -4H-1,4-Benzothiazines(III a-g)

To the stirred suspension of 3,4-dichlorobenzoyl acetone II (10mmoles) in DMSO (5ml) was added 3-methyl/3-ethoxy/5-methyl/5-ethoxy/5-methoxy/5-chloro/3,5-dimethyl-2-amino benzenethiol I (10mmoles) and mixture was



refluxed for 30-40mins. The reaction mixture was concentrated and cooled down to room temperature. The solid separated out was filtered, washed with petroleum ether and crystallized from methanol (Scheme- 1).



Scheme-1

**(ii) Preparation of 3-methyl-5/7- substituted- 2- (3,4-dichlorobenzoyl) -4(N-propyl amido) - 1,4-benzothiazines (IVa-g)**

A mixture of 3-methyl-5/7- substituted- 2- (3,4-dichlorobenzoyl) -4H-1,4-Benzothiazines III (10mmoles), 10 ml of absolute alcohol and propyl isocyanate (10mmoles) was refluxed on hot plate for 2 hrs. Then the solvent was removed under vacuum rotatory evaporator. The product 3-methyl-5/7-substituted-2-(3,4-dichlorobenzoyl)-4-(N-propyl amido)-1,4-benzothiazines was crystallised from ethanol (Scheme 2).





Scheme -2

**(iii) Preparation of 3-methyl-5/7- substituted-4- (N-propyl-N-nitrosoamido)- 2- (3,4-dichloro benzoyl) -4H-1,4-Benzothiazines (Va-g)**

3-Methyl-5/7-substituted-2-(3,4-dichlorobenzoyl)-4-(N-propylamido)-1,4-benzothiazines IV (3mmoles) was dissolved in 50 ml of acetic acid, sodium nitrite (5mmoles) was added portion wise with stirring. The mixture was stirred for 30mins at room temperature and for one hour at 50° C. Acetic acid was evaporated under reduced pressure in vacuum rotatory evaporator. The residue was treated with water. The resulting precipitate of 3-methyl-5/7-substituted-4- (N-propyl-N-nitrosoamido)- 2- (3,4-dichloro benzoyl) -4H-1,4-Benzothiazines was collected and crystallized from methanol.

(Scheme 3)



Scheme-3

**Table 1: Physical data of 3-methyl-5/7- substituted-4- (N-propyl-N-nitrosoamido)- 2- (3,4-dichloro benzoyl) -4H-1,4-Benzothiazines**

| Compound | Molecular formula                                                               | M.P<br>°C | Yield<br>% | %<br>C<br>H<br>N   |                     |                    |
|----------|---------------------------------------------------------------------------------|-----------|------------|--------------------|---------------------|--------------------|
|          |                                                                                 |           |            | (Found)<br>(Calc.) | ( Found)<br>(Calc.) | (Found)<br>(Calc.) |
| A        | C <sub>21</sub> H <sub>19</sub> Cl <sub>2</sub> N <sub>3</sub> O <sub>3</sub> S | 178       | 60         | (54.32)<br>(54.10) | (4.12)<br>(4.02)    | (9.05)<br>(9.00)   |
| B        | C <sub>22</sub> H <sub>21</sub> Cl <sub>2</sub> N <sub>3</sub> O <sub>4</sub> S | viscous   | 62         | (53.45)<br>(53.24) | (4.28)<br>(4.03)    | (8.50)<br>(8.46)   |
| C        | C <sub>21</sub> H <sub>19</sub> Cl <sub>2</sub> N <sub>3</sub> O <sub>3</sub> S | 152       | 61         | (54.32)<br>(61.42) | (4.12)<br>(5.45)    | (9.05)<br>(9.65)   |
| D        | C <sub>22</sub> H <sub>21</sub> Cl <sub>2</sub> N <sub>3</sub> O <sub>4</sub> S | viscous   | 50         | (53.45)<br>(53.29) | (4.28)<br>(4.08)    | (8.50)<br>(8.42)   |
| E        | C <sub>21</sub> H <sub>19</sub> Cl <sub>2</sub> N <sub>3</sub> O <sub>4</sub> S | viscous   | 63         | (52.51)<br>(52.45) | (3.99)<br>(3.89)    | (8.75)<br>(8.65)   |
| F        | C <sub>20</sub> H <sub>16</sub> Cl <sub>3</sub> N <sub>3</sub> O <sub>3</sub> S | 166       | 61         | (49.55)<br>(49.45) | (3.33)<br>(3.22)    | (8.67)<br>(8.42)   |
| G        | C <sub>22</sub> H <sub>21</sub> Cl <sub>2</sub> N <sub>3</sub> O <sub>3</sub> S | 170       | 63         | (55.23)<br>(55.13) | (4.22)<br>(4.12)    | (8.78)<br>(8.65)   |

**Table 2: Infra red spectral data of 3-methyl-5/7- substituted-4- (N-propyl-N-nitrosoamido)- 2- (3,4-dichloro benzoyl) -4H-1,4-Benzothiazines**

| Compound | Molecular formula                                                               | C=O<br>(cm <sup>-1</sup> ) | C-Cl<br>(cm <sup>-1</sup> ) |
|----------|---------------------------------------------------------------------------------|----------------------------|-----------------------------|
| A        | C <sub>21</sub> H <sub>19</sub> Cl <sub>2</sub> N <sub>3</sub> O <sub>3</sub> S | 1585, 1645                 | -                           |
| B        | C <sub>22</sub> H <sub>21</sub> Cl <sub>2</sub> N <sub>3</sub> O <sub>4</sub> S | 1605, 1665                 | -                           |
| C        | C <sub>21</sub> H <sub>19</sub> Cl <sub>2</sub> N <sub>3</sub> O <sub>3</sub> S | 1615, 1750                 | -                           |
| D        | C <sub>22</sub> H <sub>21</sub> Cl <sub>2</sub> N <sub>3</sub> O <sub>4</sub> S | 1610, 1695                 | -                           |
| E        | C <sub>21</sub> H <sub>19</sub> Cl <sub>2</sub> N <sub>3</sub> O <sub>4</sub> S | 1605, 1685                 | -                           |
| F        | C <sub>20</sub> H <sub>16</sub> Cl <sub>3</sub> N <sub>3</sub> O <sub>3</sub> S | 1615, 1650                 | 790                         |
| G        | C <sub>22</sub> H <sub>21</sub> Cl <sub>2</sub> N <sub>3</sub> O <sub>3</sub> S | 1610, 1640                 | -                           |

**Table 3: NMR Spectral data of 3-methyl-5/7- substituted-4- (N-propyl-N-nitrosoamido)- 2- (3,4-dichloro benzoyl) -4H-1,4-Benzothiazines**

| S.No     | Molecular formula         | Solvent  | $\delta$ (ppm) | Hydrogen | Multiplicity | Assignment                               |
|----------|---------------------------|----------|----------------|----------|--------------|------------------------------------------|
| <b>A</b> | $C_{21}H_{19}Cl_2N_3O_3S$ | $CDCl_3$ | 6.72-7.79      | 6        | Multiplet    | Aromatic protons                         |
|          |                           |          | 1.66           | 3        | Singlet      | $CH_3$ protons at $C_3$                  |
|          |                           |          | 2.25           | 3        | Singlet      | $CH_3$ protons at $C_5$                  |
|          |                           |          | 0.90–0.98      | 3        | Triplet      | $CH_3$ protons at $C'_1$ of propyl group |
|          |                           |          | 1.35-1.42      | 2        | Multiplet    | $CH_2$ protons at $C'_2$ of propyl group |
|          |                           |          | 2.7–3.05       | 2        | Triplet      | $CH_2$ protons at $C'_3$ of propyl group |
| <b>B</b> | $C_{22}H_{21}Cl_2N_3O_4S$ | $CDCl_3$ | 6.63-7.68      | 6        | Multiplet    | Aromatic protons                         |
|          |                           |          | 3.4 - 4.1      | 2        | Quartet      | $CH_2$ protons of $C_2H_5$               |
|          |                           |          | 1.30-1.4       | 3        | Triplet      | $CH_3$ protons of $C_2H_5$               |
|          |                           |          | 1.73           | 3        | Singlet      | $CH_3$ protons at $C'_3$                 |
|          |                           |          | 3.4-3.61       | 2        | Triplet      | $CH_2$ protons at $C'_1$ of propyl group |
|          |                           |          | 1.48-1.55      | 2        | Multiplet    | $H_2$ protons at $C'_2$ of propyl group  |
|          |                           |          | 0.96-1.10      | 3        | Triplet      | $CH_3$ protons at $C'_3$ of propyl group |
| <b>C</b> | $C_{21}H_{19}Cl_2N_3O_3S$ | $CDCl_3$ | 6.82-7.36      | 6        | Multiplet    | Aromatic protons                         |
|          |                           |          | 1.60           | 3        | Singlet      | $CH_3$ protons at $C_3$                  |
|          |                           |          | 2.25           | 3        | Singlet      | $CH_3$ protons at $C_7$                  |
|          |                           |          | 0.91-1.0       | 3        | Triplet      | $CH_3$ protons at $C'_1$ of propyl group |
|          |                           |          | 1.35-1.48      | 2        | Multiplet    | $CH_2$ protons at $C'_2$ of propyl group |
|          |                           |          | 2.5-2.8        | 2        | Triplet      | $CH_3$ protons at $C'_3$ of propyl group |
| <b>D</b> | $C_{22}H_{21}Cl_2N_3O_4S$ | $CDCl_3$ | 6.53-7.77      | 6        | Multiplet    | Aromatic protons                         |
|          |                           |          | 3.86-4.10      | 2        | Quartet      | $CH_2$ protons of $C_2H_5$               |
|          |                           |          | 1.20-1.25      | 3        | Triplet      | $CH_3$ protons of $C_2H_5$               |

|          |                                                                              |                         |            |   |           |                                                               |
|----------|------------------------------------------------------------------------------|-------------------------|------------|---|-----------|---------------------------------------------------------------|
| <b>E</b> | <b>C<sub>21</sub>H<sub>19</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>4</sub>S</b> | <b>CDCl<sub>3</sub></b> | 1.67       | 3 | Singlet   | CH <sub>3</sub> protons at C <sub>3</sub>                     |
|          |                                                                              |                         | 2.4 - 3.1  | 2 | Triplet   | CH <sub>2</sub> protons at C' <sub>1</sub><br>of propyl group |
|          |                                                                              |                         | 1.34-1.38  | 2 | Multiplet | CH <sub>2</sub> protons at C' <sub>2</sub><br>of propyl group |
|          |                                                                              |                         | 0.86-0.9   | 3 | Triplet   | CH <sub>3</sub> protons at C' <sub>3</sub><br>of propyl group |
|          |                                                                              |                         | 6.54 -7.77 | 6 | Multiplet | Aromatic protons                                              |
|          |                                                                              |                         | 3.73       | 3 | Singlet   | CH <sub>3</sub> protons of OCH <sub>3</sub>                   |
|          |                                                                              |                         | 1.71       | 3 | Singlet   | CH <sub>3</sub> protons at C <sub>3</sub>                     |
|          |                                                                              |                         | 3.1-3.3    | 2 | Triplet   | CH <sub>2</sub> protons at C' <sub>1</sub><br>of propyl group |
|          |                                                                              |                         | 1.54-1.65  | 2 | Multiplet | CH <sub>2</sub> protons at C' <sub>2</sub><br>of propyl group |
|          |                                                                              |                         | 0.90-0.96  | 3 | Triplet   | CH <sub>3</sub> protons at C' <sub>3</sub><br>of propyl group |
| <b>F</b> | <b>C<sub>20</sub>H<sub>16</sub>Cl<sub>3</sub>N<sub>3</sub>O<sub>3</sub>S</b> | <b>CDCl<sub>3</sub></b> | 7.03-7.87  | 6 | Multiplet | Aromatic protons                                              |
|          |                                                                              |                         | 1.66       | 3 | Singlet   | CH <sub>3</sub> protons at C <sub>3</sub>                     |
|          |                                                                              |                         | 3.5-4.0    | 2 | Triplet   | CH <sub>2</sub> protons at C' <sub>1</sub><br>of propyl group |
|          |                                                                              |                         | 1.60-1.70  | 2 | Multiplet | CH <sub>2</sub> protons at C' <sub>2</sub><br>of propyl group |
|          |                                                                              |                         | 0.95-0.99  | 3 | Triplet   | CH <sub>3</sub> protons at C' <sub>3</sub><br>of propyl group |
|          |                                                                              |                         |            |   |           |                                                               |
| <b>G</b> | <b>C<sub>22</sub>H<sub>21</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>3</sub>S</b> | <b>CDCl<sub>3</sub></b> | 6.63-7.62  | 5 | Multiplet | Aromatic protons                                              |
|          |                                                                              |                         | 1.65       | 3 | Singlet   | CH <sub>3</sub> protons of C <sub>3</sub>                     |
|          |                                                                              |                         | 2.35       | 6 | Singlet   | CH <sub>3</sub> protons at C <sub>5</sub> &C <sub>7</sub>     |
|          |                                                                              |                         | 2.9-3.25   | 2 | Triplet   | CH <sub>3</sub> protons at C' <sub>1</sub><br>of propyl group |
|          |                                                                              |                         | 1.53-1.58  | 2 | Multiplet | CH <sub>2</sub> protons at C' <sub>2</sub><br>of propyl group |
|          |                                                                              |                         | 0.85-0.9   | 3 | Triplet   | CH <sub>2</sub> protons at C' <sub>3</sub><br>of propyl group |
|          |                                                                              |                         |            |   |           |                                                               |

## Results and Discussion:

The synthesis of 3-methyl-5/7- substituted-4- (N-propyl-N-nitrosoamido)- 2- (3,4-dichloro benzoyl) -4H-1,4-benzothiazines is based on the synthesis of substituted 3-methyl -5/7- substituted- 2- (3,4-dichlorobenzoyl) -4H-1,4-Benzothiazines reported elsewhere<sup>1</sup>. 4H-1,4-Benzothiazines are analogs of phenothiazines and like phenothiazines they bear a fold along nitrogen and sulphur axis which is considered responsible to impart them biological activities. So it was considered worthwhile to incorporate the activities of benzothiazines and nitrosoureas into one molecule i.e nitrosourea derivatives of 3-methyl -5/7- substituted- 2- (3,4-dichlorobenzoyl) -4H-1,4-Benzothiazines. 4H-1,4-Benzothiazines are key compounds to synthesize the above mentioned compounds. Here the 4H-1,4-benzothiazines were allowed to undergo isocyanation at 4-position, thereby giving 3-methyl -5/7- substituted- 2- (3,4-dichlorobenzoyl) -4-(N-propyl amido) - 1,4-benzothiazines. These were then let to undergo nitrosation with sodium nitrite in acetic acid.

## References:

1. R R Gupta(Ed.) " Phenothiazines and 4H-1,4-Benzothiazines – Chemical and Biomedical Aspects" Elsevier, Amsterdam, 1988.
2. H. Keyzer, G. M. Eckert, I. S. forrest, R. R. Gupta, F. Gutmann, J. Molnar (Eds.) "Thiazines and Structurally related compounds"(Proceedings of Sixth International Conference on Phenothiazines and Structurallyrelated Psychotropic compounds, Pasadena, California, Sep.11-14(1990). Krieger Publishing Company,Malabar, Florida, USA(1992).
3. H.Y.Kogyo, Japanese patent, 59, 187,170(1984);Chem. Abstr., 102, 45965(1985).
4. Kalpana Gupta, VandanaGupta, Rajni gupta, and M. Kumar, Hetercycl.Comm., 8 (5) 485-458(2002).
5. R R Gupta, M L-Sharma, C M Rajoria, Archana Gupta, and M Nyati, Anticancer Drugs, 4, 589(1993).
6. Gulshan Kumar, Vandana Gupta, D C Gautam and R R Gupta, Heterocycl.Comm.,8(5) 381-384(2002).
7. M. Myehlstaedt, R Widera, H. Meinhold and K. Hollmann, German(East)Patent 214,128(1984); Chem Abstr.,102 203979(1985).
8. J. A. Montgomery and T.P.Johnston," Nitrosoureas" in D.E.V Wilman(Ed.) Chemistry of Anticancer agents,"Blakie ,Glassgow and London, 199, pp.131-158 and cited reference therein.
9. Mukesh Kumar Nyati, Dinesh Rai, Radha Raman Gupta and Prashant Kumar Dev. In Vivo II: 95-100(1997).
10. C. Bodea and I .Silberg" Recent advances in the Chemistry of Phenothiazines" in A. R Katrikzky and A J Boulton (Eds.)" Advances in Heterocyclic Chemistry" Vol 9 ,Academic Press , New York and London(1968), pp392.

Received on February 22,2010.