

Sulfamic acid as an efficient and cost-effective catalyst for the synthesis of 2,4,5-triarylimidazoles and 2-arylphenanthrimidazoles

Hua-Nan Peng^{1,*}, Xiao-Ming Peng², Da-Gui Zheng¹, Fei Yu¹ and Min Rao¹

¹ Key Laboratory of Applied Organic Chemistry, Higher Institutions of Jiangxi Province, Shangrao Normal University, Shangrao 334001, China

² Department of Chemistry, Nanchang University, Xuefu Road 999, Nanchang 330031, China

*Corresponding author
e-mail: huananpeng@126.com

Abstract

A simple and efficient synthesis of 2,4,5-triarylimidazoles and 2-arylphenanthrimidazoles was developed using the one-pot, three-component condensation reaction of an aromatic aldehyde, benzil or 9,10-phenanthroquinone and ammonium acetate in refluxing ethanol in the presence of the catalyst sulfamic acid.

Keywords: one-pot condensation; phenanthrimidazoles; sulfamic acid; 2,4,5-triarylimidazoles.

Introduction

Substituted imidazoles have gained remarkable importance as herbicidal (Viktor et al., 1975), fungicidal (Heeres et al., 1979), anti-inflammatory (Lombardino and Wiseman, 1974) and antibacterial agents (Sharma et al., 2009) and as inhibitors of P38 MAP kinase (Laufer et al., 2010). In addition, imidazole derivatives have been widely used as precursors to *N*-heterocyclic carbenes in organometallic catalysis and as ionic liquids in green chemistry (Kim et al., 2005; Jiménez et al., 2008). Several substituted imidazoles also show cardiovascular and antitrypanosomal activities (Malhotra et al., 2011; Samant and Sukthankar, 2011). In light of these properties, the synthesis of substituted imidazole derivatives has received considerable attention in recent years.

2,4,5-Triphenylimidazole was first synthesized more than a century ago by the three-component condensation of a 1,2-dicarbonyl compound, aldehyde and ammonia (Japp and Robinson, 1882; Radziszewski, 1882). Recently, numerous synthetic methods for the preparation of analogous compounds have been developed using similar three-component condensation in the presence of different catalysts. These catalysts include molecular iodine (Kidwai et al., 2007; Behmadi et al., 2009), trifluoroacetic acid (Mohammadizadeha et al., 2009),

scolecite (Gadekara et al., 2009), L-proline (Samai et al., 2009), ytterbium perfluorooctanesulfonate (Shen et al., 2008), tetrabutylammonium bromide (Chary et al., 2008), silica sulfuric acid (Shaabani et al., 2007), indium trichloride (Sharma et al., 2008), potassium dodecatungstocobaltate (Nagarapu et al., 2007) and silica perchloric acid (Kantevari et al., 2007). The reaction can also be performed in acetic acid under reflux conditions (Puratchikody and Doble, 2007), under microwave irradiation (Chauveau et al., 2010; Damavandi, 2011) and in ionic liquid (Zang et al., 2010).


In recent years, sulfamic acid ($\text{NH}_2\text{SO}_3\text{H}$) has successfully been used as a solid acid catalyst for various organic synthetic transformations (An et al., 2007; Heravi et al., 2007; Heydari et al., 2007; Kamal et al., 2007; Li et al., 2007; Upadhyaya et al., 2007; Mitragotri et al., 2008). The benefits of its use include low cost, eco-friendliness, ease of handling, low toxicity, excellent product yields and high reactivity. These features prompted us to explore the three-component condensation reactions of aldehydes, benzil or 9,10-phenanthroquinone and ammonium acetate in the presence of a catalytic amount of $\text{NH}_2\text{SO}_3\text{H}$ to synthesize the corresponding 2,4,5-triarylimidazoles and 2-arylphenanthrimidazoles.

Results and discussion

Different acidic catalysts in the reaction of benzaldehyde (2 mmol), benzil (2 mmol) and ammonium acetate (6 mmol) were compared (Table 1). The catalysts included F_3CCOOH , H_3BO_3 , $\text{CH}_3\text{SO}_3\text{H}$, $p\text{-CH}_3\text{C}_6\text{H}_5\text{SO}_3\text{H}$, $\text{NH}_2\text{SO}_3\text{H}$ and $p\text{-NH}_2\text{C}_6\text{H}_5\text{SO}_3\text{H}$, each at an amount equal to 10 mmol%. As can be seen from entries 1–6, $\text{NH}_2\text{SO}_3\text{H}$ is the most effective catalyst in terms of product **4a** yield (85%). Next, the amount of $\text{NH}_2\text{SO}_3\text{H}$ was varied and it was found that 10 mmol% of the catalyst was the optimal quantity for catalysis. The yield was further improved by increasing the amount of NH_4OAc from 6 mmol to 10 mmol (entry 11).

Next, a wide range of substituted benzaldehydes were tested under the optimized conditions. The results are given in Table 2. Interestingly, the electronic properties of the substituents have virtually no effect on yields. More specifically, the reaction of a benzaldehyde with an electron-donating or an electron-withdrawing group furnishes product **4** in high yield.

In a similar way, phenanthraquinone **5** was allowed to react with five substituted benzaldehydes. As shown in Table 3, these reactions produce the desired 2-arylphenanthrimidazoles **6** in excellent yields, and the substituent on the aldehyde,

Table 1 Optimizing the reaction conditions^a.


Entry	Catalyst	Catalyst (mmol%)	NH ₄ OAc (mmol)	Time (h)	Yield (%) ^b
1	F ₃ CCOOH	10	6	5	65
2	H ₃ BO ₃	10	6	5	69
3	CH ₃ SO ₃ H	10	6	5	73
4	<i>p</i> -CH ₃ C ₆ H ₄ SO ₃ H	10	6	5	77
5	<i>p</i> -NH ₂ C ₆ H ₄ SO ₃ H	10	6	5	80
6	NH ₂ SO ₃ H	10	6	5	85
7	NH ₂ SO ₃ H	2.5	6	5	75
8	NH ₂ SO ₃ H	5	6	5	80
9	NH ₂ SO ₃ H	15	6	5	85
10	NH ₂ SO ₃ H	10	8	5	89
11	NH ₂ SO ₃ H	10	10	4	93

^aThe amount of benzaldehyde was 2 mmol and benzil was 2 mmol;
^bisolated yields.

whether electron-donating or -withdrawing, has no effect on the yield.

Conclusions

We have developed a simple and efficient three-component condensation reaction for the synthesis of 2,4,5-triaryl-imidazoles and 2-arylphenanthrimidazoles using sulfamic acid as an efficient, cost-effective and eco-friendly catalyst.

Experimental section

General


Melting points were determined on X4 melting point apparatus and are uncorrected. Infrared (IR) spectra were recorded using KBr discs on a Nicolet 6700 fourier transform infrared (FT-IR) spectrophotometer. ¹H-NMR spectra were recorded on the Varian Inova-300 MHz spectrometer using tetramethylsilane (TMS) as internal standard and DMSO-*d*₆ as a solvent. Mass spectra (EI, 70 eV) were obtained on a Shimadzu SIL-10A Auto Injector liquid chromatography-mass spectrometry (LC-MS). Thin layer chromatography (TLC) was carried out on GF254 silica gel plates.

General procedure for the preparation of 2,4,5-triaryl-imidazoles **4** and 2-arylphenanthrimidazoles **6**

A mixture of benzil or phenanthraquinone (2 mmol), a benzaldehyde (2 mmol), ammonium acetate (10 mmol) and sulfamic acid (10 mmol%) in ethanol (10 ml) was stirred and heated under reflux until the reaction was completed as monitored by TLC. Then, the mixture was poured onto crushed ice and the resulting solid was filtered and crystallized from ethanol. Known compounds **4a–k** (Table 2), **6a** and **6g** (Table 3) showed ¹H nuclear magnetic resonance (NMR) spectra that were virtually identical with those reported.


2-(4-Trifluoromethylphenyl)-4,5-diphenyl-1H-imidazole (4l) White crystals; ¹H NMR: δ 8.27 (d, *J*=8 Hz, 2H), 7.83 (d, *J*=8 Hz, 2H), 7.59–7.18 (m, 10H); IR: 3402, 3057, 1602, 1588, 1496, 1441 cm⁻¹; MS: *m/z* 365 ([*M*+1]⁺, 100).

2-(2,4-Dichlorophenyl)-1H-phenanthro[9,10-*d*]imidazole (6b) White crystals; ¹H NMR: δ 8.84 (d, *J*=8 Hz, 2H), 8.59 (d, *J*=8 Hz, 2H), 7.94 (d, *J*=8 Hz, 1H), 7.87 (s, 1H), 7.73–7.60 (m, 5H); IR: 3440, 3083, 1614, 1588, 1515 cm⁻¹; MS: *m/z* 363 (*M*⁺, 100).

Table 2 Sulfamic acid catalyzed synthesis of 2,4,5-triaryl-imidazoles **4a–l**.


Entry	2	Aryl	Time (h)	Product	Yield (%) ^a	Mp (°C) ^b (found)	Mp (°C) (reported) ^{Literature}
1	2a	C ₆ H ₅	4	4a	93	277–279	274–276 ^c
2	2b	2,4-Cl ₂ C ₆ H ₃	5	4b	94	177–179	175–178 ^d
3	2c	4-MeC ₆ H ₄	4	4c	95	235–238	233–235 ^d
4	2d	4-BrC ₆ H ₄	4	4d	91	267–268	265 ^e
5	2e	2-ClC ₆ H ₄	3	4e	92	199–201	195–196 ^c
6	2f	3-NO ₂ C ₆ H ₄	3	4f	95	>300	>300 ^d
7	2g	4-OMeC ₆ H ₄	4	4g	93	231–232	227–228 ^c
8	2h	4-ClC ₆ H ₄	3	4h	95	263–265	262–264 ^d
9	2i	4-N(Me) ₂ C ₆ H ₄	4	4i	92	256–257	256–259 ^d
10	2j	2-Furyl	4	4j	89	241–243	242 ^e
11	2k	4-OHC ₆ H ₄	4	4k	90	265–267	268–269 ^c
12	2l	4-F ₃ CC ₆ H ₄	5	4l	90	273–275	

^aIsolated yields; ^bafter crystallization; ^c(Gadekara et al., 2009); ^d(Samai et al., 2009); ^e(Chary et al., 2008).

Table 3 Sulfamic acid catalyzed synthesis of 2-arylphenanthrimidazoles **6**.


Entry	2	Aryl	Time	Product	Yield (%) ^a	Mp (°C) ^b (found)	Mp (°C) (reported) ^c Literature
1	2a	C ₆ H ₅	5	6a	93	>300	314–315°
2	2b	2,4-Cl ₂ C ₆ H ₃	4	6b	90	245–246	
3	2d	4-BrC ₆ H ₄	4	6d	91	264–267	
4	2g	4-OMeC ₆ H ₄	4	6g	92	250–253	254–255°
5	2l	4-F ₃ CC ₆ H ₄	5	6l	93	241–243	

^aIsolated yields; ^bafter recrystallization; ^c(Behmadi et al., 2009).

2-(4-Bromophenyl)-1H-phenanthro[9,10-d]imidazole (6d) White crystals; ¹H NMR: δ 8.83 (d, *J*=8 Hz, 2H), 8.53 (d, *J*=8 Hz, 2H), 8.24 (d, *J*=8 Hz, 2H), 7.79 (d, *J*=8 Hz, 2H, Ar-H), 7.72 (t, *J*=8 Hz, 2H), 7.62 (t, *J*=8 Hz, 2H); IR: 3425, 3069, 1653, 1616, 1539, 1475, 1457 cm⁻¹; MS: *m/z* 373 (M⁺, 90), 375 ([M+2]⁺, 100).

2-(4-Trifluoromethylphenyl)-1H-phenanthro[9,10-d]imidazole (6l) White crystals, ¹H NMR: δ 8.93 (t, *J*=8 Hz, 2H), 8.65 (t, *J*=8 Hz, 2H), 8.59 (d, *J*=8 Hz, 2H), 8.04 (d, *J*=8 Hz, 2H), 7.88–7.68 (m, 4H); IR: 3458, 3077, 1619, 1588, 1529, 1454 cm⁻¹. MS: *m/z* 363 ([M+1]⁺, 100).

Acknowledgments

We acknowledge financial support from the Research Program of Science and Technology from the Bureau of Education of Jiangxi Province, China (No.: GJJ10736) and the Youth Science Foundation of Jiangxi Educational Committee, China (No.: GJJ11232).

References

- An, L. T.; Zou, J. P.; Zhang, L. L.; Zhang, Y. Sulfamic acid-catalyzed Michael addition of indoles and pyrrole to electron-deficient nitroolefins under solvent-free condition. *Tetrahedron Lett.* **2007**, *48*, 4297–4300.
- Behmadi, H.; Roshani, M.; Saadati, S. M. Synthesis of phenanthrimidazole from 9,10-phenanthraquinone and aldehydes by molecular iodine as catalyst. *Chinese Chem. Lett.* **2009**, *20*, 5–8.
- Chary, M. V.; Keerthysri, N. C.; Vupallapati, S. V.; Lingaiah, N.; Kantevari, S. Tetrabutylammonium bromide (TBAB) in isopropanol: an efficient, novel, neutral and recyclable catalytic system for the synthesis of 2,4,5-trisubstituted imidazoles. *Catal. Commun.* **2008**, *9*, 2013–2017.
- Chauveau, E.; Marestin, C.; Schiets, F.; Mercier, R. Synthesis of 2,4,5-triarylimidazoles in aqueous solution, under microwave irradiation. *Green Chem.* **2010**, *12*, 1018–1022.
- Damavandi, S. New approach to the multicomponent one-pot synthesis of 2-aryl-1H-phenanthro[9,10-d]imidazoles. *Heterocycl. Commun.* **2011**, *17*, 79–81.
- Gadekara, L. S.; Maneb, S. R.; Katkara, S. S.; Arbada, B. R.; Landea, M. K. Sulfamic acid as an efficient heterogeneous catalyst for the synthesis of 2,4,5-triarylimidazoles. *Cent. Eur. J. Chem.* **2009**, *7*, 550–554.
- Heeres, J.; Backx, L. J.; Mostmans, J. H.; Van Cutsem, J. Antimycotic imidazoles. Part 4. synthesis and antifungal activity of ketoconazole, a new potent orally active broad-spectrum antifungal agent. *J. Med. Chem.* **1979**, *22*, 1003–1005.
- Heravi, M. M.; Ranjbar, L.; Derikvand, F.; Bamoharram, F. F. Sulfamic acid as a cost-effective catalyst instead of metal-containing acids for the one-pot synthesis of β-acetamido ketones. *J. Mol. Catal. A: Chem.* **2007**, *276*, 226–229.
- Heydari, A.; Khaksar, S.; Pourayoubi, M.; Mahjoub, A. R. Sulfamic acid: an efficient, cost-effective and recyclable solid acid catalyst for the three-component synthesis of α-amino nitriles. *Tetrahedron Lett.* **2007**, *48*, 4059–4060.
- Japp, F. R.; Robinson, H. H.; Constitution des lophins und des amwins. *Chem. Ber.* **1882**, *15*, 1268–1270.
- Jiménez, M. V.; Pérez-Torrente, J. J.; Bartolomé, M. I.; Gierz, V.; Lahoz, F. J.; Oro, L. A. Rhodium(I) complexes with hemilabile N-heterocyclic carbenes: efficient alkyne hydrosilylation catalysts. *Organometallics* **2008**, *27*, 224–234.
- Kamal, A.; Prasad, B. R.; Reddy, A. M.; Khan, M. N. Sulfamic acid as an efficient and recyclable catalyst for the ring opening of epoxides with amines and anilines: an easy synthesis of β-amino alcohols under solvent-free conditions. *Catal. Commun.* **2007**, *8*, 1876–1880.
- Kantevari, S.; Vuppapapati, S. V.; Biradar, D. O.; Nagarapu, L. Highly efficient, one-pot, solvent-free synthesis of tetrasubstituted imidazoles using HClO₄-SiO₂ as novel heterogeneous catalyst. *J. Mol. Catal. A: Chem.* **2007**, *266*, 109–113.
- Kidwai, M.; Mothsra, P.; Bansal, V.; Somvanshi, R. K.; Ethayathulla, A. S.; Dey, S.; Singh, T. P. One-pot synthesis of highly substituted imidazoles using molecular iodine: a versatile catalyst. *J. Mol. Catal. A: Chem.* **2007**, *265*, 177–182.
- Kim, E. J.; Koa, S. Y.; Dziadulewicz, E. K. Mitsunobu alkylation of imidazole: a convenient route to chiral ionic liquids. *Tetrahedron Lett.* **2005**, *46*, 631–633.
- Laufer, S.; Hauser, D.; Stegmüller, T.; Bracht, C.; Ruff, K.; Schattel, V.; Albrecht, W.; Koch, P. Tri- and tetrasubstituted imidazoles as p38α mitogen-activated protein kinase inhibitors. *Bioorg. Med. Chem. Lett.* **2010**, *20*, 6671–6675.

- Li, Z. J.; Sun, Y. J.; Ren, X. H.; Wei, P.; Shi, Y. H.; Ouyang, P. K. Sulfamic acid catalyzed direct condensation of aldehydes, amines, and TMSCN to α -aminonitriles at ambient temperature. *Synlett*. **2007**, 5, 803–805.
- Lombardino, J. G.; Wiseman, E. H. Preparation and antiinflammatory activity of some nonacidic trisubstituted imidazoles. *J. Med. Chem.* **1974**, 17, 1182–1188.
- Malhotra, V.; Pathak, S. R.; Nath, R.; Mukherjee, D.; Shanker, K. Substituted imidazole derivatives as novel cardiovascular agents. *Bioorg. Med. Chem. Lett.* **2011**, 21, 936–939.
- Mitragotri, S. D.; Pore, D. M.; Desai, U. V.; Wadgaonkar, P. P. Sulfamic acid: an efficient and cost-effective solid acid catalyst for the synthesis of α -aminophosphonates at ambient temperature. *Catal. Commun.* **2008**, 9, 1822–1826.
- Mohammadzadeha, M. R.; Hasaninejada, A.; Bahramzadeha, M. Trifluoroacetic acid as an efficient catalyst for one-pot, four-component synthesis of 1,2,4,5-tetrasubstituted imidazoles under microwave-assisted, solvent-free conditions. *Synth. Commun.* **2009**, 39, 3232–3242.
- Nagarapu, L.; Apuri, S.; Kantevari, S. Potassium dodecatungstocobaltate trihydrate ($K_5CoW_{12}O_{40} \cdot 3H_2O$): a mild and efficient reusable catalyst for the one-pot synthesis of 1,2,4,5-tetrasubstituted imidazoles under conventional heating and microwave irradiation. *J. Mol. Catal. A. Chem.* **2007**, 266, 104–108.
- Puratchikody, A.; Doble, M. Antinociceptive and antiinflammatory activities and QSAR studies on 2-substituted-4,5-diphenyl-1H-imidazoles. *Bioorg. Med. Chem.* **2007**, 15, 1083–1090.
- Radziszewski, B. Ueber die Constitution des Lophins und verwandter Verbindungen. *Chem. Ber.* **1882**, 15, 1493–1496.
- Samai, S.; Nandi, G. C.; Singh, P.; Singh, M. S. L-Proline: an efficient catalyst for the one-pot synthesis of 2,4,5-trisubstituted and 1,2,4,5-tetrasubstituted imidazoles. *Tetrahedron* **2009**, 65, 10155–10161.
- Samant, B. S.; Sukthankar, M. G. Compounds containing 2-substituted imidazole ring for treatment against human African trypanosomiasis. *Bioorg. Med. Chem. Lett.* **2011**, 21, 1015–1018.
- Shaabani, A.; Rahmati, A.; Farhangi, E.; Badri, Z. Silica sulfuric acid promoted the one-pot synthesis of trisubstituted imidazoles under conventional heating conditions or using microwave irradiation. *Catal. Commun.* **2007**, 8, 1149–1152.
- Sharma, S. D.; Hazarika, P.; Konwar, D. An efficient and one-pot synthesis of 2,4,5-trisubstituted and 1,2,4,5-tetrasubstituted imidazoles catalyzed by $InCl_3 \cdot 3H_2O$. *Tetrahedron Lett.* **2008**, 49, 2216–2220.
- Sharma, D.; Narasimhan, B.; Kumar, P.; Judge, V.; Narang, R.; Clercq, E. D.; Balzarini, J. Synthesis, antimicrobial and antiviral evaluation of substituted imidazole derivatives. *Eur. J. Med. Chem.* **2009**, 44, 2347–2353.
- Shen, M. G.; Cai, C.; Yi, W. B. Ytterbium perfluorooctanesulfonate as an efficient and recoverable catalyst for the synthesis of trisubstituted imidazoles. *J. Fluorine Chem.* **2008**, 129, 541–544.
- Upadhyaya, D. J.; Barge, A.; Stefaniab, R.; Cravottoa, G. Efficient, solventless N-Boc protection of amines carried out at room temperature using sulfamic acid as recyclable catalyst. *Tetrahedron Lett.* **2007**, 48, 8318–8322.
- Viktor, A.; Katalin, G.; Gyorgyi, B.; Zsuzsanna, N.; Miklos, H.; Bela, R.; Erzsebet, G.; Terez, S.; Josef, D.; Szilagyi, G.; et al. Substituted imidazole pesticides. GB Patent 1407723, 1975; Chem. Abstr. **1973**, 84, 17351.
- Zang, H. J.; Su, Q. H.; Mo, Y. M.; Cheng, B. W.; Jun, S. Ionic liquid [EMIM]OAc under ultrasonic irradiation towards the first synthesis of trisubstituted imidazoles. *Ultrason. Sonochem.* **2010**, 17, 749–751.

Received August 30, 2011; accepted October 19, 2011