

Research Article

Tadele Mengesha, Armin Schikorra*, Adisak Seesanea and Sasikarn Yeepo

Calderón–Zygmund theory for strongly coupled linear system of nonlocal equations with Hölder-regular coefficient

<https://doi.org/10.1515/acv-2024-0005>

Received January 5, 2024; accepted July 18, 2024

Abstract: We extend the Calderón–Zygmund theory for nonlocal equations to strongly coupled system of linear nonlocal equations $\mathcal{L}_A^s u = f$, where the operator $\mathcal{L}_A^s$ is formally given by

$$\mathcal{L}_A^s u = \int_{\mathbb{R}^n} \frac{A(x, y)}{|x - y|^{n+2s}} \frac{(x - y) \otimes (x - y)}{|x - y|^2} (u(x) - u(y)) dy.$$

For $0 < s < 1$ and $A : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ taken to be symmetric and serving as a variable coefficient for the operator, the system under consideration is the fractional version of the classical Navier–Lamé linearized elasticity system. The study of the coupled system of nonlocal equations is motivated by its appearance in nonlocal mechanics, primarily in peridynamics. Our regularity result states that if $A(\cdot, y)$ is uniformly Hölder continuous and $\inf_{x \in \mathbb{R}^n} A(x, x) > 0$, then for $f \in L_{loc}^p$, for $p \geq 2$, the solution vector $u \in H_{loc}^{2s-\delta, p}$ for some $\delta \in (0, s)$.

Keywords: Regularity theory, Calderon–Zygmund, nonlocal PDE, elliptic systems

MSC 2020: 35R11, 35J99, 46E35, 47G30

Communicated by: Juha Kinnunen

1 Introduction

1.1 Motivation and notations

The goal of this work is to obtain Sobolev regularity estimates for solutions of the strongly coupled system of linear nonlocal equations $\mathcal{L}_A^s u = f$, where the operator $\mathcal{L}_A^s$ is formally given by

$$\mathcal{L}_A^s u = \int_{\mathbb{R}^n} \frac{A(x, y)}{|x - y|^{n+2s}} \frac{(x - y) \otimes (x - y)}{|x - y|^2} (u(x) - u(y)) dy.$$

Here we take $n \geq 1$, $0 < s < 1$, and $A : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ is taken to be symmetric and serves as a variable coefficient for the operator $\mathcal{L}_A^s$. For vectors $a = (a_1, \dots, a_n)$ and $b = (b_1, \dots, b_n)$ in $\mathbb{R}^n$, the tensor product $a \otimes b$ is the rank one matrix with its (ij) th entry being $a_i b_j$.

Coupled systems of linear nonlocal equations of the above type appear in applications. In fact, the operator $\mathcal{L}_A^s$ is related to the bond-based linearized peridynamic equation [31, 32]. To briefly describe where the operator

***Corresponding author: Armin Schikorra**, Department of Mathematics, University of Pittsburgh, 301 Thackeray Hall, Pittsburgh, PA 15260, USA, e-mail: armin@pitt.edu

Tadele Mengesha, Department of Mathematics, The University of Tennessee, Knoxville, 204 Ayres Hall, 1403 Circle Drive Knoxville, TN 37996, USA, e-mail: mengesha@utk.edu

Adisak Seesanea, Sasikarn Yeepo, Sirindhorn International Institute of Technology, Thammasat University, Khlong Nueng Pathum Thani 12120, Thailand, e-mail: adisak.see@siit.tu.ac.th, sasikarn.y@siit.tu.ac.th

comes from, consider a heterogeneous elastic solid occupying the domain Ω in $\mathbb{R}^n$, $n = 1, 2$, or 3 , that is linearly deforming when subjected to an external force field f . In the framework of the peridynamic model, a bounded domain hosting an elastic material is conceptualized as a sophisticated mass-spring system. Here, any pair of points x and y within the material is considered to interact through the bond vector $x - y$. When external load f is applied, the material undergoes a deformation, mapping a point x in the domain to the point $x + u(x) \in \mathbb{R}^n$, where the vector field u represents the displacement field. Adhering to the principles of uniform small strain theory [32], the strain of the bond $x - y$ is given by the nonlocal linearized strain

$$s[u](x, y) = \frac{u(x) - u(y)}{|x - y|} \cdot \frac{x - y}{|x - y|}.$$

The linearized bond-based peridynamic static model relates the displacement field u and the external load f by the equation (see [8, 18])

$$\int_{\mathbb{R}^n} \mathcal{C}(s[u](x, y), x, y) dy = f(x), \quad x \in \Omega,$$

where the vector-valued pairwise force density function $\mathcal{C}$ is given by

$$\mathcal{C}(s[u](x, y), x, y) = A(x, y) \rho(x - y) s[u](x, y) \frac{x - y}{|x - y|}.$$

In the above $A(x, y)$ serves as a “spring constant” for the bond joining x and y and the function ρ is the interaction kernel that is radial and describes the force strength between material points. After noting that

$$s[u](x, y) \frac{x - y}{|x - y|} = \frac{(x - y) \otimes (x - y)}{|x - y|^2} \frac{u(x) - u(y)}{|x - y|},$$

$\mathcal{L}_A^s$ is precisely the linearized bond-based peridynamic operator corresponding to the kernel of interaction

$$\rho(x - y) = \frac{1}{|x - y|^{n+2(s-1)}}.$$

To describe the problem we study, let us first introduce some standard notations and define relevant function spaces. For $t \in (0, 2)$ the fractional Laplacian $(-\Delta)^{\frac{t}{2}}$ is, defined via the Fourier transform,

$$(-\Delta)^{\frac{t}{2}} u = \mathcal{F}^{-1}(2\pi|\xi|^t \hat{u}),$$

where the Fourier transform is defined as $\mathcal{F}(u)(\xi) = \hat{u}(\xi) = \int_{\mathbb{R}^n} e^{-2\pi i x \cdot \xi} u(x) dx$. It also has a useful integral representation and for any vector field u in the Schwartz class

$$(-\Delta)^{\frac{t}{2}} u(x) = c_{t,n} \text{p.v.} \int_{\mathbb{R}^n} \frac{u(x) - u(y)}{|x - y|^{n+t}} dy,$$

where p.v. stands for the *principal value*, whose mentioning we will suppress. The inverse operator of the fractional Laplacian is the Riesz potential whose integral representation is

$$(-\Delta)^{-\frac{t}{2}} v(x) \equiv I^t v(x) = c \int_{\mathbb{R}^n} \frac{v(y)}{|x - y|^{n-t}} dy$$

for a vector field v in the Schwartz class. We introduce types of fractional Sobolev spaces that we need to state the main result: Bessel potential spaces $H^{s,p}$ and Besov spaces $W^{s,p}$. For $1 < p < \infty$ and $s \in (0, 1)$, the Bessel potential spaces $H^{s,p}(\mathbb{R}^n)$ are defined as follows: $f \in H^{s,p}(\mathbb{R}^n)$ if $f \in L^p(\mathbb{R}^n)$ and $(-\Delta)^{\frac{s}{2}} f \in L^p(\mathbb{R}^n)$. The associated norm is

$$\|f\|_{H^{s,p}(\mathbb{R}^n)} := \|f\|_{L^p(\mathbb{R}^n)} + \|(-\Delta)^{\frac{s}{2}} f\|_{L^p(\mathbb{R}^n)}.$$

For any open subset $\Omega \subset \mathbb{R}^n$, the Besov spaces $W^{s,p}(\Omega)$, for $s \in (0, 1)$, are induced by the semi-norm (called Sobolev–Slobodeckij or Gagliardo norm)

$$[f]_{W^{s,p}(\Omega)} = \left(\int_{\Omega} \int_{\Omega} \frac{|f(x) - f(y)|^p}{|x - y|^{n+sp}} dx dy \right)^{\frac{1}{p}},$$

and $\|\cdot\|_{W^{s,p}(\Omega)} = \|\cdot\|_{L^p(\Omega)} + [\cdot]_{W^{s,p}(\Omega)}$ serves as a norm. For $p = 2$, $W^{s,2}(\mathbb{R}^n) = H^{s,2}(\mathbb{R}^n)$, which we denote by $H^s(\mathbb{R}^n)$. For $p < 2$, we have $W^{s,p}(\mathbb{R}^n) \subsetneq H^{s,p}(\mathbb{R}^n)$ and for $p > 2$, $H^{s,p}(\mathbb{R}^n) \subsetneq W^{s,p}(\mathbb{R}^n)$. These spaces are particular examples of the more general Triebel–Lizorkin spaces and $F_{pp}^s(\mathbb{R}^n) = W^{s,p}(\mathbb{R}^n)$ and $F_{p,2}^s(\mathbb{R}^n) = H^{s,p}(\mathbb{R}^n)$, see [24].

Given $s \in (0, 1)$, $A \in L^\infty(\mathbb{R}^n \times \mathbb{R}^n)$ and $u \in L_{\text{loc}}^1(\mathbb{R}^n, \mathbb{R}^n)$, we understand $\mathcal{L}_A^s u$ as a distribution defined as

$$\langle \mathcal{L}_A^s u, \varphi \rangle := \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{A(x, y)}{|x - y|^{n+2s}} (u(y) - u(x)) \cdot \frac{(y - x)}{|y - x|} (\varphi(y) - \varphi(x)) \cdot \frac{(y - x)}{|y - x|} dy dx$$

for all $\varphi \in C_c^\infty(\mathbb{R}^n, \mathbb{R}^n)$. Moreover, if $u \in H^s(\mathbb{R}^n, \mathbb{R}^n)$, then from the above definition, $\mathcal{L}_A^s u \in H^{-s}(\mathbb{R}^n, \mathbb{R}^n)$ with the estimate that

$$\|\mathcal{L}_A^s u\|_{H^{-s}} \leq \frac{1}{\lambda} \|u\|_{H^s},$$

where $H^{-s}(\mathbb{R}^n, \mathbb{R}^n)$ represents the dual of $H^s(\mathbb{R}^n, \mathbb{R}^n)$.

Our interest is to address the question of regularity of solutions u to $\mathcal{L}_A^s u = f$ relative to the data f . To that end, we require the coefficient A to satisfy some continuity and boundedness assumptions. First, we say A satisfies a uniform Hölder continuity assumption if for some $\alpha \in (0, 1)$ and $\Lambda > 0$,

$$\sup_{z \in \mathbb{R}^n} |A(z, x) - A(z, y)| \leq \Lambda |x - y|^\alpha. \quad (1.1)$$

Given $\lambda, \Lambda > 0$ and $\alpha \in (0, 1)$, we define the coefficient class

$$\mathcal{A}(\alpha, \lambda, \Lambda) = \left\{ A : A(x, y) = A(y, x), \inf_{x \in \mathbb{R}^n} K(x, x) > \lambda, \|A\|_{L^\infty} \leq \frac{1}{\lambda}, \text{ and satisfies (1.1)} \right\}.$$

We note that members of the coefficient class $\mathcal{A}(\alpha, \lambda, \Lambda)$ can be negative off diagonal. Indeed, as indicated in [19], the coefficient $A(x, y) = \frac{2+|x|^\alpha+|y|^\alpha}{1+|x|^\alpha+|y|^\alpha} + 10(\sin x + \sin y) \frac{|x-y|^\alpha}{1+|x-y|^\alpha}$ belongs to $\mathcal{A}(\alpha, \lambda, \Lambda)$ for some λ, Λ and yet can be negative off diagonal.

Now for an open set $\Omega \subset \mathbb{R}^n$ and $f \in H^{-s}(\mathbb{R}^n, \mathbb{R}^n)$, a vector field $u \in H^s(\mathbb{R}^n, \mathbb{R}^n)$ is a solution to $\mathcal{L}_A^s u = f$ in Ω if

$$\langle \mathcal{L}_A^s u, \varphi \rangle = \langle f, \varphi \rangle \quad \text{for all } \varphi \in C_c^\infty(\Omega, \mathbb{R}^n). \quad (1.2)$$

In the event, $A = 1$, then operator agrees with the integral operator defined as

$$(-\mathring{\Delta})^s u(x) := \text{p.v.} \int_{\mathbb{R}^n} \frac{1}{|z|^{n+2s}} \left(\frac{z \otimes z}{|z|^2} \right) (u(x) - u(x+z)) dz,$$

where the integral converges in the sense of principal value for smooth vector fields. For vector fields u in the Schwarz space $\mathcal{S}(\mathbb{R}^n, \mathbb{R}^n)$, we have

$$\mathcal{F}((-\mathring{\Delta})^s u) = (2\pi|\xi|)^{2s} \left(\ell_1 \mathbb{I} + \ell_2 \frac{\xi \otimes \xi}{|\xi|^2} \right) \mathcal{F}(u) \quad (1.3)$$

for some positive constants ℓ_1 and ℓ_2 depending only on n and s . As a consequence, as shown in [20] for any $\tau > 0$ and $f \in L^p(\mathbb{R}^n, \mathbb{R}^n)$ with $1 < p < \infty$, then the solution u to

$$(-\mathring{\Delta})^s u + \tau u = f$$

lives in $H^{2s,p}(\mathbb{R}^n, \mathbb{R}^n)$. For the nonlocal equation of variable coefficient (1.2), we would like to obtain a Sobolev regularity of the above type for solutions in the event that the right hand side f has additional regularity. We begin by noting that for some λ and Λ , $A \in \mathcal{A}(\alpha, \lambda, \Lambda)$, and $f \in H^{-s}(\mathbb{R}^n, \mathbb{R}^n)$, a solution to (1.2) exists under some volumetric condition on u and on the domain Ω . Indeed, if Ω is a Lipschitz domain, a minimizer of the energy

$$E(u) = \frac{1}{2} \langle \mathcal{L}_A^s u, u \rangle - \langle f, u \rangle$$

over the space $\tilde{H}^s(\Omega) = \{u \in H^s(\mathbb{R}^n, \mathbb{R}^n) : u = 0 \text{ on } \mathbb{R}^n \setminus \Omega\}$ will satisfy equation (1.2). For a Lipschitz domain Ω , $\tilde{H}^s(\Omega)$ is precisely the closure of $C_c(\Omega, \mathbb{R}^n)$ in $H^s(\mathbb{R}^n, \mathbb{R}^n)$, and therefore $\tilde{H}^s(\Omega)$ is a Hilbert space with obvious inner product, [17]. The existence of a minimizer for the quadratic functional E over $\tilde{H}^s(\Omega)$, with a possible sign

changing $A \in \mathcal{A}(\alpha, \lambda, \Lambda)$ will be shown later using Lax–Milgram theorem. As has been demonstrated in [18], with a proper multiplicative constant $c(s, n)$, in terms of the nonlocality parameter s , the operator $c(s, n)\mathcal{L}_A^s u$ that corresponds to $A(x, y) = 1/2(a(x) + a(y))$ will converge in an appropriate sense to the Lamé differential operator

$$L_0 u(x) = \operatorname{div}(a(x)\nabla u) + 2\nabla(a(x) \operatorname{div} u(x)).$$

This operator is strongly elliptic in the sense of Legendre–Hadamard but not uniformly elliptic. One can then view (1.2) as a fractional analogue of the classical Navier–Lamé system of linearized elasticity equation.

1.2 Statement of the main result

The main result of the paper is the following interior regularity estimate which is the version of the regularity result proved in [19] for the coupled system of nonlocal equations under discussion.

Theorem 1.1. *Let $s \in (0, 1)$ and $s \leq t < \min\{2s, 1\}$. Let $\Omega \subset \mathbb{R}^n$ be an open bounded set. If for $2 \leq q < \infty$, $f_1, f_2 \in L^q(\Omega, \mathbb{R}^n) \cap L^2(\mathbb{R}^n, \mathbb{R}^n)$, and $u \in H^s(\mathbb{R}^n, \mathbb{R}^n)$ is a distributional solution of $\mathcal{L}_A^s u = (-\Delta)^{\frac{2s-t}{2}} f_1 + f_2$ in Ω , in the sense that*

$$\langle \mathcal{L}_A^s u, \varphi \rangle = \int_{\mathbb{R}^n} \langle f_1, (-\Delta)^{\frac{2s-t}{2}} \varphi \rangle dx + \int_{\mathbb{R}^n} \langle f_2, \varphi \rangle dx \quad \text{for all } \varphi \in C_c^\infty(\Omega; \mathbb{R}^n),$$

with $\mathcal{L}_A^s$ corresponding to $A \in \mathcal{A}(\alpha, \lambda, \Lambda)$ for some given $\alpha \in (0, 1)$ and $\lambda, \Lambda > 0$, then we have $(-\Delta)^{\frac{t}{2}} u \in L_{\text{loc}}^q(\Omega, \mathbb{R}^n)$ and for any $\Omega' \subset\subset \Omega$ we have

$$\|(-\Delta)^{\frac{t}{2}} u\|_{L^q(\Omega')} \leq C \left(\|u\|_{W^{s,2}(\mathbb{R}^n)} + \sum_{i=1}^2 \|f_i\|_{L^q(\Omega)} + \|f_i\|_{L^2(\mathbb{R}^n)} \right).$$

The constant C depends only $s, t, q, \alpha, \lambda, \Lambda, \Omega$, and Ω' .

Remark 1.2. Since $q \geq 2$, by the embedding theorems on Triebel–Lizorkin spaces, we have $F_{q,2}^s \subset F_{qq}^s$, cf. [24]. As a consequence, under the same assumption of the theorem, Theorem 1.1 implies that the solution $u \in W_{\text{loc}}^{t,q}(\Omega)$ and from the estimate we obtain

$$\|u\|_{W^{t,q}(\Omega')} = \left(\iint_{\Omega' \times \Omega'} \frac{|u(x) - u(y)|^q}{|x - y|^{n+ tq}} dx dy \right)^{\frac{1}{q}} \leq C \left(\|u\|_{W^{s,2}(\mathbb{R}^n)} + \sum_{i=1}^2 \|f_i\|_{L^q(\Omega)} + \|f_i\|_{L^2(\mathbb{R}^n)} \right).$$

The proof of Theorem 1.1 parallels the approach used in [19]. Namely, we compare the operator $\mathcal{L}_A^s$ with the simpler operator $\tilde{\mathcal{L}}_{A_D}^{s_1, s_2}$, where $s_1 + s_2 = 2s$, and is defined as, for $u \in H^s(\mathbb{R}^n, \mathbb{R}^n)$ and $\varphi \in C_c^\infty(\mathbb{R}^n, \mathbb{R}^n)$,

$$\langle \tilde{\mathcal{L}}_{A_D}^{s_1, s_2} u, \varphi \rangle = \int_{\mathbb{R}^n} A_D(z) \langle (c_1 \mathbb{I} + c_2 \mathcal{R} \otimes \mathcal{R})(-\Delta)^{\frac{s_1}{2}} u(z), (-\Delta)^{\frac{s_2}{2}} \varphi(z) \rangle dz \quad (1.4)$$

for constants c_1 and c_2 that will be determined as a function of s and n . In the above definition, the operator $\mathcal{R} = (\mathcal{R}_1, \mathcal{R}_2, \dots, \mathcal{R}_n)$ is the vector of Riesz transforms, and $A_D(z) = A(z, z)$, the restriction of the coefficient A on the diagonal. Notice that for constant coefficients the two operators $\mathcal{L}_A^s$ and $\tilde{\mathcal{L}}_{A_D}^{s,s}$ coincide. Indeed, if $A(x, y) = A$, constant, then by using (1.3), for vector fields in the Schwarz space

$$\mathcal{L}_A^s u = A(-\Delta)^s u = A(\ell_1(-\Delta)^{\frac{s}{2}} u + \ell_2(\mathcal{R} \otimes \mathcal{R})(-\Delta)^{\frac{s}{2}} u) = \tilde{\mathcal{L}}_A^{s,s} u$$

with $c_1 = \ell_1$ and $c_2 = \ell_2$. We will prove an optimal regularity result for solutions of the strongly coupled equation

$$\langle \tilde{\mathcal{L}}_{A_D}^{s_1, s_2} u, \varphi \rangle = \langle g, \varphi \rangle \quad \text{for all } \varphi \in C_c^\infty(\mathbb{R}^n, \mathbb{R}^n) \quad (1.5)$$

and use those solutions as approximations of the solution to the original system of equations. The mechanism we accomplish this is via perturbation argument where we show that the difference operator

$$\mathcal{D}_{s,t} u = \mathcal{L}_A^s u - \tilde{\mathcal{L}}_{A_D}^{s,t} u$$

can be understood as a lower order term in the event that A is Hölder continuous.

While our work studies solutions to strongly coupled linear nonlocal PDEs, there has been a number of results in the literature that studied the regularity of solutions to scalar nonlocal PDEs. To name a few, opti-

mal local regularity results are obtained in [2] for weak solutions to the Dirichlet problem associated with the fractional Laplacian. Similar results are obtained for the fractional heat equation in [12, 13]. Almost optimal regularity results are obtained in [5] for weak solutions to nonlocal equations with Hölder regular coefficients. Optimal Sobolev regularity are proved in [7] for strong solutions to nonlocal equations with translation invariant coefficients ($A(x, y) = A(x - y)$). For equations with less regular coefficients, higher integrability and higher differentiability results are obtained in [21, 23] for nonlocal equations with variable coefficients that have small mean oscillations. See also [9, 10] for related results. For elliptic, measurable, and bounded coefficients, solutions to nonlocal PDEs are proved in [15] to have a self-improvement property where higher integrability and higher differentiability are obtained without any smoothness assumption on the coefficients, see also [27]. Similar results are also verified in [4, 30] for solutions to nonlocal double phase problems. For a concise description of the results of the above mentioned manuscripts, we refer to [19]. See also [1, 3, 22].

The paper is organized as follows. In the next section, we will discuss some preliminary results we need in the sequel. In Section 3, we estimate $\langle \mathcal{D}_{s,t} u, \varphi \rangle$ in terms of the Riesz potential $I^s = (-\Delta)^{-\frac{s}{2}}$. In Section 4, we will develop the optimal regularity result for a solution of equation (1.5). In Section 5, we prove the main result of the paper by using an iterative argument making use of the commutator estimate we prove in Section 3 and the optimal regularity result obtained in Section 4.

Throughout the paper, we work under the convention that domains of integrals are always open sets and we use the symbol $\llcorner$ to say compactly contained, e.g., $\Omega_1 \llcorner \Omega_2$ if $\overline{\Omega_1}$ is compact and $\overline{\Omega_1} \subset \Omega_2$. Constants change from line to line, and unless it is important we may not detail their dependence on various parameters. We will make frequent use of $\lesssim$, $\gtrsim$ and $\asymp$, which denotes inequalities with multiplicative constants (depending on non-essential data). For example we say $A \lesssim B$ if for some constant $C > 0$ we have $A \leq CB$. We will use the angle bracket $\langle \cdot, \cdot \rangle$ to represent the standard inner product or the duality pairing depending on the context.

2 Some preliminaries

Our arguments below make use of the various definition and properties of fractional Laplacian operators, accompanying Sobolev spaces, Sobolev inequalities, and various embedding that can be found in [24]. See [6, 11], or monographs [25] for more on fractional operators.

Sobolev inequalities needed for this paper are proved in [19, Proposition 2.1] (see also [33]) and we summarize them as follows.

Lemma 2.1. *The following statements hold:*

(a) *If $sp < n$, then there exists a constant $C = C(s, p, n) > 0$ such that*

$$\|I^s v\|_{L^{\frac{np}{n-sp}}(\mathbb{R}^n)} \leq C \|v\|_{L^p(\mathbb{R}^n)} \quad \text{for any } v \in L^p(\mathbb{R}^n, \mathbb{R}^n). \quad (2.1)$$

In addition, if $\Omega \subset \mathbb{R}^n$ is bounded, then corresponding to any $q \in [1, \frac{np}{n-sp}]$, there is a constant $C = C(s, p, n, \Omega) > 0$ such that

$$\|I^s v\|_{L^q(\Omega)} \leq C \|v\|_{L^p(\mathbb{R}^n)} \quad \text{for any } v \in L^p(\mathbb{R}^n, \mathbb{R}^n). \quad (2.2)$$

(b) *If $sp \geq n$ and $\Omega \subset \mathbb{R}^n$ is bounded domain, then for any $q \in [1, \infty)$, and $r \in [1, \frac{n}{s})$, there exists a constant $C = C(s, p, n, \Omega) > 0$ such that for any $v \in L^p(\mathbb{R}^n, \mathbb{R}^n)$,*

$$\|I^s v\|_{L^q(\Omega)} \leq C (\|v\|_{L^p(\mathbb{R}^n)} + \|v\|_{L^r(\mathbb{R}^n)}). \quad (2.3)$$

The above Sobolev estimates together with the relationship between the fractional Laplacian and the Riesz potentials yield the following result that is also stated and proved in [19, Proposition 2.4]. We state it here in a slightly different way to suit our setting.

Lemma 2.2 ([19]). *Suppose that $\eta_1, \eta_2 \in C_c^\infty(\mathbb{R}^n)$, and $\eta_2 = 1$ in the neighborhood of the support of η_1 . Then for any $\psi \in C_c^\infty(\mathbb{R}^n)$ such that $\text{supp}(\psi) \subset \{x : \eta_1(x) = 1\}$, and any $q, p \in (1, \infty)$ and $\tau \in (0, 2)$ we have*

$$\|(1 - \eta_2)(-\Delta)^{\frac{\tau}{2}}((1 - \eta_1)I^\tau \psi)\|_{L^q(\mathbb{R}^n)} \leq C \|\psi\|_{L^p(\mathbb{R}^n)}. \quad (2.4)$$

Moreover, if $r > \frac{np}{n+\tau p} > 1$ for $\tau < 1$, then for any bounded set $\Omega \subset \mathbb{R}^n$, there exists a constant $C(\Omega)$ such that

$$\|(-\Delta)^{\frac{s}{2}}((1-\eta_1)I^\tau\psi)\|_{L^{p'}(\Omega)} \leq C\|\psi\|_{L^{p'}(\mathbb{R}^n)}. \quad (2.5)$$

In either case the constant C may also depend on r, q, τ, p, n , and on η_1, η_2 , but not on ψ .

Notice that because of the strict inclusion of the support of $1 - \eta_2$ into the support of $1 - \eta_1$, inequality (2.4) holds for any $p, q \in (1, \infty)$. The way it is written here, the inequality is slightly different from part (a) of [19, Proposition 2.4] but the same proof can be repeated for the proof of (2.4).

We also mention the dual definition of $(-\Delta)^{\frac{s}{2}}$ operator. Indeed, for vector fields u and v in the Schwartz class, the L^2 -inner product of $(-\Delta)^{\frac{s}{2}}u(x)$ and $v(x)$ can be represented as, for $s \in (0, 2)$,

$$\int_{\mathbb{R}^n} (-\Delta)^{\frac{s}{2}}u(x) \cdot v(x) dx = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{(u(y) - v(x)) \cdot (v(y) - v(x))}{|x - y|^{n+s}} dy dx. \quad (2.6)$$

The proof can be found [26, Proposition 2.36.] or [6].

The Riesz transform, $\mathcal{R} = (\mathcal{R}_1, \dots, \mathcal{R}_n) := \nabla I^1$, plays a central role in this work. First, $\mathcal{R}$ has the Fourier symbol $c|\xi|^{-1}$, and can also be represented as

$$\mathcal{R}f(x) = \int_{\mathbb{R}^n} \frac{\frac{x-y}{|x-y|}}{|x-y|^n} f(y) dy.$$

Second, we will use the fact that they are Calderón–Zygmund operators and for $1 < p < \infty$, there exists a constant $C = C(n, p) > 0$ such that

$$\|\mathcal{R}f\|_{L^p} \leq C\|f\|_{L^p} \quad \text{for all } f \in L^p.$$

Finally, we state and prove existence of a solution to the nonlocal system (1.2). We recall the space

$$\tilde{H}^s(\Omega) = \{u \in H^s(\mathbb{R}^n, \mathbb{R}^n) : u = 0 \text{ on } \mathbb{R}^n \setminus \Omega\},$$

which is equal to the closure of $C_c(\Omega, \mathbb{R}^n)$ in $H^s(\mathbb{R}^n, \mathbb{R}^n)$ when Ω is a Lipschitz domain. As noted earlier, and therefore, $\tilde{H}^s(\Omega)$ is a Hilbert space with obvious inner product, [17].

Proposition 2.3. Suppose that $\Omega \subset \mathbb{R}^n$ is an open bounded set with Lipschitz boundary. Then for any, $\alpha, s \in (0, 1)$, there exists $C > 0$ so that if $\frac{\lambda}{\Lambda} > C$, $A \in \mathcal{A}(\alpha, \lambda, \Lambda)$, and $f \in H^{-s}(\mathbb{R}^n, \mathbb{R}^n)$, then there exists a unique $u \in \tilde{H}^s(\Omega)$ such that

$$\langle \mathcal{L}_A^s u, v \rangle = \langle f, v \rangle \quad \text{for all } v \in \tilde{H}^s(\Omega).$$

Moreover, the solution is a minimizer of the functional

$$v \mapsto \frac{1}{2} \langle \mathcal{L}_A^s v, v \rangle - \langle f, v \rangle \quad \text{for all } v \in \tilde{H}^s(\Omega).$$

Proof. We will apply the Lax–Milgram theorem to show existence of a unique solution corresponding to any f in the dual of the Hilbert space $\tilde{H}^s(\Omega)$. To that end, we introduce the bilinear form $B_A : \tilde{H}^s(\Omega) \times \tilde{H}^s(\Omega) \rightarrow \mathbb{R}$ by

$$B_A(u, v) = \langle \mathcal{L}_A^s u, v \rangle.$$

By applying Hölder's inequality, we see that B is continuous. In fact, for any $u, v \in \tilde{H}^s(\mathbb{R}^n, \mathbb{R}^n)$,

$$|B_A(u, v)| \leq \frac{1}{\lambda} |u|_{H^s} |v|_{H^s} \leq \frac{1}{\lambda} \|u\|_{H^s} \|v\|_{H^s}.$$

Next, we show that this bilinear form is coercive in $\tilde{H}^s(\Omega)$. Now for any $v \in \tilde{H}^s(\Omega)$,

$$\begin{aligned} B_A(v, v) &= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} A(x, y) \frac{|(v(y) - v(x)) \cdot \frac{(y-x)}{|y-x|}|^2}{|y-x|^{n+2s}} dy dx \\ &= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} A_D(x, x) \frac{|(v(y) - v(x)) \cdot \frac{(y-x)}{|y-x|}|^2}{|y-x|^{n+2s}} dy dx + E_{A-A_D}(v), \end{aligned} \quad (2.7)$$

where

$$E_{A-A_D}(v) = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} [A(x, y) - A_D(x, x)] \frac{|(v(y) - v(x)) \cdot \frac{(y-x)}{|y-x|}|^2}{|y-x|^{n+2s}} dy dx.$$

Since $\inf_{x \in \mathbb{R}^n} A_D(x, x) \geq \lambda > 0$ and using the fractional Korn's inequality in [29] which proves that there is a constant $\tilde{c} > 0$ such that for any $s \in (0, 1)$ and $v \in H^s(\mathbb{R}^n, \mathbb{R}^n)$,

$$\tilde{c} |v|_{H^s(\mathbb{R}^n)}^2 \leq \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|(v(y) - v(x)) \cdot \frac{(y-x)}{|y-x|}|^2}{|y-x|^{n+2s}} dy dx \leq |v|_{H^s(\mathbb{R}^n)}^2,$$

we can estimate the first term of (2.7) as

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} A_D(x, x) \frac{|(v(y) - v(x)) \cdot \frac{(y-x)}{|y-x|}|^2}{|y-x|^{n+2s}} dy dx \geq \tilde{c} \lambda |v|_{H^s(\mathbb{R}^n)}^2.$$

Applying the fractional Poincaré inequality on $\tilde{H}^s(\Omega)$, there exists $C = C(\Omega, s, n) > 0$ such that for any $v \in \tilde{H}^s(\Omega)$, $|v|_{H^s(\mathbb{R}^n)}^2 \geq C \|v\|_{H^s(\mathbb{R}^n)}^2$, and therefore combining with above estimate we have

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} A_D(x, x) \frac{|(v(y) - v(x)) \cdot \frac{(y-x)}{|y-x|}|^2}{|y-x|^{n+2s}} dy dx \geq \tilde{C} \lambda \|v\|_{H^s(\mathbb{R}^n)}^2. \quad (2.8)$$

To estimate the second term of (2.7), we first notice that if $A \in \mathcal{A}(\alpha, \lambda, \Lambda)$, then for any $\beta \in (0, \alpha)$, we have $A \in \mathcal{A}(\beta, \lambda, \tilde{\Lambda}_\beta)$ for some $\tilde{\Lambda}_\beta > 0$. This follows from the estimate that for any $\beta \in (0, \alpha)$ and any $x, y, z \in \mathbb{R}^n$,

$$|A(z, x) - A(z, y)| = |A(z, x) - A(z, y)|^{\frac{\beta}{\alpha}} |A(z, x) - A(z, y)|^{1-\frac{\beta}{\alpha}} \leq (2\|A\|_{L^\infty})^{1-\frac{\beta}{\alpha}} \Lambda^{\frac{\beta}{\alpha}} |x-y|^\beta.$$

Thus, without loss of generality we may assume that $0 < \alpha < 2s$. It then follows that

$$E_{A-A_D}(v) \geq -\Lambda \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|(v(y) - v(x)) \cdot \frac{(y-x)}{|y-x|}|^2}{|y-x|^{n+2(s-\frac{\alpha}{2})}} dy dx \geq -\Lambda |v|_{H^{(s-\frac{\alpha}{2})}(\mathbb{R}^n)}^2.$$

We now use the continuous embedding $H^s(\mathbb{R}^n, \mathbb{R}^n) \subset H^{(s-\frac{\alpha}{2})}(\mathbb{R}^n, \mathbb{R}^n)$, to conclude that there is a positive constant $c = c(s, \alpha, n)$ such that

$$E_{A-A_D}(v) \geq -\Lambda c \|v\|_{H^s(\mathbb{R}^n)}^2. \quad (2.9)$$

Combining (2.8) and (2.9), we obtain that for any $v \in H^s(\Omega)$,

$$B(v, v) \geq (\tilde{c}\lambda - c\Lambda) \|v\|_{H^s(\mathbb{R}^n)}^2.$$

For λ, Λ satisfying $(\tilde{c}\lambda - c\Lambda > 0)$, the bilinear form is coercive. $\square$

3 Commutator estimates

In this section, we obtain estimates for the quantity $\mathcal{D}_{s_1, s_2} u$ defined in the previous section. To be precise, for $s \in (0, 1)$ such that $s_1 + s_2 = 2s$, $u \in H^s(\mathbb{R}^n, \mathbb{R}^n)$, and $\varphi \in C_c(\mathbb{R}^n, \mathbb{R}^n)$, we recall that

$$\mathcal{D}_{s_1, s_2}(u, \varphi) = \langle \mathcal{L}_A^s u, \varphi \rangle - \langle \tilde{\mathcal{L}}_{A_D}^{s_1, s_2} u, \varphi \rangle. \quad (3.1)$$

Before we begin estimating this difference, let us first find a different characterization of the operator $\mathcal{L}_A^s u$ that uses Riesz potentials. To that end, for any $x, y \in \mathbb{R}^n$ and $\varphi \in C_c(\mathbb{R}^n, \mathbb{R}^n)$,

$$\begin{aligned} \varphi(x) - \varphi(y) &= c_2 (I^{s_2}(-\Delta)^{\frac{s_2}{2}} \varphi(x) - I^{s_2}(-\Delta)^{\frac{s_2}{2}} \varphi(y)) \\ &= c_2 \int_{\mathbb{R}^n} (-\Delta)^{\frac{s_2}{2}} \varphi(z_2) (|x - z_2|^{s_2-n} - |y - z_2|^{s_2-n}) dz_2 \end{aligned} \quad (3.2)$$

for a constant c_2 that depends only on s_2 and n . This identity remains valid for $\varphi \in H^{s_2}(\mathbb{R}^n, \mathbb{R}^n)$. Similarly, for any $0 \leq \varepsilon < s_2$, we can write

$$\begin{aligned}\varphi(x) - \varphi(y) &= c_2(\varepsilon)(I^{s_2-\varepsilon}(-\Delta)^{\frac{s_2-\varepsilon}{2}}\varphi(x) - I^{s_2-\varepsilon}(-\Delta)^{\frac{s_2-\varepsilon}{2}}\varphi(y)) \\ &= c_2(\varepsilon) \int_{\mathbb{R}^n} (-\Delta)^{\frac{s_2-\varepsilon}{2}}\varphi(z_2)(|x - z_2|^{s_2-\varepsilon-n} - |y - z_2|^{s_2-\varepsilon-n}) dz_2,\end{aligned}\quad (3.3)$$

where $c_2(\varepsilon)$ depends on ε , in addition to s_2 and n . We note that $c_2(0) = c_2 > 0$. Now we plug $\varphi(x) - \varphi(y)$ and $u(x) - u(y)$ in

$$\langle \mathcal{L}_A^s u, \varphi \rangle = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{A(x, y)}{|x - y|^{n+2s}} \left\langle \frac{(x - y) \otimes (x - y)}{|x - y|^2} (u(x) - u(y)), \varphi(x) - \varphi(y) \right\rangle dx dy$$

and apply Fubini's theorem to obtain the expression that for any $\varepsilon \in [0, s_2)$,

$$\langle \mathcal{L}_A^s u, \varphi \rangle = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \langle \mathbb{K}_A^\varepsilon(z_1, z_2)(-\Delta)^{\frac{s_1}{2}} u(z_1), (-\Delta)^{\frac{s_2-\varepsilon}{2}} \varphi(z_2) \rangle dz_1 dz_2, \quad (3.4)$$

where for any set function $B(x, y)$, and $0 \leq \varepsilon < s_2$,

$$\mathbb{K}_B^\varepsilon(z_1, z_2) = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{B(x, y)}{|x - y|^{n+2s}} \kappa_{s_1}^{s_2-\varepsilon}(x, y, z_1, z_2) \frac{(x - y) \otimes (x - y)}{|x - y|^2} dx dy \quad (3.5)$$

with, $c(\varepsilon) = c_1 \cdot c_2(\varepsilon)$,

$$\kappa_{s_1}^{s_2-\varepsilon}(x, y, z_1, z_2) = c(\varepsilon)(|x - z_1|^{s_1-n} - |y - z_1|^{s_1-n})(|x - z_2|^{s_2-\varepsilon-n} - |y - z_2|^{s_2-\varepsilon-n}). \quad (3.6)$$

See [19, Lemma 3.6.] for a rigorous justification of the above calculations. With this at hand, we introduce an intermediate operator $A_D \mathcal{L}_1^s$ given by

$$\langle A_D \mathcal{L}_1^s u, \varphi \rangle := \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} A_D(z_1) \langle \mathbb{K}_1^0(z_1, z_2)(-\Delta)^{\frac{s_1}{2}} u(z_1), (-\Delta)^{\frac{s_2}{2}} \varphi(z_2) \rangle dz_1 dz_2,$$

where $\mathbb{K}_1^0$ is as given in (3.5) with $\varepsilon = 0$ and $B = 1$. It then follows from (3.3) that for $\varepsilon \in [0, s_2)$, and integrating in the z_2 variable that

$$\langle A_D \mathcal{L}_1^s u, \varphi \rangle = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} A_D(z_1) \langle \mathbb{K}_1^\varepsilon(z_1, z_2)(-\Delta)^{\frac{s_1}{2}} u(z_1), (-\Delta)^{\frac{s_2-\varepsilon}{2}} \varphi(z_2) \rangle dz_1 dz_2.$$

Now, for a given $\varepsilon \in [0, s_2)$, we first write the difference $\mathcal{D}_{s_1, s_2}(u, \varphi)$ defined in (3.1) as

$$\begin{aligned}\mathcal{D}_{s_1, s_2}(u, \varphi) &= \langle \mathcal{L}_A^s u - (A_D \mathcal{L}_1^s)u, \varphi \rangle + \langle (A_D \mathcal{L}_1^s)u, \varphi \rangle - \langle \tilde{\mathcal{L}}_{A_D}^{s_1, s_2} u, \varphi \rangle \\ &= \mathcal{D}_1^s(u, \varphi) + \mathcal{D}_2^{s_1, s_2}(u, \varphi).\end{aligned}$$

We then have that

$$\langle \mathcal{L}_A^s u, \varphi \rangle = \langle \tilde{\mathcal{L}}_{A_D}^{s_1, s_2} u, \varphi \rangle + \mathcal{D}_1^s(u, \varphi) + \mathcal{D}_2^{s_1, s_2}(u, \varphi). \quad (3.7)$$

The next two propositions estimate the last two terms of (3.7). First, we estimate $\mathcal{D}_1^s(u, \varphi) = \langle \mathcal{L}_A^s u - (A_D \mathcal{L}_1^s)u, \varphi \rangle$.

Proposition 3.1. *Let $s \in (0, 1)$ with $s_1 + s_2 = 2s$, $\alpha \in (0, 1)$ and $\Lambda > 0$. Then there exist constants $\sigma_0 \in (0, \alpha]$ and a constant $c > 0$ such that the decomposition (3.7) holds and for any $A \in \mathcal{A}(\alpha, \Lambda) = \{A : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R} : |A(x, y)| \leq \Lambda, (1.1) \text{ holds}\}$, any $\sigma \in (0, \sigma_0)$, and any $\varepsilon \in (0, \frac{\sigma}{4})$, we have*

$$|\mathcal{D}_1^s(u, \varphi)| \leq \int_{\mathbb{R}^n} I^{\sigma-\varepsilon} |(-\Delta)^{\frac{s_1}{2}} u|(x) |(-\Delta)^{\frac{s_2-\varepsilon}{2}} \varphi|(x) dx$$

and

$$|\mathcal{D}_1^s(u, \varphi)| \leq \int_{\mathbb{R}^n} I^{\sigma-\varepsilon} |(-\Delta)^{\frac{s_1-\varepsilon}{2}} u|(x) |(-\Delta)^{\frac{s_2}{2}} \varphi|(x) dx$$

for all $u \in H^{s_1, p}(\mathbb{R}^n, \mathbb{R}^n)$ and $\varphi \in C_c^\infty(\mathbb{R}^n, \mathbb{R}^n)$.

Proof. From the definition of the operator $A_D \mathcal{L}_1^s$ and integrating in the z_2 variable that for any $\varepsilon \in [0, s_2]$,

$$\langle A_D \mathcal{L}_1^s u, \varphi \rangle = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} A_D(z_1) \langle \mathbb{K}_1^\varepsilon(z_1, z_2) (-\Delta)^{\frac{s_1}{2}} u(z_1), (-\Delta)^{\frac{s_2-\varepsilon}{2}} \varphi(z_2) \rangle dz_1 dz_2,$$

and so using the (3.4) and the definition of $A_D \mathcal{L}_1^s$, we have that

$$\mathcal{D}_1^s(u, \varphi) = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \langle \mathbb{M}^\varepsilon(z_1, z_2) (-\Delta)^{\frac{s_1}{2}} u(z_1), (-\Delta)^{\frac{s_2-\varepsilon}{2}} \varphi(z_2) \rangle dz_1 dz_2, \quad (3.8)$$

where

$$\begin{aligned} \mathbb{M}^\varepsilon(z_1, z_2) &= \mathbb{K}_A^\varepsilon(z_1, z_2) - A_D(z_1) \mathbb{K}_1^\varepsilon(z_1, z_2) \\ &= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{A(x, y) - A_D(z_1)}{|x - y|^{n+2s}} \kappa_{s_1}^{s_2-\varepsilon}(x, y, z_1, z_2) \frac{(x - y) \otimes (x - y)}{|x - y|^2} dx dy \end{aligned}$$

and $\kappa_{s_1}^{s_2-\varepsilon}$ as defined in (3.6). It then follows that

$$|\mathbb{M}^\varepsilon(z_1, z_1)| \leq \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} |A(x, y) - A_D(z_1)| \frac{|\kappa_{s_1}^{s_2-\varepsilon}(x, y, z_1, z_1)|}{|x - y|^{n+2s}} dx dy,$$

and as a consequence,

$$|\mathcal{D}_1^s(u, \varphi)| \leq \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} |\mathbb{M}^\varepsilon(z_1, z_2)| |(-\Delta)^{\frac{s_1}{2}} u(z_1)| |(-\Delta)^{\frac{s_2-\varepsilon}{2}} \varphi(z_2)| dz_1 dz_2.$$

We observe that the upper bound of $|\mathbb{M}^\varepsilon(z_1, z_2)|$ is exactly the quantity that appear in [19, Lemma 3.5], and so for $\sigma > 0$ small enough, the inequality

$$|\mathcal{D}_1^s(u, \varphi)| \lesssim \int_{\mathbb{R}^n} I^{\sigma-\varepsilon} |(-\Delta)^{\frac{s_1}{2}} u|(x) |(-\Delta)^{\frac{s_2-\varepsilon}{2}} \varphi|(x) dx.$$

follows from [19, Theorem 3.1]. The other estimate follows the same way by reversing the role of u and φ , and s_1 and s_2 . $\square$

Next we estimate $\mathcal{D}_2^{s_1, s_2}(u, \varphi) = \langle A_D \mathcal{L}_1^s u, \varphi \rangle - \langle \tilde{\mathcal{L}}_{A_D}^{s_1, s_2} u, \varphi \rangle$. Recall that the operator $A_D \mathcal{L}_1^s$ is defined as follows:

$$\langle A_D \mathcal{L}_1^s u, \varphi \rangle = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} A_D(z_1) \langle \mathbb{K}_1^0(z_1, z_2) (-\Delta)^{\frac{s_1}{2}} u(z_1), (-\Delta)^{\frac{s_2}{2}} \varphi(z_2) \rangle dz_1 dz_2.$$

Let us obtain a compact form of the action of the operator $A_D \mathcal{L}_1^s$ on vector fields. Denoting $U := (-\Delta)^{\frac{s_1}{2}} u$ and $V := (-\Delta)^{\frac{s_2}{2}} \varphi$, it follows that

$$\begin{aligned} \langle A_D \mathcal{L}_1^s u, \varphi \rangle &= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} A_D(z_1) \langle \mathbb{K}_1^0(z_1, z_2) U(z_1), V(z_2) \rangle dz_1 dz_2 \\ &= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \left\langle \frac{(x - y) \otimes (x - y)}{|x - y|^{n+2(s+1)}} (I^{s_1}(A_D U)(x) - I^{s_1}(A_D U)(y)), I^{s_2} V(x) - I^{s_2} V(y) \right\rangle dx dy. \end{aligned}$$

Observe the following for $\gamma_1 = (n + 2s - 2)(n + 2s)$ and $\gamma_2 = (n + 2s - 2)$:

$$\nabla^2 \left(\frac{1}{|x - y|^{n+2s-2}} \right) = \gamma_1 \frac{(x - y) \otimes (x - y)}{|x - y|^2} \frac{1}{|x - y|^{n+2s}} - \gamma_2 \frac{1}{|x - y|^{n+2s}} \mathbb{I}. \quad (3.9)$$

It then follows that

$$\begin{aligned} \langle A_D \mathcal{L}_1^s u, \varphi \rangle &\stackrel{(3.9)}{=} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{\tilde{\gamma}_2}{|x - y|^{n+2s}} (I^{s_1}(A_D U)(x) - I^{s_1}(A_D U)(y)) \cdot (I^{s_2} V(x) - I^{s_2} V(y)) dx dy \\ &\quad + \tilde{\gamma}_1 \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \left\langle \nabla^2 \left(\frac{1}{|x - y|^{n+2s-2}} \right) (I^{s_1}(A_D U)(x) - I^{s_1}(A_D U)(y)), I^{s_2} V(x) - I^{s_2} V(y) \right\rangle \\ &= \int_{\mathbb{R}^n} \tilde{\gamma}_2 A_D(z) U(z) \cdot V(z) + \tilde{\gamma}_1 \langle \nabla^2 I^2(A_D U)(z), V(z) \rangle dz \\ &= \int_{\mathbb{R}^n} (\tilde{\gamma}_2 A_D(z) U(z) + \tilde{\gamma}_1 \mathcal{R} \otimes \mathcal{R}(A_D U)(z)) \cdot V(z) dz. \end{aligned}$$

The precise value of constants $\tilde{\gamma}_1$ and $\tilde{\gamma}_2$ are computed in [28] and verify that $\tilde{\gamma}_1 \neq \tilde{\gamma}_2$. For the argument to follow, the exact value is not as important, but these are the constants that appear in the regular operator $\tilde{\mathcal{L}}_{A_D}^{s_1, s_2}$. As a consequence, the expression for $\mathcal{D}_2^{s_1, s_2}(u, \varphi)$ simplifies to

$$\begin{aligned}\mathcal{D}_2^{s_1, s_2}(u, \varphi) &= \tilde{\gamma}_1 \int_{\mathbb{R}^n} \langle (\mathcal{R} \otimes \mathcal{R}(A_D(-\Delta)^{\frac{s_1}{2}} u)(z) - A_D \mathcal{R} \otimes \mathcal{R}((-\Delta)^{\frac{s_1}{2}} u)(z)), (-\Delta)^{\frac{s_2}{2}} \varphi(z) \rangle dz \\ &= \tilde{\gamma}_1 \int_{\mathbb{R}^n} \langle [\mathcal{R} \otimes \mathcal{R}, A_D]((-\Delta)^{\frac{s_1}{2}} u)(z), (-\Delta)^{\frac{s_2}{2}} \varphi(z) \rangle dz,\end{aligned}$$

where we used the commutator notation

$$[T, b](f) = T(bf) - bTf.$$

We normalize the constant and assume that $\tilde{\gamma}_1 = 1$. The next proposition estimates $\mathcal{D}_2^{s_1, s_2}(u, \varphi)$.

Proposition 3.2. *Let $s \in (0, 1)$ with $s_1 + s_2 = 2s$ and $\alpha > 0$. For $u \in H^{s_1, p}(\mathbb{R}^n)$ and $\varphi \in C_c^\infty(\mathbb{R}^n)$, let*

$$\mathcal{D}_2^{s_1, s_2}(u, \varphi) = \int_{\mathbb{R}^n} \langle [\mathcal{R} \otimes \mathcal{R}, A_D]((-\Delta)^{\frac{s_1}{2}} u)(z), (-\Delta)^{\frac{s_2}{2}} \varphi(z) \rangle dz.$$

Then there exists $c > 0$ such that (3.7) holds and for any $\varepsilon \in (0, \alpha)$ and for any $A \in \mathcal{A}(\alpha, \Lambda) = \{A : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R} : |A(x, y)| \leq \Lambda, (1.1) \text{ holds}\}$ we have the estimates

$$|\mathcal{D}_2^{s_1, s_2}(u, \varphi)| \leq \int_{\mathbb{R}^n} I^{\alpha-\varepsilon} |(-\Delta)^{\frac{s_1-\varepsilon}{2}} u|(z) |(-\Delta)^{\frac{s_2}{2}} \varphi|(z) dz$$

and

$$|\mathcal{D}_2^{s_1, s_2}(u, \varphi)| \leq \int_{\mathbb{R}^n} I^{\alpha-\varepsilon} |(-\Delta)^{\frac{s_1}{2}} u|(z) |(-\Delta)^{\frac{s_2-\varepsilon}{2}} \varphi|(z) dz$$

for all $u \in H^{s_1, p}(\mathbb{R}^n)$ and $\varphi \in C_c^\infty(\mathbb{R}^n)$.

Proof. By applying an integration by parts, we get

$$\begin{aligned}\mathcal{D}_2^{s_1, s_2}(u, \varphi) &= \int_{\mathbb{R}^n} \langle (-\Delta)^{\frac{\varepsilon}{2}} ([\mathcal{R} \otimes \mathcal{R}, A_D]((-\Delta)^{\frac{s_1}{2}} u))(z), (-\Delta)^{\frac{s_2-\varepsilon}{2}} \varphi(z) \rangle dz \\ &= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \left\langle \frac{[\mathcal{R} \otimes \mathcal{R}, A_D]((-\Delta)^{\frac{s_1}{2}} u)(z) - [\mathcal{R} \otimes \mathcal{R}, A_D]((-\Delta)^{\frac{s_1}{2}} u)(\tilde{z})}{|z - \tilde{z}|^{n+\varepsilon}}, (-\Delta)^{\frac{s_2-\varepsilon}{2}} \varphi(z) \right\rangle d\tilde{z} dz.\end{aligned}$$

Notice that

$$\begin{aligned}[\mathcal{R} \otimes \mathcal{R}, A_D]((-\Delta)^{\frac{s_1}{2}} u)(z) &= \mathcal{R} \otimes \mathcal{R}(A_D(-\Delta)^{\frac{s_1}{2}} u)(z) - A_D \mathcal{R} \otimes \mathcal{R}((-\Delta)^{\frac{s_1}{2}} u)(z) \\ &= \int_{\mathbb{R}^n} \frac{\Omega(y-z)}{|y-z|^n} (A(y, y) - A(z, z)) (-\Delta)^{\frac{s_1}{2}} u(y) dy,\end{aligned}$$

where we are using the notation $\Omega(\xi) = \frac{\xi \otimes \xi}{|\xi|^2}$. We recall that $\frac{\Omega(\xi)}{|\xi|^n}$ is the Calderón–Zygmund kernel for the second order matrix of Riesz transform $\mathcal{R} \otimes \mathcal{R}$, see (3.9). Thus we have

$$\mathcal{D}_2^{s_1, s_2}(u, \varphi) = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \langle \mathbb{W}^\varepsilon(y, z) (-\Delta)^{\frac{s_1}{2}} u(y), (-\Delta)^{\frac{s_2-\varepsilon}{2}} \varphi(z) \rangle dy dz,$$

where

$$\mathbb{W}^\varepsilon(y, z) := \int_{\mathbb{R}^n} \frac{1}{|z - \tilde{z}|^{n+\varepsilon}} \left(\frac{\Omega(y-z)}{|y-z|^n} (A(y, y) - A(z, z)) - \frac{\Omega(y-\tilde{z})}{|y-\tilde{z}|^n} (A(y, y) - A(\tilde{z}, \tilde{z})) \right) d\tilde{z}.$$

We claim that

$$|\mathbb{W}^\varepsilon(y, z)| \leq |y-z|^{\alpha-\varepsilon-n}.$$

Assume the claim is proved for now. We then have

$$\begin{aligned} |\mathcal{D}_2^{s_1, s_2}(u, \varphi)| &\leq \int_{\mathbb{R}^n} \left(\int_{\mathbb{R}^n} |z-y|^{\alpha-\varepsilon-n} |(-\Delta)^{\frac{s_1}{2}} u|(y) dy \right) |(-\Delta)^{\frac{s_2-\varepsilon}{2}} \varphi|(z) dz \\ &= \int_{\mathbb{R}^n} I^{\alpha-\varepsilon} |(-\Delta)^{\frac{s_1}{2}} u|(z) |(-\Delta)^{\frac{s_2-\varepsilon}{2}} \varphi|(z) dz. \end{aligned}$$

Hence, we obtain the second estimate for $\mathcal{D}_2^{s_1, s_2}(u, \varphi)$. By reversing the role of u and φ , and s_1 and s_2 , the first estimate follows the same way.

What remains is to prove the claim. To that end, we divide the domain into three cases.

Case 1: $|z - \tilde{z}| < \frac{1}{2}|y - z|$ or $|z - \tilde{z}| < \frac{1}{2}|y - \tilde{z}|$. We first consider

$$\begin{aligned} &\left| \frac{\Omega(y-z)}{|y-z|^n} (A(y, y) - A(z, z)) - \frac{\Omega(y-\tilde{z})}{|y-\tilde{z}|^n} (A(y, y) - A(\tilde{z}, \tilde{z})) \right| \\ &\leq \left| \left(\frac{\Omega(y-z)}{|y-z|^n} - \frac{\Omega(y-\tilde{z})}{|y-\tilde{z}|^n} \right) (A(y, y) - A(z, z)) \right| + \left| \frac{\Omega(y-\tilde{z})}{|y-\tilde{z}|^n} (A(\tilde{z}, \tilde{z}) - A(z, z)) \right|. \end{aligned}$$

Since in this case we have $|y - z| \asymp |y - \tilde{z}|$, we can use the application of the fundamental theorem of calculus (see [19, Lemma 3.2]) to obtain

$$\begin{aligned} |\mathbb{W}^\varepsilon(y, z)| &\leq \int_{\mathbb{R}^n} \frac{|z - \tilde{z}|}{|z - \tilde{z}|^{n+\varepsilon} |y - z|^{n+1}} |A(y, y) - A(z, z)| d\tilde{z} + \int_{\mathbb{R}^n} \frac{1}{|z - \tilde{z}|^{n+\varepsilon} |y - z|^n} |A(\tilde{z}, \tilde{z}) - A(z, z)| d\tilde{z} \\ &\leq \int_{\mathbb{R}^n} \frac{|z - \tilde{z}|}{|z - \tilde{z}|^{n+\varepsilon} |y - z|^{n+1}} |y - z|^\alpha d\tilde{z} + \int_{\mathbb{R}^n} \frac{1}{|z - \tilde{z}|^{n+\varepsilon} |y - z|^n} |z - \tilde{z}|^\alpha d\tilde{z}, \end{aligned}$$

where the second inequality above follows from the α -Hölder continuous of A . Then we integrate with respect to $\tilde{z}$ and get

$$\begin{aligned} |\mathbb{W}^\varepsilon(y, z)| &\leq \int_{|z-\tilde{z}| \leq |y-z|} \frac{|z - \tilde{z}|^{1-\varepsilon-n}}{|y - z|^{n+1-\alpha}} d\tilde{z} + \int_{|z-\tilde{z}| \leq |y-z|} \frac{|z - \tilde{z}|^{\alpha-\varepsilon-n}}{|y - z|^n} d\tilde{z} \\ &\leq |y - z|^{1-\varepsilon} |y - z|^{\alpha-1-n} + |y - z|^{\alpha-\varepsilon-n} \leq |y - z|^{\alpha-\varepsilon-n}. \end{aligned}$$

Case 2: $|z - \tilde{z}| \geq \frac{1}{2}|y - z|$ and $|z - \tilde{z}| \geq \frac{1}{2}|y - \tilde{z}|$ and $|y - z| < |y - \tilde{z}|$. Since A is α -Hölder continuous, we have

$$\begin{aligned} |\mathbb{W}^\varepsilon(y, z)| &\leq \int_{|z-\tilde{z}| \geq |y-z|} \frac{1}{|z - \tilde{z}|^{n+\varepsilon}} (|y - z|^{\alpha-n} + |y - \tilde{z}|^{\alpha-n}) d\tilde{z} \\ &\leq |y - z|^{\alpha-n} \int_{|z-\tilde{z}| \geq |y-z|} \frac{1}{|z - \tilde{z}|^{n+\varepsilon}} d\tilde{z} \asymp |y - z|^{\alpha-\varepsilon-n}. \end{aligned}$$

Case 3: $|z - \tilde{z}| \geq \frac{1}{2}|y - z|$ and $|z - \tilde{z}| \geq \frac{1}{2}|y - \tilde{z}|$ and $|y - z| \geq |y - \tilde{z}|$. We have

$$\begin{aligned} |\mathbb{W}^\varepsilon(y, z)| &\leq \int_{|z-\tilde{z}| \geq |y-z|} \frac{1}{|z - \tilde{z}|^{n+\varepsilon}} (|y - z|^{\alpha-n} + |y - \tilde{z}|^{\alpha-n}) d\tilde{z} \\ &\leq |y - z|^{-\varepsilon-n} \int_{|y-\tilde{z}| \leq |y-z|} |y - \tilde{z}|^{\alpha-n} d\tilde{z} \asymp |y - z|^{\alpha-\varepsilon-n}. \end{aligned}$$

That completes the proof of the proposition. $\square$

4 The weighted fractional Lamé system

In this section, we prove an optimal regularity result for the system of equations

$$\tilde{\mathcal{L}}_A^{t, 2s-t} u = (-\Delta)^{\frac{2s-t}{2}} f_1 + f_2, \quad (4.1)$$

where $\bar{A}$ is a positive, measurable function that is bounded from below and above by positive constants, and $\bar{\mathcal{L}}_{\bar{A}}^{t, 2s-t}$ is as defined in (1.4), and can be understood as the operator

$$(-\Delta)^{\frac{2s-t}{2}} (\bar{A}(z)[(-\Delta)^{\frac{t}{2}} u(z) + c\mathcal{R} \otimes \mathcal{R}(-\Delta)^{\frac{t}{2}} u(z)]),$$

where after scaling in (1.4), we assume that $c \neq 1$. The following is an a priori regularity estimate that we will use as an iterative device to obtain the optimal regularity result for the (4.1).

Proposition 4.1. *Let $s \in (0, 1)$ and $t \in (0, 2s)$ such that $2s - t < 1$. Suppose that $\bar{A} : \mathbb{R}^n \rightarrow \mathbb{R}$ is a positive, measurable, and bounded from above and below, i.e.,*

$$\Lambda^{-1} \leq \bar{A}(z) \leq \Lambda \quad \text{for a.e. } z \in \mathbb{R}^n.$$

Assume that for some $q \in (1, \infty)$, $u \in H^{t,q}(\mathbb{R}^n, \mathbb{R}^n)$ is a distributional solution to

$$\begin{aligned} & \int_{\mathbb{R}^n} \bar{A}(z) \langle (-\Delta)^{\frac{t}{2}} u(z) + c\mathcal{R} \otimes \mathcal{R}(-\Delta)^{\frac{t}{2}} u(z), (-\Delta)^{\frac{2s-t}{2}} \varphi(z) \rangle dz \\ &= \int_{\mathbb{R}^n} \langle f_1(z), (-\Delta)^{\frac{2s-t}{2}} \varphi(z) \rangle dz + \int_{\mathbb{R}^n} \langle f_2(z), \varphi(z) \rangle dz \quad \text{for all } \varphi \in C_c^\infty(\Omega, \mathbb{R}^n), \end{aligned} \quad (4.2)$$

where $c \neq 1$. Suppose now that $\Omega_1 \subset \subset \Omega_2 \subset \subset \Omega \subset \mathbb{R}^n$. Then:

(a) *There exists $\bar{q}$ such that $\bar{q} > q > \frac{n\bar{q}}{n+(2s-t)\bar{q}} > 1$ such that if $f_1, f_2 \in L^q(\mathbb{R}^n, \mathbb{R}^n) \cap L^{\bar{q}}(\Omega_2, \mathbb{R}^n)$, then*

$$(-\Delta)^{\frac{t}{2}} u \in L^{\bar{q}}(\Omega_1, \mathbb{R}^n)$$

and

$$\|(-\Delta)^{\frac{t}{2}} u\|_{L^{\bar{q}}(\Omega_1)} \leq \sum_{j=1}^2 (\|f_j\|_{L^q(\Omega_2)} + \|f_j\|_{L^q(\mathbb{R}^n)}) + \|(-\Delta)^{\frac{t}{2}} u\|_{L^q(\mathbb{R}^n)}.$$

(b) *For any $p > q$, and $r \in (1, p)$ such that $r > \frac{np}{n+(2s-t)p} > 1$, if $f_1, f_2 \in L^q(\mathbb{R}^n, \mathbb{R}^n) \cap L^p(\Omega_2, \mathbb{R}^n)$, then the L^p norm of $(-\Delta)^{\frac{t}{2}} u$ can be estimated as*

$$\|(-\Delta)^{\frac{t}{2}} u\|_{L^p(\Omega_1)} \leq \sum_{j=1}^2 (\|f_j\|_{L^p(\Omega_2)} + \|f_j\|_{L^q(\mathbb{R}^n)}) + \|(-\Delta)^{\frac{t}{2}} u\|_{L^r(\Omega_2)} + \|(-\Delta)^{\frac{t}{2}} u\|_{L^q(\mathbb{R}^n)}. \quad (4.3)$$

For $c = 0$, part (b) of Proposition 4.1 is precisely [19, Proposition 4.1]. For the case when $c \neq 1$, the proof of the proposition uses arguments that parallel the proof of [19, Proposition 4.1]. Notice also that the estimate for part (a) follows from part (b) after we made sure $\bar{q}$ exists and taking $p = \bar{q}$ and $r = q$. For the existence, given $q > 1$, we can choose

$$\bar{q} \in \left(\frac{nq}{n - (2s - t)q}, \frac{n}{n - (2s - t)} \right).$$

The interval is nonempty because $q > 1$. Below we will sketch the proof of part (b). First, we state and prove the following observation, see also [28].

Lemma 4.2. *Assume $c \neq 1$. Then for any $U : \mathbb{R}^d \rightarrow \mathbb{R}^d$, and $1 < p < \infty$, we have*

$$\|U\|_{L^p(\mathbb{R}^d)} \leq \|U + c(\mathcal{R} \otimes \mathcal{R})U\|_{L^p(\mathbb{R}^d)}. \quad (4.4)$$

Moreover, we have for any open set $\Omega_1 \subset \subset \Omega_2$ and any $\tau \in [0, 1]$ such that $\frac{np}{n+\tau p} > 1$ and $q \in [1, \infty)$,

$$\|U\|_{L^p(\Omega_1)} \leq \|U + c(\mathcal{R} \otimes \mathcal{R})U\|_{L^p(\Omega_2)} + \|U\|_{L^{\frac{np}{n+\tau p}}(\Omega_2)} + \|U\|_{L^q(\mathbb{R}^n)}$$

with the constant depending on $\Omega_1, \Omega_2, \tau, q$.

Proof. The first estimate of lemma is known, see, e.g., [28], but for the convenience of the reader we sketch the argument. After recalling that the Riesz-transform $\mathcal{R}$ is the operator with Fourier symbol $i \frac{\xi}{|\xi|}$, we may take the Fourier transform to obtain

$$\mathcal{F}(U + c(\mathcal{R} \otimes \mathcal{R})U) = C\mathcal{D}(\xi)\mathcal{F}U(\xi),$$

where

$$\mathbb{D}(\xi) = \mathbb{I}_n - c \frac{\xi \otimes \xi}{|\xi|^2}.$$

We observe that $\mathbb{D}(\xi)$ is a symmetric matrix with eigenvalues 1 (eigenspace: $\xi^\perp$ which is $(d-1)$ -dimensional) and $1-c \neq 0$ (eigenspace $\text{span}(\xi)$, whenever $\xi \in \mathbb{R}^n \setminus \{0\}$). In particular, for any $\xi \in \mathbb{R}^n \setminus \{0\}$, $\mathbb{D}^{-1}(\xi)$ exists, and is given by

$$\mathbb{D}^{-1}(\xi) = \mathbb{I}_n + \frac{1}{1-c} \frac{\xi \otimes \xi}{|\xi|^2}.$$

It then follows that $|\mathbb{D}^{-1}(\xi)| \lesssim \frac{2-c}{1-c}$. Then we may write

$$U = \mathcal{F}^{-1}(\mathbb{D}^{-1} \mathcal{F}(U + c(\mathcal{R} \otimes \mathcal{R})U)).$$

The claim about the L^p estimate follows from Mikhlin- or Hörmander multiplier theorem, [20] or [28] for detailed calculation.

For the second inequality, take $\eta_1, \eta_2 \in C_c^\infty(\Omega_2)$, $0 \leq \eta_1, \eta_2 \leq 1$, $\eta_1 \equiv 1$ in Ω_1 , $\eta_2 \equiv 1$ in $\text{supp } \eta_1$. Then we apply inequality (4.4) to $\eta_1 U$ to obtain that

$$\|U\|_{L^p(\Omega_1)} \leq \|\eta_1 U\|_{L^p(\mathbb{R}^n)} \leq \|\eta_1 U + c(\mathcal{R} \otimes \mathcal{R})(\eta_1 U)\|_{L^p(\mathbb{R}^n)}.$$

It then follows that

$$\|U\|_{L^p(\Omega_1)} \leq \|U + c(\mathcal{R} \otimes \mathcal{R})U\|_{L^p(\Omega_2)} + \|[\mathcal{R} \otimes \mathcal{R}, \eta_1](U)\|_{L^p(\mathbb{R}^n)},$$

where here we used the commutator notation $[T, \eta_1](f) := T(\eta_1 f) - \eta_1 T(f)$. To estimate the last term in the previous inequality, we use the identity $\eta_1(1 - \eta_2) \equiv 0$ to write

$$[\mathcal{R} \otimes \mathcal{R}, \eta_1](U) = [\mathcal{R} \otimes \mathcal{R}, \eta_1](\eta_2 U) + \eta_1 \mathcal{R} \otimes \mathcal{R}((1 - \eta_2)U)$$

and therefore

$$\|[\mathcal{R} \otimes \mathcal{R}, \eta_1](U)\|_{L^p(\mathbb{R}^n)} \leq \|[\mathcal{R} \otimes \mathcal{R}, \eta_1](\eta_2 U)\|_{L^p(\mathbb{R}^n)} + \|\eta_1 \mathcal{R} \otimes \mathcal{R}((1 - \eta_2)U)\|_{L^p(\mathbb{R}^n)}.$$

For the first term, in view of commutator estimates, say in [16, Theorem 6.1.] or [14], for any $\tau \in [0, 1]$ denoting by $I^\tau = (-\Delta)^{-\frac{\tau}{2}}$ the Riesz potential, and then using Sobolev inequality (if $\frac{np}{n+\tau p} \in (1, \infty)$)

$$\|[\mathcal{R} \otimes \mathcal{R}, \eta_1](\eta_2 U)\|_{L^p(\mathbb{R}^n)} \leq \|\eta_1\|_{\text{Lip}} \|I^\tau(\eta_2 U)\|_{L^p(\mathbb{R}^n)} \leq \|\eta_2 U\|_{L^{\frac{np}{n+\tau p}}(\mathbb{R}^n)} \lesssim \|U\|_{L^{\frac{np}{n+\tau p}}(\Omega_2)}.$$

Here, as usual,

$$\|\eta_1\|_{\text{Lip}} := \|\eta_1\|_{L^\infty} + \|\nabla \eta_1\|_{L^1}.$$

For the other term, we observe that for any $x \in \mathbb{R}^n$,

$$|\eta_1(x) \mathcal{R} \otimes \mathcal{R}((1 - \eta_2)U)(x)| \lesssim \int_{\mathbb{R}^n} \eta_1(x) \frac{1}{|x-y|^n} (1 - \eta_2(y)) |U(y)| \lesssim \kappa * |U|(x),$$

where

$$\kappa(z) := \frac{1}{|z|^n} \chi_{|z| \geq 1},$$

where the constant in $\gtrsim$ depends on the distance of the support of $(1 - \eta_2)$ to the support of η_1 . We observe that $\kappa \in L^q(\mathbb{R}^n)$ for any $q \in (1, \infty]$. It then follows from Young's convolution inequality that for any $q \in [1, \infty)$ (observing that κ is integrable to any power)

$$\|\eta_1 \mathcal{R} \otimes \mathcal{R}((1 - \eta_2)U)\|_{L^p(\mathbb{R}^n)} \lesssim \|\kappa * |U|\|_{L^\infty(\mathbb{R}^n)} \lesssim \|\kappa\|_{L^{q'}(\mathbb{R}^n)} \|U\|_{L^q(\mathbb{R}^n)} \lesssim \|U\|_{L^q(\mathbb{R}^n)}.$$

Putting the inequalities together we complete the proof of the lemma. $\square$

We are now ready to sketch the proof of Proposition 4.1.

Sketch of the proof of Proposition 4.1. Instead of Ω_1 and Ω_2 we are going to prove the statement for Ω_1 and Ω_4 and some choice of Ω_2, Ω_3 such that $\Omega_1 \subset \subset \Omega_2 \subset \subset \Omega_3 \subset \subset \Omega_4$. From the previous lemma, Lemma 4.2, we have

$$\|(-\Delta)^{\frac{t}{2}} v\|_{L^p(\Omega_1)} \lesssim \|(-\Delta)^{\frac{t}{2}} v + c \mathcal{R} \otimes \mathcal{R}(-\Delta)^{\frac{t}{2}} v\|_{L^p(\Omega_2)} + \|(-\Delta)^{\frac{t}{2}} u\|_{L^{q'}(\Omega_2)} + \|(-\Delta)^{\frac{t}{2}} u\|_{L^q(\mathbb{R}^n)}.$$

By the ellipticity of $\bar{A}$ and duality

$$\begin{aligned} \|(-\Delta)^{\frac{t}{2}} v + c\mathcal{R} \otimes \mathcal{R}(-\Delta)^{\frac{t}{2}} v\|_{L^p(\Omega_2)} &\leq \|\bar{A}((-\Delta)^{\frac{t}{2}} v + c\mathcal{R} \otimes \mathcal{R}(-\Delta)^{\frac{t}{2}} v)\|_{L^p(\Omega_2)} \\ &\leq \sup_{\psi} \int_{\mathbb{R}^n} \langle \bar{A}(x)((-\Delta)^{\frac{t}{2}} v + c\mathcal{R} \otimes \mathcal{R}(-\Delta)^{\frac{t}{2}} v)(x), \psi(x) \rangle dx, \end{aligned}$$

where the supremum is taken over all $\psi \in C_c^\infty(\Omega_2; \mathbb{R}^n)$ with $\|\psi\|_{L^{p'}(\Omega_2)} \leq 1$. To finish the proof of the proposition, it suffices to prove that for any $\psi \in C_c^\infty(\Omega_2; \mathbb{R}^n)$,

$$\begin{aligned} &\int_{\mathbb{R}^n} \langle \bar{A}(x)((-\Delta)^{\frac{t}{2}} v + c\mathcal{R} \otimes \mathcal{R}(-\Delta)^{\frac{t}{2}} v)(x), \psi(x) \rangle dx \\ &\leq (\|f_1\|_{L^p(\Omega_2)} + \|f_2\|_{L^p(\Omega_3)} + \|f_1\|_{L^{q_+}(\mathbb{R}^n)} + \|(-\Delta)^{\frac{t}{2}} v\|_{L^r(\Omega_4)} + \|(-\Delta)^{\frac{t}{2}} v\|_{L^q(\mathbb{R}^n)}) \|\psi\|_{L^{p'}(\Omega_3)}. \end{aligned}$$

To that end, pick $\eta_1, \eta_2 \in C_c^\infty(\Omega_3)$ with $\eta_1 \equiv 1$ in a neighborhood of Ω_2 , and $\eta_2 \equiv 1$ on the support of η_1 . For any $\psi \in C_c^\infty(\Omega_2; \mathbb{R}^n)$, write $\psi = (-\Delta)^{\frac{2s-t}{2}}(I^{2s-t}\psi)$ and

$$\begin{aligned} \psi &= (-\Delta)^{\frac{2s-t}{2}}(\eta_1 I^{2s-t}\psi) + (-\Delta)^{\frac{2s-t}{2}}((1-\eta_1)I^{2s-t}\psi) \\ &= (-\Delta)^{\frac{2s-t}{2}}(\eta_1 I^{2s-t}\psi) + \eta_2(-\Delta)^{\frac{2s-t}{2}}((1-\eta_1)I^{2s-t}\psi) + (1-\eta_2)(-\Delta)^{\frac{2s-t}{2}}((1-\eta_1)I^{2s-t}\psi). \end{aligned} \quad (4.5)$$

Then we have

$$\begin{aligned} &\int_{\mathbb{R}^n} \langle \bar{A}((-\Delta)^{\frac{t}{2}} v + c\mathcal{R} \otimes \mathcal{R}(-\Delta)^{\frac{t}{2}} v)(x), \psi(x) \rangle dx \\ &= \int_{\mathbb{R}^n} \langle \bar{A}((-\Delta)^{\frac{t}{2}} v + c\mathcal{R} \otimes \mathcal{R}(-\Delta)^{\frac{t}{2}} v)(x), (-\Delta)^{\frac{2s-t}{2}}(\eta_1 I^{2s-t}\psi)(x) \rangle dx \\ &\quad + \int_{\mathbb{R}^n} \langle \bar{A}((-\Delta)^{\frac{t}{2}} v + c\mathcal{R} \otimes \mathcal{R}(-\Delta)^{\frac{t}{2}} v)(x), \eta_2(x)(-\Delta)^{\frac{2s-t}{2}}((1-\eta_1)I^{2s-t}\psi)(x) \rangle dx \\ &\quad + \int_{\mathbb{R}^n} \langle \bar{A}((-\Delta)^{\frac{t}{2}} v + c\mathcal{R} \otimes \mathcal{R}(-\Delta)^{\frac{t}{2}} v)(x), (1-\eta_2)(-\Delta)^{\frac{2s-t}{2}}((1-\eta_1)I^{2s-t}\psi)(x) \rangle dx \\ &= I + II + III. \end{aligned}$$

We estimate each of the above integrals. To estimate I , we set $\varphi := \eta_1 I^{2s-t}\psi \in C_c^\infty(\Omega_3)$. Then we notice that φ is an admissible test function in equation (4.2) and, thus, we can use it in the equation

$$\begin{aligned} I &= \int_{\mathbb{R}^n} \langle f_1(z), (-\Delta)^{\frac{2s-t}{2}} \varphi(z) \rangle dz + \int_{\mathbb{R}^n} \langle f_2(z), \varphi(z) \rangle dz \\ &= \int_{\mathbb{R}^n} \langle f_1(z), \psi(z) \rangle dz + \int_{\mathbb{R}^n} \langle f_2(z), \varphi(z) \rangle dz - \int_{\mathbb{R}^n} \langle f_1(z), (-\Delta)^{\frac{2s-t}{2}}(1-\eta_1)I^{2s-t}\psi \rangle dz, \end{aligned}$$

where the latter is obtained using the decomposition (4.5). Now the first two terms can be estimated as follows:

$$\begin{aligned} \int_{\mathbb{R}^n} \langle f_1(z), \psi(z) \rangle dz + \int_{\mathbb{R}^n} \langle f_2(z), \varphi(z) \rangle dz &\leq \|f_1\|_{L^p(\Omega_2)} \|\psi\|_{L^{p'}(\Omega_2)} + \|f_2\|_{L^p(\Omega_3)} \|\varphi\|_{L^{p'}(\Omega_3)} \\ &\leq (\|f_1\|_{L^p(\Omega_2)} + \|f_2\|_{L^p(\Omega_3)}) \|\psi\|_{L^{p'}(\Omega_2)}, \end{aligned}$$

where we used Sobolev inequalities (2.2) and (2.3) and the fact that ψ is compactly supported to estimate

$$\|\varphi\|_{L^{p'}(\Omega_3)} \leq \|I^{2s-t}\psi\|_{L^{p'}(\Omega_3)} \leq \|\psi\|_{L^{p'}(\Omega_2)}.$$

To estimate the last term of I , first we write it as

$$\begin{aligned} \int_{\mathbb{R}^n} \langle f_1(z), (-\Delta)^{\frac{2s-t}{2}}(1-\eta_1)I^{2s-t}\psi \rangle dz &= \int_{\mathbb{R}^n} \langle f_1(z), \eta_2(-\Delta)^{\frac{2s-t}{2}}(1-\eta_1)I^{2s-t}\psi \rangle dz \\ &\quad + \int_{\mathbb{R}^n} \langle f_1(z), (1-\eta_2)(-\Delta)^{\frac{2s-t}{2}}(1-\eta_1)I^{2s-t}\psi \rangle dz. \end{aligned}$$

Then while application of (2.5) of Lemma 2.2 (or [19, Proposition 2.4 part b)]) yields

$$\begin{aligned} \int_{\mathbb{R}^n} \langle f_1(z), \eta_2(-\Delta)^{\frac{2s-t}{2}} ((1-\eta_1)I^{2s-t}\psi) \rangle dz &\leq \|f_1\|_{L^p(\Omega_3)} \|(-\Delta)^{\frac{2s-t}{2}} (1-\eta_1)I^{2s-t}\psi\|_{L^{p'}(\Omega_3)} \\ &\lesssim \|f_1\|_{L^p(\Omega_3)} \|\psi\|_{L^{p'}(\Omega_2)} \end{aligned}$$

and application of (2.4) of Lemma 2.2 which holds for any $r > 1$ implies that

$$\begin{aligned} \int_{\mathbb{R}^n} \langle f_1(z), (1-\eta_2)(-\Delta)^{\frac{2s-t}{2}} (1-\eta_1)I^{2s-t}\psi \rangle dz &\leq \|f_1\|_{L^q(\mathbb{R}^n)} \|(1-\eta_2)(-\Delta)^{\frac{2s-t}{2}} (1-\eta_1)I^{2s-t}\psi\|_{L^{q'}(\mathbb{R}^n)} \\ &\lesssim \|f_1\|_{L^q(\mathbb{R}^n)} \|\psi\|_{L^{p'}(\Omega_2)}. \end{aligned}$$

That finishes the estimate for I . To estimate II , we again apply (2.5) of Lemma 2.2 to obtain

$$\begin{aligned} II &\leq \int_{\mathbb{R}^n} |\bar{A}((-\Delta)^{\frac{t}{2}}v + c\mathcal{R} \otimes \mathcal{R}(-\Delta)^{\frac{t}{2}}v)(x)| |\eta_2(x)(-\Delta)^{\frac{2s-t}{2}} ((1-\eta_1)I^{2s-t}\psi)(x)| dx \\ &\leq \|\bar{A}((-\Delta)^{\frac{t}{2}}v + c\mathcal{R} \otimes \mathcal{R}(-\Delta)^{\frac{t}{2}}v)\|_{L^r(\Omega_3)} \|(-\Delta)^{\frac{2s-t}{2}} ((1-\eta_1)I^{2s-t}\psi)\|_{L^{r'}(\Omega_3)} \\ &\lesssim \|(-\Delta)^{\frac{t}{2}}v\|_{L^r(\Omega_4)} \|\psi\|_{L^{p'}(\Omega_3)}. \end{aligned}$$

Finally, the estimate III follows from (2.4) of Lemma 2.2 as

$$\begin{aligned} III &\leq \int_{\mathbb{R}^n} |\bar{A}((-\Delta)^{\frac{t}{2}}v + c\mathcal{R} \otimes \mathcal{R}(-\Delta)^{\frac{t}{2}}v)(x)| |(1-\eta_2)(x)(-\Delta)^{\frac{2s-t}{2}} ((1-\eta_1)I^{2s-t}\psi)(x)| dx \\ &\lesssim \|(-\Delta)^{\frac{t}{2}}v\|_{L^q(\Omega_4)} \|\psi\|_{L^{p'}(\Omega_3)}. \end{aligned}$$

That concludes the proof of the proposition. $\square$

We are now ready to state and prove the optimal regularity result for the weighted fractional Lamé equation given in (4.1). The result follows from Proposition 4.1 by iterating the result on successive subdomains. We sketch its proof below.

Theorem 4.3. *Let $s \in (0, 1)$, $t \in (0, 2s)$ such that $2s - t < 1$. Assume that for some $q \in (1, \infty)$, $(-\Delta)^{\frac{t}{2}}u \in L^q(\mathbb{R}^n)$ is a distributional solution to*

$$\begin{aligned} &\int_{\mathbb{R}^n} \langle \bar{A}(z)((-\Delta)^{\frac{t}{2}}u + c\mathcal{R} \otimes \mathcal{R}(-\Delta)^{\frac{t}{2}}u)(z), (-\Delta)^{\frac{2s-t}{2}}\varphi(z) \rangle dz \\ &= \int_{\mathbb{R}^n} \langle f_1(z), (-\Delta)^{\frac{2s-t}{2}}\varphi(z) \rangle dz + \int_{\mathbb{R}^n} \langle f_2(z), \varphi(z) \rangle dz \quad \text{for all } \varphi \in C_c^\infty(\Omega). \end{aligned}$$

Here $\bar{A} : \mathbb{R}^n \rightarrow \mathbb{R}$ is a positive, measurable, and bounded from above and below, i.e.,

$$\Lambda^{-1} \leq \bar{A}(z) \leq \Lambda \quad \text{for a.e. } x \in \mathbb{R}^n.$$

Then for any $\Omega' \subset\subset \Omega \subset\subset \mathbb{R}^n$, $p \in (1, \infty)$, if $f_1, f_2 \in L^q(\mathbb{R}^n) \cap L^p(\Omega)$, then $(-\Delta)^{\frac{t}{2}}u \in L^p(\Omega')$ with

$$\|(-\Delta)^{\frac{t}{2}}u\|_{L^p(\Omega')} \lesssim \|f_1\|_{L^p(\Omega)} + \|f_2\|_{L^p(\Omega)} + \|f_1\|_{L^q(\mathbb{R}^n)} + \|(-\Delta)^{\frac{t}{2}}u\|_{L^q(\mathbb{R}^n)}.$$

If, in addition, $\bar{A}$ is γ -Hölder continuous uniformly, that is,

$$\sup_{x, y \in \mathbb{R}^n} \frac{|\bar{A}(x) - \bar{A}(y)|}{|x - y|^\gamma} \leq \Lambda,$$

then for any $\beta \in (0, \min\{\gamma, 2s - t\})$ and any $\Omega' \subset\subset \Omega \subset\subset \mathbb{R}^n$,

$$\|(-\Delta)^{\frac{t+\beta}{2}}u\|_{L^p(\Omega')} \lesssim \|(-\Delta)^{\frac{t}{2}}u\|_{L^q(\mathbb{R}^n)} + \|(-\Delta)^{\frac{\beta}{2}}f_1\|_{L^q(\mathbb{R}^n)} + \|f_2\|_{L^q(\Omega)}.$$

Proof. Suppose that $p > q$, and $2s - t < 1$. We consider a sequence of pairs (Ω_i, p_i) for $i = 1, 2, \dots, L$ such that

$$\Omega_1 = \Omega', \quad p_1 = p, \quad \Omega_i \subset \subset \Omega_{i+1},$$

and

$$p_{i+1} \in [q, p_i] \quad \text{such that } p_{i+1} > \frac{np_i}{n + (2s - t)p} > 1$$

for some L and $p_{L+1} = q$, $p_L = \bar{q}$, where $\bar{q}$ is obtained in part (a) of Proposition 4.1. A finite L depending on t, s, p, q exists. We then apply part (b) of Proposition 4.1 to obtain the inequality that

$$\|(-\Delta)^{\frac{t}{2}} u\|_{L^{p_i}(\Omega_i)} \leq \sum_{j=1}^2 (\|f_j\|_{L^{p_i}(\Omega_{i+1})} + \|f_j\|_{L^q(\mathbb{R}^n)}) + \|(-\Delta)^{\frac{t}{2}} u\|_{L^{p_{i+1}}(\Omega_{i+1})} + \|(-\Delta)^{\frac{t}{2}} u\|_{L^q(\mathbb{R}^n)}.$$

We now iterate to get the desired inequality.

The second part of the theorem can be proved in exactly the same way as [19, Proposition 4.2]. $\square$

5 The regularity theorem: Proof of Theorem 1.1

Theorem 1.1 will be proved by an iteration argument that is explained in detail in [19]. In short, it follows the following steps. First, we obtain a localized small incremental improvement for a solution to a globally posed problem. Second, via a cutoff argument, extend the solution with locally improved regularity to be globally defined and also at the same time solve a globally posed problem. This extension is accompanied by essential controlled estimates. We now iterate and get a localized small improved regularity further increasing the regularity of the solution, and so on. The localizing estimate can be done in exactly the same way as [19, Theorem 5.1]. The only component missing is the “small localized improvement” that replaces [19, Theorem 6.1]. In the remaining, we will only prove this missing regularity result and refer the execution of the iterative argument to [19].

Theorem 5.1. Fix $s \in (0, 1)$, $t \in [s, 2s)$, $t < 1$. For given $\alpha \in (0, 1)$, $\lambda, \Lambda > 0$, let $A \in \mathcal{A}(\alpha, \lambda, \Lambda)$. Suppose also that for any $2 \leq p < \infty$, $u \in H^{s,2}(\mathbb{R}^n, \mathbb{R}^n) \cap H^{t,p}(\mathbb{R}^n, \mathbb{R}^n) \cap H^{t,2}(\mathbb{R}^n, \mathbb{R}^n)$ with $\text{supp } u \subset \Omega \subset \subset \mathbb{R}^n$ is a solution to

$$\langle \mathcal{L}_A^s u, \varphi \rangle = \int_{\mathbb{R}^n} \langle f_1, (-\Delta)^{\frac{2s-t}{2}} \varphi \rangle dz + \int_{\mathbb{R}^n} \langle f_2, \varphi \rangle dz \quad \text{for all } \varphi \in C_c^\infty(\mathbb{R}^n). \quad (5.1)$$

Then there exists $\bar{\varepsilon} > 0$ such that if $r \in [p, p + \bar{\varepsilon})$ and $f_1, f_2 \in L^r(\mathbb{R}^n) \cap L^p(\mathbb{R}^n)$, then

$$\|(-\Delta)^{\frac{t}{2}} u\|_{L^r(\Omega)} \leq \sum_{i=1}^2 \|f_i\|_{L^r(\mathbb{R}^n)} + \|f_i\|_{L^p(\mathbb{R}^n)} + \|(-\Delta)^{\frac{t}{2}} u\|_{L^p(\mathbb{R}^n)}.$$

In addition, if $\beta \in [0, \bar{\varepsilon}]$, $(-\Delta)^{\frac{\beta}{2}} f_1 \in L^p(\mathbb{R}^n)$, and $f_1, f_2 \in L^p(\mathbb{R}^n)$, then $(-\Delta)^{\frac{t+\beta}{2}} u \in L_{\text{loc}}^p(\mathbb{R}^n)$ and

$$\|(-\Delta)^{\frac{t+\beta}{2}} u\|_{L^p(\Omega)} \leq \|(-\Delta)^{\frac{\beta}{2}} f_1\|_{L^p(\mathbb{R}^n)} + \|f_1\|_{L^p(\mathbb{R}^n)} + \|f_2\|_{L^p(\mathbb{R}^n)} + \|(-\Delta)^{\frac{t}{2}} u\|_{L^p(\mathbb{R}^n)}. \quad (5.2)$$

Here, $\bar{\varepsilon} > 0$ is uniform in the following sense: $\bar{\varepsilon}$ depends only on α and the number $\theta \in (0, 1)$ which is such that

$$\theta < s, \quad t, \quad 2s - t < 1 - \theta \quad \text{and} \quad 2 \leq p < \frac{1}{\theta}.$$

Proof. We proceed very similar to [19], by reformulating the system of equations to the weighted fractional Lamé system studied in Proposition 4.1 – up to the commutators introduced in Section 3. Set

$$F[\varphi] := \int_{\mathbb{R}^n} \langle f_1, (-\Delta)^{\frac{2s-t}{2}} \varphi \rangle dz + \int_{\mathbb{R}^n} \langle f_2, \varphi \rangle dz.$$

Then for $t \in [s, 2s)$, if u solves (5.1), then recalling the decomposition (3.7), we have up to a constant multiple

$$\langle \tilde{\mathcal{L}}_{A_D}^{t,2s-t} u, \varphi \rangle = F[\varphi] - \mathcal{D}_1^s(u, \varphi) - \mathcal{D}_2^{t,2s-t}(u, \varphi) \quad \text{for all } \varphi \in C_c(\mathbb{R}^n; \mathbb{R}^n), \quad (5.3)$$

where the linear operator $\tilde{\mathfrak{L}}_{A_D}^{t,2s-t}$ is the weighted fractional Lamé operator introduced in equation (1.4) with $A_D(z) = A(z, z)$ is bounded from below and above by positive constants. The functionals $\mathcal{D}_1^s$ and $\mathcal{D}_2^{s_1, s_2}$ are as defined in Section 3. We now define the two operators

$$T_1[\varphi] = \mathcal{D}_1^s(u, \varphi) \quad \text{and} \quad T_2[\varphi] = \mathcal{D}_2^{s_1, s_2}(u, \varphi),$$

which are linear in $\varphi \in C_c^\infty(\mathbb{R}^n)$. Given θ as in the theorem, we can choose ε sufficiently small so that if we take $\sigma = 8\varepsilon$, we have

$$\frac{np'}{n + \sigma p'} \in (1, \infty) \quad \text{for all } p \in \left(2, \frac{1}{\theta}\right)$$

and that Proposition 3.1 and Proposition 3.2 hold. Applying Proposition 3.1 with this σ and Sobolev inequalities (see Lemma 2.1) we see that for any $\beta \in [0, \varepsilon]$,

$$\begin{aligned} |T_1[\varphi]| &\leq \int_{\mathbb{R}^n} |(-\Delta)^{\frac{t}{2}} u|(x) I^{\sigma-\varepsilon} |(-\Delta)^{\frac{2s-t-\varepsilon}{2}} \varphi|(x) dx \\ &\leq \|(-\Delta)^{\frac{t}{2}} u\|_{L^p} \|(-\Delta)^{\frac{2s-t-\varepsilon}{2}} \varphi\|_{L^{\frac{np'}{n+(\sigma-\varepsilon)p'}}(\mathbb{R}^n)} \\ &\leq \|(-\Delta)^{\frac{t}{2}} u\|_{L^p} \|(-\Delta)^{\frac{2s-t-\varepsilon}{2}} \varphi\|_{L^{\frac{np'}{n+(\sigma-\beta)p'}}(\mathbb{R}^n)}. \end{aligned}$$

As a consequence, we have that for any $\beta \in [0, \varepsilon]$

$$T_1 \in \left(\dot{H}^{2s-t-\beta, \frac{np'}{n+(\sigma-\beta)p'}}(\mathbb{R}^n) \right)^*.$$

By representation of the dual elements [19, Proposition 2.2], there exists $g_\beta^1 \in L^{\frac{np}{n-(\sigma-\beta)p}}(\mathbb{R}^n, \mathbb{R}^n)$ such that

$$T_1[\varphi] = \int_{\mathbb{R}^n} \langle g_\beta^1(x), (-\Delta)^{\frac{2s-t-\beta}{2}} \varphi(x) \rangle dx.$$

Similarly, applying Proposition 3.2 and repeating the above calculation for T_2 , for any $\beta \in [0, \varepsilon]$, and $\varepsilon < \alpha$, we can get from the representation of dual elements that a vector field $g_\beta^2 \in L^{\frac{np}{n-(\sigma-\beta)p}}(\mathbb{R}^n, \mathbb{R}^n)$ such that

$$T_2[\varphi] = \int_{\mathbb{R}^n} \langle g_\beta^2(x), (-\Delta)^{\frac{2s-t-\beta}{2}} \varphi(x) \rangle dx.$$

After denoting $g_\beta := g_\beta^1 + g_\beta^2$, we can now rewrite (5.3) as

$$\langle \tilde{\mathfrak{L}}_{A_D}^{t,2s-t} u, \varphi \rangle = \int_{\mathbb{R}^n} \langle (-\Delta)^{\frac{\beta}{2}} f_1 + g_\beta, (-\Delta)^{\frac{2s-t-\beta}{2}} \varphi \rangle dx + \int_{\mathbb{R}^n} \langle f_2, \varphi \rangle$$

for all $\beta \in [0, \varepsilon]$ and $\varphi \in C_c(\mathbb{R}^n, \mathbb{R}^n)$.

Now if $\beta = 0$, then we may apply Theorem 4.3 to conclude that for any $\Omega \subset \subset \mathbb{R}^n$ and $r \in [p, \frac{np}{n-\sigma p}]$ we have

$$\|(-\Delta)^{\frac{t}{2}} u\|_{L^r(\Omega)} \leq \sum_{i=1}^2 \|f_i\|_{L^r(\mathbb{R}^n)} + \|f_i\|_{L^p(\mathbb{R}^n)} + \|(-\Delta)^{\frac{t}{2}} u\|_{L^p(\mathbb{R}^n)}.$$

We notice that there exists an $\bar{\varepsilon} > 0$ such that $\frac{np}{n-\sigma p} \geq p + \bar{\varepsilon}$ for all $p \in [0, \frac{2}{\theta}]$.

Also, if $\beta \in (0, \varepsilon)$, since A_D is α -Hölder continuous uniformly, we may apply the second part of Theorem 4.3 to obtain (5.2). $\square$

Funding: The authors acknowledge funding as follows: Tadele Mengesha: National Science Foundation (NSF), grant no. DMS-2206252; Adisak Seesanea: Thammasat University Research Unit in Gait Analysis and Intelligent Technology (GaitTech); Armin Schikorra: National Science Foundation (NSF), NSF Career DMS-2044898; Sasikarn Yeepo: Thammasat Postdoctoral Fellowship. The research that led to this work was partially carried out while Armin Schikorra was visiting Chulalongkorn University. Armin Schikorra is a Humboldt fellow.

References

- [1] P. Auscher, S. Bortz, M. Egert and O. Saari, Nonlocal self-improving properties: A functional analytic approach, *Tunis. J. Math.* **1** (2019), no. 2, 151–183.
- [2] U. Biccari, M. Warma and E. Zuazua, Local elliptic regularity for the Dirichlet fractional Laplacian, *Adv. Nonlinear Stud.* **17** (2017), no. 2, 387–409.
- [3] S.-S. Byun and K. Kim, L^q estimates for nonlocal p -Laplacian type equations with BMO kernel coefficients in divergence form, preprint (2023), <https://arxiv.org/abs/2303.08517>.
- [4] S.-S. Byun, K. Kim and D. Kumar, Regularity results for a class of nonlocal double phase equations with VMO coefficients, *Publ. Mat.* **68** (2024), no. 2, 507–544.
- [5] M. Cozzi, Interior regularity of solutions of non-local equations in Sobolev and Nikol'skii spaces, *Ann. Mat. Pura Appl. (4)* **196** (2017), no. 2, 555–578.
- [6] E. Di Nezza, G. Palatucci and E. Valdinoci, Hitchhiker's guide to the fractional Sobolev spaces, *Bull. Sci. Math.* **136** (2012), no. 5, 521–573.
- [7] H. Dong and D. Kim, On L_p -estimates for a class of non-local elliptic equations, *J. Funct. Anal.* **262** (2012), no. 3, 1166–1199.
- [8] Q. Du, M. Gunzburger, R. B. Lehoucq and K. Zhou, Analysis of the volume-constrained peridynamic Navier equation of linear elasticity, *J. Elasticity* **113** (2013), no. 2, 193–217.
- [9] M. M. Fall, Regularity results for nonlocal equations and applications, *Calc. Var. Partial Differential Equations* **59** (2020), no. 5, Paper No. 181.
- [10] M. M. Fall, T. Mengesha, A. Schikorra and S. Yeepo, Calderón–Zygmund theory for non-convolution type nonlocal equations with continuous coefficient, *Partial Differ. Equ. Appl.* **3** (2022), no. 2, Paper No. 24.
- [11] N. Garofalo, Fractional thoughts, in: *New Developments in the Analysis of Nonlocal Operators*, Contemp. Math. 723, American Mathematical Society, Providence (2019), 1–135.
- [12] G. Grubb, Fractional-order operators: Boundary problems, heat equations, in: *Mathematical Analysis and Applications—Plenary Lectures*, Springer Proc. Math. Stat. 262, Springer, Cham (2018), 51–81.
- [13] G. Grubb, Regularity in L_p Sobolev spaces of solutions to fractional heat equations, *J. Funct. Anal.* **274** (2018), no. 9, 2634–2660.
- [14] J. Ingmanns, Estimates for commutators of fractional differential operators via harmonic extension, preprint (2020), <https://arxiv.org/abs/2012.12072>.
- [15] T. Kuusi, G. Mingione and Y. Sire, Nonlocal self-improving properties, *Anal. PDE* **8** (2015), no. 1, 57–114.
- [16] E. Lenzmann and A. Schikorra, Sharp commutator estimates via harmonic extensions, *Nonlinear Anal.* **193** (2020), Article ID 111375.
- [17] W. McLean, *Strongly Elliptic Systems and Boundary Integral Equations*, Cambridge University, Cambridge, 2000.
- [18] T. Mengesha and Q. Du, Nonlocal constrained value problems for a linear peridynamic Navier equation, *J. Elasticity* **116** (2014), no. 1, 27–51.
- [19] T. Mengesha, A. Schikorra and S. Yeepo, Calderón–Zygmund type estimates for nonlocal PDE with Hölder continuous kernel, *Adv. Math.* **383** (2021), Article ID 107692.
- [20] T. Mengesha and J. M. Scott, The solvability of a strongly-coupled nonlocal system of equations, *J. Math. Anal. Appl.* **486** (2020), no. 2, Article ID 123919.
- [21] S. Nowak, $H^{s,p}$ regularity theory for a class of nonlocal elliptic equations, *Nonlinear Anal.* **195** (2020), Article ID 111730.
- [22] S. Nowak, Improved Sobolev regularity for linear nonlocal equations with VMO coefficients, *Math. Ann.* **385** (2023), no. 3–4, 1323–1378.
- [23] S. Nowak, Regularity theory for nonlocal equations with VMO coefficients, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **40** (2023), no. 1, 61–132.
- [24] T. Runst and W. Sickel, *Sobolev Spaces of Fractional Order, Nemytskij Operators, and Nonlinear Partial Differential Equations*, De Gruyter Ser. Nonlinear Anal. Appl. 3, Walter de Gruyter, Berlin, 1996.
- [25] S. G. Samko, *Hypersingular Integrals and Their Applications*, Anal. Methods Spec. Funct. 5, Taylor & Francis, London, 2002.
- [26] A. Schikorra, Regularity of $n/2$ -harmonic maps into spheres, *J. Differential Equations* **252** (2012), no. 2, 1862–1911.
- [27] A. Schikorra, Nonlinear commutators for the fractional p -Laplacian and applications, *Math. Ann.* **366** (2016), no. 1–2, 695–720.
- [28] J. M. Scott, The fractional Lamé–Navier operator: Appearances, properties and applications, preprint (2022), <https://arxiv.org/abs/2204.12029>.
- [29] J. M. Scott and T. Mengesha, A fractional Korn-type inequality, *Discrete Contin. Dyn. Syst.* **39** (2019), no. 6, 3315–3343.
- [30] J. M. Scott and T. Mengesha, Self-improving inequalities for bounded weak solutions to nonlocal double phase equations, *Commun. Pure Appl. Anal.* **21** (2022), no. 1, 183–212.
- [31] S. A. Silling, Reformulation of elasticity theory for discontinuities and long-range forces, *J. Mech. Phys. Solids* **48** (2000), no. 1, 175–209.
- [32] S. A. Silling, Linearized theory of peridynamic states, *J. Elasticity* **99** (2010), no. 1, 85–111.
- [33] E. M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton Math. Ser. 30, Princeton University, Princeton, 1970.