Research Article

lie Xiao*

A trace problem for associate Morrey potentials

DOI: 10.1515/anona-2016-0069

Received March 27, 2016; accepted August 18, 2016

Abstract: As a solution to the restriction question for associate Morrey potentials (Question 1.1), this paper characterizes a Radon measure μ on \mathbb{R}^n such that the Riesz operator I_{α} maps the associate Morrey spaces $H_1^{p,\kappa} \subset H^{p,\kappa}$ to the μ -induced Morrey spaces $L_{\mu}^{q,\lambda} \subset L_{\mu,\infty}^{q,\lambda}$ continuously. The discovered restriction/trace principle (Theorem 1.2) is brand-new, and its demonstration is non-trivial.

Keywords: Trace, associate Morrey potential, embedding

MSC 2010: 31B15, 42B35, 46E35

1 Introduction

1.1 Description of Question 1.1

Stemming from an extensive investigation of [2, 7, 9, 10, 45], this paper attempts to answer the comingup-next restriction question, i.e., a trace problem for associate Morrey potentials. Namely, the Riesz images $I_{\alpha}H_1^{p,\kappa} \subset I_{\alpha}H^{p,\kappa}$ of the spaces $H_1^{p,\kappa} \subset H^{p,\kappa}$ associated with the Morrey spaces $L_{\infty}^{\frac{p}{p-1},\kappa} \supset L^{\frac{p}{p-1},\kappa}$ that have been proven to be of independent interest in geometric potential analysis (see, e.g., [5]) but also of great value in treating the local behaviour of an appropriate solution to certain elliptic partial differential equation or the incompressible Navier-Stokes system (see, e.g., [11, 14–16, 20–22, 28, 29, 31–35, 38, 39, 42, 43]):

Question 1.1. Let μ be a Radon measure on the Euclidean space \mathbb{R}^n . What ball condition must μ have so that the Riesz mapping $f \mapsto I_{\alpha}f$ is continuous (i.e., the potential $I_{\alpha}f$ is restricted to the support of μ) in the chain

$$H_1^{p,\kappa} \subset H^{p,\kappa} \xrightarrow{I_\alpha} L_\mu^{q,\lambda} \subset L_{\mu,\infty}^{q,\lambda}$$

for the associate Morrey spaces $H_1^{p,\kappa} \subset H^{p,\kappa}$ and the μ -based Morrey spaces $L_\mu^{q,\lambda} \subset L_{\mu,\infty}^{q,\lambda}$?

A resolution of the above question relies on the following fundamentals.

First of all, from a geometric measure theory perspective, a Radon measure on \mathbb{R}^n is defined as a Borel regular measure assigning finite mass to any compact subset of \mathbb{R}^n (cf. [18, p. 5]). For a given Radon measure μ on \mathbb{R}^n and a nonnegative number β , denote by

$$\|\|\mu\|\|_{\beta} \equiv \sup_{(x,r)\in\mathbb{R}^n\times(0,\infty)} r^{-\beta}\mu\big(B(x,r)\big)$$

the β -th order ball variation of μ . Trivially, $\|\|\mu\|\|_0$ is the total variation of μ . Obviously, this variation $\|\|\mu\|\|_{\beta}$ is finite whenever $\beta \in \{1, \ldots, n\}$ and μ is the β -dimensional Hausdorff measure \mathcal{H}^{β} restricted on a smooth manifold $\mathbb{M} \subset \mathbb{R}^n$ with $\mathcal{H}^{\beta}(\mathbb{M}) < \infty$.

^{*}Corresponding author: Jie Xiao: Department of Mathematics and Statistics, Memorial University, St. John's, NL A1C 5S7, Canada, e-mail: jxiao@mun.ca

Next, given $\alpha \in (0, n)$ the Riesz operator I_{α} acting on a Lebesgue ν -measurable function f in \mathbb{R}^n is determined by

$$I_{\alpha}f(x) = \int_{\mathbb{R}^n} |y - x|^{\alpha - n} f(y) \, d\nu(y),$$

whose constant multiple

$$u = \left(\frac{\Gamma(\frac{n-\alpha}{2})}{\pi^{\frac{n}{2}}2^{\alpha}\Gamma(\frac{\alpha}{2})}\right)I_{\alpha}f$$

defines an inverse for a power of the Laplace operator on \mathbb{R}^n (cf. [5, p. 9]), i.e., such an singular integral u solves the $\frac{\alpha}{2}$ -th order Laplace equation $(-\Delta)^{\frac{\alpha}{2}}u = f$ using the Fourier transform.

After that, for $(q, \lambda) \in [1, \infty) \times (0, n]$, let $L_{\mu, \infty}^{q, \lambda}$ be the weak Morrey space of all μ -measurable functions f on \mathbb{R}^n with

$$\|f\|_{L^{q,\lambda}_{\mu,\infty}} = \sup_{(x,r,t) \in \mathbb{R}^n \times (0,\infty) \times (0,\infty)} \left(r^{\lambda-n} t^q \mu \left(\{ y \in B(x,r) : |f(y)| > t \} \right) \right)^{\frac{1}{q}} < \infty,$$

where B(x, r) is the Euclidean open ball with centre x and radius r. Naturally, as a proper subspace of $L_{\mu,\infty}^{q,\lambda}$, the symbol $L_{\mu}^{q,\lambda}$ is employed to represent the Morrey space of all μ -measurable functions f with

$$||f||_{L^{q,\lambda}_{\mu}} = \sup_{(x,r)\in\mathbb{R}^n\times(0,\infty)} \left(r^{\lambda-n}\int\limits_{B(x,r)} |f|^q d\mu\right)^{\frac{1}{q}} < \infty.$$

In particular, $L_{\mu,\infty}^q = L_{\mu,\infty}^{q,n}$ and $L_{\mu}^q = L_{\mu}^{q,n}$. Moreover, if $\mu = \nu$ (the n-dimensional Lebesgue measure on \mathbb{R}^n) and $(q,\lambda) = (p,\kappa) \in [1,\infty) \times (0,n]$, then $L_{\mu,\infty}^{q,\lambda}$ and $L_{\mu}^{q,\lambda}$ will be simply written as $L_{\infty}^{p,\kappa}$ and $L^{p,\kappa}$ respectively, and the endpoint spaces $L_{\infty}^{p,n}$ and $L^{p,n}$ coincide with the weak- L^p space L_{∞}^p and L^p , respectively; but if $q = \infty$, then $L_{\infty}^{\infty,\kappa} = L^{\infty,\kappa} = L^{\infty}$. Furthermore, for $p \in [1,\infty]$ and $p' \equiv p/(p-1)$ with $\infty \equiv 1/(1-1)$ and $1 \equiv \infty/(\infty-1)$, set

$$\begin{split} H_1^{p,\kappa} &= \left\{ \nu - \text{measurable functions } f : \|f\|_{H_1^{p,\kappa}} \equiv \sup_{\|g\|_{L^{p',\kappa}_{\infty}} \le 1} \left| \int\limits_{\mathbb{R}^n} fg \ d\nu \right| < \infty \right\} \equiv [L^{p',\kappa}_{\infty}]^{\star}, \\ H^{p,\kappa} &= \left\{ \nu - \text{measurable functions } f : \|f\|_{H^{p,\kappa}} \equiv \sup_{\|g\|_{L^{p',\kappa}} \le 1} \left| \int\limits_{\mathbb{R}^n} fg \ d\nu \right| < \infty \right\} \equiv [L^{p',\kappa}]^{\star}. \end{split}$$

Naturally, $H_1^{p,\kappa}$ and $H^{p,\kappa}$ are called the associate spaces $[L_\infty^{p',\kappa}]^*$ and $[L^{p',\kappa}]^*$ of the Morrey spaces $L_\infty^{p',\kappa}$ and $L^{p',\kappa}$, respectively (cf. [36]). Especially, one has $H_1^{1,\kappa} = H^{1,\kappa} = L^1$ (for $0 < \kappa \le n$) and $H_1^{\infty,n} \subsetneq H^{\infty,n} = L^\infty$ (due to $L^1 = L^{1,n} \subsetneq L_\infty^{1,n}$, cf. [17]).

Last but not least, note that $x \mapsto |x|^{-\frac{\kappa}{p}}$ is an element of $L^{p,\kappa}$ for $0 \le \kappa < n$ (cf. [46, p. 587]) and that $x \mapsto (1+|x|^{\eta})^{-1}$ belongs to $H^{p,\kappa}$ for $\eta > \frac{\kappa}{p}$ (cf. [24, Example 2.5]). So, the next implication seems to be very natural and motivative (cf. [26, p.85] and [6, Theorem 3.3]):

$$||f||_{H^{p,\kappa}_1} \gtrsim ||f||_{H^{p,\kappa}} \gtrsim ||f||_{L^q} \gtrsim ||f||_{L^q_\infty} \quad \text{ for } q = \frac{pn}{n + (n - \kappa)(p - 1)} \le p.$$

In the above and below, $X \gtrsim Y$ stands for that there exists a constant c > 0 such that $X \ge cY$, $X \gtrsim Y \gtrsim X$ will be simply written as $X \approx Y$, and 1_E represents the characteristic function of a set $E \subseteq \mathbb{R}^n$.

1.2 Answering Question 1.1

Working on the above question plus the last implication one discovers the following trace and embedding principle for the associate Morrey potentials $I_{\alpha}H^{p,\kappa}$ (regarded as the space of tempered distributions having derivatives of order α in the associate Morrey space $H^{p,\kappa}$).

Theorem 1.2. Assume $(\alpha, \beta, \lambda) \in (0, n) \times (0, n] \times (0, n]$.

(i) *I*i

$$1 < q = \frac{\beta + \lambda - n}{n - \alpha} < \infty, \quad 0 < \kappa \le n < \beta + \lambda \le 2n,$$

then $I_{\alpha} \colon H^{1,\kappa} \to L^{q,\lambda}_{\mu}$ is continuous if and only if

$$\|\|\mu\|\|_{\alpha,\beta,\lambda} \equiv \sup_{K \subset \mathbb{R}^n \ compact \ with \ \nu(K) > 0} \frac{\|I_\alpha 1_K\|_{L^{q,\lambda}_\mu}}{\nu(K)} < \infty,$$

while $I_{\alpha} \colon H^{1,\kappa} \to L^{q,\lambda}_{\mu,\infty}$ is continuous if and only if $\|\|\mu\|\|_{\beta} < \infty$.

(ii) If

$$1 < q = \frac{p(\beta + \lambda - n)}{\kappa + p(n - \alpha - \kappa)} \le p < \infty, \quad 0 < \kappa < n < \beta + \lambda \le 2n,$$

then $I_{\alpha}\colon H_{1}^{p,\kappa}\subset H^{p,\kappa}\to L_{\mu}^{q,\lambda}\subset L_{\mu,\infty}^{q,\lambda}$ is continuous if and only if $\|\|\mu\|_{eta}<\infty$.

(iii) If

$$1 < q = \frac{\beta + \lambda - n}{n - \alpha - \kappa} < \infty, \quad 0 < \kappa < n < \beta + \lambda \le 2n,$$

then $I_{\alpha}\colon H^{\infty,\kappa}\to L^{q,\lambda}_{\mu}$ or $I_{\alpha}\colon H^{\infty,\kappa}\to L^{q,\lambda}_{\mu,\infty}$ is continuous if and only if $\|\|\mu\|_{\beta}<\infty$.

Interestingly, as an immediate application of Theorem 1.2, the next corollary completely describes the embedding of the k-th order associate Morrey–Sobolev space into $L_{\mu}^{q,\lambda}$. More precisely, if

$$\nabla^k f = \begin{cases} (-\Delta)^{\frac{k}{2}} f & \text{for } k \text{ even,} \\ \nabla (-\Delta)^{\frac{k-1}{2}} f & \text{for } k \text{ odd,} \end{cases}$$

then

$$|f| \leq I_k(|\nabla^k f|)$$

for all C^{∞} -functions f with compact support in \mathbb{R}^n (see, e.g., [4, p. 393] or [30, p. 19]), and hence applying Theorem 1.2, together with a small modification of the argument used in the proof of [30, p. 71, Theorem] as p = 1, produces the following corollary.

Corollary 1.3. *If* $(k, \beta, \lambda) \in \{1, ..., n-1\} \times (0, n] \times (0, n], 1 \le p \le \infty$,

$$1 < q = \frac{\beta + \lambda - n}{p^{-1}\kappa + n - k - \kappa}, \quad 0 < \kappa < n < \beta + \lambda \le 2n,$$

then

$$\sup_{0<|\nabla^k f|\in H^{p,\kappa}}\|f\|_{L^{q,\lambda}_{\mu}}\||\nabla^k f|\|_{H^{p,\kappa}}^{-1}<\infty\iff \|\|\mu\|_{\beta}<\infty.$$

Through exploiting the local nature of some indicator-like functions, introducing Lemmas 2.1, 3.1 and 4.1, and utilising the dual structure of a local L_{μ}^{q} -space, the non-trivial arguments for (i), (ii) and (iii) of Theorem 1.2 and their crucial Remarks 2.2, 3.2 and 4.2 will be arranged in Sections 2, 3 and 4, respectively. It is believed that the techniques developed in the forthcoming three sections can be applied to investigate a similar trace problem for the following Cordes–Nirenberg potentials (see, e.g., [16, 26, 27, 34]):

$$\left\{I_{\alpha}f: \sup_{x\in\mathbb{R}^n}\int_{\mathbb{D}^n}|f(y)|^p|x-y|^{-\gamma}\,d\nu(y)<\infty\right\},\quad 1\leq p<\infty,\,0<\gamma< n.$$

Nevertheless, this investigation will be the subject of a future article.

2 Demonstration of Theorem 1.2 (i)

2.1 A lemma for Theorem 1.2(i)

To prove Theorem 1.2 (i), one needs the following result which improves [2, Theorem 5.1].

Lemma 2.1. If $0 < \alpha < n$, $0 < \lambda \le \kappa \le n$,

$$1 \leq p < \frac{\kappa}{\alpha}, \quad n - \alpha p < \beta \leq n, \quad 1 \leq q = \frac{p(\beta + \lambda - n)}{\kappa - \alpha p} \leq \frac{pn}{\kappa - \alpha p},$$

then

$$\|\|\mu\|\|_{\beta}<\infty \Leftrightarrow \sup_{0\neq f\in L^{p,\kappa}}\|I_{\alpha}f\|_{L^{q,\lambda}_{\mu,\infty}}\|f\|_{L^{p,\kappa}}^{-1}<\infty.$$

Proof. Suppose

$$\sup_{0\neq f\in L^{p,\kappa}}\|I_{\alpha}f\|_{L^{q,\lambda}_{\mu,\infty}}\|f\|_{L^{p,\kappa}}^{-1}<\infty.$$

For a given ball B(x, r), we compute (cf. [6])

$$\|1_{B(x,r)}\|_{L^{p,\kappa}}\approx r^{\frac{\kappa}{p}},\quad r^{\alpha}\lesssim \inf_{y\in B(x,r)}I_{\alpha}1_{B(x,r)}(y),\quad \mu\big(B(x,r)\big)\lesssim \mu\big(\big\{y\in B(x,r):\ r^{\alpha}\lesssim I_{\alpha}1_{B(x,r)}(y)\big\}\big).$$

Note that $1_{B(x,r)}$ belongs to $L^{p,\kappa}$, so

$$\mu(B(x,r))r^{\alpha q + \lambda - n} \leq \mu(\{y \in B(x,r) : r^{\alpha} \leq I_{\alpha}1_{B(x,r)}(y)\})r^{\alpha q + \lambda - n} \leq \|1_{B(x,r)}\|_{I^{p,x}}^{q} \leq r^{\frac{\kappa q}{p}}.$$

This inequality ensures that $\|\|\mu\|\|_{\beta} < \infty$ due to

$$0 < \beta = (\kappa p^{-1} - \alpha)q + n - \lambda \le n.$$

Next, suppose $\|\mu\|_{\beta} < \infty$. Without loss of generality, one may assume $0 \le f \in L^{p,\kappa}$. For a positive number t and a given ball B(x,r) with $B(x,r)^c = \mathbb{R}^n \setminus B(x,r)$ let $0 \le f \in L^{p,\kappa}$, $f_1 = f1_{B(x,r)}$, $f_2 = f1_{B(x,r)^c}$,

$$F_t = \{ y \in B(x, r) : I_{\alpha}f(y) > t \}, \quad F_{t,1} = \{ y \in B(x, r) : I_{\alpha}f_1(y) > t \}, \quad F_{t,2} = \{ y \in B(x, r) : I_{\alpha}f_2(y) > t \}.$$

Then

$$\mu(F_t) \leq \mu(F_{t/2,1}) + \mu(F_{t/2,2}).$$

It remains to control the last two terms separately.

Step 1: In order to prove the desired estimate

$$\mu(F_{t/2,1}) \leq \|\|\mu\|\|_{\beta} t^{-\frac{p(\kappa+\beta-n)}{\kappa-\alpha p}} r^{n-\kappa} \quad \text{for all } (x,r,t) \in \mathbb{R}^n \times (0,\infty) \times (0,\infty),$$

one utilises the Fubini theorem to obtain

$$\begin{split} t\mu(F_{t,1}) &\leq \int\limits_{F_{t,1}} I_{\alpha}f_{1}(z) \, d\mu(z) \\ &= \int\limits_{F_{t,1}} \left(\int\limits_{\mathbb{R}^{n}} |y-z|^{\alpha-n} f_{1}(y) \, d\nu(y) \right) d\mu(z) \\ &\leq \int\limits_{F_{t,1}} \left(\int\limits_{0}^{\infty} \left(\int\limits_{B(z,s)} f_{1}(y) \, d\nu(y) \right) s^{\alpha-n-1} \, ds \right) d\mu(z) \\ &\leq \int\limits_{0}^{\infty} \left(\int\limits_{F_{t,1}} \int\limits_{B(z,s)} f_{1}(y) \, d\nu(y) \, d\mu(z) \right) s^{\alpha-n-1} \, ds \\ &\leq \int\limits_{0} (\delta) + \int\limits_{\infty} (\delta), \end{split}$$

where $\delta > 0$ will be determined later with

$$\mathsf{J}_{j}(\delta) = \begin{cases} \int\limits_{0}^{\delta} \left(\int\limits_{F_{t,1}} \int\limits_{B(z,s)} f_{1}(y) \, d\nu(y) \, d\mu(z) \right) s^{\alpha-n-1} \, ds & \text{for } j = 0, \\ \int\limits_{0}^{\infty} \left(\int\limits_{F_{t,1}} \int\limits_{B(z,s)} f_{1}(y) \, d\nu(y) \, d\mu(z) \right) s^{\alpha-n-1} \, ds & \text{for } j = \infty. \end{cases}$$

First, if p > 1, then an application of the Hölder inequality with p' = p/(p-1) plus $f \in L^{p,\kappa}$ derives

$$\begin{split} \int_{F_{t,1}} \int_{B(z,s)} f_{1}(y) \, d\nu(y) \, d\mu(z) &\leq \int_{\mathbb{R}^{n}} \mu(B(y,s) \cap F_{t,1}) (1_{B(x,r)} f)(y) \, d\nu(y) \\ &\leq \left(\int_{B(x,r)} (f(y))^{p} \, d\nu(y) \right)^{\frac{1}{p}} \left(\int_{B(x,r)} \left(\mu(B(y,s) \cap F_{t,1}) \right)^{p'} \, d\nu(y) \right)^{\frac{1}{p'}} \\ &\leq r^{\frac{n-\kappa}{p}} \|f\|_{L^{p,\kappa}} \left(\int_{B(x,r)} \left(\mu(B(y,s) \cap F_{t,1}) \right) \left(\mu(B(y,s) \cap F_{t,1}) \right)^{p'-1} \, d\nu(y) \right)^{\frac{1}{p'}} \\ &\leq r^{\frac{n-\kappa}{p}} \|f\|_{L^{p,\kappa}} s^{\frac{\beta}{p}} \|\mu\|_{\beta}^{\frac{1}{p}} \left(\int_{B(x,r)} \mu(B(y,s) \cap F_{t,1}) \, d\nu(y) \right)^{\frac{1}{p'}} \\ &\leq r^{\frac{n-\kappa}{p}} \|f\|_{L^{p,\kappa}} s^{\frac{\beta}{p}} \|\mu\|_{\beta}^{\frac{1}{p}} \left(\int_{B(x,r)} 1_{B(y,s)}(z) \, d\nu(z) \right)^{\frac{1}{p'}} \left(\mu(F_{t,1}) \right)^{\frac{1}{p'}} \\ &\leq r^{\frac{n-\kappa}{p}} \|f\|_{L^{p,\kappa}} s^{\frac{\beta}{p}+\frac{n}{p'}} \|\mu\|_{\beta}^{\frac{1}{p}} \left(\mu(F_{t,1}) \right)^{\frac{1}{p'}}. \end{split}$$

At the same time, if p = 1, then a slight modification of the above estimation gives

$$\int_{F_{t,1}} \int_{B(z,s)} f_1(y) \, d\nu(y) \, d\mu(z) \leq r^{n-\kappa} \|f\|_{L^{1,\kappa}} s^{\beta} \|\|\mu\|_{\beta}.$$

Second, for $J_0(\delta)$ one uses the foregoing inequalities to gain that if p > 1, then

$$\begin{split} \mathsf{J}_{0}(\delta) &= \int\limits_{0}^{\delta} \bigg(\int\limits_{F_{t,1}} \int\limits_{B(z,s)} f_{1}(y) \, d\nu(y) \, d\mu(z) \bigg) s^{\alpha-n-1} \, ds \\ &\lesssim r^{\frac{n-\kappa}{p}} \|f\|_{L^{p,\kappa}} \|\|\mu\|_{\beta}^{\frac{1}{p}} \big(\mu(F_{t,1})\big)^{\frac{1}{p'}} \int\limits_{0}^{\delta} s^{\frac{\beta}{p} + \frac{n}{p'} + \alpha - n - 1} \, ds \\ &\approx r^{\frac{n-\kappa}{p}} \delta^{\alpha + \frac{\beta-n}{p}} \|f\|_{L^{p,\kappa}} \|\|\mu\|_{\beta}^{\frac{1}{p}} \big(\mu(F_{t,1})\big)^{\frac{1}{p'}}, \end{split}$$

and if p = 1, then

$$\mathsf{J}_0(\delta) \lesssim r^{n-\kappa} \delta^{\alpha+\beta-n} \|f\|_{L^{1,\kappa}} \|\|\mu\|_{\beta}.$$

Of course, the condition $\beta > n - \alpha p$ is needed for the treatment of $J_0(\delta)$. Meanwhile, for $J_{\infty}(\delta)$ one utilises $f \in L^{p,\kappa}$ to deduce

$$\mathsf{J}_{\infty}(\delta) \leq \|f\|_{L^{p,\kappa}} \int\limits_{s}^{\infty} s^{\alpha-\frac{\kappa}{p}-1} \mu(F_{t,1}) \ ds \approx \delta^{\alpha-\frac{\kappa}{p}} \mu(F_{t,1}) \|f\|_{L^{p,\kappa}}.$$

Here the condition $\alpha p < \kappa$ has been used.

Third, putting the estimates for $J_0(\delta)$ and $J_{\infty}(\delta)$ together gives

$$t\mu(F_{t,1}) \lesssim \begin{cases} r^{\frac{n-\kappa}{p}} \delta^{\alpha+\frac{\beta-n}{p}} \|f\|_{L^{p,\kappa}} \|\mu\|_{\beta}^{\frac{1}{p}} \left(\mu(F_{t,1})\right)^{\frac{1}{p'}} + \delta^{\alpha-\frac{\kappa}{p}} \mu(F_{t,1}) \|f\|_{L^{p,\kappa}} & \text{for } p > 1, \\ r^{n-\kappa} \delta^{\alpha+\beta-n} \|f\|_{L^{1,\kappa}} \|\mu\|_{\beta} + \delta^{\alpha-\kappa} \mu(F_{t,1}) \|f\|_{L^{1,\kappa}} & \text{for } p = 1. \end{cases}$$

Now, choosing

$$\delta = \left(\frac{\mu(F_{t,1})}{r^{n-\kappa} \|\|\mu\|\|_{\beta}}\right)^{\frac{1}{\beta+\kappa-n}}$$

gives

$$\mu(F_{t,1}) \lesssim \|\|\mu\|\|_{\beta} r^{n-\kappa} (t^{-1} \|f\|_{L^{p,\kappa}})^{\frac{p(n-\beta-\kappa)}{\alpha p-\kappa}}.$$

Step 2: In order to prove the estimate

$$\mu(F_{t/2,2}) \leq \|\|\mu\|\|_{\beta} r^{n-\kappa} (t^{-1} \|f\|_{L^{p,\kappa}})^{\frac{p(\kappa+\beta-n)}{\kappa-ap}} \quad \text{for all } (x,r,t) \in \mathbb{R}^n \times (0,\infty) \times (0,\infty),$$

one can borrow the idea used in Step 1 to get

$$t\mu(F_{t,2}) \leq \mathsf{K}_0(\delta) + \mathsf{K}_{\infty}(\delta),$$

where

$$\mathsf{K}_{j}(\delta) = \begin{cases} \int\limits_{0}^{\delta} \left(\int\limits_{F_{t,2}} \int\limits_{B(z,s)} f_{2}(y) \, d\nu(y) \, d\mu(z) \right) s^{\alpha-n-1} \, ds & \text{for } j = 0, \\ \int\limits_{0}^{\infty} \left(\int\limits_{F_{t,2}} \int\limits_{B(z,s)} f_{2}(y) \, d\nu(y) \, d\mu(z) \right) s^{\alpha-n-1} \, ds & \text{for } j = \infty. \end{cases}$$

Notice that

$$y \in B(x, r)^c$$
 and $z \in B(y, s) \cap F_{t,2} \implies y \in B(x, r + s)$.

So, if p > 1, then

$$\begin{split} \int\limits_{F_{t,2}} \int\limits_{B(z,s)} f_2(y) \, d\nu(y) \, d\mu(z) & \leq \int\limits_{B(x,r+s)} f(y) \Biggl(\int\limits_{\mathbb{R}^n} \mathbf{1}_{B(y,s)}(z) \mathbf{1}_{F_{t,2}}(z) \, d\mu(z) \Biggr) \, d\nu(y) \\ & \leq \Biggl(\int\limits_{B(x,r+s)} (f(y))^p \, d\nu(y) \Biggr)^{\frac{1}{p}} \Biggl(\int\limits_{B(x,r+s)} \Biggl(\int\limits_{\mathbb{R}^n} \mathbf{1}_{B(y,s)}(z) \mathbf{1}_{F_{t,2}}(z) \, d\mu(z) \Biggr)^{p'} \, d\nu(y) \Biggr)^{\frac{1}{p'}} \\ & \leq \|f\|_{L^{p,\kappa}} (r+s)^{\frac{n-\kappa}{p}} \bigl(\mu(F_{t,2}) \bigr)^{\frac{1}{p'}} s^{n+\frac{\beta-n}{p}} \| \mu \|_{\frac{p}{p}}^{\frac{1}{p}}, \end{split}$$

and if p = 1, then

$$\int\limits_{F_{t,2}} \int\limits_{B(z,s)} f_2(y) \, d\nu(y) \, d\mu(z) \lesssim \|f\|_{L^{1,\kappa}} (r+s)^{n-\kappa} s^{\beta} \|\|\mu\|_{\beta},$$

whence

$$\mathsf{K}_{0}(\delta) \lesssim \begin{cases} \|f\|_{L^{p,\kappa}} \delta^{\alpha + \frac{\beta - n}{p}} (\delta + r)^{\frac{n - \kappa}{p}} (\mu(F_{t,2}))^{\frac{1}{p'}} \|\mu\|_{\beta}^{\frac{1}{p}} & \text{for } p > 1, \\ \|f\|_{L^{1,\kappa}} \delta^{\alpha + \beta - n} (\delta + r)^{n - \kappa} \|\mu\|_{\beta} & \text{for } p = 1. \end{cases}$$

Similarly, one has

$$\mathsf{K}_{\infty}(\delta) \lesssim \|f\|_{L^{p,\kappa}} \int_{\delta}^{\infty} s^{\alpha - \frac{\kappa}{p} - 1} \mu(F_{t,2}) \, ds \approx \delta^{\alpha - \frac{\kappa}{p}} \mu(F_{t,2}) \|f\|_{L^{p,\kappa}}.$$

Putting together the estimates for $K_0(\delta)$ and $K_{\infty}(\delta)$ yields

$$t\mu(F_{t,2}) \lesssim \begin{cases} \|f\|_{L^{p,\kappa}} \big(\delta^{\alpha+\frac{\beta-n}{p}}(\delta+r)^{\frac{n-\kappa}{p}} \big(\mu(F_{t,2})\big)^{\frac{1}{p'}} \|\mu\|_{\beta}^{\frac{1}{p}} + \delta^{\alpha-\frac{\kappa}{p}} \mu(F_{t,2}) \big) & \text{for } p>1, \\ \|f\|_{L^{1,\kappa}} \big(\delta^{\alpha+\beta-n}(\delta+r)^{n-\kappa} \|\mu\|_{\beta} + \delta^{\alpha-\kappa} \mu(F_{t,2}) \big) & \text{for } p=1. \end{cases}$$

Below is a treatment of two cases: $\delta \ge r$ and $\delta \le r$.

Case δ ≥ r . Under this situation, one has

$$\begin{cases} \delta^{\alpha+\frac{\beta-n}{p}}(\delta+r)^{\frac{n-\kappa}{p}}\big(\mu(F_{t,2})\big)^{\frac{1}{p'}}\|\mu\|_{\beta}^{\frac{1}{p}}+\delta^{\alpha-\frac{\kappa}{p}}\mu(F_{t,2}) \lesssim \delta^{\alpha+\frac{\beta-\kappa}{p}}\big(\mu(F_{t,2})\big)^{\frac{1}{p'}}\|\mu\|_{\beta}^{\frac{1}{p}}+\delta^{\alpha-\frac{\kappa}{p}}\mu(F_{t,2}) & \text{for } p>1,\\ \delta^{\alpha+\beta-n}(\delta+r)^{n-\kappa}\|\mu\|_{\beta}+\delta^{\alpha-\kappa}\mu(F_{t,2}) \lesssim \delta^{\alpha+\beta-\kappa}\|\mu\|_{\beta}+\delta^{\alpha-\kappa}\mu(F_{t,2}) & \text{for } p=1, \end{cases}$$

and hence selecting

$$\delta \approx \left(\frac{\mu(F_{t,2})}{\|\|\mu\|\|_{\beta}}\right)^{\frac{1}{\beta}}$$

yields

$$\left(\frac{\mu(F_{t,2})}{\|\mu\|_{\beta}}\right)^{\frac{\kappa-\alpha p}{\beta p}} \lesssim t^{-1} \|f\|_{L^{p,\kappa}}.$$

Taking the $p(\beta + \kappa - n)/(\kappa - \alpha p)$ -th power on this last estimate gives

$$\mu(F_{t,2}) \lesssim \|\|\mu\|\|_{\beta}(t^{-1}\|f\|_{L^{p,\kappa}})^{\frac{p(\beta+\kappa-n)}{\kappa-ap}} \left(\frac{\mu(F_{t,2})}{\|\|\mu\|\|_{\beta}}\right)^{\frac{n-\kappa}{\beta p}} \lesssim (t^{-1}\|f\|_{L^{p,\kappa}})^{\frac{p(\beta+\kappa-n)}{\kappa-ap}} \|\|\mu\|\|_{\beta} r^{n-\kappa}.$$

Case δ ≤ r . Under this situation, one has

$$\begin{cases} \delta^{\alpha+\frac{\beta-n}{p}}(\delta+r)^{\frac{n-\kappa}{p}}\big(\mu(F_{t,2})\big)^{\frac{1}{p'}}\|\mu\|_{\beta}^{\frac{1}{p}}+\delta^{\alpha-\frac{\kappa}{p}}\mu(F_{t,2})\leq \bigg(\frac{r^{\frac{n-\kappa}{p}}}{\delta^{\frac{n-\beta}{p}-\alpha}}\bigg)\big(\mu(F_{t,2})\big)^{\frac{1}{p'}}\|\mu\|_{\beta}^{\frac{1}{p}}+\frac{\mu(F_{t,2})}{\delta^{\frac{\kappa}{p}-\alpha}} & \text{for } p>1, \\ \delta^{\alpha+\beta-n}(\delta+r)^{n-\kappa}\|\mu\|_{\beta}+\delta^{\alpha-\kappa}\mu(F_{t,2})\leq \delta^{\alpha+\beta-n}r^{n-\kappa}\|\mu\|_{\beta}+\delta^{\alpha-\kappa}\mu(F_{t,2}) & \text{for } p=1. \end{cases}$$

By selecting

$$\delta \approx \left(\frac{r^{n-\kappa}|||\mu|||_{\beta}}{\mu(F_{t,2})}\right)^{\frac{1}{n-\kappa-\beta}},$$

one achieves

$$t\lesssim \|f\|_{L^{p,\kappa}} \left(\frac{r^{n-\kappa}\|\|\mu\|\|_{\beta}}{\mu(F_{t,2})}\right)^{\frac{\alpha-\kappa}{n-\kappa-\beta}},$$

thereby arriving at

$$\mu(F_{t,2}) \leq \|\|\mu\|\|_{\beta} (t^{-1} \|f\|_{L^{p,\kappa}})^{\frac{p(\kappa+\beta-n)}{\kappa-\alpha p}} r^{n-\kappa}.$$

Now, a combination of the preceding two steps gives

$$\mu(F_t) \lesssim \|\|\mu\|\|_{\beta} (t^{-1} \|f\|_{L^{p,\kappa}})^{\frac{p(\kappa+\beta-n)}{\kappa-ap}} r^{n-\kappa} \quad \text{for all } (x,r,t) \in \mathbb{R}^n \times (0,\infty) \times (0,\infty).$$

Since $\lambda \leq \kappa$, one gets

$$q=\frac{p(\beta+\lambda-n)}{\kappa-\alpha p}\leq \frac{p(\beta+\kappa-n)}{\kappa-\alpha p}=\tilde{q},$$

thereby implying

$$(\mu(F_t))^{\frac{1}{q}} = (\mu(F_t))^{\frac{1}{q}} (\mu(F_t))^{\frac{1}{q} - \frac{1}{q}} \ge (\mu(F_t))^{\frac{1}{q}} r^{\beta(\frac{1}{q} - \frac{1}{q})} \|\mu\|_{\dot{q}}^{\frac{1}{q} - \frac{1}{q}}.$$

As a result, one establishes

$$\mu(F_t) \lesssim \|\|\mu\|\|_{\mathcal{B}} t^{-q} r^{n-\lambda} \|f\|_{L^{p,\kappa}}^q \quad \text{for all } (x,r,t) \in \mathbb{R}^n \times (0,\infty) \times (0,\infty),$$

as desired.

2.2 Proof of Theorem 1.2 (i)

On the one hand, since

$$||1_K||_{H^{1,\kappa}} = ||1_K||_{L^1} = \nu(K)$$
 for all compact $K \subset \mathbb{R}^n$,

one has

$$\sup_{0 \not\equiv f \in H^{1,\kappa}} \|I_{\alpha}f\|_{L^{q,\lambda}_{\mu}} \|f\|_{H^{1,\kappa}}^{-1} < \infty \ \Rightarrow \ \|\|\mu\|_{\alpha,\beta,\lambda} < \infty.$$

Conversely, assume $\|\|\mu\|\|_{\alpha,\beta,\lambda} < \infty$. Without loss of generality, one may assume $0 \le f \in H^{1,\kappa} = L^1$. Then such a function f can be written as $f = \lim_{j\to\infty} f_j$ (cf. [44, p. 88]), where

$$f_j(y) = \sum_{i=1}^{j2^j} (j-1)2^{-j} \mathbf{1}_{K_{i,j}}(y) + j \mathbf{1}_{K_j}(y), \quad K_{i,j} = \{ y \in \mathbb{R}^n : (i-1)2^{-j} \le f(y) < i2^{-j} \}, \quad K_j = \{ y \in \mathbb{R}^n : f(y) \ge j \}.$$

Thus, applying the Minkowski inequality for $L^q_\mu(B(x,r))$ (all L^q_μ -integrable functions on a given ball B(x,r)) derives

$$\begin{split} \|I_{\alpha}f_{j}\|_{L^{q}_{\mu}(B(x,r))} &\leq \sum_{i=1}^{j2^{j}} (j-1)2^{-j} \|I_{\alpha}1_{K_{i,j}}\|_{L^{q}_{\mu}(B(x,r))} + j \|I_{\alpha}1_{K_{j}}\|_{L^{q}_{\mu}(B(x,r))} \\ &\leq r^{\frac{n-\lambda}{q}} \left(\sum_{i=1}^{j2^{j}} (j-1)2^{-j} \|I_{\alpha}1_{K_{i,j}}\|_{L^{q,\lambda}_{\mu}} + j \|I_{\alpha}1_{K_{j}}\|_{L^{q,\lambda}_{\mu}} \right) \\ &\leq r^{\frac{n-\lambda}{q}} \|\|\mu\|\|_{\alpha,\beta,\lambda} \left(\sum_{i=1}^{j2^{j}} (j-1)2^{-j} \nu(K_{i,j}) + j \nu(K_{j}) \right) \\ &\leq r^{\frac{n-\lambda}{q}} \|\|\mu\|\|_{\alpha,\beta,\lambda} \|f_{j}\|_{L^{1}}. \end{split}$$

Now, sending j to ∞ leads to

$$||I_{\alpha}f||_{L^{q}_{u}(B(x,r))} \leq r^{\frac{n-\lambda}{q}} |||\mu|||_{\alpha,\beta,\lambda} ||f||_{L^{1}},$$

whence

$$\|I_{\alpha}f\|_{L^{q,\lambda}_{\alpha}} \leq \|\|\mu\|\|_{\alpha,\beta,\lambda} \|f\|_{H^{1,\kappa}} \approx \|\|\mu\|\|_{\alpha,\beta,\lambda} \|f\|_{H^{1,\kappa}}.$$

On the other hand, an application of Lemma 2.1 with $0 \le \lambda \le \kappa = n$, p = 1 and $q = (\beta + \lambda - n)/(n - \alpha) \ge 1$ gives

$$\|\|\mu\|\|_{\beta} < \infty \iff \sup_{0 \neq f \in L^1} \|I_{\alpha}f\|_{L^{q,\lambda}_{\mu,\infty}} \|f\|_{L^1}^{-1} < \infty.$$

So, if $\|\|\mu\|\|_{\beta} < \infty$, then $I_{\alpha} : H^{1,\kappa} \to L^{q,\lambda}_{\mu,\infty}$ is continuous, and vice versa.

Remark 2.2. In the first equivalence of Theorem 1.2 (i), $\|\mu\|_{\alpha,\beta,\lambda} < \infty$ implies easily $\|\mu\|_{\beta} < \infty$ (cf. the beginning of the argument for the proof of Lemma 2.1). However, if $\lambda = n$, $\mu = \nu$ (ensuring $\beta = n$) and $q = n/(n-\alpha)$, then

$$\|\mu\|_{\alpha,\beta,\lambda} \ge \|I_{\alpha}1_{B(0,1)}\|_{L^{n/(n-\alpha)}}(\nu(B(0,1)))^{-1} = \infty,$$

and hence $I_{\alpha}L^{1}$ is not subset of $L^{\frac{n}{n-\alpha}}$ according to the first equivalence of Theorem 1.2 (i).

Moreover, as a key tool of verifying the second equivalence of Theorem 1.2 (i), Lemma 2.1 can be used, along with the well-known layer-cake formula, to imply the following. If $0 < \alpha < n$, $0 < \lambda \le \kappa \le n$,

$$1 \leq p < \frac{\kappa}{\alpha}, \quad n - \alpha p < \beta \leq n, \quad 1 \leq \hat{q} < q = \frac{p(\beta + \lambda - n)}{\kappa - \alpha p}, \quad \|\|\mu\|\|_{\beta} < \infty, \quad f \in L^{p,\kappa},$$

then

$$\int\limits_{B(x,r)} |I_{\alpha}f(y)|^{\hat{q}} \ d\mu(y) = \left(\int\limits_{0}^{\delta} + \int\limits_{\delta}^{\infty}\right) \mu\left(\left\{y \in B(x,r) : |I_{\alpha}f(y)| > t\right\}\right) dt^{\hat{q}} \leq \|\|\mu\|_{\beta} \left(r^{\beta}\delta^{\hat{q}} + \|f\|_{L^{p,\kappa}}^{q} r^{n-\lambda}\delta^{\hat{q}-q}\right),$$

and hence

$$\int\limits_{B(x,r)}|I_{\alpha}f(y)|^{\hat{q}}\;d\mu(y)\leq \|\|\mu\|_{\beta}\|f\|_{L^{p,\kappa}}^{\hat{q}}r^{\beta+\frac{\hat{q}(n-\lambda-\beta)}{q}},$$

through choosing $\delta = (r^{n-\lambda-\beta} \|f\|_{L^{p,x}}^q)^{\frac{1}{q}}$. Consequently,

$$I_{\alpha}f \in L_{\mu}^{\hat{q},\hat{\lambda}} \quad \text{with } \hat{\lambda} = n - \beta + \frac{\hat{q}(\kappa - \alpha p)}{p} < n.$$

In other words,

$$\|\|\mu\|\|_{\beta}<\infty \Rightarrow \sup_{0\neq f\in L^{p,\kappa}}\|I_{\alpha}f\|_{L^{\hat{q},\lambda}_{\mu}}\|f\|_{L^{p,\kappa}}^{-1}<\infty.$$

Naturally, if $\hat{q} = p$ in the last implication, then

$$\|\|\mu\|_{\beta}<\infty \Leftrightarrow \sup_{0\neq f\in L^{p,\kappa}}\|I_{\alpha}f\|_{L^{p,\lambda}_{\mu}=n-\beta+\kappa-\alpha p}\|f\|_{L^{p,\kappa}}^{-1}<\infty,$$

whose speciality $\mu = \nu$ recovers the second imbedding of [8, Theorem 16 (i)] which in turn yields the following new k-th order Morrey–Poincare inequality:

$$||f||_{L^{p,\kappa-kp}} \lesssim ||\nabla^k f||_{L^{p,\kappa}}$$
 for all $||\nabla^k f||_{L^{p,\kappa}} > 0$ with $kp < \kappa \le n$.

3 Demonstration of Theorem 1.2 (ii)

3.1 A lemma for Theorem 1.2 (ii)

To demonstrate Theorem 1.2(ii), one needs an equivalent characterisation of $\|\mu\|_{\mathcal{B}}$.

Lemma 3.1. *For* $(\beta, \gamma) \in [0, n] \times [0, n]$ *, let*

$$\|\|\mu\|\|_{\beta,\gamma} \equiv \sup_{(x,y,r)\in\mathbb{R}^n\times\mathbb{R}^n\times(0,\infty)} r^{-\beta} \int\limits_{B(x,r)} |z-y|^{-\gamma} d\mu(z)$$

be the (β, γ) -order ball variation of μ . If $\gamma \leq \beta$, then

$$\|\|\mu\|\|_{\beta} \leq \|\|\mu\|\|_{\beta-\gamma,\gamma}.$$

Moreover, if $y < \beta$ *, then*

$$\|\|\mu\|\|_{\beta} \gtrsim \|\|\mu\|\|_{\beta-\nu,\nu}.$$

But, if $0 < y = \beta$, then there exists a Radon measure μ_0 such that

$$\|\|\mu_0\|\|_{\beta} < \infty$$
, $\|\|\mu_0\|\|_{\beta-\gamma,\gamma} = \infty$.

Proof. On the one hand, one utilises the above definition to discover that if $y \le \beta$, then

$$\|\|\mu\|\|_{\beta-\gamma,\gamma} \ge \sup_{(x,r)\in\mathbb{R}^n\times(0,\infty)} r^{-(\beta-\gamma)} \int_{B(x,r)} |z-x|^{-\gamma} d\mu(z) \ge \sup_{(x,r)\in\mathbb{R}^n\times(0,\infty)} r^{-\beta} \mu(B(x,r)) = \|\|\mu\|\|_{\beta}.$$

On the other hand, if $y < \beta$, then [29, Lemma 1.27] is employed to estimate

$$\begin{aligned} \|\mu\|_{\beta-\gamma,\gamma} &\approx \sup_{(x,y,r)\in\mathbb{R}^n\times\mathbb{R}^n\times(0,\infty)} r^{-(\beta-\gamma)} \bigg(\int\limits_0^r + \int\limits_r^\infty \bigg) \mu \big(B(x,r)\cap B(y,s)\big) s^{-\gamma-1} \, ds \\ &\leq \|\mu\|_{\beta} \sup_{(x,y,r)\in\mathbb{R}^n\times\mathbb{R}^n\times(0,\infty)} r^{-(\beta-\gamma)} \bigg(\int\limits_0^r s^{\beta-\gamma-1} \, ds + \int\limits_r^\infty r^{\beta} s^{-\gamma-1} \, ds \bigg) \approx \|\mu\|_{\beta}. \end{aligned}$$

Unfortunately, this last estimation is not valid for $\gamma = \beta = j$ (a natural number less than n). In fact, for the origin $\{0\}^{n-j}$ of \mathbb{R}^{n-j} , the Radon measure $d\mu_j = 1_{\mathbb{R}^{j} \times \{0\}^{n-j}} d\nu$ satisfies

$$\|\mu_i\|_i < \infty, \quad \|\mu_i\|_{0,i} = \infty.$$

3.2 Proof of Theorem 1.2 (ii)

The argument comprises two parts.

Part 1: Verify that I_{α} maps $H^{p,\kappa}$ to either $L^{q,\lambda}_{\mu}$ or $L^{q,\lambda}_{\mu,\infty}$ continuously if and only if $\|\mu\|_{\beta} < \infty$. Since $L^{q,\lambda}_{\mu} \subset L^{q,\lambda}_{\mu,\infty}$, it is enough to check

$$\sup_{0 \neq f \in H^{p,\kappa}} \|I_{\alpha}f\|_{L^{q,\lambda}_{\mu,\infty}} \|f\|_{H^{p,\kappa}}^{-1} < \infty \ \Rightarrow \ \|\mu\|_{\beta} < \infty \ \Rightarrow \ \sup_{0 \neq f \in H^{p,\kappa}} \|I_{\alpha}f\|_{L^{q,\lambda}_{\mu}} \|f\|_{H^{p,\kappa}}^{-1} < \infty.$$

To this end, recall that (cf. [6, Theorem 3.3]) if $f \in H^{p,\kappa}$, then

$$||f||_{H^{p,\kappa}} \approx \inf \left\{ \sum_j |c_j| : f = \sum_j c_j a_j \right\},$$

where a_i is a $(p, n - \kappa)$ -atom, that is, a_i is supported on a ball $B_i \in \mathbb{R}^n$ and satisfies

$$||a_j||_{L^p} \leq \left(\nu(B_j)\right)^{\frac{\kappa-n}{np^j}}, \quad \text{where } p' = \frac{p}{p-1}.$$

So, for a given ball B(x, r), the function

$$f_{x,r} = (\nu(B(x,r)))^{\frac{\kappa-n}{np'}-\frac{1}{p}} 1_{B(x,r)}$$

belongs to $H^{p,\kappa}$. Consequently,

$$\|f_{x,r}\|_{H^{p,x}}\approx 1, \quad r^{\alpha-n+\frac{\kappa}{p'}}\lesssim \inf_{y\in B(x,r)}I_{\alpha}f_{x,r}(y), \quad \mu\big(B(x,r)\big)\lesssim \mu\big(\big\{y\in B(x,r): r^{\alpha-n+\frac{\kappa}{p'}}\lesssim I_{\alpha}f_{x,r}(y)\big\}\big).$$

Now, if

$$\sup_{0\neq f\in H^{p,\kappa}} \lVert I_{\alpha}f\rVert_{L^{q,\lambda}_{\mu,\infty}} \lVert f\rVert_{H^{p,\kappa}}^{-1} < \infty,$$

then $||I_{\alpha}f_{x,r}||_{L^{q,\lambda}_{u,\infty}} \leq 1$, and hence

$$r^{\lambda-n+q(\alpha-n+\frac{\kappa}{p'})}\mu\big(B(x,r)\big)\lesssim r^{\lambda-n+q(\alpha-n+\frac{\kappa}{p'})}\mu\big(\big\{y\in B(x,r): r^{\alpha-n+\frac{\kappa}{p'}}\lesssim I_{\alpha}f_{x,r}(y)\big\}\big)\lesssim \|I_{\alpha}f_{x,r}\|_{L^{q,\lambda}_u}^q\lesssim 1.$$

This in turn validates $\|\|\mu\|\|_{\beta} < \infty$.

Next, suppose $\|\mu\|_{\beta} < \infty$. The desired continuity follows from a consideration of the forthcoming two cases.

Case $\lambda < n$. Since $0 < n - \lambda < \beta \le 2n - \lambda$, Lemma 3.1 is utilised to ensure $\|\mu\|_{n-\lambda,\beta-(n-\lambda)} < \infty$. Let $f = \sum_j c_j a_j$ belong to $H^{p,\kappa}$. Given a ball B(x,r) and the conjugate index q' = q/(q-1), let $L^{q'}_{\mu}(B(x,r))$ be the μ -measurable functions g with

$$\|g\|_{L^{q'}_{\mu}(B(x,r))} = \left(\int\limits_{B(x,r)} |g|^{q'} d\mu\right)^{\frac{1}{q'}} < \infty.$$

Then, from duality and the Fubini theorem it follows that

$$\left(\int_{B(x,r)} |I_{\alpha}f|^{q} d\mu\right)^{\frac{1}{q}} = \sup_{\|g\|_{L_{\mu}^{q'}(B(x,r))} \le 1} \left|\int_{B(x,r)} gI_{\alpha}f d\mu\right|
= \sup_{\|g\|_{L_{\mu}^{q'}(B(x,r))} \le 1} \left|\int_{B(x,r)} g(z) \int_{\mathbb{R}^{n}} f(y)|z - y|^{\alpha - n} d\nu(y) d\mu(z)\right|
= \sup_{\|g\|_{L_{\mu}^{q'}(B(x,r))} \le 1} \left|\int_{\mathbb{R}^{n}} \left(\int_{B(x,r)} g(z)|z - y|^{\alpha - n} d\mu(z)\right) f(y) d\nu(y)\right|
\le \sup_{\|g\|_{L_{\mu}^{q'}(B(x,r))} \le 1} \sum_{j} |c_{j}| \int_{\mathbb{R}^{n}} \left(\int_{B(x,r)} |g(z)||z - y|^{\alpha - n} d\mu(z)\right) |a_{j}(y)| d\nu(y).$$

Next, choose the parameter pair $\{\alpha_1, \alpha_2\}$ such that

$$n-\beta < \alpha_1 = 2n-\beta - \lambda < n$$
, $0 < \alpha_2 < n$, $\alpha q = \alpha_1 + (q-1)\alpha_2$.

Then the Hölder inequality on the supporting ball B_i of a_i , with $q \le p$, yields

$$\begin{split} & \int\limits_{\mathbb{R}^{n}} \left(\int\limits_{B(x,r)} |g(z)| |z-y|^{\alpha-n} \, d\mu(z) \right) |a_{j}(y)| \, d\nu(y) \\ & \leq \int\limits_{B_{j}} \left(\int\limits_{B(x,r)} |g(z)| |z-y|^{\alpha-n} \, d\mu(z) \right) |a_{j}(y)| \, d\nu(y) \\ & \leq \left(\int\limits_{B_{j}} \left(\int\limits_{B(x,r)} |g(z)| |z-y|^{\alpha-n} \, d\mu(z) \right)^{q'} \, d\nu(y) \right)^{\frac{1}{q'}} \left(\int\limits_{B_{j}} |a_{j}|^{q} \, d\nu \right)^{\frac{1}{q}} \\ & \leq \left(\nu(B_{j}) \right)^{\frac{1}{q} - \frac{1}{p}} \left(\int\limits_{B_{j}} |a_{j}|^{p} \, d\nu \right)^{\frac{1}{p}} \left(\int\limits_{B_{j}} \left(\int\limits_{B(x,r)} |g(z)| |z-y|^{\alpha-n} \, d\mu(z) \right)^{q'} \, d\nu(y) \right)^{\frac{1}{q'}} \\ & \leq \left(\nu(B_{j}) \right)^{\frac{1}{q} - 1 + \frac{\kappa}{p'n}} \left(\int\limits_{B_{j}} \left(\int\limits_{B(x,r)} |g(z)| |z-y|^{\alpha-n} \, d\mu(z) \right)^{q'} \, d\nu(y) \right)^{\frac{1}{q'}}. \end{split}$$

Another application of the Hölder inequality, along with $\|\mu\|_{n-\lambda,\beta-(n-\lambda)} < \infty$, gives

$$\begin{split} \left(\int\limits_{B(x,r)} |g(z)| |z-y|^{\alpha-n} \, d\mu(z) \right)^{q'} & \leq \left(\int\limits_{B(x,r)} \frac{|g(z)|^{q'}}{|z-y|^{n-\alpha_2}} \, d\mu(z) \right) \left(\int\limits_{B(x,r)} \frac{d\mu(z)}{|z-y|^{n-\alpha_1}} \right)^{q'-1} \\ & \leq \left(r^{n-\lambda} \| \|\mu\|_{n-\lambda,\beta-(n-\lambda)} \right)^{q'-1} \int\limits_{B(x,r)} \frac{|g(z)|^{q'}}{|z-y|^{n-\alpha_2}} \, d\mu(z). \end{split}$$

Now, the foregoing two groups of estimates are put together, and using the Fubini theorem and [29, Lemma 1.30], as well as the equalities

$$q=\frac{p(\beta+\lambda-n)}{\kappa-p(\alpha+\kappa-n)}, \quad \alpha_2=q'\bigg(\alpha-\frac{2n-\beta-\lambda}{q}\bigg), \quad \frac{1}{q}+\frac{\alpha_2}{nq'}=1-\frac{\kappa}{np'},$$

gives

$$\begin{split} & \int\limits_{\mathbb{R}^{n}} \left(\int\limits_{B(x,r)} |g(z)| |z-y|^{\alpha-n} \ d\mu(z) \right) |a_{j}(y)| \ d\nu(y) \\ & \lesssim \left(\nu(B_{j}) \right)^{\frac{1}{q}-1+\frac{\kappa}{p'n}} r^{\frac{n-\lambda}{q}} \left\| \mu \right\|_{n-\lambda,\beta-(n-\lambda)}^{\frac{1}{q}} \left(\int\limits_{B_{j}} \int\limits_{B(x,r)} \frac{|g(z)|^{q'}}{|z-y|^{n-\alpha_{2}}} \ d\mu(z) \ d\nu(y) \right)^{\frac{1}{q'}} \\ & \approx \left(\nu(B_{j}) \right)^{\frac{1}{q}-1+\frac{\kappa}{p'n}} r^{\frac{n-\lambda}{q}} \left\| \mu \right\|_{n-\lambda,\beta-(n-\lambda)}^{\frac{1}{q}} \left(\int\limits_{B(x,r)} \left(\int\limits_{B_{j}} \frac{d\nu(y)}{|z-y|^{n-\alpha_{2}}} \right) |g(z)|^{q'} \ d\mu(z) \right)^{\frac{1}{q'}} \\ & \lesssim \left(\nu(B_{j}) \right)^{\frac{1}{q}-1+\frac{\kappa}{p'n}+\frac{\alpha_{2}}{nq'}} r^{\frac{n-\lambda}{q}} \left\| \mu \right\|_{n-\lambda,\beta-(n-\lambda)}^{\frac{1}{q}} \left(\int\limits_{B(x,r)} |g(z)|^{q'} \ d\mu(z) \right)^{\frac{1}{q'}} \\ & \approx r^{\frac{n-\lambda}{q}} \left\| \mu \right\|_{n-\lambda,\beta-(n-\lambda)}^{\frac{1}{q}} \left(\int\limits_{B(x,r)} |g(z)|^{q'} \ d\mu(z) \right)^{\frac{1}{q'}}. \end{split}$$

This in turn yields

$$\left(\int\limits_{B(x,r)}|I_{\alpha}f|^{q}\ d\mu\right)^{\frac{1}{q}}\lesssim r^{\frac{n-\lambda}{q}}\|\|\mu\|\|_{n-\lambda,\beta-(n-\lambda)}^{\frac{1}{q}}\sum_{j}|c_{j}|,$$

and so by Lemma 3.1,

$$\|I_{\alpha}f\|_{L^{q,\lambda}} \lesssim \|\|\mu\|_{n-\lambda,\beta-(n-\lambda)}^{\frac{1}{q}} \|f\|_{H^{p,\kappa}} \lesssim \|\|\mu\|_{\beta}^{\frac{1}{q}} \|f\|_{H^{p,\kappa}}.$$

Case $\lambda=n$. Under this situation, one has $L_{\mu}^{q,\lambda}=L_{\mu}^q$. So, a particular care is needed. However, the initial issue is to take such a parameter pair (α_1, α_2) such that

$$n-\beta < \alpha_1 < n$$
, $0 < \alpha_2 < n$, $\alpha q = \alpha_1 + (q-1)\alpha_2$.

Fix a point x_0 in the supporting ball B of a given $(p, n-\kappa)$ -atom a and select $B(x_0, r_0)$ so that $v(B(x_0, r_0)) = v(B)$ and $B \subseteq B(x_0, 2r_0)$. One can use [29, Lemma 1.30] to get

$$\int_{B} |y-z|^{\alpha_2-n} d\nu(z) \leq (\nu(B))^{\frac{\alpha_2}{n}},$$

and then use the Fubini theorem, [29, Lemma 1.27] and the Hölder inequality with $q \le p$ to compute

$$\begin{split} \int\limits_{B(x_0,2r_0)} \left(\int\limits_{B} |a(z)|^q |y-z|^{\alpha_1-n} \, dv(z) \right) d\mu(y) &= \int\limits_{B} \left(\int\limits_{B(x_0,2r_0)} |y-z|^{\alpha_1-n} \, d\mu(y) \right) |a(z)|^q \, dv(z) \\ &\leq \int\limits_{B} \left(\int\limits_{0}^{2r_0} + \int\limits_{2r_0}^{\infty} \mu \big(B(x_0,2r_0) \cap B(z,s) \big) s^{\alpha_1-n-1} \, ds \right) |a(z)|^q \, dv(z) \end{split}$$

$$\leq \|\|\mu\|_{\beta} \left(\int_{B} |a|^{q} dv \right) r_{0}^{\beta + \alpha_{1} - n}$$

$$\leq \|\|\mu\|_{\beta} \left(\int_{B} |a|^{p} dv \right)^{\frac{q}{p}} (v(B))^{1 - \frac{q}{p}} r_{0}^{\beta + \alpha_{1} - n}$$

$$\approx \|\|\mu\|_{\beta} (v(B))^{\frac{\beta + \alpha_{1}}{n} + \frac{q\kappa}{n}(1 - \frac{1}{p}) - q}.$$

Now, one utilises $\alpha = \alpha_1/q + (q-1)\alpha_2/q$, the Hölder inequality and $q = \beta p/(p(n-\alpha) - (p-1)\kappa)$ to find the following estimation:

$$\int_{B(x_0,2r_0)} (I_{\alpha}|a|)^q d\mu \leq \int_{B(x_0,2r_0)} \left(\int_{B} \frac{|a(y)|^q dv(y)}{|y-z|^{n-\alpha_1}} \right) \left(\int_{B} \frac{dv(y)}{|y-z|^{n-\alpha_2}} \right)^{q-1} d\mu(z) \\
\leq \|\mu\|_{\beta} (v(B))^{\frac{\beta+\alpha_1}{n} + \frac{q\kappa}{n}(1-\frac{1}{p}) - q + \frac{\alpha_2(q-1)}{n}} \\
\approx \|\mu\|_{\beta}.$$

Meanwhile, a combination of the Minkowski inequality, [29, Lemma 1.27], the Hölder inequality and the fact that $q = \beta p/(p(n-\alpha) - (p-1)\kappa) > \beta/(n-\alpha)$ yields

$$\left(\int_{B(x_{0},2r_{0})^{c}} (I_{\alpha}|a|)^{q} d\mu\right)^{\frac{1}{q}} \leq \int_{B} \left(\int_{B(x_{0},2r_{0})^{c}} |y-z|^{(\alpha-n)q} d\mu(y)\right)^{\frac{1}{q}} |a(z)| d\nu(z)$$

$$\leq \int_{B} \left(\int_{2r_{0}}^{\infty} s^{(\alpha-n)q-1} \mu(B(x_{0},2r_{0})^{c} \cap B(z,s)) ds\right)^{\frac{1}{q}} |a(z)| d\nu(z)$$

$$\leq \left(\int_{B} |a|^{p} d\nu\right)^{\frac{1}{p}} (\nu(B))^{1-\frac{1}{p}} (\nu(B))^{\frac{\alpha q+\beta}{nq}-1} \|\mu\|_{\beta}^{\frac{1}{q}}$$

$$\leq (\nu(B))^{\frac{\alpha q+\beta}{nq}-\frac{1}{p}+\frac{\kappa}{np'}-\frac{1}{p'}} \|\mu\|_{\beta}^{\frac{1}{q}}$$

$$\approx \|\mu\|_{\beta}^{\frac{1}{q}}.$$

Putting the integrals $\int_{B(x_0,2r_0)}(\cdots) d\mu$ and $\int_{B(x_0,2r_0)^c}(\cdots) d\mu$ together implies

$$||I_{\alpha}|a||_{L^{q}_{\mu}} \leq |||\mu||_{\beta}^{\frac{1}{q}}.$$

Upon expressing $f \in H^{p,\kappa}$ as $f = \sum_i c_i a_i$, one uses the foregoing estimate and the Minkowski inequality to establish the following inequality:

$$\|I_{\alpha}f\|_{L^{q}_{\mu}} \leq \sum_{j} |c_{j}| \|I_{\alpha}|a_{j}|\|_{L^{q}_{\mu}} \lesssim \||\mu||_{\beta}^{\frac{1}{q}} \sum_{j} |c_{j}|.$$

Consequently, one finds

$$||I_{\alpha}f||_{L^{q}_{\mu}} \leq |||\mu||_{\beta}^{\frac{1}{q}} ||f||_{H^{p,\kappa}}.$$

Part 2: Verify that I_{α} maps $H_{1}^{p,\kappa}$ to either $L_{\mu}^{q,\lambda}$ or $L_{\mu,\infty}^{q,\lambda}$ continuously if and only if $\|\mu\|_{\beta} < \infty$. Thanks to both $H_{1}^{p,\kappa} \in H^{p,\kappa}$ and $L_{\mu}^{q,\lambda} \in L_{\mu,\infty}^{q,\lambda}$, using Part 1, one has that $\|\mu\|_{\beta} < \infty$ implies that I_{α} maps $H_{1}^{p,\kappa}$ to either $L_{\mu}^{q,\lambda}$ or $L_{\mu,\infty}^{q,\lambda}$ continuously. So, it remains to check that if $I_{\alpha} : H_{1}^{p,\kappa} \to L_{\mu,\infty}^{q,\lambda}$ is continuous, then $\|\mu\|_{\beta} < \infty$. To this end, utilising [20, Lemma 3.1] one gets that $f \in H_{1}^{p,\kappa}$ if and only if there exist a sequence of numbers $\{c_i\}$ and a sequence of functions $\{a_i\}$ such that

$$f = \sum_{j} c_j a_j$$
 with $||f||_{H_1^{p,\kappa}} \approx \sum_{j} |c_j| < \infty$,

for which there exists a ball B_i containing the support of a_i such that

$$\left(\nu(B_j)\right)^{\frac{n-\kappa}{p^ln}}\int\limits_0^\infty \left(\nu\left(\left\{x\in\mathbb{R}^n: 1_{B_j}(x)|a_j(x)|>t\right\}\right)\right)^{\frac{1}{p}}dt\leq 1.$$

Upon choosing

$$f_{x,r} = (\nu(B(x,r)))^{\frac{\kappa-n}{np'} - \frac{1}{p}} 1_{B(x,r)}$$

for a given ball B(x, r), one gets

$$||f_{x,r}||_{H^{p,\kappa}} \lesssim 1.$$

Now, using the induced condition

$$||I_{\alpha}f_{x,r}||_{L^{q,\lambda}_{1}} \lesssim ||f_{x,r}||_{H^{p,\kappa}_{1}}$$

and the calculation (with $f_{x,r}$) done at the beginning of Part 1, one can readily reach $\|\mu\|_{\beta} < \infty$.

Remark 3.2. If $H^{p,0}$ or $H^{p,n}$ is defined as the space of all ν -measurable functions f with

$$||f||_{H^{p,\kappa}} \approx \inf \left\{ \sum_{j} |c_j| : f = \sum_{j} c_j a_j \right\} < \infty,$$

and a_j is a (p, n)-atom or a (p, 0)-atom, i.e., a_j is supported on a ball $B_j \subset \mathbb{R}^n$ and satisfies $\|a_j\|_{L^p} \leq (\nu(B_j))^{-\frac{1}{p'}}$ or $\|a_j\|_{L^p} \leq 1$, then the above argument is still valid for $H^{p,0}$ and $H^{p,n}$, and hence Theorem 1.2 can be extended to $H^{p,0\leq \kappa\leq n}$. Therefore, not only this extension gives a new approach to [1, Theorem 2], but also the limiting case $\alpha \to 0$ of Theorem 1.2 (ii) for $\lambda = n$ and $\mu = \nu$ actually gives (cf. [26, p.85])

$$H^{p,\kappa} \subset L^q$$
 with $q = \frac{pn}{\kappa + p(n - \kappa)}$.

Furthermore, one has the following cases:

- (a) If $\lambda = n$ and $0 < \kappa < n$, then Theorem 1.2 (ii) recovers [10, Theorem 2.4 (ii)].
- (b) If $\mu = \nu$ (forcing $\beta = n$) and $\lambda = \kappa$ in Theorem 1.2 (ii), then

$$||I_{\alpha}f||_{L^{\tilde{q}=p\kappa/(\kappa+p(n-\alpha-\kappa)),\kappa}} \lesssim ||f||_{H^{p,\kappa}} \quad \text{for all } f \in H^{p,\kappa},$$

which may be regarded as a sort of the associate representation of the first embedding of [8, Theorem 16 (ii)] (cf. [2, Theorem 3.1])

$$||I_{\alpha}f||_{L^{\bar{q}=p\kappa/(\kappa-\alpha p),\kappa}} \lesssim ||f||_{L^{p,\kappa}} \quad \text{for all } f \in L^{p,\kappa}$$

due to $1/\tilde{q} + 1/\bar{q} + \kappa/n = 1$.

(c) If $p = q = (\beta + \lambda - n - \kappa)/(n - \alpha - \kappa)$, then Theorem 1.2 (ii) is utilised to deduce

$$\|\|\mu\|\|_{\beta}<\infty \ \Leftrightarrow \ \|I_{\alpha}f\|_{L^{p,\bar{\lambda}=n+\kappa-\beta+p(n-\alpha-\kappa)}_{\nu}}\lesssim \|f\|_{H^{p,\kappa}} \quad \text{ for all } f\in H^{p,\kappa},$$

which may be treated as a kind of the associate form of the last equivalence in Remark 2.2 thanks to $(\bar{\lambda} - \tilde{\lambda})/(pn) + \kappa/n = 1$.

4 Demonstration of Theorem 1.2 (iii)

4.1 A lemma for Theorem 1.2 (iii)

In accordance with [6, Theorem 2.3], [36, Theorem 4.1] and [12, Theorem 2.7], if $p \in (1, \infty)$ and $\gamma \in [0, n)$, then

$$H^{p,\gamma} = \left\{ f \in L^p_{\text{loc}} : \|f\|_{H^{p,\gamma}} \approx \inf_{w \in B_1^{(n-\gamma)}} \left(\int_{\mathbb{D}^n} |f|^p w^{1-p} \ dv \right)^{\frac{1}{p}} < \infty \right\},\,$$

where $B_1^{(n-\gamma)}$ comprises all nonnegative functions w with

$$\int_{\mathbb{R}^n} w \, d\Lambda_{n-y}^{(\infty)} \equiv \int_0^\infty \Lambda_{n-\beta}^{(\infty)} \big(\big\{ y \in \mathbb{R}^n : w(y) > t \big\} \big) \, dt \le 1,$$

and, for $E \subset \mathbb{R}^n$, the symbol $\Lambda_{n-v}^{(\infty)}(E)$ expresses the Hausdorff capacity of E with order $n-\beta$, i.e.,

$$\Lambda_{n-\gamma}^{(\infty)}(E)=\inf\sum_i r_j^{n-\gamma},$$

where the infimum ranges over all countable coverings of E by open balls of radius r_i .

To validate Theorem 1.2 (iii), it is required to push the above description of $H^{p,\gamma}$ to the limiting case $p=\infty$, that is, to characterise $H^{\infty,\gamma}$ by means of $\Lambda_{n-\gamma}^{(\infty)}$ as seen below.

Lemma 4.1. For $\gamma \in (0, n)$, let $L^{1,\gamma}$ be the Morrey space of all signed Radon (locally finite regular signed Borel) measures μ whose total variation measures $\|\mu\| \equiv \tilde{\mu}$ obey $\|\|\tilde{\mu}\|\|_{\beta} < \infty$, and set $L^{1}_{\Lambda_{\gamma}^{(\infty)}}$ be the class of all $\Lambda_{\gamma}^{(\infty)}$ -quasicontinuous functions f on \mathbb{R}^n for which

$$||f||_{L^{1}_{\Lambda_{y}^{(\infty)}}} \equiv \int_{\mathbb{R}^{n}} |f| \, d\Lambda_{y}^{(\infty)} < \infty.$$

Then

$$\mathsf{L}^{1,\gamma} = \left\{ \mu \ signed \ Radon \ measure : \|\mu\|_{[L^1_{\Lambda^{(\infty)}_y}]^\star} \equiv \sup_{\|g\|_{L^1_{\Lambda^{(\infty)}_y} \le 1}} \left| \int\limits_{\mathbb{R}^n} g \ d\mu \right| < \infty \right\} \equiv [L^1_{\Lambda^{(\infty)}_y}]^\star.$$

Moreover, if L_0^{∞} is the class of L^{∞} -functions with compact support in \mathbb{R}^n , then

$$H^{\infty,\gamma} = \left\{ f \in L_0^{\infty} : \|f\|_{\mathsf{H}^{\infty,\gamma}} \equiv \inf_{w \in B_1^{(n-\gamma)}} \|fw^{-1}\|_{L^{\infty}} < \infty \right\} \equiv \mathsf{H}^{\infty,\gamma}.$$

Consequently, $L^1_{\Lambda^{(\infty)}}$ exists as a subspace of $H^{\infty,\gamma}$.

Proof. The first identity immediately follows from [3, Proposition 1]. The second needs a demonstration.

Suppose $f \in H^{\infty,\gamma}$. Then $f \in L_0^{\infty}$ and $||f||_{H^{\infty,\gamma}} < \infty$. Now for any $g \in L^{1,\kappa}$ with $||g||_{L^{1,\kappa}} \le 1$, one has (cf. [6, Theorem 2.2])

$$||g||_{L^{1,\gamma}} = \sup_{w \in B_1^{(n-\gamma)}} \int_{\mathbb{R}^n} |f| w \, dv \le 1.$$

This in turn implies

$$\left| \int_{\mathbb{R}^n} fg \, dv \right| \leq \int_{\mathbb{R}^n} |g| w |f| w^{-1} \, dv \leq \|g\|_{L^{1,\gamma}} \|fw^{-1}\|_{L^{\infty}} \quad \text{for all } w \in B_1^{(n-\gamma)},$$

and so $f \in H^{\infty,\gamma}$. As a consequence, $H^{\infty,\gamma} \subset H^{\infty,\gamma}$.

Since $L^{1,\gamma}$ is just the associate space of $H^{\infty,\gamma}$, in short, $L^{1,\gamma} = [H^{\infty,\gamma}]^*$, in order to prove $H^{\infty,\gamma} \in H^{\infty,\gamma}$ it is enough to check

$$[\mathsf{H}^{\infty,y}]^* \equiv \left\{ f \text{ ν-measurable function} : \|f\|_{[\mathsf{H}^{\infty,y}]^*} \equiv \sup_{\|g\|_{\mathsf{H}^{\infty,y}} \le 1} \left| \int_{\mathbb{R}^n} fg \, d\nu \right| < \infty \right\} \subset L^{1,y}.$$

Now, if f lies in $[H^{\infty,y}]^*$, then for a given ball $B(x_0, r_0)$ and any function g supported in $B(x_0, r_0)$ one has

$$\|g\|_{\mathsf{H}^{\infty,y}} = \inf_{w \in B_1^{(n-y)}} \|gw^{-1}\|_{L^{\infty}(B(x_0,r_0))}.$$

Since *g* vanishes in $\mathbb{R}^n \setminus B(x_0, r_0)$, one can choose

$$w_0 = r_0^{\beta-n} 1_{B(x_0, r_0)} \in B_1^{(n-\beta)},$$

and then get

$$||g||_{\mathsf{H}^{\infty,\gamma}} \lesssim ||g||_{L^{\infty}(B(x_0,r_0))} r_0^{n-\gamma}$$

Upon taking

$$g_0(x) = \begin{cases} 1_{B(x_0, r_0)} |f(x)| f(x)^{-1} & \text{for } f(x) \neq 0, \\ 0 & \text{for } f(x) = 0, \end{cases}$$

one achieves

$$\int_{B(x_0, r_0)} |f| \, d\nu = \left| \int_{\mathbb{R}^n} f g_0 \, d\nu \right| \le \|f\|_{[H^{\infty, y}]^*} \|g_0\|_{H^{\infty, y}} \lesssim \|f\|_{[H^{\infty, y}]^*} r_0^{n-y},$$

thereby reaching $f \in L^{1,\gamma}$, and thus $[H^{\infty,\gamma}]^* \subset L^{1,\gamma}$, as desired.

To see that $L^1_{\Lambda^{(\infty)}_{n-\nu}}$ is contained in $H^{\infty,\gamma}$, suppose $f \in L^1_{\Lambda^{(\infty)}_{n-\nu}}$. Then $w = |f| \|f\|^{-1}_{L^1_{\Lambda^{(\infty)}}}$ satisfies

$$\int_{\mathbb{R}^n} w \, d\Lambda_{n-y}^{(\infty)} = \int_0^\infty \Lambda_{n-y}^{(\infty)} \big(\big\{ y \in \mathbb{R}^n : |f(y)| \|f\|_{L_{\Lambda_{n-y}}^{-1}}^{-1} > t \big\} \big) \, dt \le 1,$$

i.e., $w \in B_1^{(n-\gamma)}$, and hence

$$||f||_{H^{\infty,\gamma}} \leq ||fw^{-1}||_{L^{\infty}} \leq ||f||_{L^{1}(\infty)},$$

that is, $f \in H^{\infty,\gamma}$, as desired.

4.2 Proof of Theorem 1.2 (iii)

For a given nonnegative Radon measure μ , according to Lemma 4.1 it is better to begin with proving that

$$I_{\alpha} \colon L^{1}_{\Lambda_{n-\kappa}^{(\infty)}} \to L^{q,\lambda}_{\mu}$$

is continuous if and only if

$$I_{\alpha} \colon L^{1}_{\Lambda^{(\infty)}} \to L^{q,\lambda}_{\mu,\infty}$$

is continuous if and only if $\mu \in L^{1,\beta}$. As a matter of fact, it is enough to check

$$\|\|\mu\|\|_{\beta} < \infty \Rightarrow \|I_{\alpha}f\|_{L^{q,\lambda}_u} \lesssim \|f\|_{L^1_{\Lambda^{(\infty)}_{n-\kappa}}} \quad \text{for all } f \in L^1_{\Lambda^{(\infty)}_{n-\kappa}}.$$

To this end, assume $\|\|\mu\|\|_{\beta} < \infty$. In accordance with [6, Remark 3.4], one has

$$||f||_{L^1_{\Lambda^{(\infty)}_{n-\kappa}}} \approx \inf\left\{\sum_j |c_j| : f = \sum_j c_j a_j\right\} < \infty,$$

where a_j is a $(\infty, n - \kappa)$ -atom, i.e., a_j is supported on a ball $B_j \subset \mathbb{R}^n$ and satisfies $\|a_j\|_{L^\infty} \leq (\nu(B_j))^{\frac{\kappa-n}{n}}$.

Next, for the given parameter $q = (\beta + \lambda - n)/(n - \alpha - \kappa) > 1$ and a function $f = \sum_i c_i a_i \in L^1_{\Lambda^{(\infty)}}$, one can slightly modify the second part of the argument used for the proof of Theorem 1.2 (ii) (with not only putting p'=1 when $p=\infty$ but also noticing that $\|a_i\|_{L^\infty}\leq (\nu(B_i))^{\frac{\kappa-n}{n}}$ to obtain

$$||I_{\alpha}a_{j}||_{L_{\mu}^{q,\lambda}} \lesssim |||\mu|||_{\beta}^{\frac{1}{q}},$$

which, by using the Minkowski inequality, implies

$$||I_{\alpha}f||_{L^{q,\lambda}_{\mu}} \leq \sum_{j} |c_{j}||I_{\alpha}a_{j}||_{L^{q,\lambda}_{\mu}} \lesssim |||\mu|||_{\beta}^{\frac{1}{q}} \sum_{j} |c_{j}|.$$

Consequently, one finds

$$||I_{\alpha}f||_{L^{q,\lambda}_{\mu}} \leq |||\mu|||_{\beta}^{\frac{1}{q}} ||f||_{L^{1}_{\Lambda_{n-\kappa}}}^{1}.$$

Finally, if $I_{\alpha} \colon H^{\infty,\kappa} \to L^{q,\lambda}_{\mu}$ is continuous, then $I_{\alpha} \colon H^{\infty,\kappa} \to L^{q,\lambda}_{\mu,\infty}$ is continuous, and hence $\|\mu\|_{\beta} < \infty$ follows: lows from choosing $1_{B(x,r)}$ as the test function in $H^{\infty,\kappa}$. Now, if $\|\|\mu\|\|_{\beta} < \infty$, then the previous analysis ensures that $I_{\alpha}:L^1_{\Lambda_{n-\kappa}^{(\infty)}}\to L^{q,\lambda}_{\mu}$ is continuous. Hence, for a given ball B(x,r), any $w\in B_1^{(n-\kappa)}$ and any $g\in L^{q'}_{\mu}(B(x,r))$, one applies the Fubini theorem to compute

$$\begin{split} \left(\int\limits_{B(x,r)} |I_{\alpha}f|^{q} \, d\mu \right)^{\frac{1}{q}} &= \sup_{\|g\|_{L_{\mu}^{q'}(B(x,r))} \le 1} \left| \int\limits_{B(x,r)} (I_{\alpha}f)g \, d\mu \right| \\ &\leq \sup_{\|g\|_{L_{\mu}^{q'}(B(x,r))} \le 1} \int\limits_{B(x,r)} \left(\int\limits_{\mathbb{R}^{n}} |f(y)| |y - z|^{\alpha - n} \, dv(y) \right) |g(z)| \, d\mu(z) \\ &= \sup_{\|g\|_{L_{\mu}^{q'}(B(x,r))} \le 1} \int\limits_{\mathbb{R}^{n}} \left(\int\limits_{B(x,r)} |g(z)| |y - z|^{\alpha - n} \, d\mu(z) \right) |f(y)| \, dv(y) \\ &\leq \|fw^{-1}\|_{L^{\infty}} \sup_{\|g\|_{L_{\mu}^{q'}(B(x,r))} \le 1} \int\limits_{\mathbb{R}^{n}} \left(\int\limits_{B(x,r)} |g(z)| |y - z|^{\alpha - n} \, d\mu(z) \right) w(y) \, dv(y) \\ &= \|fw^{-1}\|_{L^{\infty}} \sup_{\|g\|_{L_{\mu}^{q'}(B(x,r))} \le 1} \int\limits_{B(x,r)} (I_{\alpha}w) |g| \, d\mu \\ &= \|fw^{-1}\|_{L^{\infty}} \|I_{\alpha}w\|_{L_{\mu}^{q}(B(x,r))} \\ &\leq \|fw^{-1}\|_{L^{\infty}} r^{\frac{n-\lambda}{q}} \|I_{\alpha}w\|_{L_{\mu}^{q,\lambda}} \\ &\leq \|fw^{-1}\|_{L^{\infty}} r^{\frac{n-\lambda}{q}} \|\mu\|_{\beta}^{\frac{1}{q}} \|w\|_{L_{\lambda_{n-x}}^{1/2}} \\ &\leq r^{\frac{n-\lambda}{q}} \|fw^{-1}\|_{L^{\infty}} \|\mu\|_{\beta}^{\frac{1}{q}}. \end{split}$$

Therefore, one utilises Lemma 4.1 to achieve

$$\|I_{\alpha}f\|_{L^{q,\lambda}_{\mu}} \lesssim \|\|\mu\|\|_{\beta}^{\frac{1}{q}} \|f\|_{\mathsf{H}^{\infty,\kappa}} \lesssim \|\|\mu\|\|_{\beta}^{\frac{1}{q}} \|f\|_{H^{\infty,\kappa}}.$$

That is, $I_{\alpha} : H^{\infty,\kappa} \to L^{q,\lambda}_{\mu}$ is continuous.

Remark 4.2. Perhaps, it is worth pointing out that $\kappa = 0$ is suitably allowable in Theorem 1.2 (iii). To see this nature more clearly, see [3, Proposition 5] which says, as Fefferman–Stein's predual of BMO (John Nirenberg's space of functions of bounded mean oscillation on \mathbb{R}^n ; see, e.g., [19, 25, 40]), that if H^1 stands for the real 1-Hardy space comprising all L^1 -functions f for which the $\{1, \ldots, n\} \ni j$ -Riesz transform

$$y \mapsto R_j f(y) = \lim_{\epsilon \to 0} \Gamma\left(\frac{n+1}{2}\right) \pi^{-\frac{n+1}{2}} \int_{B(0,\epsilon)^c} z_j |z|^{-n-1} f(y-z) \, d\nu(z)$$

belongs to L^1 , then

$$||I_{\alpha}f||_{L^{1}_{\Lambda_{n-\alpha}}} \leq ||f||_{H^{1}} = ||f||_{L^{1}} + \sum_{i=1}^{n} ||R_{i}f||_{L^{1}} \quad \text{ for all } f \in H^{1},$$

and hence, by [3, Corollary],

$$\left| \int_{\mathbb{R}^n} f \, d\mu \right| \lesssim \|\|\mu\|\|_{n-\alpha} \|f\|_{H^1} \quad \text{for all } (\mu, f) \in \mathsf{L}^{1, n-\alpha}_+ \times H^1,$$

where $\mathsf{L}^{1,n-\alpha}_+$ consists of all nonnegative elements in $\mathsf{L}^{1,n-\alpha}$. Also, recall that (cf., e.g., [23, Theorem 6.6.10]) $f \in H^1$ if and only if there exist a sequence of (∞, n) -atoms $\{a_j\}$ with $\int_{\mathbb{R}^n} a \, d\nu = 0$ and a sequence of numbers $\{c_j\}$ such that

$$f=\sum_{j}c_{j}a_{j},\quad \|f\|_{H^{1}}\approx\sum_{j}|c_{j}|<\infty.$$

So, this leads to an investigation of the endpoint $\kappa=0$ of Theorem 1.2 (iii) for $q=(\beta+\lambda-n)/(n-\alpha)>1$. On the one hand, if $I_\alpha\colon H^1\to L^{q,\lambda}_\mu$ is continuous, then

$$||I_{\alpha}a||_{L^{q,\lambda}_{\mu}} \lesssim ||a||_{H^1} \quad \text{for all } a \in H^1.$$

For $B(x_-, r) = B(x, r)$, $B(x_+, r)$ (whose distance to B(x, r) is 3r, i.e., $|x - x_+| = 5r$) and

$$a = \frac{1_{B(x_-,r)} - 1_{B(x_+,r)}}{\nu(B(x,6r))} \in H^1 \quad \text{with } ||a||_{H^1} \le 1,$$

via a simple geometric computation, one gets

$$I_{\alpha}a(y) = \left(\nu(B(x,6r))\right)^{-1} \left(\int\limits_{B(x_{-},r)} \frac{d\nu(z)}{|z-y|^{n-\alpha}} - \int\limits_{B(x_{+},r)} \frac{d\nu(z)}{|z-y|^{n-\alpha}}\right) \geq r^{\alpha-n} \quad \text{for all } y \in B(x,r),$$

whence finding

$$1 \geq \|I_\alpha a\|_{L^{q,\lambda}_\alpha}^q \geq r^{q(\alpha-n)+\lambda-n} \mu\big(B(x,r)\big) \geq r^{-\beta} \mu\big(B(x,r)\big) \, \Rightarrow \, \|\|\mu\|_\beta < \infty.$$

On the other hand, assume $\|\|\mu\|\|_{\beta} < \infty$. Upon writing $f = \sum_{j=1}^{\infty} c_j a_j \in H^1$ and observing that the argument for the proof of Theorem 1.2 (iii) actually reveals

$$\|I_{\alpha}a_j\|_{L^{q,\lambda}_u} \lesssim \|\|\mu\|\|_{\beta}^{\frac{1}{q}},$$

one utilises the Minkowski inequality to achieve

$$\|I_{\alpha}f\|_{L^{q,\lambda}_{\mu}} \leq \sum_{j} |c_{j}| \|I_{\alpha}a_{j}\|_{L^{q,\lambda}_{\mu}} \leq \||\mu||_{\beta}^{\frac{1}{q}} \sum_{j} |c_{j}| \approx \||\mu||_{\beta}^{\frac{1}{q}} \|f\|_{H^{1}}.$$

Surprisingly, this last estimation for $\mu = \nu$, i.e., $I_{\alpha}H^1 \subset L^{\frac{\lambda}{n-\alpha},\lambda}$, is an associate form of [26, Theorem 2], which implies $I_{\alpha}H^1 \subset H^{p,p'\alpha}$, since $\lambda = p'\alpha$ amounts to $p = \lambda/(\lambda - \alpha)$. When $\lambda = n$, both go back to the wellknown Stein-Weiss embedding $I_{\alpha}H^1 \subset L^{\frac{n}{n-\alpha}}$, see, e.g., [13, 37, 40, 41].

Funding: The author is in part supported by NSERC of Canada (202979463102000).

References

- D. R. Adams, Traces of potentials arising from translation invariant operators, Ann. Sc. Norm, Super. Pisa Cl. Sci. (3) 25 (1971), 203-217.
- [2] D. R. Adams, A note on Riesz potentials, Duke Math. J. 42 (1975), 765-778.
- [3] D. R. Adams, A note on Choquet integrals with respect to Hausdorff capacity, in: Function Spaces and Applications (Lund 1986), Lecture Notes in Math. 1302, Springer, Berlin (1988), 115-124.
- [4] D. R. Adams, A sharp inequality of J. Moser for higher order derivatives, Ann. of Math. (2) 128 (1988), 385-398.
- D. R. Adams and L. I. Hedberg, Function Spaces and Potential Theory, Springer, Berlin, 1996.
- [6] D. R. Adams and J. Xiao, Nonlinear analysis on Morrey spaces and their capacities, Indiana Univ. Math. J. 53 (2004), 1629-1663.
- [7] D. R. Adams and J. Xiao, Morrey potentials and harmonic maps, Comm. Math. Phys. 308 (2011), 439-456.
- [8] D. R. Adams and J. Xiao, Morrey spaces in harmonic analysis, Ark. Mat. 50 (2012), 201–230.
- [9] D. R. Adams and J. Xiao, Erratum to: "Morrey potentials and harmonic maps", Comm. Math. Phys. 339 (2015), 769-771.
- [10] D. R. Adams and J. Xiao, Restrictions of Riesz-Morrey potentials, Ark. Mat. 54 (2016), 201-231.
- [11] H. Beirão da Veiga, On nonlinear potential theory, and regular boundary points, for the p-Laplacian in N space variables, Adv. Nonlinear Anal. 3 (2014), 45-67.
- [12] C. Bennett and R. Sharpley, Interpolation of Operators, Academic Press, New York, 1988.
- [13] J. Bourgain and H. Brezis, Sur l'équation divu = f, C. R. Math. Acad. Sci. Paris **334** (2002), 973–976.
- [14] S. Campanato, Equazioni ellittiche non variazionali a coefficienti continui, Ann. Mat. Pura Appl. (4) 86 (1970), 125-154.
- [15] S. Campanato, A Cordes type condition for nonlinear non-variational systems, Rend. Accad. Naz. Sci. XL Mem. Mat. Appl. (5) **13** (1989), 307–321.
- [16] H. O. Cordes, Über die erste Randwertaufgabe bei quasilinearen Differentialgleichungen zweiter Ordnung in mehr als zwei Variablen, Math. Ann. 131 (1956), 278-312.
- [17] M. Cwikel and C. Fefferman, The canonical seminorm on weak L¹, Studia Math. **78** (1984), 275–278.
- [18] L. C. Evans and R. F. Gariepy, Measure Theory and Fine Properties of Functions, CRC Press, Boca Raton, 1992.
- [19] C. Fefferman and E. M. Stein, H^p spaces of several variables, Acta Math. 129 (1972), 137–193.

- [20] L. C. F. Ferreira, On a bilinear estimate in weak-Morrey spaces and uniquenness for Navier-Stokes equations, J. Math. Pures Appl. (9) 105 (2016), 228-247.
- [21] M. Giaquinta, Multiple Integrals in the Calculus of Variations and Nonlinear Elliptic Systems, Ann. of Math. Stud., Princeton University Press, Princeton, 1983.
- [22] D. Gilbarg and N. S. Trudinger, Elliptic Partial Differential Equations of Second Order, Springer, Berlin, 2001.
- [23] L. Grafakos, Classical and Modern Fourier Analysis, Pearson Education, Upper Saddle River, 2004.
- [24] T. Izumi, Y. Sawano and H. Tanaka, Littlewood-Paley theory for Morrey spaces, Rev. Mat. Complut. 28 (2015), 411-447.
- [25] F. John and L. Nirenberg, On functions of bounded mean oscillation, Comm. Pure Appl. Math. 14 (1961), 415-426.
- [26] E. A. Kalita, Dual Morrey spaces, Dokl. Math. 58 (1998), 85–87.
- [27] E. A. Kalita, Morrey regularity of solutions of nonlinear elliptic systems of arbitrary order under restrictions on the modulus of ellipticity, Ukrainian Math. J. 54 (2002), 1760-1777.
- [28] P. G. Lemarié-Rieusset, On some classes of time-periodic solutions for the Navier-Stokes equations in the whole space, SIAM J. Math. Anal. 47 (2015), 1022-1043.
- [29] J. Malý and W. P. Ziemer, Fine Regularity of Solutions of Elliptic Partial Differential Equations, Math. Surveys Monogr. 51, American Mathematical Society, Providence, 1997.
- [30] V. Maz'ya, Sobolev Spaces with Applications to Elliptic Partial Differential Equations, Springer, Berlin, 2011.
- [31] Y. Meyer, Wavelets, paraproducts and Navier-Stokes equations, in: Current Developments in Mathematics, International Press, Cambridge (1996), 105-212.
- [32] C. Miao and B. Yuan, Weak Morrey spaces and strongsolutions to the Navier-Stokes equations, Sci. China Ser. A 50 (2007), 1401-1417.
- [33] C. B. Morrey, Multiple Integrals in the Calculus of Variations, Springer, Berlin, 1966.
- [34] L. Nirenberg, On nonlinear elliptic partial differential equations and Hölder continuity, Comm. Pure Appl. Math. 6 (1953), 103-156.
- [35] V. Rădulescu and D. Repovš, Partial Differential Equations with Variable Exponents: Variational Methods and Qualitative Analysis, Monogr. Res. Notes Math., CRC Press, Boca Raton, 2015.
- [36] Y. Sawano and H. Tanaka, The Fatou property of block spaces, J. Math. Sci. Univ. Tokyo 22 (2015), no. 3, 663-683.
- [37] A. Schikorra, D. Spector and J. V. Schaftingen, An L¹-type estimate for Riesz potentials, preprint (2014), http://arxiv.org/abs/1411.2318.
- [38] R. Schoen and S. T. Yau, Lectures on Harmonic Maps, Conf. Proc. Lecture Notes Geom. Topol. II, International Press, Somerville, 1997.
- [39] J. Serrin, A remark on the Morrey potential, in: Control Methods in PDE-Dynamical Systems (Snowbird 2005), Contemp. Math. 426, American Mathematical Society, Providence (2007), 307-315.
- [40] E. M. Stein and G. Weiss, On the theory of harmonic functions of several variables. I: Theory of H^p -spaces, Acta Math. 103 (1960), 25-62.
- [41] R. S. Strichartz, H^p Sobolev spaces, Collog. Math. **60–61** (1990), 129–139.
- [42] T. Tao and G. Tian, A singularity removal theorem for Yang-Mills fields in higher dimensions, J. Amer. Math. Soc. 17 (2004), 557-593.
- [43] M. E. Tayler, Tools for PDE. Pseudodifferential Operators, Paradifferential Operators, and Layer Potentials, Math. Surveys Monogr. 81, American Mathematical Society, Providence, 2000.
- [44] A. Torchinsky, Real Variables, Addison-Wesley, Redwood City, 1988.
- [45] J. Xiao, A new perspective on the Riesz potential, Adv. Nonlinear Anal. (2016), DOI 10.1515/anona-2015-0183.
- [46] C. T. Zorko, Morrey spaces, Proc. Amer. Math. Soc. 98 (1986), 586-592.