

## Research Article

## Special Issue: In honor of David Jerison

Rupert L. Frank\*

# Sharp inequalities for coherent states and their optimizers

<https://doi.org/10.1515/ans-2022-0050>

received September 27, 2022; accepted January 20, 2023

**Abstract:** We are interested in sharp functional inequalities for the coherent state transform related to the Wehrl conjecture and its generalizations. This conjecture was settled by Lieb in the case of the Heisenberg group, Lieb and Solovej for  $SU(2)$ , and Kulikov for  $SU(1, 1)$  and the affine group. In this article, we give alternative proofs and characterize, for the first time, the optimizers in the general case. We also extend the recent Faber-Krahn-type inequality for Heisenberg coherent states, due to Nicola and Tilli, to the  $SU(2)$  and  $SU(1, 1)$  cases. Finally, we prove a family of reverse Hölder inequalities for polynomials, conjectured by Bodmann.

**Keywords:** functional inequalities, coherent states, inequalities for analytic functions, representations of Lie groups, isoperimetric inequality

**MSC 2020:** Primary 39B62, Secondary 22E70, 30H10, 30H20, 81R30

## 1 Introduction and main results

Coherent states appear in various areas of pure and applied mathematics, including mathematical physics, signal and image processing, and semiclassical and microlocal analysis. Some background can be found, for instance, in [33, 39, 40]. Here, we are interested in sharp functional inequalities for coherent state transforms.

To motivate the questions we are interested in, let us recall Wehrl's conjecture [44] and its resolution by Lieb [23]. Following Schrödinger, Bargmann, Segal, Glauber, Klauder, and others, we consider a certain family of normalized Gaussian functions  $\psi_{p,q} \in L^2(\mathbb{R})$ , parametrized by  $p, q \in \mathbb{R}$ . Explicitly,

$$\psi_{p,q}(x) := (\pi\hbar)^{-\frac{1}{4}} e^{-\frac{1}{2\hbar}(x-q)^2 + \frac{1}{\hbar}ipx} \quad \text{for all } x \in \mathbb{R},$$

where  $\hbar > 0$  is a fixed constant. For a nonnegative operator  $\rho$  in  $L^2(\mathbb{R})$  with  $\text{Tr}\rho = 1$ , one considers the function

$$(p, q) \mapsto \langle \psi_{p,q}, \rho \psi_{p,q} \rangle,$$

known as the Husimi function, the covariant symbol, or the lower symbol. Thus, to a quantum state  $\rho$  in  $L^2(\mathbb{R})$ , one associates a function defined on the classical phase space  $\mathbb{R}^2$ . Wehrl [44] was interested in the entropy-like quantity

---

To David Jerison, in admiration, on the occasion of his 70th birthday.

---

\* **Corresponding author: Rupert L. Frank**, Mathematisches Institut, Ludwig-Maximilians Universität München, Theresienstr. 39, 80333 München, Germany; Munich Center for Quantum Science and Technology, Schellingstr. 4, 80799 München, Germany; Mathematics 253-37, Caltech, Pasadena, CA 91125, USA, e-mail: r.frank@lmu.de

$$- \iint_{\mathbb{R} \times \mathbb{R}} \langle \psi_{p,q}, \rho \psi_{p,q} \rangle \ln \langle \psi_{p,q}, \rho \psi_{p,q} \rangle dp dq,$$

showed that it is positive, and conjectured that its minimum value occurs when  $\rho = |\psi_{p_0,q_0}\rangle\langle\psi_{p_0,q_0}|$  for some  $p_0, q_0 \in \mathbb{R}$ . That this is indeed the case was shown in a celebrated article by Lieb [23]. Lieb's proof was based on the sharp forms of the Young and the Hausdorff-Young inequalities and showed more generally that, for power functions  $\Phi(s) = s^r$  with  $r \geq 1$ , the quantity

$$\iint_{\mathbb{R} \times \mathbb{R}} \Phi(\langle \psi_{p,q}, \rho \psi_{p,q} \rangle) dp dq \quad (1)$$

is maximal for  $\rho$  as above. The result for  $\Phi(s) = s \ln s$  then follows by differentiating at  $r = 1$ , noting that the value for  $\Phi(s) = s$  is a constant independent of  $\rho$ .

In [10], Carlen gave an alternative proof of Lieb's result, both for  $\Phi(s) = s^r$ ,  $r \geq 1$ , and  $\Phi(s) = s \ln s$ , and characterized the cases of equality. He also extended the result to  $\Phi(s) = -s^r$  with  $0 < r < 1$ , again including a characterization of cases of equality. Carlen's proof is based on the logarithmic Sobolev inequality and an identity for analytic functions. For yet another proof in the logarithmic case, see [27]. For an interesting recent generalization of Lieb's result, see [13].

In [24], Lieb and Solovej extended the earlier results and showed what they called the generalized Wehrl conjecture. Namely, for *any* convex function  $\Phi$  on  $[0, 1]$ , the quantity (1) is maximal if  $\rho = |\psi_{p_0,q_0}\rangle\langle\psi_{p_0,q_0}|$  for some  $p_0, q_0 \in \mathbb{R}$ . The Lieb-Solovej proof proceeds by a limiting argument, based on sharp inequalities for  $SU(2)$  coherent states discussed below. Because of the limiting process, it does not provide a characterization of the cases of equality. Carlen's analysis is based on differentiating the power function  $\Phi(s) = s^r$  with respect to the exponent  $r$  and using the logarithmic Sobolev inequality for the resulting quantity. We do not know how to adopt this method to deal with general convex functions  $\Phi$ .

Our first main result in this article gives an alternative proof of the theorem of Lieb and Solovej and includes a new characterization of the cases of equality.

**Theorem 1.** *Let  $\Phi : [0, 1] \rightarrow \mathbb{R}$  be convex. Then,*

$$\sup \left\{ \iint_{\mathbb{R} \times \mathbb{R}} \Phi(|\langle \psi_{p,q}, \psi \rangle|^2) dp dq : \psi \in L^2(\mathbb{R}), \|\psi\|_{L^2(\mathbb{R})} = 1 \right\} = 2\pi\hbar \int_0^1 \Phi(s) \frac{ds}{s}$$

*and the supremum is attained for  $\psi = e^{i\theta}\psi_{p_0,q_0}$  with some  $p_0, q_0 \in \mathbb{R}$ ,  $\theta \in \mathbb{R}/2\pi\mathbb{Z}$ . If  $\Phi$  is not linear and if the supremum is finite, then it is attained only for such  $\psi$ .*

Note that the value of the double integral with  $\psi = e^{i\theta}\psi_{p_0,q_0}$  does not depend on  $p_0, q_0 \in \mathbb{R}$ ,  $\theta \in \mathbb{R}/2\pi\mathbb{Z}$ . It may or may not be finite, depending on  $\Phi$ . For finiteness, it is necessary that  $\lim_{s \rightarrow 0^+} \Phi(s) = 0$ .

Under a slightly stronger assumption on  $\Phi$ , we can extend the characterization of cases of equality to density matrices.

**Corollary 2.** *Let  $\Phi : [0, 1] \rightarrow \mathbb{R}$  be convex. Then,*

$$\sup \left\{ \iint_{\mathbb{R} \times \mathbb{R}} \Phi(\langle \psi_{p,q}, \rho \psi_{p,q} \rangle) dp dq : \rho \geq 0 \text{ on } L^2(\mathbb{R}), \text{Tr} \rho = 1 \right\} = 2\pi\hbar \int_0^1 \Phi(s) \frac{ds}{s}$$

*and the supremum is attained for  $\rho = |\psi_{p_0,q_0}\rangle\langle\psi_{p_0,q_0}|$  with some  $p_0, q_0 \in \mathbb{R}$ . If  $\Phi$  is strictly convex and if the supremum is finite, then it is attained only for such  $\rho$ .*

**Remark 3.** The statement and proof of Theorem 1 and Corollary 2 extend, with minor changes, to the case of higher dimensions. We omit the details.

Coherent states are often closely related to representations of an underlying Lie group. The coherent states discussed so far are related to the Heisenberg group. In his article, containing the proof of Wehrl's conjecture, Lieb conjectured that the analog of Wehrl's conjecture also holds for Bloch coherent states, that is, for a family of coherent states related to  $SU(2)$ . After some partial results in [9,38], this conjecture was finally solved by Lieb and Solovej in [24]; see also [25] for a partially alternate proof. Again, they prove a generalized version of Lieb's conjecture involving general convex functions  $\Phi$ . However, they employ a limiting argument, and therefore, their article does not characterize the cases of equality. Our second main result settles this open question by showing that, indeed, equality is only attained by rank one projections onto a coherent state.

Let us be more specific. As is well known (see, e.g., [19, Chapter II] and [41, Section VIII.4]), the nontrivial irreducible representations of  $SU(2)$  are labeled by  $J \in \frac{1}{2}\mathbb{N}$ , where  $2J + 1$  is the dimension of the representation and  $\mathbb{N} = \{1, 2, 3, \dots\}$ . Let  $\mathcal{H}$  be a  $(2J + 1)$ -dimensional representation space. Then, there are operators  $S_1, S_2$ , and  $S_3$  on  $\mathcal{H}$  satisfying  $[S_1, S_2] = iS_3$  and cyclically, representing the generators of  $SU(2)$ . For any  $\omega = (\omega_1, \omega_2, \omega_3) \in \mathbb{S}^2 \subset \mathbb{R}^3$ , the operator  $\omega \cdot S = \omega_1 S_1 + \omega_2 S_2 + \omega_3 S_3$  has minimal eigenvalue  $-J$ , and this eigenvalue is nondegenerate. We choose  $\psi_\omega \in \mathcal{H}$  as a corresponding normalized eigenvector. It is unique up to a phase, but since we are only interested in the state  $|\psi_\omega\rangle\langle\psi_\omega|$ , this choice of the phase is irrelevant for us. This defines the Bloch coherent states. (We follow here the convention in [33]; other definitions are based on the *maximal* eigenvalue  $+J$ , but this leads to the same family of coherent states, just interchanging  $\omega$  and  $-\omega$ .)

**Theorem 4.** Let  $J \in \frac{1}{2}\mathbb{N}$  and consider an irreducible  $(2J + 1)$ -dimensional representation of  $SU(2)$  on  $\mathcal{H}$ . Let  $\Phi : [0, 1] \rightarrow \mathbb{R}$  be convex. Then,

$$\sup \left\{ \int_{\mathbb{S}^2} \Phi(|\langle\psi_\omega, \psi\rangle|^2) d\omega : \psi \in \mathcal{H}, \|\psi\|_{\mathcal{H}} = 1 \right\} = \frac{4\pi}{2J} \int_0^1 \Phi(s) s^{\frac{1}{2J}-1} ds$$

and the supremum is attained for  $\psi = e^{i\theta}\psi_{\omega_0}$  with some  $\omega_0 \in \mathbb{S}^2$ ,  $\theta \in \mathbb{R}/2\pi\mathbb{Z}$ . If  $\Phi$  is not affine linear, then it is attained only for such  $\psi$ .

Note that the value of the integral with  $\psi = e^{i\theta}\psi_{\omega_0}$  does not depend on  $\omega_0 \in \mathbb{S}^2$ ,  $\theta \in \mathbb{R}/2\pi\mathbb{Z}$ . Since  $\Phi$  is bounded, the supremum in the theorem is always finite, in contrast to the situation of Theorem 1.

**Corollary 5.** Let  $J \in \frac{1}{2}\mathbb{N}$  and consider an irreducible  $(2J + 1)$ -dimensional representation of  $SU(2)$  on  $\mathcal{H}$ . Let  $\Phi : [0, 1] \rightarrow \mathbb{R}$  be convex. Then,

$$\sup \left\{ \int_{\mathbb{S}^2} \Phi(\langle\psi_\omega, \rho\psi_\omega\rangle) d\omega : \rho \geq 0 \text{ on } \mathcal{H}, \text{Tr}\rho = 1 \right\} = \frac{4\pi}{2J} \int_0^1 \Phi(s) s^{\frac{1}{2J}-1} ds$$

and the supremum is attained for  $\rho = |\psi_{\omega_0}\rangle\langle\psi_{\omega_0}|$  with some  $\omega_0 \in \mathbb{S}^2$ . If  $\Phi$  is strictly convex, then it is attained only for such  $\rho$ .

Our third main result concerns coherent states for certain representations of  $SU(1, 1)$ . After initial results in [3,6,26], the analog of Wehrl's conjecture was recently settled by Kulikov in [20], again for general convex functions  $\Phi$ . (In fact, slightly less than convexity is required in [20].) Kulikov [20, Remark 4.3] also characterized optimizers in the case where  $\Phi$  is strictly convex. We extend this to the case where  $\Phi$  is not linear.

All nontrivial representations of  $SU(1, 1)$  are infinite-dimensional. Its nontrivial irreducible unitary representations consist of discrete, principal, and complementary series, as well as limits of the discrete series; see, e.g., [19, Chapters II and XVI; also (2.20)]. Here, we are only interested in one of the two discrete series. The results for the other one can be deduced from the results below by applying complex conjugation at the appropriate places.

Following the notation in [5], the discrete series representation under consideration is labeled by  $K \in \frac{1}{2}\mathbb{N} \setminus \{\frac{1}{2}\} = \{1, \frac{3}{2}, 2, \dots\}$ . Let  $\mathcal{H}$  be a corresponding representation space. The generators of the Lie algebra of  $SU(1, 1)$  give rise to operators  $K_0, K_1$ , and  $K_2$  in  $\mathcal{H}$  satisfying

$$[K_1, K_2] = -iK_0, \quad [K_2, K_0] = iK_1, \quad [K_0, K_1] = iK_2.$$

Moreover, one has

$$K_0^2 - K_1^2 - K_2^2 = K(K - 1),$$

where  $K$  is the number labeling the representations. (There are also representations of  $SU(1, 1)$  corresponding to  $K = \frac{1}{2}$ , called limits of the discrete series, but their coherent state transforms are in some sense degenerate; see Section 4.4. We also briefly discuss the case of arbitrary real  $K > \frac{1}{2}$  after Corollary 7.)

For any  $(n_0, n_1, n_2) \in \mathbb{R}^3$  with  $n_0^2 - n_1^2 - n_2^2 = 1$  and  $n_0 > 0$ , the operator  $n_0 K_0 - n_1 K_1 - n_2 K_2$  has minimal eigenvalue  $K$  and this eigenvalue is simple. Therefore, we can choose a corresponding normalized eigenvector (which is unique up to a phase). It is convenient to label this vector not by  $(n_0, n_1, n_2)$  but rather by  $z \in \mathbb{D}$ , the open unit disk in  $\mathbb{C}$ , using the parametrization

$$(n_0, n_1 + in_2) = \left( \frac{1 + |z|^2}{1 - |z|^2}, \frac{2z}{1 - |z|^2} \right).$$

In this way, we obtain a family of vectors  $\psi_z$ ,  $z \in \mathbb{D}$ , giving rise to coherent states for a discrete series representation of  $SU(1, 1)$ .

In what follows, we denote by  $dA(z) = dx dy$  the two-dimensional Lebesgue measure on  $\mathbb{C}$ .

**Theorem 6.** *Let  $K \in \frac{1}{2}\mathbb{N} \setminus \{\frac{1}{2}\}$  and consider the irreducible discrete series representation of  $SU(1, 1)$  on  $\mathcal{H}$  corresponding to  $K$ . Let  $\Phi : [0, 1] \rightarrow \mathbb{R}$  be convex. Then,*

$$\sup \left\{ \int_{\mathbb{D}} \Phi(|\langle \psi_z, \psi \rangle|^2) \frac{dA(z)}{(1 - |z|^2)^2} : \psi \in \mathcal{H}, \|\psi\|_{\mathcal{H}} = 1 \right\} = \frac{\pi}{2K} \int_0^1 \Phi(s) s^{-\frac{1}{2K}-1} ds$$

and the supremum is attained for  $\psi = e^{i\theta} \psi_{z_0}$  with some  $z_0 \in \mathbb{D}$ ,  $\theta \in \mathbb{R}/2\pi\mathbb{Z}$ . If  $\Phi$  is not linear and if the supremum is finite, then it is attained only for such  $\psi$ .

Note that the value of the integral with  $\psi = e^{i\theta} \psi_{z_0}$  does not depend on  $z_0 \in \mathbb{D}$ ,  $\theta \in \mathbb{R}/2\pi\mathbb{Z}$ . It may or may not be finite, depending on  $\Phi$ . For finiteness, it is necessary that  $\lim_{s \rightarrow 0^+} \Phi(s) = 0$ .

Theorem 6 proves the uniqueness part of a conjecture of Lieb and Solovej [26, Conjecture 5.2]. As we mentioned before, the inequality part is due to Kulikov [20].

**Corollary 7.** *Let  $K \in \frac{1}{2}\mathbb{N} \setminus \{\frac{1}{2}\}$  and consider the irreducible discrete series representation of  $SU(1, 1)$  on  $\mathcal{H}$  corresponding to  $K$ . Let  $\Phi : [0, 1] \rightarrow \mathbb{R}$  be convex. Then,*

$$\sup \left\{ \int_{\mathbb{D}} \Phi(\langle \psi_z, \rho \psi_z \rangle) \frac{dA(z)}{(1 - |z|^2)^2} : \rho \geq 0 \text{ on } \mathcal{H}, \text{Tr} \rho = 1 \right\} = \frac{\pi}{2K} \int_0^1 \Phi(s) s^{-\frac{1}{2K}-1} ds$$

and the supremum is attained for  $\rho = |\psi_{z_0}\rangle\langle\psi_{z_0}|$  with some  $z_0 \in \mathbb{D}$ . If  $\Phi$  is strictly convex and if the supremum is finite, then it is attained only for such  $\rho$ .

For every real  $K > \frac{1}{2}$ , there is an irreducible representation of the Lie algebra generated by  $K_0, K_1$ , and  $K_2$  satisfying the above relations. For  $K \notin \frac{1}{2}\mathbb{N}$ , such a representation does not come from the Lie group  $SU(1, 1)$  – recall that this group is not simply connected. However, it comes from a representation of the covering group of  $SU(1, 1)$  [32], and we could prove sharp inequalities for the corresponding coherent states. From an

analytic point of view, this would lead to the same problem as coherent states for the affine group, which we discuss next.

The affine group (in one space dimension), also known as the  $(aX + b)$ -group, has two nontrivial irreducible unitary representations [2,12]. Again, we focus on a single one since the results for the other one can be obtained by appropriate complex conjugations. What distinguishes the affine group from the above cases of the Heisenberg group,  $SU(2)$ , and  $SU(1, 1)$  is that different choices of an extremal weight vector lead to inequivalent coherent state transforms.

We fix a parameter  $\beta > \frac{1}{2}$ , emphasizing that this parameter does not label the representation, but rather the choice of the extremal weight vector. We consider the following family of normalized functions  $\psi_{a,b} \in L^2(\mathbb{R}_+)$ ,  $\mathbb{R}_+ = (0, \infty)$ , parametrized by  $a \in \mathbb{R}_+$ ,  $b \in \mathbb{R}$ ,

$$\psi_{a,b}(x) := 2^\beta \Gamma(2\beta)^{-\frac{1}{2}} a^\beta x^{\beta-\frac{1}{2}} e^{-ax+ibx} \quad \text{for all } x \in \mathbb{R}_+.$$

(Here, we follow the convention in [12]. What we call  $\beta$  is called  $\alpha - \frac{1}{2}$  in [26], and they do not normalize  $\psi_{a,b}$  in  $L^2(\mathbb{R}_+)$  but choose a different, natural normalization.)

**Theorem 8.** *Let  $\beta > \frac{1}{2}$  and let  $\Phi : [0, 1] \rightarrow \mathbb{R}$  be convex. Then,*

$$\sup \left\{ \iint_{\mathbb{R}_+ \times \mathbb{R}} \Phi(|\langle \psi_{a,b}, \psi \rangle|^2) \frac{da db}{a^2} : \psi \in L^2(\mathbb{R}_+), \|\psi\|_{L^2(\mathbb{R}_+)} = 1 \right\} = \frac{2\pi}{\beta} \int_0^1 \Phi(s) s^{-\frac{1}{2\beta}-1} ds$$

*and the supremum is attained for  $\psi = e^{i\theta} \psi_{a_0, b_0}$  with some  $a_0 \in \mathbb{R}_+$ ,  $b_0 \in \mathbb{R}$ ,  $\theta \in \mathbb{R}/2\pi\mathbb{Z}$ . If  $\Phi$  is not linear and if the supremum is finite, then it is attained only for such  $\psi$ .*

Note that the value of the double integral with  $\psi = e^{i\theta} \psi_{a_0, b_0}$  does not depend on  $a_0 \in \mathbb{R}_+$ ,  $b_0 \in \mathbb{R}$ , and  $\theta \in \mathbb{R}/2\pi\mathbb{Z}$ . It may or may not be finite, depending on  $\Phi$ . For finiteness, it is necessary that  $\lim_{s \rightarrow 0^+} \Phi(s) = 0$ .

Theorem 8 settles the equality part of a conjecture of Lieb and Solovej [26, Conjecture 3.1]. For strictly convex  $\Phi$ , it had been settled earlier in [20, Remark 4.3]. Clearly, the assumption that  $\Phi$  is not linear is optimal, because otherwise the supremum is attained for any  $\psi \in L^2(\mathbb{R}_+)$ .

We note that Theorem 8 has a version for  $\beta = \frac{1}{2}$ ; see Remark 20.

**Corollary 9.** *Let  $\beta > \frac{1}{2}$  and let  $\Phi : [0, 1] \rightarrow \mathbb{R}$  be convex. Then,*

$$\sup \left\{ \iint_{\mathbb{R}_+ \times \mathbb{R}} \Phi(\langle \psi_{a,b}, \rho \psi_{a,b} \rangle) \frac{da db}{a^2} : \rho \geq 0 \text{ on } L^2(\mathbb{R}), \text{Tr} \rho = 1 \right\} = \frac{2\pi}{\beta} \int_0^1 \Phi(s) s^{-\frac{1}{2\beta}-1} ds$$

*and the supremum is attained for  $\rho = |\psi_{a_0, b_0}\rangle \langle \psi_{a_0, b_0}|$  with some  $a_0 \in \mathbb{R}_+$ ,  $b_0 \in \mathbb{R}$ . If  $\Phi$  is strictly convex and if the supremum is finite, then it is attained only for such  $\rho$ .*

This concludes the description of our main results, but we would like to draw the reader's attention also to Sections 3 and 5 where we prove, respectively, sharp reverse Hölder inequalities for analytic functions, thereby settling a conjecture of Bodmann [9], and optimal Faber-Krahn-type inequalities for coherent state transforms.

The method that we will be using is that from a recent, beautiful article by Kulikov [20]. He developed this method to solve the Lieb-Solovej conjectures for  $SU(1, 1)$  and the affine group. Here, we show that it can be adapted to deal with the Heisenberg and the  $SU(2)$  cases. We also push the characterization of optimizers a bit further than in [20], thus leading to the optimal results in Theorems 1, 4, 6, and 8.

Kulikov's article in turn seems to be inspired by an equally beautiful recent article by Nicola and Tilli [29]. They were the first, as far as we know, to use the isoperimetric inequality in connection with the coherent state transform to obtain optimal functional inequalities. (Talenti [43] used a closely related method for comparison theorems for solutions of PDEs.) Kulikov proved his results by instead using the

isoperimetric inequality in hyperbolic space, and we will prove Theorem 4 by using that on the sphere. While it is tempting to try to use the same method for more general groups, an obstacle will have to be overcome (see Section 4.6).

Nicola and Tilli proved Faber-Krahn-type inequalities for the Heisenberg coherent states. We will show that their main result (at least without the characterization of the cases of equality) follows from Theorem 1, and we will use this idea to prove analogs of their results for coherent states based on  $SU(2)$ ,  $SU(1, 1)$ , and the affine group (see Section 5). For further developments started by [29], see, for instance, [18, 30, 35].

After this article was submitted for publication, we learned that Kulikov et al. had independently obtained similar results with similar techniques. These results have appeared in preprint form [21].

Thanks are due to Eric Carlen, Elliott Lieb, and Jan Philip Solovej for many discussions on the topics of this article.

It is my pleasure to dedicate this article to David Jerison on the occasion of his 70th birthday. His articles have been an inspiration for me, those on sharp inequalities [17] and others. I am particularly indebted to him for his remarks in the fall of 2008, which indirectly were a great motivation for work that eventually led to [14].

## 2 Inequalities for analytic functions

The main ingredient behind the results in the previous section are sharp inequalities for analytic functions and the characterization of their optimizers, which we discuss in this section.

### 2.1 Definitions and main result

There are three different types of inequalities, corresponding to the cases of the Heisenberg group,  $SU(2)$ , and  $SU(1, 1)$ . We refer to these different scenarios as Cases 1, 2, and 3. In Cases 2 and 3, there is a parameter  $J \in \frac{1}{2}\mathbb{N}$  and  $\alpha > 1$ , respectively, that is fixed in what follows.

In Case 1, we consider functions from the Fock space  $\mathcal{F}^2(\mathbb{C})$ , that is, entire functions  $f$  satisfying

$$\|f\|_{\mathcal{F}^2} := \left( \int_{\mathbb{C}} |f(z)|^2 e^{-\pi|z|^2} dA(z) \right)^{1/2} < \infty.$$

We recall that we write  $dA(z) = dx dy$  for the two-dimensional Lebesgue measure. In Case 2, we consider functions in  $\mathcal{P}_{2J}$ , that is, polynomials  $f$  of degree  $\leq 2J$  endowed with the norm

$$\|f\|_{\mathcal{P}_{2J}} := \left( \frac{2J+1}{\pi} \int_{\mathbb{C}} |f(z)|^2 (1 + |z|^2)^{-2J-2} dA(z) \right)^{1/2}.$$

This norm is finite for any  $f \in \mathcal{P}_{2J}$ . In Case 3, we consider functions from the weighted Bergman space  $A_{\alpha}^2(\mathbb{D})$ , that is, analytic functions  $f$  on the disk  $\mathbb{D}$  satisfying

$$\|f\|_{A_{\alpha}^2} := \left( \frac{\alpha-1}{\pi} \int_{\mathbb{D}} |f(z)|^2 (1 - |z|^2)^{\alpha-2} dA(z) \right)^{1/2} < \infty.$$

To treat all cases simultaneously, we set

$$\mathcal{H} := \begin{cases} \mathcal{F}^2(\mathbb{C}) & \text{in Case 1,} \\ \mathcal{P}_{2J} & \text{in Case 2,} \\ \mathcal{A}_\alpha^2(\mathbb{D}) & \text{in Case 3,} \end{cases}$$

and denote the norm in  $\mathcal{H}$  by  $|||\cdot|||$ .

Thus, the set on which the relevant functions are defined is

$$\Omega := \begin{cases} \mathbb{C} & \text{in Cases 1 and 2,} \\ \mathbb{D} & \text{in Case 3,} \end{cases}$$

and the relevant measure is

$$dm(z) := \begin{cases} dA(z) & \text{in Case 1,} \\ \pi^{-1}(1 + |z|^2)^{-2}dA(z) & \text{in Case 2,} \\ \pi^{-1}(1 - |z|^2)^{-2}dA(z) & \text{in Case 3.} \end{cases}$$

To a function  $f \in \mathcal{H}$ , we associate a function  $u_f$  on  $\Omega$ , defined by

$$u_f(z) := \begin{cases} |f(z)|e^{-\frac{\pi}{2}|z|^2} & \text{in Case 1,} \\ |f(z)|(1 + |z|^2)^{-J} & \text{in Case 2,} \\ |f(z)|(1 - |z|^2)^{\alpha/2} & \text{in Case 3.} \end{cases} \quad (2)$$

The problem that we are interested in is to maximize, given a convex function  $\Phi$ , the quantity

$$\int_{\Omega} \Phi(u_f(z)^2) dm(z)$$

over all  $f \in \mathcal{H}$  with  $|||f||| = 1$ . Our main result will characterize the set of  $f$ 's for which this supremum is attained. This set  $\mathcal{M} \subset \{f \in \mathcal{H} : |||f||| = 1\}$  is defined as follows. In Case 1, we consider the functions  $F_w$ , parametrized by  $w \in \mathbb{C}$ , given as

$$F_w(z) := e^{-\frac{\pi}{2}|w|^2 + \pi \bar{w}z} \quad \text{for all } z \in \mathbb{C}$$

and set

$$\mathcal{M} := \{e^{i\theta} F_w : w \in \mathbb{C}, \theta \in \mathbb{R}/2\pi\mathbb{Z}\}.$$

In Case 2, we consider the functions  $F_w$ , parametrized by  $w \in \mathbb{C} \cup \{\infty\}$ , given as

$$F_w(z) := \frac{(1 + \bar{w}z)^{2J}}{(1 + |w|^2)^J}, \quad w \neq 0, \quad F_\infty(z) := z^{2J} \quad \text{for all } z \in \mathbb{C}$$

and set

$$\mathcal{M} := \{e^{i\theta} F_w : w \in \mathbb{C} \cup \{\infty\}, \theta \in \mathbb{R}/2\pi\mathbb{Z}\}.$$

In Case 3, we consider the function  $F_w$ , parametrized by  $w \in \mathbb{D}$ , given as

$$F_w(z) := \frac{(1 - |w|^2)^{\frac{\alpha}{2}}}{(1 - \bar{w}z)^\alpha} \quad \text{for all } z \in \mathbb{D}$$

and set

$$\mathcal{M} := \{e^{i\theta} F_w : w \in \mathbb{D}, \theta \in \mathbb{R}/2\pi\mathbb{Z}\}.$$

In each case, it can be verified that  $F_w \in \mathcal{H}$  and that  $|||F_w||| = 1$  for all  $w$  in the respective index set. Indeed, this can be seen by a direct computation for  $w = 0$ . For a general  $w$ , we use the fact that the functions  $u_{F_w}$  are equimeasurable, that is, for every  $\kappa > 0$ , the measure  $m(\{u_{F_w} > \kappa\})$  is independent of  $w$ . As a consequence of the equimeasurability, the norm in  $\mathcal{H}$  is independent of  $w$ . The equimeasurability in turn is a consequence

of the fact that  $F_w$  is obtained from, say,  $F_0$  by the action of the Heisenberg group,  $SU(2)$ , or  $SU(1, 1)$  in the respective cases, and of the invariance of the measure  $m$  under this action.

The following is the main result of this section.

**Theorem 10.** *Let  $\Phi : [0, 1] \rightarrow \mathbb{R}$  be convex. Then,*

$$\sup \left\{ \int_{\Omega} \Phi(u_f(z)^2) dm(z) : f \in \mathcal{H}, \|f\| = 1 \right\} = \begin{cases} \int_0^1 \Phi(s) s^{-1} ds & \text{in Case 1,} \\ (2J)^{-1} \int_0^1 \Phi(s) s^{\frac{1}{2J}-1} ds & \text{in Case 2,} \\ \alpha^{-1} \int_0^1 \Phi(s) s^{-\frac{1}{\alpha}-1} ds & \text{in Case 3,} \end{cases}$$

and the supremum is attained in  $\mathcal{M}$ . If  $\Phi$  is not affine linear and if the supremum is finite, then it is attained only in  $\mathcal{M}$ .

We will prove this theorem in Subsection 2.2, after establishing some lemmas.

## 2.2 Proof of Theorem 10

We begin the proof of Theorem 10 by recalling a simple and well-known bound on the supremum of  $u_f$ . This bound shows that  $u_f \leq 1$  for  $\|f\| = 1$ , so  $\Phi(u_f^2)$  appearing in the theorem is well defined for  $\Phi$  defined on  $[0, 1]$ . The characterization of the cases of equality in the inequality  $u_f \leq 1$  will eventually lead to the corresponding characterization in Theorem 10.

**Lemma 11.** *Let  $f \in \mathcal{H}$ . Then*

$$\|u_f\|_{L^\infty(\Omega)} \leq \|f\|$$

with equality if and only if either  $f = 0$  or  $\|f\|^{-1}f \in \mathcal{M}$ .

**Proof.** In Case 1, this is essentially [29, Proposition 2.1]. Indeed, there the inequality in the lemma is proved, and it is shown that  $u_f$  tends to zero at infinity. The latter fact, together with continuity, implies that there is a  $z \in \mathbb{C}$  such that  $\|u_f\|_{L^\infty(\mathbb{C})} = u_f(z)$ , and then [29, Proposition 2.1] implies  $\|f\|^{-1}f \in \mathcal{M}$ , provided  $f \neq 0$ .

In Case 2, the inequality is mentioned in [9, Paragraph after Remark 3.2]. Since the function  $u_f$  extends continuously to a function on the Riemann sphere  $\mathbb{C} \cup \{\infty\}$ , there is a  $z \in \mathbb{C} \cup \{\infty\}$  such that  $\|u_f\|_{L^\infty(\mathbb{C})} = u_f(z)$ , and then one obtains the equality condition from that in the Cauchy-Schwarz inequality.

In Case 3, the inequality and the fact that  $u_f$  tends to zero as  $|z| \rightarrow 1$  is mentioned in [20, (1.1) and the paragraph thereafter]. As in the other cases, from the latter fact, one can deduce the equality condition. Let us add some details concerning the facts mentioned in [20]. To carry out the proof of the inequality, one can, for instance, use [6, (5)] and argue as in [29, Proposition 2.1]. To deduce the vanishing of  $u_f$  on  $\partial D$ , one can observe that this is true when  $f$  is a polynomial and that those are dense in  $A_\alpha^2(D)$  by [6, (5)].  $\square$

We now come to the core of the proof of Theorem 10, which concerns a certain monotonicity of the measure of superlevel sets of  $u_f$ . In Case 3, the following lemma and its proof are a special case of [20, Theorem 2.1]. Our contribution is to extend the reasoning to Cases 1 and 2. (Indeed, similar arguments in the setting of Case 1 have already appeared in [29, Theorem 3.1]; in particular, inequality (3) is the same as [29, (3.17) combined with (3.10)]. Note, however, that the authors of [29] use Lieb's inequality as an ingredient



and do not reprove it using their method, in contrast to what we do here; see, in particular, [29, Theorem 5.2].) We also point out that arguments of this type are reminiscent of those of Talenti in [43].

**Lemma 12.** *Let  $f \in \mathcal{H}$  and  $\mu(\kappa) := m(\{u_f > \kappa\})$  for  $\kappa > 0$ . Then, the function*

$$\kappa \mapsto \begin{cases} -2 \ln \kappa - \mu(\kappa) & \text{in Case 1,} \\ \kappa^{-\frac{1}{\beta}}(1 - \mu(\kappa)) & \text{in Case 2,} \\ \kappa^{\frac{2}{\alpha}}(-1 - \mu(\kappa)) & \text{in Case 3} \end{cases}$$

*is nondecreasing on  $(0, \|u_f\|_\infty)$ . Moreover, if  $f \in \mathcal{M}$ , then this function is constant.*

**Proof.** Writing  $M := \|u_f\|_{L^\infty(\Omega)}$  for brevity, we observe that, by Sard's theorem (noting that  $u_f$  is real analytic on  $\Omega$  viewed as a subset of  $\mathbb{R}^2$ ), for almost every  $\kappa \in (0, M)$ ,  $\{u_f = \kappa\}$  is a smooth curve (or, possibly, a union thereof). Denote the one-dimensional Hausdorff measure on this curve by  $|dz|$ , we have, by the co-area formula,

$$\int_{\Omega} g(x, y) |\nabla u_f| dA(z) = \int_0^{t_0} \int_{\{u_f = \kappa\}} g(x, y) |dz| d\tau.$$

Let

$$\omega(z) := \frac{dm(z)}{dA(z)} = \begin{cases} 1 & \text{in Case 1,} \\ \pi^{-1}(1 + |z|^2)^{-2} & \text{in Case 2,} \\ \pi^{-1}(1 - |z|^2)^{-2} & \text{in Case 3.} \end{cases}$$

Taking  $g = |\nabla u_f|^{-1} \omega \mathbb{1}_{\{u_f > \kappa\}} \mathbb{1}_{\{|\nabla u_f| \neq 0\}}$  and noting that  $m(\{|\nabla u_f| = 0\}) = 0$  by real analyticity, we find, for any  $\kappa \in (0, M)$ ,

$$\mu(\kappa) = \int_{\kappa}^M \int_{\{u_f = \kappa\}} |\nabla u_f|^{-1} \omega |dz| d\tau.$$

Thus,  $\mu$  is absolutely continuous on compact subintervals of  $(0, M]$  and, for almost every  $\kappa \in (0, M)$ ,

$$\mu'(\kappa) = - \int_{\{u_f = \kappa\}} |\nabla u_f|^{-1} \omega |dz|.$$

For a curve  $\gamma$  in  $\Omega$  let us set

$$\ell(\gamma) := \int_{\gamma} \sqrt{\omega} |dz|.$$

In particular, for the level set  $\{u_f = \kappa\}$  we obtain, by the Schwarz inequality,

$$\ell(\{u_f = \kappa\})^2 \leq \int_{\{u_f = \kappa\}} |\nabla u_f|^{-1} \omega |dz| \int_{\{u_f = \kappa\}} |\nabla u_f| |dz|.$$

As we argued before, the first term on the right side is  $-\mu'(\kappa)$ . Let us consider the second term. Since the outer unit normal vector field  $\nu$  to  $\{u_f > \kappa\}$  on the boundary  $\{u_f = \kappa\}$  is given by  $-\nabla u_f / |\nabla u_f|$ , we have  $|\nabla u_f| = -\kappa \nu \cdot \nabla(\ln u_f)$ , and therefore, by Green's theorem,

$$\int_{\{u_f = \kappa\}} |\nabla u_f| |dz| = -\kappa \int_{\{u_f > \kappa\}} \Delta \ln u_f dA(z).$$

To compute the Laplacian of  $\ln u_f$ , we recall that  $f$  is a positive weight times the absolute value of an analytic function. On the set  $\{u_f > \kappa\}$ , the analytic function does not have zeros, so the logarithm of its absolute value is harmonic there. Thus, the Laplacian of  $\ln u_f$  coincides with the Laplacian of the logarithm of the weight. Explicitly,

$$\Delta \ln u_f = \begin{cases} -\frac{\pi}{2} \Delta |z|^2 = -2\pi & \text{in Case 1,} \\ -j \Delta \ln(1 + |z|^2) = -4j(1 + |z|^2)^{-2} & \text{in Case 2,} \\ \frac{\alpha}{2} \Delta \ln(1 - |z|^2) = -2\alpha(1 - |z|^2)^{-2} & \text{in Case 3.} \end{cases}$$

Note that the right side is equal to a constant multiple of  $\omega$ , and therefore,

$$\int_{\{u_f = \kappa\}} |\nabla u_f| |dz| = \begin{cases} 2\pi\kappa\mu(\kappa) & \text{in Case 1} \\ 4\pi j\kappa\mu(\kappa) & \text{in Case 2,} \\ 2\pi\alpha\kappa\mu(\kappa) & \text{in Case 3.} \end{cases}$$

To summarize, we have shown that

$$\ell(\{u_f = \kappa\})^2 \leq \begin{cases} -2\pi\kappa\mu'(\kappa)\mu(\kappa) & \text{in Case 1} \\ -4\pi j\kappa\mu'(\kappa)\mu(\kappa) & \text{in Case 2,} \\ -2\pi\alpha\kappa\mu'(\kappa)\mu(\kappa) & \text{in Case 3.} \end{cases}$$

We now use the isoperimetric inequality to bound the left side from below; for references in the spherical and hyperbolic case, see, for instance, [31, (4.23)], as well as [8], [22, Third part, Chapter IV], [34,36,37]. We have

$$\ell(\partial A)^2 \geq \begin{cases} 4\pi m(A) & \text{in Case 1,} \\ 4\pi m(A)(1 - m(A)) & \text{in Case 2,} \\ 4\pi m(A)(1 + m(A)) & \text{in Case 3.} \end{cases}$$

Using these inequalities with  $A = \{u_f > \kappa\}$ , dividing by  $\mu(\kappa)$  (which is nonzero for  $\kappa < M$ ) and combining the resulting inequality with the above upper bound on  $\ell(\{u_f = \kappa\})^2$ , we obtain

$$\begin{cases} 2 \leq -\kappa\mu'(\kappa) & \text{in Case 1,} \\ 1 - \mu(\kappa) \leq -j\kappa\mu'(\kappa) & \text{in Case 2,} \\ 1 + \mu(\kappa) \leq -\frac{\alpha}{2}\kappa\mu'(\kappa) & \text{in Case 3.} \end{cases} \quad (3)$$

These inequalities are equivalent to the monotonicity assertions in the lemma.

It remains to verify that this function is constant if  $f \in \mathcal{M}$ . By the equimeasurability discussed before the statement of Theorem 10, it suffices to prove this for  $f = F_0 \in \mathcal{M}$ . For all  $\kappa \leq 1$ , we have

$$m(\{u_{F_0} > \kappa\}) = \begin{cases} \int_{\mathbb{C}} \mathbb{1}(e^{-\frac{\alpha}{2}|z|^2} > \kappa) dA(z) = -2\ln \kappa & \text{in Case 1,} \\ \pi^{-1} \int_{\mathbb{C}} \mathbb{1}((1 + |z|^2)^{-j} > \kappa) \frac{dA(z)}{(1 + |z|^2)^2} = 1 - \kappa^{\frac{1}{j}} & \text{in Case 2,} \\ \pi^{-1} \int_{\mathbb{D}} \mathbb{1}((1 - |z|^2)^{\frac{\alpha}{2}} > \kappa) \frac{dA(z)}{(1 - |z|^2)^2} = \kappa^{-\frac{2}{\alpha}} - 1 & \text{in Case 3.} \end{cases}$$

It follows that for  $f = F_0$ , the function in Lemma 12 is, indeed, constant.  $\square$

The last ingredient in the proof of Theorem 10 is an inequality due to Chebyshev [11] (see also [16, Theorems 43 and 236]). For a proof of the following lemma, with a slightly weaker assumption than monotonicity of one of the functions, see [20, Lemma 4.1].

**Lemma 13.** Let  $t_0 > 0$  and let  $w$  and  $h$  be nondecreasing functions on  $[0, t_0]$ . Then

$$\int_0^{t_0} h(t)w(t)dt \geq t_0^{-1} \int_0^{t_0} h(t)dt \int_0^{t_0} w(t)dt.$$

We are finally in position to prove the main result of this section.

**Proof of Theorem 10.** We begin with some preliminary remarks concerning convex functions  $\Phi$  on  $[0, 1]$ . We first argue that, without loss of generality, we may assume that  $\Phi$  is continuous on  $[0, 1]$ . By convexity, it is continuous on  $(0, 1)$ , so we only need to discuss the endpoints. It is elementary that  $\Phi(0^+) := \lim_{s \rightarrow 0^+} \Phi(s)$  and  $\Phi(1^-) := \lim_{s \rightarrow 1^-} \Phi(s)$  exist and are finite. (Note that these limits are  $\leq \Phi(0)$  and  $\leq \Phi(1)$ , respectively, and thus are not  $+\infty$ .) By analyticity,  $m(\{u_f = 0\}) = m(\{f = 0\}) = 0$ , so on this set, we may replace  $\Phi(0)$  by  $\Phi(0^+)$  without changing the value of the integral. Similarly, by Lemma 11 and its proof,  $\{u_f = 1\}$  consists at most of one point, so on this set, we may replace  $\Phi(1)$  by  $\Phi(1^-)$  without changing the value of the integral. Thus, we may assume that  $\Phi$  is continuous on  $[0, 1]$ .

Next, we argue that we may assume that  $\Phi(0) = 0$ . In Case 2,  $m$  is a finite measure, so this can be accomplished by replacing  $\Phi$  by  $\Phi - \Phi(0)$ , which has a trivial effect on the supremum. In Cases 1 and 3, we take  $f \in \mathcal{M}$  and see from the explicit form that  $u_f(z) \rightarrow 0$  as  $|z| \rightarrow \infty$  in Case 1 and  $|z| \rightarrow 1$  in Case 3. (In fact, this holds for any  $f \in \mathcal{H}$ , as discussed in the proof of Lemma 11, but this is not needed here.) It follows that, if  $\Phi(0) \neq 0$ , then the supremum is equal to  $+\infty$ , and this value is achieved by all  $f \in \mathcal{M}$ , so the assertion of the theorem is true in this case. Thus, in what follows, we may assume that  $\Phi(0) = 0$ .

After these preliminaries, we begin with the main part of the argument. Let  $f \in \mathcal{H}$  with  $\|f\| = 1$ . We define  $u_f$  by (2) and set

$$s_0 := \|u_f\|_{L^\infty(\Omega)}^2.$$

Then, the quantity we are interested in can be written as follows:

$$\int_{\Omega} \Phi(u_f(z)^2) dm(z) = \int_0^{s_0} m(\{u_f^2 > s\}) \Phi'(s) ds.$$

Here,  $\Phi'$  denotes either the left- or the right-sided derivative of  $\Phi$ , which are known to exist everywhere and to coincide outside of a countable set [42, Theorem 1.26]. We also used the facts that  $\Phi$  is absolutely continuous [42, Theorem 1.28] and that  $\Phi(0) = 0$ .

We now write the quantity on the right side as follows:

$$\begin{cases} \int_0^{s_0} \left( -\ln s - g\left(s^{\frac{1}{2}}\right) \right) \Phi'(s) ds & \text{in Case 1,} \\ \int_0^{s_0} \left( 1 - s^{\frac{1}{2}} g\left(s^{\frac{1}{2}}\right) \right) \Phi'(s) ds & \text{in Case 2,} \\ \int_0^{s_0} \left( -1 - s^{\frac{1}{2}} g\left(s^{\frac{1}{2}}\right) \right) \Phi'(s) ds & \text{in Case 3,} \end{cases}$$

where, according to Lemma 12,  $\kappa \mapsto g(\kappa)$  is nondecreasing on  $(0, s_0^{\frac{1}{2}})$ . In particular, when  $\Phi$  is the identity, we obtain, in view of the normalization of  $f$ ,

$$1 = \|f\|^2 = \begin{cases} \int_0^{s_0} \left(-\ln s - g\left(s^{\frac{1}{2}}\right)\right) ds & \text{in Case 1,} \\ (2J+1) \int_0^{s_0} \left(1 - s^{\frac{1}{2J}} g\left(s^{\frac{1}{2}}\right)\right) ds & \text{in Case 2,} \\ (\alpha-1) \int_0^{s_0} \left(-1 - s^{-\frac{1}{\alpha}} g\left(s^{\frac{1}{2}}\right)\right) ds & \text{in Case 3.} \end{cases} \quad (4)$$

Let us set

$$t_0 := \begin{cases} s_0 & \text{in Case 1,} \\ s_0^{\frac{2J+1}{2J}} & \text{in Case 2,} \\ s_0^{\frac{\alpha-1}{\alpha}} & \text{in Case 3,} \end{cases}$$

and for  $0 \leq t \leq t_0$ ,

$$h(t) := \begin{cases} g(t^{\frac{1}{2}}) & \text{in Case 1,} \\ g(t^{\frac{1}{2J+1}}) & \text{in Case 2,} \\ g(t^{\frac{\alpha}{2(\alpha-1)}}) & \text{in Case 3.} \end{cases}$$

Then, the normalization (4) can be equivalently written as follows:

$$1 = \begin{cases} \int_0^{t_0} (-\ln t - h(t)) dt & \text{in Case 1,} \\ 2J \int_0^{t_0} \left(t^{-\frac{1}{2J+1}} - h(t)\right) dt & \text{in Case 2,} \\ \alpha \int_0^{t_0} \left(-t^{\frac{1}{\alpha-1}} - h(t)\right) dt & \text{in Case 3,} \end{cases}$$

while the quantity to be maximized is

$$\int_{\Omega} \Phi(u_f(z)^2) dm(z) = \begin{cases} \int_0^{t_0} (-\ln t - h(t)) w(t) dt & \text{in Case 1,} \\ 2J \int_0^{t_0} \left(t^{-\frac{1}{2J+1}} - h(t)\right) w(t) dt & \text{in Case 2,} \\ \alpha \int_0^{t_0} \left(-t^{\frac{1}{\alpha-1}} - h(t)\right) w(t) dt & \text{in Case 3,} \end{cases} \quad (5)$$

where, for  $0 \leq t \leq t_0$ ,

$$w(t) := \begin{cases} \Phi'(t) & \text{in Case 1,} \\ \Phi'(t^{\frac{2J}{2J+1}}) & \text{in Case 2,} \\ \Phi'(t^{\frac{\alpha}{\alpha-1}}) & \text{in Case 3.} \end{cases}$$

Since  $\Phi$  is convex,  $\Phi'$  is nondecreasing, and therefore,  $w$  is nondecreasing as well. Also,  $h$  is nondecreasing since  $g$  is. Thus, Lemma 13 is applicable, and, for given  $t_0$ , an upper bound on the right side of (5) is obtained by replacing the function  $h$  by the constant  $t_0^{-1} \int_0^{t_0} h(t) dt$ . According to the normalization, we have

$$t_0^{-1} \int_0^{t_0} h(t) dt = C(t_0),$$

where, for  $\tau \in (0, 1]$ ,

$$C(\tau) := \begin{cases} -\tau^{-1} - \tau^{-1} \int_0^\tau \ln t dt & \text{in Case 1,} \\ -(2J)^{-1} \tau^{-1} + \tau^{-1} \int_0^\tau t^{-\frac{1}{2J+1}} dt & \text{in Case 2,} \\ -\alpha^{-1} \tau^{-1} - \tau^{-1} \int_0^\tau t^{\frac{1}{\alpha-1}} dt & \text{in Case 3.} \end{cases}$$

Thus, we have shown the upper bound as follows:

$$\int_{\Omega} \Phi(u_f(z)^2) dm(z) \leq A(t_0), \quad (6)$$

where, for  $\tau \in (0, 1]$ ,

$$A(\tau) := \begin{cases} \int_0^\tau (-\ln t - C(\tau)) w(t) dt & \text{in Case 1,} \\ 2J \int_0^\tau \left( t^{-\frac{1}{2J+1}} - C(\tau) \right) w(t) dt & \text{in Case 2,} \\ \alpha \int_0^\tau \left( -t^{\frac{1}{\alpha-1}} - C(\tau) \right) w(t) dt & \text{in Case 3.} \end{cases}$$

Our goal now is to show that  $A$  is nondecreasing in  $(0, 1]$ . Since, by Lemma 11,  $t_0 \leq 1$ , inserting this into (6) gives us the upper bound  $A(1)$ . Later, we will argue that this is the claimed optimal bound and discuss the cases of equality.

In order to prove the monotonicity of  $A$ , we first compute

$$C(\tau) = \begin{cases} -\tau^{-1} - \ln \tau + 1 & \text{in Case 1,} \\ -(2J)^{-1} \tau^{-1} + \frac{2J+1}{2J} \tau^{-\frac{1}{2J+1}} & \text{in Case 2,} \\ -\alpha^{-1} \tau^{-1} - \frac{\alpha-1}{\alpha} \tau^{\frac{1}{\alpha-1}} & \text{in Case 3.} \end{cases}$$

From these expressions, one can easily deduce that  $C' > 0$  in  $(0, 1)$ . Another consequence that we will use soon is that

$$0 = \begin{cases} -\ln \tau - C(\tau) - C'(\tau)\tau & \text{in Case 1,} \\ \tau^{-\frac{1}{2J+1}} - C(\tau) - C'(\tau)\tau & \text{in Case 2,} \\ -\tau^{\frac{1}{\alpha-1}} - C(\tau) - C'(\tau)\tau & \text{in Case 3.} \end{cases} \quad (7)$$

We now compute

$$A'(\tau) = \begin{cases} (-\ln \tau - C(\tau))w(\tau) - C'(\tau) \int_0^\tau w(t)dt & \text{in Case 1,} \\ 2J\left(\tau^{-\frac{1}{2J+1}} - C(\tau)\right)w(\tau) - 2JC'(\tau) \int_0^\tau w(t)dt & \text{in Case 2,} \\ \alpha\left(-\tau^{\frac{1}{\alpha-1}} - C(\tau)\right)w(\tau) - \alpha C'(\tau) \int_0^\tau w(t)dt & \text{in Case 3.} \end{cases}$$

Since  $w$  is nondecreasing, we have  $\int_0^\tau w(t)dt \leq \tau w(\tau)$ . This, together with  $C'(\tau) \geq 0$ , implies

$$A'(\tau) \geq \begin{cases} (-\ln \tau - C(\tau))w(\tau) - C'(\tau)\tau w(\tau) & \text{in Case 1,} \\ 2J\left(\tau^{-\frac{1}{2J+1}} - C(\tau)\right)w(\tau) - 2JC'(\tau)\tau w(\tau) & \text{in Case 2,} \\ \alpha\left(-\tau^{\frac{1}{\alpha-1}} - C(\tau)\right)w(\tau) - \alpha C'(\tau)\tau w(\tau) & \text{in Case 3.} \end{cases}$$

According to (7), the right side is equal to zero in all cases. This proves that  $A' \geq 0$  in  $(0, 1]$ .

As mentioned before, the monotonicity of  $A$  allows us to replace  $A(t_0)$  by  $A(1)$  in (6). We claim that this bound is optimal. Indeed, if  $f \in \mathcal{M}$ , then, by the second part of Lemma 12,  $g$  is constant. Thus, also  $h$  is constant and nothing was lost when applying Lemma 13. This proves that, in this case, (6) is an equality and, since  $t_0 = 1$  by Lemma 11, we have shown the claimed optimality.

Finally, assume that  $A(1) < \infty$  and that  $\Phi$  is not affine linear. Then,  $\Phi'$  is not constant and neither is  $w$ . We deduce that there is an  $\varepsilon > 0$  such that the inequality  $\int_0^\tau w(t)dt \leq \tau w(\tau)$  is strict for all  $\tau \in (1 - \varepsilon, 1]$ . This, together with the fact that  $C'(\tau) > 0$  for all  $\tau \in (0, 1)$  implies that  $A'(\tau) > 0$  for all  $\tau \in (1 - \varepsilon, 1)$ . In particular,  $A(\tau) < A(1)$  if  $\tau \in [0, 1)$ .

As a consequence, if  $f \in \mathcal{H}$  with  $\|f\| = 1$  attains the supremum in Theorem 10, then necessarily  $t_0 = 1$ . Then, by Lemma 11,  $f \in \mathcal{M}$ , as claimed. This completes the proof of Theorem 10, except for the explicit value of the supremum.

To compute the latter, we may choose an arbitrary element in  $\mathcal{M}$ , and it is convenient to take  $f = F_0 = 1$ . Then, we obtain, by integrating in radial coordinates,

$$\int_{\Omega} \Phi(u_{F_0}(z)^2) dz = \begin{cases} 2\pi \int_0^\infty \Phi(e^{-\pi r^2}) r dr & \text{in Case 1,} \\ 2 \int_0^\infty \Phi((1+r^2)^{-2J})(1+r^2)^{-2} r dr & \text{in Case 2,} \\ 2 \int_0^1 \Phi((1-r^2)^\alpha)(1-r^2)^{-2} r dr & \text{in Case 3.} \end{cases}$$

Changing variables  $s = e^{-\pi r^2}$ ,  $s = (1+r^2)^{-2J}$ , and  $s = (1+r^2)^\alpha$  in the three cases, we easily arrive at the claimed formulas.  $\square$

## 2.3 Extension to density matrices

In this subsection, we generalize the inequality in Theorem 10, and under a slightly stronger assumption on  $\Phi$ , we characterize the cases of equality. We use an argument similar to [23, Lemma 2].

Given an operator  $\rho \geq 0$  with  $\text{Tr} \rho = 1$  on one of the Hilbert spaces  $\mathcal{H}$ , we define a function  $u_\rho$  on  $\Omega$  as follows. We can write

$$\rho = \sum_n p_n |f_n\rangle\langle f_n| \quad \text{with} \quad \sum_n p_n = 1, p_n \geq 0, \langle f_n, f_m \rangle = \delta_{n,m}.$$

We then set

$$u_\rho(z) := \left( \sum_n p_n u_{f_n}(z)^2 \right)^{\frac{1}{2}}.$$

It is easily checked that this is well-defined. (Note, in particular, the nonuniqueness of the above decomposition of  $\rho$  in the case of a degenerate eigenvalue.) Moreover, for  $\rho = |f\rangle\langle f|$ , this definition of  $u_\rho$  coincides with the earlier one of  $u_f$ .

**Corollary 14.** *Let  $\Phi : [0, 1] \rightarrow \mathbb{R}$  be convex. Then,*

$$\sup \left\{ \int_{\Omega} \Phi(u_\rho(z)^2) dm(z) : \rho \geq 0 \text{ on } \mathcal{H}, \text{Tr} \rho = 1 \right\} \\ = \begin{cases} \int_0^1 \Phi(s) s^{-1} ds & \text{in Case 1,} \\ (2J)^{-1} \int_0^1 \Phi(s) s^{\frac{1}{2J}-1} ds & \text{in Case 2,} \\ (2K)^{-1} \int_0^1 \Phi(s) s^{-\frac{1}{2K}-1} ds & \text{in Case 3,} \end{cases}$$

and the supremum is attained for  $\rho = |F\rangle\langle F|$  with  $F \in \mathcal{M}$ . If, moreover,  $\Phi$  is strictly convex and if the supremum is finite, then it is attained only for such  $\rho$ .

**Proof.** We use the above expansion of  $\rho$ . By convexity of  $\Phi$ , for any  $z \in \Omega$ ,

$$\Phi(u_\rho(z)^2) = \Phi\left(\sum_n p_n u_{f_n}(z)^2\right) \leq \sum_n p_n \Phi(u_{f_n}(z)^2).$$

Thus, with  $S$  denoting the supremum in Theorem 10,

$$\int_{\Omega} \Phi(u_\rho(z)^2) dm(z) \leq \sum_n p_n \int_{\Omega} \Phi(u_{f_n}(z)^2) dm(z) \leq \sum_n p_n S = S.$$

Since, by Theorem 10,  $S$  is attained for  $\rho = |F\rangle\langle F|$  with  $F \in \mathcal{M}$ , we obtain the first assertion in the corollary.

Now assume that  $S < \infty$  and that equality is achieved for some  $\rho$ . If  $\Phi$  is not linear, then, by Theorem 10,  $f_n \in \mathcal{M}$  for each  $n$ . (Throughout we restrict ourselves to values of  $n$  for which  $p_n > 0$ .) Moreover,

$$\Phi\left(\sum_n p_n u_{f_n}(z)^2\right) = \sum_n p_n \Phi(u_{f_n}(z)^2) \quad \text{for a.e. } z \in \Omega.$$

Assuming now that  $\Phi$  is strictly convex, we deduce that  $u_{f_n}(z)^2 = u_{f_1}(z)^2$  for a.e.  $z \in \Omega$  and every  $n$ . Thus, by continuity,  $|f_n(z)| = |f_1(z)|$  for all  $z \in \Omega$  and all  $n$ . By analyticity, there are  $\theta_n \in \mathbb{R}/2\pi\mathbb{Z}$  such that  $f_n(z) = e^{i\theta_n} f_1(z)$  for all  $z \in \Omega$ . (Indeed, by the maximum modulus principle,  $f_n/f_1$  is equal to a constant in  $\Omega$  without the zeros of  $f_1$  and then by continuity in all of  $\Omega$ .) Since  $\langle f_n, f_1 \rangle = \delta_{n,1}$ , we conclude that there is only a single index  $n$ , namely,  $n = 1$ .  $\square$

## 2.4 Another inequality of Kulikov

For later purposes, in this subsection, we record another inequality from [20], which corresponds, in some sense, to the limiting case  $\alpha = 1$  in Theorem 10.

The underlying Hilbert space is the Hardy space  $H^2(\mathbb{D})$  consisting of all analytic functions  $f$  in  $\mathbb{D}$  such that

$$\|f\|_{H^2(\mathbb{D})} := \left( \sup_{0 < r < 1} (2\pi)^{-1} \int_{-\pi}^{\pi} |f(re^{i\varphi})|^2 d\varphi \right)^{1/2} < \infty.$$

To emphasize the analogy with Theorem 10, we denote this space by  $\mathcal{H}$  and its norm by  $|||\cdot|||$ . We also use the same notation  $\Omega$  and  $dm(z)$  as in Case 3. The function  $u_f$  is defined by (2) with  $\alpha = 1$ . The functions  $F_w$  are defined as in Case 3 with  $\alpha = 1$ , and one easily checks that they are normalized. The set  $\mathcal{M}$  is defined as before.

**Proposition 15.** *Let  $\Phi : [0, 1] \rightarrow \mathbb{R}$  be nondecreasing. Then,*

$$\sup \left\{ \int_{\Omega} \Phi(u_f(z)^2) dm(z) : f \in \mathcal{H}, |||f||| = 1 \right\} = \int_0^1 \Phi(s) s^{-2} ds$$

and the supremum is attained in  $\mathcal{M}$ . If  $\Phi$  is strictly increasing near 1 and if the supremum is finite, then it is attained only in  $\mathcal{M}$ .

By  $\Phi$  being “strictly increasing near 1,” we mean that  $\Phi(s) < \Phi(1^-)$  for all  $s < 1$ , where  $\Phi(1^-) := \lim_{s \rightarrow 1^-} \Phi(s)$ . We will see in the proof that  $u_f \leq 1$ , so the quantity in the supremum is well defined.

**Proof.** The first part is a special case of [20, Theorem 1.1]. The second part can be obtained by an inspection of the proof of the first part, but since this is not explicitly stated in [20], we provide some details. As in the proof of Theorem 10, we may assume that  $\Phi(0) = \lim_{s \rightarrow 0^+} \Phi(s) = 0$ . Then,

$$\int_{\Omega} \Phi(u_f(z)^2) dm(z) = \int_0^{s_0} \mu\left(s^{\frac{1}{2}}\right) d\Phi(s),$$

where  $s_0 := \|u_f\|_{L^\infty(\Omega)}^2$ . By an analog of Lemma 11, we have  $s_0 \leq 1$  [20, (1.2)] with equality if and only if  $f \in \mathcal{M}$ . The argument for the latter assertion is essentially the same as in Lemma 11, using the fact that  $u_f(z) \rightarrow 0$  as  $|z| \rightarrow 1$ , stated in [20, paragraph after (1.2)], and the cases of equality in the Schwarz inequality for a reproducing kernel (see also the proof of Proposition 19).

As shown in [20, Theorem 3.1], we have  $\mu(\kappa) \leq (\kappa^{-2} - 1)_+$  for all  $\kappa > 0$ , and this bound is an equality if  $f \in \mathcal{M}$ . Thus,

$$\int_0^{s_0} \mu\left(s^{\frac{1}{2}}\right) d\Phi(s) \leq \int_0^1 (s^{-1} - 1) d\Phi(s) - \int_{s_0}^1 (s^{-1} - 1) d\Phi(s),$$

where the first term on the right side corresponds to the value of the supremum. Thus, if this term is finite and  $f$  attains the supremum, then the second term on the right side has to vanish. If  $\Phi$  is strictly increasing near 1, then the measure  $d\Phi(s)$  does not vanish on any interval  $(1 - \varepsilon, 1)$  with  $\varepsilon > 0$ , and therefore, necessarily,  $s_0 = 1$ . By the above-mentioned analog of Lemma 11, this means  $f \in \mathcal{M}$ , as claimed.

To compute the value of the supremum, we can proceed exactly as in Case 3 of Theorem 10, setting  $\alpha = 1$  in that calculation. This proves the proposition.  $\square$



### 3 Reverse Hölder inequalities for analytic functions

The material in this section is an extension of that in the previous section. It is not relevant for the proof of the results in Section 1.

In Theorem 10, we were working under a constraint on a Hilbertian norm. It turns out that this is an unnecessary restriction. We will now prove a generalization of Theorem 10 with a constraint on a more general norm or quasinorm. This will allow us to settle a conjecture by Bodmann [9, Conjecture 3.5].

We continue to use the notation of Section 2. For  $0 < p < \infty$ , we define

$$|||f|||_p := \begin{cases} \left( \frac{p}{2} \int_{\mathbb{C}} |f(z)|^p e^{-\frac{p}{2}|z|^2} dA(z) \right)^{1/p} & \text{in Case 1,} \\ \left( \frac{pJ+1}{\pi} \int_{\mathbb{C}} |f(z)|^p (1+|z|^2)^{-pJ-2} dA(z) \right)^{1/p} & \text{in Case 2,} \\ \left( \frac{\alpha p-2}{2\pi} \int_{\mathbb{D}} |f(z)|^p (1-|z|^2)^{\frac{\alpha}{2}-2} dA(z) \right)^{1/p} & \text{in Case 3.} \end{cases}$$

This is a norm for  $p \geq 1$  and a quasinorm for  $p < 1$ . The prefactors are chosen such that  $|||1|||_p = 1$ . We still assume that  $J \in \frac{1}{2}\mathbb{N}$  in Case 2 and now  $\alpha > \frac{2}{p}$  in Case 3. We denote by  $\mathcal{X}^p$  the space of all analytic functions  $f$  on  $\Omega$  such that  $|||f|||_p < \infty$ . In Case 2, we require, in addition, that  $f$  is a polynomial of degree  $\leq 2J$ . The function  $u_f$  for  $f \in \mathcal{X}^p$  is defined as before. We note that every  $F \in \mathcal{M}$  satisfies  $|||F|||_p = 1$ . Indeed, we have already noted that this holds for  $F = F_0 = 1$ , and for general  $F \in \mathcal{M}$ , it follows from the equimeasurability of  $u_{F_w}$  for different  $w$ , discussed before Theorem 10.

**Theorem 16.** *Let  $0 < p < \infty$  and let  $\Phi : [0, 1] \rightarrow \mathbb{R}$  be convex. Then,*

$$\sup \left\{ \int_{\Omega} \Phi(u_f(z)^p) dm(z) : f \in \mathcal{X}^p, |||f|||_p = 1 \right\} = \begin{cases} \frac{2}{p} \int_0^1 \Phi(s) s^{-1} ds & \text{in Case 1,} \\ (pJ)^{-1} \int_0^1 \Phi(s) s^{\frac{1}{pJ}-1} ds & \text{in Case 2,} \\ \frac{2}{\alpha p} \int_0^1 \Phi(s) s^{-\frac{2}{\alpha p}-1} ds & \text{in Case 3,} \end{cases}$$

and the supremum is attained in  $\mathcal{M}$ . If  $\Phi$  is not affine linear and if the supremum is finite, then it is attained only in  $\mathcal{M}$ .

For  $p = 2$ , this theorem reduces to Theorem 10. In Case 3, it reduces to [20, Theorem 1.2 and Remark 4.3], except that our equality statement allows for more general  $\Phi$ . In Cases 1 and 2, the theorem seems to be new.

Taking  $\Phi(s) = s^{\frac{q}{p}}$  with  $q > p$ , we obtain the following reverse Hölder inequalities.

**Corollary 17.** *Let  $0 < p < q < \infty$ . Then, for any  $f \in \mathcal{X}^p$ ,*

$$|||f|||_q \leq |||f|||_p$$

with equality if and only if  $f = 0$  or  $|||f|||_p^{-1} f \in \mathcal{M}$ .

This corollary in Case 1 is due to Carlen [10, Theorem 2]. In fact, Carlen proves a more general inequality including an additional parameter. Carlen's method of proof depends on the logarithmic Sobolev inequality

and an identity for analytic functions. It is different from ours. Corollary 17 in Case 2 has been conjectured by Bodmann [9, Conjecture 3.5], who proved it in the special case where  $q = p + J^{-1}n$  with  $n \in \mathbb{N}$  and  $p > J^{-1}$ . Bodmann's proof relies on a sharp Sobolev inequality and an analog of Carlen's identity. Corollary 17 in Case 3 is due to Kulikov [20, Corollary 1.3]. The special case  $q = p + 2\alpha^{-1}$  with  $p \geq 2$ , and  $\alpha p > 4$  was proved earlier by Bandyopadhyay in [3, Corollary 3.3] using the method of Carlen and Bodmann. (Note that in [3], it is assumed that  $\alpha \in \mathbb{N} \setminus \{1\}$ , in her notation  $\alpha = 2k$ , but this seems to be irrelevant for [3, Section 3].)

We now turn to the proof of Theorem 16. The main new ingredient is the following generalization of Lemma 11. In Case 3, this is well known [20, (1.1)] and probably also in Case 1, but in Case 2 it might be new.

**Lemma 18.** *Let  $0 < p < \infty$  and let  $f \in X^p$ . Then,*

$$\|u_f\|_{L^\infty(\Omega)} \leq \|f\|_p$$

*with equality if and only if either  $f = 0$  or  $\|f\|_p^{-1}f \in \mathcal{M}$ .*

**Proof.** We begin by showing that

$$u_f(0) \leq \|f\|_p \quad (8)$$

with equality if and only if  $\|f\|_p^{-1}f = e^{i\theta}F_0$  for some  $\theta \in \mathbb{R}/2\pi\mathbb{Z}$  (provided  $f \neq 0$ ). Since  $\ln|f|$  is subharmonic in  $\Omega$ , we have, for any  $r \in (0, R)$ , where  $R := \infty$  in Cases 1 and 2 and  $R := 1$  in Case 3,

$$\ln|f(0)| \leq (2\pi)^{-1} \int_{-\pi}^{\pi} \ln|f(re^{i\varphi})| d\varphi.$$

We multiply by  $re^{-\frac{p}{2}r^2}$ ,  $r(1+r^2)^{-pJ-2}$ , and  $r(1-r^2)^{\frac{p}{2}-2}$  in the different cases and integrate with respect to  $r \in (0, R)$ . In this way, we obtain

$$\ln|f(0)| \leq \int_{\Omega} \ln|f(z)| w(z) dA(z),$$

where

$$w(z) := \begin{cases} \frac{p}{2} e^{-\frac{p}{2}|z|^2} & \text{in Case 1,} \\ \frac{pJ+1}{\pi} (1+|z|^2)^{-pJ-2} & \text{in Case 2,} \\ \frac{\alpha p-2}{2\pi} (1-|z|^2)^{\frac{p}{2}-2} & \text{in Case 3.} \end{cases}$$

The measure  $w(z)dA(z)$  is a probability measure. Multiplying the inequality by  $p$ , we can write it as follows:

$$|f(0)|^p \leq \exp \left( \int_{\Omega} \ln(|f(z)|^p) w(z) dA(z) \right) \leq \int_{\Omega} |f(z)|^p w(z) dA(z) = \|f\|_p^p,$$

where the second inequality comes from Jensen's inequality. Since the exponential function is strictly convex, Jensen's inequality is strict unless  $\ln(|f|^p)$  is almost everywhere constant. Since  $f$  is continuous, this happens if and only if  $|f|$  is constant and, by the maximum modulus principle, if and only if  $f$  is constant. This proves the claim.

We now claim that, for any  $z \in \Omega$ ,

$$u_f(z) \leq \|f\|_p \quad (9)$$

if and only if  $\|f\|_p^{-1}f = e^{i\theta}F_z$  for some  $\theta \in \mathbb{R}/2\pi\mathbb{Z}$  (provided  $f \neq 0$ ). In Case 2, the same inequality remains valid for  $z = \infty$ , recalling that  $u_f$  extends continuously to this point. Indeed, inequality (9) and its equality statement follow from the corresponding assertions concerning (8), by applying an element of the

Heisenberg group,  $SU(2)$ , or  $SU(1, 1)$  to move the point  $z$  to the point 0 and by noting that  $|||\cdot|||_p$  is invariant under this group action. The latter fact follows from the equimeasurability property discussed before Theorem 10.

Inequality (9) implies the inequality in the lemma. Now assume that  $f \neq 0$  achieves equality in this inequality. We claim that  $u_f(z) \rightarrow 0$  as  $|z| \rightarrow \infty$  or  $|z| \rightarrow 1$  in Cases 1 and 3. This, together with the continuity of  $u_f$  in  $\Omega$  in Cases 1 and 3 and in  $\mathbb{C} \cup \{\infty\}$  in Case 2, implies that there is a  $z$  such that  $u_f(z) = \|u_f\|_{L^\infty(\Omega)}$ . The equality statement in (9) then implies the equality statement in the lemma.

Thus, it suffices to prove the asymptotic vanishing of  $u_f$  in Cases 1 and 3. In both cases, this is clear when  $f$  is a polynomial and follows, in the general case, from the fact that polynomials are dense with respect to  $|||\cdot|||_p$  and the inequality in the lemma. This completes the proof.  $\square$

**Proof of Theorem 16.** Given Lemma 18, which replaces Lemma 11, the proof is a minor variation of that of Theorem 10. We only sketch the major steps. The task is to maximize

$$\int_0^{s_0} m(\{u_f^p > s\}) \Phi'(s) ds = \int_0^{s_0} \mu(s^{\frac{1}{p}}) \Phi'(s) ds$$

under the constraint

$$\int_0^{s_0} m(\{u_f^p > s\}) ds = \int_0^{s_0} \mu(s^{\frac{1}{p}}) ds = \begin{cases} \frac{2}{p} & \text{in Case 1,} \\ \frac{1}{pJ+1} & \text{in Case 2,} \\ \frac{2}{\alpha p - 2} & \text{in Case 3,} \end{cases}$$

with  $s_0 := \|u_f\|_{L^\infty(\Omega)}^p$  and  $\mu$  as in Lemma 12. The latter lemma allows us to write  $\mu(s^{\frac{1}{p}})$  as the sum of a fixed piece and one that involves the nondecreasing function  $g(s^{\frac{1}{p}})$ . We pass from the variable  $s$  to a variable  $t$  so that the constraint for the resulting nondecreasing function  $h$  can be written as an integral with respect to the unweighted measure  $dt$ . Then, we can use Chebyshev's bound (Lemma 13) to replace  $h$  by its average. This leads to a certain bound  $A(t_0)$ , and a computation, similarly as for  $p = 2$ , shows that  $A$  is nondecreasing. Moreover, if  $\Phi$  is not affine linear, then  $A$  is strictly increasing. This concludes the sketch of the proof of Theorem 16.  $\square$

## 4 Proof of the main results

In this section, we prove the main results stated in Section 1. In each case, we will work with a concrete representation of the group action that involves analytic functions. The inequalities will then be deduced from Theorem 10.

### 4.1 Proof of Theorem 1

By scaling, it suffices to prove the theorem for a single value of  $\hbar$ , and it is convenient to choose  $\hbar = (2\pi)^{-1}$ . Then, given  $\psi \in L^2(\mathbb{R})$ , we can write

$$\langle \psi_{p,q}, \psi \rangle = e^{-\frac{\pi}{2}(q^2+p^2)} e^{i\pi q p f(q - ip)}$$

with

$$f(z) := 2^{\frac{1}{4}} \int_{\mathbb{R}} e^{2\pi izx - \frac{\pi}{2} z^2 - \pi x^2} \psi(x) dx \quad \text{for all } z \in \mathbb{C}.$$

It is well known and easy to see that  $f$  is entire and that

$$\|f\|_{\mathcal{F}^2}^2 = \iint_{\mathbb{R} \times \mathbb{R}} |\langle \psi_{p,q}, \psi \rangle|^2 dp dq = \|\psi\|_{L^2(\mathbb{R})}^2,$$

where the last identity is the completeness relation of the coherent states. In particular,  $f \in \mathcal{F}^2(\mathbb{C})$ . Moreover,

$$u_f(q - ip) = |\langle \psi_{p,q}, \psi \rangle|,$$

so Theorem 1 follows immediately from Theorem 10 in Case 1. Similarly, Corollary 2 follows from Corollary 14.

## 4.2 Proof of Theorem 4

Let  $J \in \frac{1}{2}\mathbb{N}$ . We consider the representation of  $SU(2)$  on functions  $f$  on  $\mathbb{C}$  given as follows:

$$\pi_U(f)(z) := (\beta z + \bar{\alpha})^{2J} f\left(\frac{\alpha z - \bar{\beta}}{\beta z + \bar{\alpha}}\right) \quad \text{for all } z \in \mathbb{C},$$

where

$$U = \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix} \in SU(2), \quad \text{that is, } \alpha, \beta \in \mathbb{C} \quad \text{with } |\alpha|^2 + |\beta|^2 = 1.$$

This representation restricted to  $\mathcal{P}_{2J}$  is irreducible and unitary for the norm defined above. In this representation,

$$S_1 = \frac{1}{2} \left( (-z^2 + 1) \frac{d}{dz} + 2Jz \right), \quad S_2 = \frac{1}{2i} \left( (-z^2 - 1) \frac{d}{dz} + 2Jz \right), \quad S_3 = z \frac{d}{dz} - J.$$

We may choose the space  $\mathcal{H}$  in Theorem 4 as  $\mathcal{P}_{2J}$ . By an explicit computation, one sees that the functions  $F_w$  are eigenvectors of the operator  $S(w) \cdot S$  corresponding to the eigenvalue  $-J$ , where we used the stereographic projection  $S : \mathbb{C} \mapsto \mathbb{S}^2$ , which is given as follows:

$$S_1(w) + iS_2(w) := \frac{2w}{1 + |w|^2}, \quad S_3(w) := \frac{1 - |w|^2}{1 + |w|^2}.$$

Consequently, the phases of  $\psi_w$ ,  $w \in \mathbb{S}^2$ , can be chosen such that these functions coincide with the functions  $F_w$ ,  $w \in \mathbb{C} \cup \{\infty\}$ . Thus, using the explicit form of  $F_w$ ,

$$\langle \psi_{S(w)}, \psi \rangle = (1 + |w|^2)^{-J} f(w)$$

with

$$f(w) := \frac{2J+1}{\pi} \int_{\mathbb{C}} (1 + w\bar{z})^{2J} \psi(z) (1 + |z|^2)^{-2J-2} dA(z).$$

Since  $\psi$  is a polynomial of degree  $\leq 2J$ , the reproducing property of the kernel implies that  $f(w) = \psi(w)$  for all  $w \in \mathbb{C}$ . Moreover,

$$u_f(w) = |\langle \psi_{S(w)}, \psi \rangle|,$$

and so, by a change of variables,

$$\pi^{-1} \int_{\mathbb{C}} \Phi(u_f(w)^2) \frac{dA(w)}{(1 + |w|^2)^2} = (4\pi)^{-1} \int_{\mathbb{S}^2} \Phi(|\langle \psi_w, \psi \rangle|^2) d\omega.$$

Thus, Theorem 4 follows immediately from Theorem 10 in Case 2. Similarly, Corollary 5 follows from Corollary 14.

### 4.3 Proof of Theorem 6

Let  $K \in \frac{1}{2}\mathbb{N} \setminus \{\frac{1}{2}\}$ . We consider the representation of  $SU(1, 1)$  on functions  $f$  on  $\mathbb{D}$  given as follows:

$$\pi_U(f)(z) := (\beta z + \bar{\alpha})^{-2K} f\left(\frac{\alpha z + \bar{\beta}}{\beta z + \bar{\alpha}}\right) \quad \text{for all } z \in \mathbb{C},$$

where

$$U = \begin{pmatrix} \alpha & \beta \\ \bar{\beta} & \bar{\alpha} \end{pmatrix} \in SU(1, 1), \quad \text{that is, } \alpha, \beta \in \mathbb{C} \quad \text{with } |\alpha|^2 - |\beta|^2 = 1.$$

This representation restricted to  $A_{2K}^2(\mathbb{D})$  is irreducible and unitary for the norm defined above. In this representation,

$$K_0 = z \frac{d}{dz} + K, \quad K_1 = \frac{1}{2i} \left( (z^2 - 1) \frac{d}{dz} + 2Kz \right), \quad K_2 = -\frac{1}{2} \left( (z^2 + 1) \frac{d}{dz} + 2Jz \right).$$

We may choose the space  $\mathcal{H}$  in Theorem 6 as  $A_{2K}^2(\mathbb{D})$ . By an explicit computation, one sees that the functions  $F_w$  are eigenvectors of the operator  $n_0 K_0 - n_1 K_1 - n_2 K_2$  corresponding to the eigenvalue  $K$ . Here,  $(n_0, n_1, n_2)$  is related to  $w$  as in the discussion before the statement of Theorem 6. Consequently, we can choose the phases in such a way that  $\psi_w = F_w$  for all  $w \in \mathbb{D}$ . Thus, using the explicit form of the  $F_w$ ,

$$\langle \psi_w, \psi \rangle = (1 + |w|^2)^K f(w)$$

with

$$f(w) := \frac{2K-1}{\pi} \int_{\mathbb{D}} (1 + w\bar{z})^{-2K} \psi(z) (1 + |z|^2)^{2K-2} dA(z).$$

Since  $\psi \in A_{2K}^2(\mathbb{D})$ , the reproducing property of the kernel implies that  $f(w) = \psi(w)$  for all  $w \in \mathbb{D}$ . Moreover,

$$u_f(w) = |\langle \psi_w, \psi \rangle|,$$

so Theorem 6 follows immediately from Theorem 10 in Case 3. Similarly, Corollary 7 follows from Corollary 14.

### 4.4 The limit of the discrete series

Two other irreducible unitary representations of  $SU(1, 1)$  are not in the discrete series, but are closely related to it, the so-called limits of discrete series [19, Chapter II]. They are typically not considered in the context of coherent states, since they are not square-integrable, but the questions discussed in this article make perfect sense for them and can be completely answered.

We restrict our attention to one of the limits of the discrete series, since the results for the other one can be deduced by appropriate complex conjugation. The construction of the coherent states is verbatim the same as for the discrete series, except that the value of  $K$  now is  $\frac{1}{2}$ .

**Proposition 19.** Consider the irreducible limit of the discrete series representation of  $SU(1, 1)$  on  $\mathcal{H}$ . Let  $\Phi : [0, 1] \rightarrow \mathbb{R}$  be nondecreasing. Then,

$$\sup \left\{ \int_{\mathbb{D}} \Phi(|\langle \psi_z, \psi \rangle|^2) \frac{dA(z)}{(1 - |z|^2)^2} : \psi \in \mathcal{H}, \|\psi\|_{\mathcal{H}} = 1 \right\} = \pi \int_0^1 \Phi(s) s^{-2} ds$$

and the supremum is attained for  $\psi = e^{i\theta} \psi_{z_0}$  with some  $z_0 \in \mathbb{D}$ ,  $\theta \in \mathbb{R}/2\pi\mathbb{Z}$ . If  $\Phi$  is strictly increasing near 1 and if the supremum is finite, then it is attained only for such  $\psi$ .

Note that the value of the integral with  $\psi = e^{i\theta} \psi_{z_0}$  does not depend on  $z_0 \in \mathbb{D}$ ,  $\theta \in \mathbb{R}/2\pi\mathbb{Z}$ . It may or may not be finite, depending on  $\Phi$ . For finiteness, it is necessary that  $\lim_{s \rightarrow 0^+} s^{-1} \Phi(s) = 0$ . In particular, the function  $\Phi(s) = s$  leads to an infinite supremum, which reflects the non-squareintegrability of the representation.

**Proof of Proposition 19.** We consider the same representation of  $SU(1, 1)$  on functions on  $\mathbb{D}$  as in the proof of Theorem 6 but with  $K = \frac{1}{2}$ . This representation is irreducible when restricted to the Hardy space  $H^2(\mathbb{D})$  and unitary for the norm defined above (see [19, Section II.6]). We choose the representation space  $\mathcal{H} = H^2(\mathbb{D})$ . The functions  $F_w$  were defined before Proposition 15, and one verifies that, by an appropriate choice of phases,  $\psi_w = F_w$ . It is well known that functions in the Hardy space have radial boundary values in  $L^2(\partial\mathbb{D})$  and that, in their norm, it suffices to consider this boundary value. Thus, using the explicit form of the  $F_w$ ,

$$\langle \psi_w, \psi \rangle = (2\pi)^{-1} \int_{-\pi}^{\pi} \overline{F_w(e^{i\varphi})} \psi(e^{i\varphi}) d\varphi = (1 + |w|^2)^{-\frac{1}{2}} f(w)$$

with

$$f(w) := (2\pi)^{-1} \int_{-\pi}^{\pi} (1 - we^{-i\varphi})^{-1} \psi(e^{i\varphi}) d\varphi.$$

By the reproducing property of the kernel (seen, for instance, by expanding both functions in the integrand into a Fourier series), we see that  $f(w) = \psi(w)$  for all  $w \in \mathbb{D}$ . Moreover,

$$u_f(w) = |\langle \psi_w, \psi \rangle|,$$

so Proposition 19 follows from Proposition 15.  $\square$

There is also an analog of Corollary 7 extending Proposition 19 (with convex  $\Phi$ ) to density matrices, but we omit it for the sake of brevity.

## 4.5 Proof of Theorem 8

Given  $\psi \in L^2(\mathbb{R}_+)$ , we can write

$$\langle \psi_{a,b}, \psi \rangle = a^\beta f(ia - b)$$

with

$$f(z) := 2^\beta \Gamma(2\beta)^{-\frac{1}{2}} \int_0^\infty x^{\beta-\frac{1}{2}} e^{izx} \psi(x) dx.$$

It is easy to see and known that  $f$  is analytic in  $\mathbb{C}_+ = \{z \in \mathbb{C} : \text{Im} z > 0\}$  and that

$$\frac{\beta - \frac{1}{2}}{2\pi} \int_{\mathbb{C}_+} |f(z)|^2 (\operatorname{Im} z)^{2\beta-2} dA(z) = \frac{\beta - \frac{1}{2}}{2\pi} \iint_{\mathbb{R}_+ \times \mathbb{R}} |\langle \psi_{a,b}, \psi \rangle|^2 \frac{dad b}{a^2} = \|\psi\|_{L^2(\mathbb{R}_+)}^2,$$

where the last identity is the completeness relation of the coherent states [12, (2.10)]. Consider the conformal map  $\Sigma : \mathbb{C}_+ \rightarrow \mathbb{D}$ ,

$$\Sigma(z) = \frac{z - i}{-iz + 1} \quad \text{for } z \in \mathbb{C}_+, \quad \Sigma^{-1}(\zeta) = \frac{\zeta + i}{i\zeta + 1} \quad \text{for } \zeta \in \mathbb{D}.$$

Setting

$$g(\zeta) := (i\zeta + 1)^{-2\beta} f\left(\frac{\zeta + i}{i\zeta + 1}\right) \quad \text{for all } \zeta \in \mathbb{D},$$

we find that  $g$  is analytic in  $\mathbb{D}$  and, using  $dA(\zeta) = |\Sigma'(z)|^2 dA(z)$  for  $\zeta = \Sigma(z)$ ,

$$\frac{\beta - \frac{1}{2}}{2\pi} \int_{\mathbb{C}_+} |f(z)|^2 (\operatorname{Im} z)^{2\beta-2} dA(z) = \frac{2\beta - 1}{\pi} \int_{\mathbb{D}} |g(\zeta)|^2 (1 - |\zeta|^2)^{2\beta-2} dA(\zeta) = \|g\|_{A_{2\beta}(\mathbb{D})}^2.$$

Moreover, after a simple computation,

$$u_g(\Sigma(ia - b)) = |\langle \psi_{a,b}, \psi \rangle|,$$

and therefore,

$$\iint_{\mathbb{R}_+ \times \mathbb{R}} \Phi(|\langle \psi_{a,b}, \psi \rangle|^2) \frac{dad b}{a^2} = 4\pi \int_{\mathbb{D}} \Phi(u_g(\zeta)^2) dm(\zeta).$$

In addition, the coherent states  $\psi_{a,b}$ ,  $(a, b) \in \mathbb{R}_+ \times \mathbb{R}$ , are in one-to-one correspondence with the functions  $F_w$ ,  $w \in \mathbb{D}$ . Indeed, a straightforward computation shows that the  $f$  corresponding to  $\psi = \psi_{1,0}$  is  $f(z) = (2i/(z + i))^{2\beta}$ , which corresponds to  $g(\zeta) = 1 = F_0(\zeta)$ . The result, in the general case, follows from the facts that every  $(a, b)$  can be moved to  $(1, 0)$  by an  $aX + b$ -action and every point  $w \in \mathbb{D}$  can be moved to 0 by the action of a subgroup of  $SU(1, 1)$  isomorphic to  $aX + b$  (see, e.g., [26, equation after (1.3)]) and that  $\Sigma$  relates these actions to each other.

Thus, Theorem 8 follows immediately from Theorem 10 in Case 3 with  $\alpha = 2\beta$  and, similarly, Corollary 9 follows from Corollary 14.

**Remark 20.** The functions  $\psi_{a,b}$  are also well defined for  $\beta \in (0, \frac{1}{2}]$ . In this case, the coherent state transform cannot be normalized to be an isometry to a subset of  $L^2(\mathbb{R}_+ \times \mathbb{R}, a^{-2}dad b)$ , but the optimization problem in Theorem 8 still makes sense. We claim that, for  $\beta = \frac{1}{2}$ , Theorem 8 remains valid, replacing the assumptions “convex” and “not linear” on  $\Phi$  by “nondecreasing” and “strictly increasing near 1,” respectively. Indeed, in this case, the function  $f$  in the previous proof belongs to the Hardy space  $H^2(\mathbb{C}_+)$  and, by Plancherel, its norm in that space is equal to  $\|\psi\|_{L^2(\mathbb{R}_+)}$ . Mapping  $\mathbb{C}_+$  to  $\mathbb{D}$  via  $\Sigma$ , we can deduce the assertion from Proposition 15. We do not know whether Theorem 8 extends to  $\beta \in (0, \frac{1}{2})$ .

## 4.6 Limitations of the method

In this article, we have discussed the cases of the Heisenberg group,  $SU(2)$ ,  $SU(1, 1)$ , and the affine group. It is a natural question, potentially of relevance for representation theory, to which extent the results can be generalized to arbitrary Lie groups.

While the method of the present article is able to treat various cases in a unified way, it will probably not be able to deal with the general case, as we argue now. One of the key ingredients in the argument is Lemma 12, whose proof uses the fact that the superlevel sets of the overlap of two coherent states are isoperimetric set. This may fail in general, as we are going to show.

Following Lieb and Solovej [25], we consider the case of symmetric representations of  $SU(N)$ . We fix  $N \geq 3$ . The relevant representations are labeled by  $M \in \mathbb{N}$  and we choose the representation space  $\mathcal{H}$  to be the symmetric subspace of the tensor product  $\otimes^M \mathbb{C}^N$ . Coherent states are defined through elements of the form  $\otimes^M z$  with  $z \in \mathbb{C}^N$ ,  $|z| = 1$ . Note that if two  $z$ 's differ by a phase, then the corresponding vectors  $\otimes^M z$  in  $\mathcal{H}$  also differ by a phase and correspond to the same state. Thus, we will label the coherent states by points  $z$  in the complex projective space

$$\mathbb{C}P^{N-1} = \{z \in \mathbb{C}^N : |z| = 1\} / \sim,$$

where  $z \sim w$  if  $z = e^{i\theta}w$  for some  $\theta \in \mathbb{R}/2\pi\mathbb{Z}$ . We denote integration with respect to the natural  $SU(N)$ -invariant probability measure on  $\mathbb{C}P^{N-1}$  by  $dz$ .

Lieb and Solovej have solved the corresponding problem and shown that, for any convex  $\Phi : [0, 1] \rightarrow \mathbb{R}$ ,

$$\sup \left\{ \int_{\mathbb{C}P^{N-1}} \Phi(|\langle \otimes^M z, \psi \rangle|^2) dz : \psi \in \mathcal{H}, \|\psi\|_{\mathcal{H}} = 1 \right\}$$

is attained for coherent states.

If we tried to reprove this through the method in the present article, we would consider the measure of the superlevel sets of the function  $z \mapsto |\langle \otimes^M z, \psi \rangle|$  and try to prove some monotonicity properties of it. This monotonicity property should be saturated if  $\psi$  is of the form  $\otimes^M z_0$ . In this special case, the level sets are of the form

$$\{z \in \mathbb{C}P^{N-1} : |z^* z_0|^M > \kappa\}.$$

These are geodesic balls (see, e.g., [15, Example 2.110]), and if we want to use the method based on an isoperimetric inequality, they should be optimizers for the isoperimetric inequality. (More precisely, this isoperimetric property should hold for all  $\kappa$  for which the measure of the corresponding set is  $\leq \frac{1}{2}$ . For  $\kappa$  such that the measure is  $\geq \frac{1}{2}$ , the complement of the corresponding set should be an isoperimetric set.) This, however, is not the case for all  $\kappa$ , at least not for  $N = 4$ , as pointed out in [4, Appendix]; see also [28, Remark 4.2]. For the solution of the isoperimetric problem in  $\mathbb{C}P^{N-1}$ , see also [28, Theorem 4.1]. The isoperimetric sets are expected to transition from geodesic balls for small volumes to tubes around some  $\mathbb{C}P^{M-1} \subset \mathbb{C}P^{N-1}$  for intermediate volumes.

## 5 Faber-Krahn-type inequalities for the coherent state transform

The main result of the recent article [29] by Nicola and Tilli states that, for any measurable set  $E \subset \mathbb{R}^2$  of finite measure,

$$\iint_E |\langle \psi_{p,q}, \psi \rangle|^2 dp dq \leq 2\pi\hbar(1 - e^{-(2\pi\hbar)^{-1}|E|}) \quad (10)$$

with equality if and only if  $\psi = e^{i\theta}\psi_{p_0,q_0}$  for some  $p_0, q_0 \in \mathbb{R}$ ,  $\theta \in \mathbb{R}/2\pi\mathbb{Z}$ , and  $E$  is equal to a ball centered at  $(p_0, q_0)$  (up to sets of measure zero). (Here, we restrict ourselves to the one-dimensional case of their result. Since the proof of Theorem 1 extends to higher dimensions, the discussion in this subsection does so as well.)

We claim that the inequality (10) follows by abstract arguments from Theorem 1. Of course, this is not too surprising, since Kulikov's arguments, which we have adapted to yield a proof of Theorem 1, are inspired by those in [29]. Nevertheless, this observation will allow us to derive an analog of the Nicola-Tilli results in the  $SU(2)$ ,  $SU(1, 1)$  and  $aX + b$  cases.



**Proof of (10) given Theorem 1.** Fixing  $p_0, q_0 \in \mathbb{R}$  and  $\psi \in L^2(\mathbb{R})$  with  $\|\psi\|_{L^2(\mathbb{R})} = 1$ , we can write the first assertion of Theorem 1 as the statement that

$$\iint_{\mathbb{R} \times \mathbb{R}} \Phi(|\langle \psi_{p,q}, \psi \rangle|^2) dp dq \leq \iint_{\mathbb{R} \times \mathbb{R}} \Phi(|\langle \psi_{p,q}, \psi_{p_0,q_0} \rangle|^2) dp dq$$

for any convex function  $\Phi$  on  $[0, 1]$ . By Hardy-Littlewood majorization theory (see, e.g., [16, Theorems 108, 249, 250], [1, Corollary 2.1], [42, Theorem 15.27], and also [7, Chapter 2, Proposition 3.3]), this is equivalent to the fact that

$$\iint_E |\langle \psi_{p,q}, \psi \rangle|^2 dp dq \leq \sup_{|F|=|E|} \iint_F |\langle \psi_{p,q}, \psi_{p_0,q_0} \rangle|^2 dp dq$$

for any measurable set  $E \subset \mathbb{R}^2$  of finite measure. By an explicit computation,

$$|\langle \psi_{p,q}, \psi_{p_0,q_0} \rangle| = e^{-\frac{1}{4\hbar}((q-q_0)^2 + (p-p_0)^2)}.$$

This is symmetric decreasing around  $(p_0, q_0)$ , and therefore, the supremum above is attained if (and only if)  $F$  is a ball centered at  $(p_0, q_0)$  (up to sets of measure zero). In this case, the right side can be computed to be  $1 - e^{-(2\pi\hbar)^{-1}|E|}$ , yielding (10).  $\square$

By carefully going through the majorization argument, it should be possible to deduce the equality statement by Nicola and Tilli from the equality statement in Theorem 1, but we omit this here.

Obviously, the above argument can be generalized to the  $SU(2)$ ,  $SU(1, 1)$ , and  $aX + b$  cases. For the sake of brevity, we leave out a statement about the cases of equality.

**Theorem 21.** Let  $J \in \frac{1}{2}\mathbb{N}$  and consider an irreducible  $(2J + 1)$ -dimensional representation of  $SU(2)$  on  $\mathcal{H}$ . Then, for any  $\psi \in \mathcal{H}$  with  $\|\psi\|_{\mathcal{H}} = 1$  and any measurable  $E \subset \mathbb{S}^2$ ,

$$\int_E |\langle \psi_\omega, \psi \rangle|^2 d\omega \leq \frac{4\pi}{2J+1} \left( 1 - \left( 1 - \frac{|E|}{4\pi} \right)^{2J+1} \right).$$

Equality is attained if  $\psi = e^{i\theta} \psi_{\omega_0}$  with some  $\omega_0 \in \mathbb{S}^2$ ,  $\theta \in \mathbb{R}/2\pi\mathbb{Z}$ , and  $E$  is a spherical cap centered at  $\omega_0$ .

**Theorem 22.** Let  $K \in \frac{1}{2}\mathbb{N} \setminus \{\frac{1}{2}\}$  and consider the irreducible discrete series representation of  $SU(1, 1)$  on  $\mathcal{H}$  corresponding to  $K$ . Then, for any  $\psi \in \mathcal{H}$  with  $\|\psi\|_{\mathcal{H}} = 1$  and any measurable  $E \subset \mathbb{D}$ ,

$$\int_E |\langle \psi_z, \psi \rangle|^2 \frac{dA(z)}{(1 - |z|^2)^2} \leq \frac{\pi}{2K-1} (1 - (1 + m(E))^{-2K+1}),$$

where  $dm(z) = \pi^{-1}(1 - |z|^2)^{-2} dA(z)$ . Equality is attained if  $\psi = e^{i\theta} \psi_{z_0}$  with some  $z_0 \in \mathbb{D}$ ,  $\theta \in \mathbb{R}/2\pi\mathbb{Z}$ , and  $E$  is a hyperbolic ball centered at  $z_0$ .

**Theorem 23.** Let  $\beta > \frac{1}{2}$ . Then, for any  $\psi \in L^2(\mathbb{R}_+)$  with  $\|\psi\|_{L^2(\mathbb{R}_+)} = 1$  and any measurable  $E \subset \mathbb{R}_+ \times \mathbb{R}$ ,

$$\int_E |\langle \psi_{a,b}, \psi \rangle|^2 \frac{da db}{a^2} \leq \frac{4\pi}{2\beta-1} (1 - (1 + (4\pi)^{-1}\mu(E))^{-2\beta+1}),$$

where  $d\mu(a, b) = a^{-2} da db$ . Equality is attained if  $\psi = e^{i\theta} \psi_{a_0, b_0}$  with some  $a_0 \in \mathbb{R}_+$ ,  $b_0 \in \mathbb{R}$ ,  $\theta \in \mathbb{R}/2\pi\mathbb{Z}$ , and  $E$  is a hyperbolic ball centered at  $(a_0, b_0)$ .

In Theorems 22 and 23, by a “hyperbolic ball,” we mean a geodesic ball with respect to the hyperbolic metric on  $\mathbb{D}$  and  $\mathbb{C}_+$  (identified with  $\mathbb{R}_+ \times \mathbb{R}$ ), respectively.

Theorem 23, including a characterization of equality cases, has recently been proved in [35] by a direct adaptation of the method in [29]. Our proof, based on Theorem 8, is different.

**Proof of Theorems 21, 22, and 23.** As above, one can show that the left sides in the theorems are bounded by the supremum of the integral of  $|\langle \psi_\alpha, \psi_{\alpha_0} \rangle|^2$  over sets  $F$  of the same measure as  $E$ . Here, the index  $\alpha$  labels  $\omega \in \mathbb{S}^2$ ,  $z \in \mathbb{D}$ , and  $(a, b) \in \mathbb{R}_+ \times \mathbb{R}$  in the three cases, respectively, and  $\alpha_0$  is a fixed such index. By the bathtub principle, the supremum over  $F$  is attained at a set of the form  $\{|\langle \psi_\alpha, \psi_{\alpha_0} \rangle| > \kappa_0\} \cup G$ , where  $G$  is a measurable subset of  $\{|\langle \psi_\alpha, \psi_{\alpha_0} \rangle| = \kappa_0\}$ .

To complete the proof, we will need some explicit knowledge about the function  $|\langle \psi_\alpha, \psi_{\alpha_0} \rangle|$ . We choose the representation space  $\mathcal{H}$  in Theorems 21 and 22 in the same way as in the proofs of Theorems 4 and 6, namely as  $\mathcal{P}_{2J}$  and  $A_{2K}^2(\mathbb{D})$ , respectively. Then, as shown there,  $|\langle \psi_\alpha, \psi_{\alpha_0} \rangle| = u_{F_\omega}(z)$ , where  $\alpha = S(z)$  and  $\alpha = z$  in the first two cases and, similarly,  $\alpha_0 = S(w)$  and  $\alpha_0 = w$ . In the third case, if  $\alpha = (a, b)$ , then  $z = \Sigma(ia - b)$  and similarly for  $\alpha_0$  and  $w$ . In particular,  $w = 0$  if we choose  $\alpha_0$  to be  $\omega_0 = (0, 0, 1)$ ,  $z = 0$ , and  $(a_0, b_0) = (1, 0)$  in different cases. The explicit definition of  $u_f$  then shows that  $\{|\langle \psi_\alpha, \psi_{\alpha_0} \rangle| > \kappa_0\}$  is a spherical cap in the first case or a hyperbolic ball in the last two cases and that, in all cases,  $\{|\langle \psi_\alpha, \psi_{\alpha_0} \rangle| = \kappa_0\}$  has measure zero. Thus, the set  $G$  above can be ignored, and we have identified the optimal set in the case of a special choice of  $\alpha_0$ .

This, in fact, yields the shape for an arbitrary choice of  $\alpha_0$ . Indeed, as discussed before Theorem 10, the functions  $\alpha \mapsto \langle \psi_\alpha, \psi_{\alpha_0} \rangle$  are equimeasurable for different  $\alpha_0$ 's and one such function can be obtained from another by the action of  $SU(2)$  or  $SU(1, 1)$ . Since this action maps spherical caps to spherical caps, or hyperbolic balls to hyperbolic balls, we obtain that the supremum is attained for any  $\alpha_0$  at such a set.

It remains to compute the supremum. It is convenient to do this in terms of the functions  $u_f$ . The second and the third cases can be treated together with the convention that  $\alpha = 2K$  in the second and  $\alpha = 2\beta$  in the third case. We have, with an arbitrary  $F \in \mathcal{M}$ ,

$$\int_{\{u_F > \kappa_0\}} u_F(z)^2 dm(z) = \begin{cases} (4\pi)^{-1} \int_{\{|\langle \psi_\omega, \psi_{\omega_0} \rangle| > \kappa_0\}} |\langle \psi_\omega, \psi_{\omega_0} \rangle|^2 d\omega, \\ \pi^{-1} \int_{\{|\langle \psi_z, \psi_{z_0} \rangle| > \kappa_0\}} |\langle \psi_z, \psi_{z_0} \rangle|^2 \frac{dA(z)}{(1 - |z|^2)^2}. \end{cases}$$

Meanwhile, by the layer cake formula,

$$\begin{aligned} \int_{\{u_F > \kappa_0\}} u_F(z)^2 dm(z) &= 2 \int_0^{\kappa_0} m(\{u_F > \kappa\}) \kappa d\kappa + 2 \int_0^{\kappa_0} m(\{u_F > \kappa\}) \kappa d\kappa \\ &= \int_{\Omega} u_F(z)^2 dm(z) - 2 \int_0^{\kappa_0} (m(\{u_F > \kappa\}) - m(\{u_F > \kappa_0\})) \kappa d\kappa. \end{aligned} \quad (11)$$

The first term on the right side is equal to

$$\int_{\Omega} u_F(z)^2 dm(z) = c \|F\|^2 = c$$

with  $c = (2J + 1)^{-1}$  and  $c = (\alpha - 1)^{-1}$  in different cases. For the second term on the right side of (11), we use the explicit expressions for  $m(\{u_F > \kappa\})$  from the proof of Lemma 12 and obtain, after a computation,

$$2 \int_0^{\kappa_0} (m(\{u_F > \kappa\}) - m(\{u_F > \kappa_0\})) \kappa d\kappa = \begin{cases} c \kappa_0^{\frac{2J+1}{J}}, \\ c \kappa_0^{\frac{2(\alpha-1)}{\alpha}}. \end{cases} \quad (12)$$

This gives the expression of the supremum in terms of  $\kappa_0$ . The parameter  $\kappa_0$  satisfies

$$m(\{u_F > \kappa_0\}) = \begin{cases} (4\pi)^{-1}|E| & \text{in the case of Theorem 21,} \\ m(E) & \text{in the case of Theorem 22,} \\ (4\pi)^{-1}\mu(E) & \text{in the case of Theorem 23.} \end{cases} \quad (13)$$

(In the last case, we used the fact that  $(4\pi)^{-1}\mu(E) = m(\Sigma(\tilde{E}))$ , where  $\Sigma$  is the conformal map from  $\mathbb{C}_+$  to  $\mathbb{D}$  from the proof of Theorem 8 and  $\tilde{E} \subset \mathbb{C}_+$  is obtained from  $E$  by identifying  $(a, b) \in \mathbb{R}_+ \times \mathbb{R}$  with  $ia - b \in \mathbb{C}_+$ .) Using (13) and the expressions from the proof of Lemma 12, we can express  $\kappa_0$  in terms of the measure of  $E$ . Inserting this into (12) gives an expression for the second term on the right side of (11). This leads to the claimed explicit form of the upper bound.  $\square$

**Funding information:** Partial support through the US National Science Foundation grant DMS-1954995, as well as through the Deutsche Forschungsgemeinschaft (German Research Foundation) through Germany's Excellence Strategy EXC-2111-390814868, is acknowledged.

**Conflict of interest:** The author states that there is no conflict of interest.

## References

- [1] A. Alvino, G. Trombetti, and P.-L. Lions, *On optimization problems with prescribed rearrangements*, Nonlinear Anal. **13** (1989), no. 2, 185–220.
- [2] E. W. Aslaksen and J. R. Klauder, *Unitary representations of the affine group*, J. Mathematical Phys. **9** (1968), 206–211.
- [3] J. Bandyopadhyay, *Optimal concentration for  $SU(1, 1)$  coherent state transforms and an analog of the Lieb-Wehrl conjecture for  $SU(1, 1)$* , Comm. Math. Phys. **285** (2009), no. 3, 1065–1086.
- [4] J. L. Barbosa, M. doCarmo, and J. Eschenburg, *Stability of hypersurfaces of constant mean curvature in Riemannian manifolds*, Math. Z. **197** (1988), no. 1, 123–138.
- [5] V. Bargmann, *Irreducible unitary representations of the Lorentz group*, Ann. Math. **48** (1947), no. 2, 568–640.
- [6] F. Bayart, O. F. Brevig, A. Haimi, J. Ortega-Cerdà, and K.-M. Perfekt, *Contractive inequalities for Bergman spaces and multiplicative Hankel forms*, Trans. Amer. Math. Soc. **371** (2019), no. 1, 681–707.
- [7] C. Bennett and R. Sharpley, *Interpolation of operators*, Pure and Applied Mathematics, vol. 129, Academic Press, Inc., Boston, MA, 1988.
- [8] F. Bernstein, *Über die isoperimetrische Eigenschaft des Kreises auf der Kugeloberfläche und in der Ebene*, Math. Ann. **60** (1905), no. 1, 117–136.
- [9] B. G. Bodmann, *A lower bound for the Wehrl entropy of quantum spin with sharp high-spin asymptotics*, Comm. Math. Phys. **250** (2004), no. 2, 287–300.
- [10] E. A. Carlen, *Some integral identities and inequalities for entire functions and their application to the coherent state transform*, J. Funct. Anal. **97** (1991), no. 1, 231–249.
- [11] P. L. Chebyshev, *On approximate expressions of some integrals in terms of others, taken within the same limits*, Proc. Math. Soc. Kharkov **2** (1882), 93–98.
- [12] I. Daubechies, J. R. Klauder, and T. Paul, *Wiener measures for path integrals with affine kinematic variables*, J. Math. Phys. **28** (1987), no. 1, 85–102.
- [13] G. De Palma, *The Wehrl entropy has Gaussian optimizers*, Lett. Math. Phys. **108** (2018), no. 1, 97–116.
- [14] R. L. Frank and E. H. Lieb, *Sharp constants in several inequalities on the Heisenberg group*, Ann. Math. (2) **176** (2012), no. 1, 349–381.
- [15] S. Gallot, D. Hulin, and J. Lafontaine, *Riemannian Geometry*, 3rd edition, Universitext., Springer-Verlag, Berlin, 2004.
- [16] G. H. Hardy, J. E. Littlewood and G. Pólya, *Inequalities*, Reprint of the 1952 edition, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1988.
- [17] D. Jerison and J. M. Lee, *Extremals for the Sobolev inequality on the Heisenberg group and the CR Yamabe problem*, J. Amer. Math. Soc. **1** (1988), no. 1, 1–13.
- [18] D. Kalaj, *Contraction Property of Differential Operator on Fock Space*, Preprint (2022), arXiv:2207.13606.
- [19] A. W. Knap, *Representation Theory of Semisimple Groups. An Overview Based on Examples*, Reprint of the 1986 Original, Princeton Landmarks in Mathematics, Princeton University Press, Princeton, NJ, 2001.
- [20] A. Kulikov, *Functionals with extrema at reproducing kernels*, Geom. Funct. Anal. **32** (2022), no. 4, 938–949.

- [21] A. Kulikov, F. Nicola, J. Ortega-Cerdà and P. Tilli, *A Monotonicity Theorem for Subharmonic Functions on Manifolds*, Preprint, 2022, arXiv:2212.14008.
- [22] P. Lévy, *Leçons d'analyse fonctionnelle*, Gauthier-Villars, Paris, 1922.
- [23] E. H. Lieb, *Proof of an entropy conjecture of Wehrl*, Comm. Math. Phys. **62** (1978), no. 1, 35–41.
- [24] E. H. Lieb and J. P. Solovej, *Proof of an entropy conjecture for Bloch coherent spin states and its generalizations*, Acta Math. **212** (2014), no. 2, 379–398.
- [25] E. H. Lieb and J. P. Solovej, *Proof of the Wehrl-type entropy conjecture for symmetric  $SU(N)$  coherent states*, Comm. Math. Phys. **348** (2016), no. 2, 567–578.
- [26] E. H. Lieb and J. P. Solovej, *Wehrl-type coherent state entropy inequalities for  $SU(1, 1)$  and its  $AX+B$  subgroup*, in: *Partial Differential Equations, Spectral Theory, and Mathematical Physics - The Ari Laptev Anniversary Volume*, EMS Ser. Congr. Rep., EMS Press, Berlin, 2021, 301–314.
- [27] S. Luo, *A simple proof of Wehrl's conjecture on entropy*, J. Phys. A **33** (2000), 3093–3096.
- [28] Y. Memarian, *The Isoperimetric Inequality on Compact Rank One Symmetric Spaces and Beyond*, Preprint, 2021, arXiv:1710.03952.
- [29] F. Nicola and P. Tilli, *The Faber-Krahn inequality for the short-time Fourier transform*, Invent. Math. **230** (2022), no. 1, 1–30.
- [30] F. Nicola and P. Tilli, *The Norm of Time-Frequency Localization Operators*, Preprint, 2022, arXiv:2207.08624.
- [31] R. Osserman, *The isoperimetric inequality*, Bull. Amer. Math. Soc. **84** (1978), no. 6, 1182–1238.
- [32] A. Perelomov, *Coherent states for arbitrary Lie group*, Comm. Math. Phys. **26** (1972), 222–236.
- [33] A. Perelomov, *Generalized coherent states and their applications*, *Texts and Monographs in Physics*. Springer-Verlag, Berlin, 1986.
- [34] T. Rado, *The isoperimetric inequality on the sphere*, Amer. J. Math. **57** (1935), no. 4, 765–770.
- [35] J. P. G. Ramos and P. Tilli, *A Faber-Krahn Inequality for Wavelet Transforms*, Preprint, 2022, arXiv:2205.07998.
- [36] E. Schmidt, *Über die isoperimetrische Aufgabe im  $n$ -dimensionalen Raum konstanter negativer Krümmung. I. Die isoperimetrischen Ungleichungen in der hyperbolischen Ebene und für Rotationskörper im  $n$ -dimensionalen hyperbolischen Raum*, Math. Z. **46** (1940), 204–230.
- [37] E. Schmidt, *Die isoperimetrischen Ungleichungen auf der gewöhnlichen Kugel und für Rotationskörper im  $n$ -dimensionalen sphärischen Raum*, Math. Z. **46** (1940), 743–794.
- [38] P. Schupp, *On Lieb's conjecture for the Wehrl entropy of Bloch coherent states*, Comm. Math. Phys. **207** (1999), no. 2, 481–493.
- [39] P. Schupp, *Wehrl entropy, coherent states and quantum channels*, in: *The Physics and Mathematics of Elliott Lieb*, Vol. II, EMS Press, Berlin, 2022, p. 329–344.
- [40] B. Simon, *The classical limit of quantum partition functions*, Comm. Math. Phys. **71** (1980), no. 3, 247–276.
- [41] B. Simon, *Representations of finite and compact groups*, *Graduate Studies in Mathematics*, vol. 10, American Mathematical Society, Providence, RI, 1996.
- [42] B. Simon, *Convexity. An analytic viewpoint*, *Cambridge Tracts in Mathematics*, vol. 187, Cambridge University Press, Cambridge, 2011.
- [43] G. Talenti, *Elliptic equations and rearrangements*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. **3** (1976), no. 4, 697–718.
- [44] A. Wehrl, *On the relation between classical and quantum-mechanical entropy*, Rep. Math. Phys. **16** (1979), no. 3, 353–358.