

Review

Susanne H. Kirsch, Wolfgang Herrmann and Rima Obeid*

Genetic defects in folate and cobalamin pathways affecting the brain

Abstract

Folate and cobalamin are necessary for early brain development and function. Deficiency of folate or cobalamin during pregnancy can cause severe malformation in the central nervous system such as neural tube defects. After birth, folate and cobalamin deficiency can cause anemia, failure to thrive, recurrent infections, psychiatric and neurological symptoms. The folate and the homocysteine metabolic pathways interact at a central step where 5-methyltetrahydrofolate donates its methyl group to homocysteine to produce methionine and tetrahydrofolate. Methyl cobalamin and folate interact at this critical step. Both nutrients have a crucial role in DNA synthesis and in delivering S-adenosylmethionine, the universal methyl donor. Severe and mild inherited disorders in folate and cobalamin pathways have been described. The two groups of disorders share some similarities, but differ in the molecular mechanism, metabolic dysregulation, and disease management. This review summarizes selected disorders, including rare and common mutations that affect folate and cobalamin absorption, transport, or dependent enzymes. When the mutations are discovered early enough, many of the described disorders are easily treatable by B vitamin supplementation, which often prevents or reverses the manifestation of the disease. Therefore, the screening for mutations is recommended and should be carried out as early as possible: after occurrence of the first symptoms or when a certain constellations of the folate and cobalamin related markers are measured, such as elevated homocysteine and/or methylmalonic acid.

Keywords: brain; cobalamin; folate.

Introduction

It is our great pleasure to congratulate *Clinical Chemistry and Laboratory Medicine* (CCLM) on its 50th anniversary and to contribute this article for the celebration issue. CCLM, an internationally recognized and leading journal in the field of clinical chemistry and laboratory medicine, also plays an important role in the scientific area of hyperhomocysteinemia (HHcy) and B vitamin deficiency. One of the first CCLM publications in this field dates back to the year 1998 [1] and examined the relationship between HHcy and cardiovascular diseases. In the period from 2000 to 2012, CCLM has published four international conferences on HHcy and B vitamin deficiency which have been organized by our group (2nd Conference on Hyperhomocysteinemia [2], 3rd Conference on Hyperhomocysteinemia [3], 4th Conference on Hyperhomocysteinemia [4], and the World Congress on Hyperhomocysteinemia [5]) with more than 100 articles. To date over 200 articles dealing with homocysteine (Hcy) and B vitamin deficiency have been published in CCLM. This reflects the contribution that CCLM has made and underlines the important role of the journal for the scientific community active in that research field. HHcy and B vitamin deficiency are still the focus of medical research but topics have changed during the last decade. Ten years ago the main interest in HHcy research was directed towards its relationship with cardiovascular diseases, however, the current focus lies in its association with neurodegenerative diseases [6, 7]. Large scale intervention has failed to show that B vitamin supplementation improved patients cardiovascular outcome [8]. A very recent meta-analysis with 47,921 participants from 19 studies (plasma Hcy was reduced in all studies) found that B vitamin supplementation has a significant protective effect on stroke, but none on the risk of cardiovascular disease, myocardial infarction, chronic heart disease, cardiovascular death, or all-cause mortality [9]. It has also been shown that HHcy and B vitamin deficiency is correlated to neurological and psychiatric diseases as well as cognitive decline [7, 10, 11]. The important role of biomarkers in neurodegenerative

*Corresponding author: Prof. Dr. Rima Obeid, Department of Clinical Chemistry and Laboratory Medicine, University Hospital of Saarland, Building 57, 66421 Homburg/Saar, Germany, Phone: +49 68411630701, Fax: +49 68411630703, E-mail: rima.obeid@uniklinikum-saarland.de

Susanne H. Kirsch and Wolfgang Herrmann: Department of Clinical Chemistry and Laboratory Medicine, University Hospital of Saarland, Homburg, Germany

diseases has been highlighted in a special issue of CCLM (Biomarkers of Neurodegenerative Diseases) [7, 12–21]. Moreover, recent publications provided convincing evidence that B vitamin supplementation significantly slowed brain atrophy and cognitive decline in patients with mild cognitive impairment after 2 years of treatment [22, 23].

Plausible pathomechanisms behind this relationship are, on the one hand, oxidative stress and other neurotoxic the mechanisms and a lowered cellular methylation potential caused by HHcy and B vitamin deficiency [10, 24, 25]. On the other hand, HHcy and B vitamin deficiency result from lifestyle factors but genetic factors also contribute significantly to dysregulated metabolic pathways in this respect. Beside the very common 5,10-methylenetetrahydrofolate reductase (MTHFR, EC 1.5.1.20) polymorphism that influences remethylation of Hcy to methionine [26] many other genetic mutations affect folate and cobalamin dependent pathways [27] and have disease modifying effects [28]. Beside disturbed DNA synthesis, DNA hypomethylation due to HHcy and B vitamin deficiency is an important epigenetic factor in gene regulation and expression and has impact on disease development [29, 30]. This review is focused on the role of genetic defects in folate and cobalamin metabolism and their impact on brain function.

Genetic defects in folate pathway

Folate metabolism and distribution

Folate is a methyl donor for Hcy methylation to methionine. It participates in the *de novo* synthesis of purines and thymidylates [deoxythymidine monophosphate (dTMP)]. Therefore, folate cycle links the DNA synthesis with the methyl group metabolism. The universal methyl donor, S-adenosylmethionine (SAM), is synthesized from methionine and it regulates the coordination between folate and methionine metabolism. Folate and cobalamin deficiency as well as HHcy at conception may negatively influence the health of the offspring [31] and have been related to early pregnancy loss and other congenital birth defects [32]. Several mutations in genes of folate-catabolizing enzymes and transporting proteins have been described. Many of these disorders affect the central nervous system and are associated with severe clinical symptoms that are manifested at early infancy.

Dietary folates are derivatives of folylpolyglutamates that are enzymatically hydrolyzed upon ingestion into monoglutamates in the brush border cells of the

duodenum and jejunum. The hydrolysis of the polyglutamate is mediated by glutamate carboxypeptidase II (GCP II, EC 3.4.17.21) or the exopeptidase γ -glutamyl hydrolase (GGH, EC 3.4.19.9) located in the lysosome [33]. The folylpoly- γ -glutamate synthase (FPGS, EC 6.3.2.17) facilitates folate polyglutamation. The anionic nature of the folate oligomers renders them unable to leave the cell and, therefore, the polyglutamate forms of reduced folates (and certain antifolates) accumulate in the cell. Only the monoglutamates can be transported across cell membranes [34].

Three independent types of membrane systems are responsible for cross membrane transport of folate monoglutamate forms. These are the membrane folate receptor (FR), the reduced folate carrier 1 (RFC1), and the proton-coupled folate transporter/heme carrier protein 1 (PCFT/HCP1). The FR has high affinity for folate (K_m approx. 1 nmol/L) and conducts the receptor-mediated endocytosis (unidirectional) across the cell membrane at neutral pH [35]. FR α (FOLR1) is expressed in certain epithelial cells, the choroid plexus, the placenta [36], and the kidney. FR β (FOLR2) is expressed in the fetal brain, kidney, placenta, spleen, and thymus [37]. RFC1 (SLC19A1) has a higher affinity for 5-methyltetrahydrofolate (5-methylTHF) than for folic acid and transports folate at an optimal pH of 7.5 [38]. The PCFT (SLC46A1) system acts at an optimal pH in the acidic range (4.5–5.5) explaining its role as the major intestinal folate transporter [39]. Transport of folates into the cerebrospinal fluid (CSF) occurs in the choroid plexus, where 5-methylTHF is transported across the blood-brain barrier by FR α in adult or FR β in the fetal brain. PCFT is ubiquitously expressed in the human brain, where it functions in concert with FR α and FR β or might export folates after FR α -mediated endocytosis [37, 40]. Biomarkers of folate and methionine metabolism in blood are important determinants of CSF levels of the metabolites [24, 41].

Tetrahydrofolate (THF) represents the active form of folate that is formed from dihydrofolate (DHF) by means of dihydrofolate reductase (DHFR, EC 1.5.1.3) (Figure 1). 5,10-MethyleneTHF is converted to 5-methylTHF by MTHFR in an irreversible reaction [42]. 5-MethylTHF donates its methyl group to Hcy and is converted into THF by means of methionine synthase (MTR, EC 2.1.1.13), a cobalamin (Cbl)-dependent enzyme. The enzyme methionine synthase reductase (MTRR, EC 1.16.1.8) is involved in the reductive regeneration of the Cbl-cofactor, which is required for MTR function. 10-FormylTHF is utilized for purine synthesis and 5,10-methyleneTHF for synthesis of dTMP and methionine. Serine hydroxymethyltransferase 1 (SHMT1, EC 2.1.2.1) utilizes serine to form 5,10-methyleneTHF [43]. The conversion




Figure 1 Folate metabolism, related enzymes, and transporters.

DHF, dihydrofolate; DHFR, dihydrofolate reductase; dTMP, deoxythymidine monophosphate; dUMP, deoxyuridine monophosphate; FR, folate receptor; MTHFD, 5,10-methylenetetrahydrofolate dehydrogenase; MTHFR, 5,10-methylenetetrahydrofolate reductase; MTR, methionine synthase; MTRR, methionine synthase reductase; PCFT, proton coupled folate transporter; RFC, reduced folate carrier; SAH, S-adenosylhomocysteine; SAM, S-adenosylmethionine; THF, tetrahydrofolate; TYMS, thymidylate synthase.

of the different forms of folate is very fast and depends on other factors like Cbl and vitamin B₆ status, SAM, and certain polymorphisms in folate-catabolizing enzymes. The 5-methylTHF is the predominant folate form in plasma that constitutes up to 90% of total folate. Folic acid, the synthetic form of the vitamin, enters the folate cycle after a two-step reduction via DHFR into DHF and then THF.

Inherited disorders of folate metabolism and transport

Several inherited disorders of folate metabolism and transport have been described: MTHFR deficiency, MTR deficiency (either caused by mutations in *MTR* gene or *MTRR* gene), cerebral folate deficiency (CFD) caused by FOLR1 mutation, hereditary folate malabsorption, and glutamate formiminotransferase (FTCD) deficiency [44] (Table 1). In addition, several putative inherited disorders related to folate metabolism are discussed in the literature like DHFR deficiency, cellular uptake defects, and the 5,10-methyleneTHF cyclohydrolase [part of the trifunctional enzyme 5,10-methyleneTHF dehydrogenase 1 (MTHFD1, EC 1.5.1.5)] deficiency [66].

Hereditary folate malabsorption

Hereditary folate malabsorption is a clinical syndrome manifested a few months after birth [67]. The pathophysiology of the disorder is attributed to impaired intestinal folate absorption and impaired folate transport into the central nervous system causing very low serum folate concentrations and CFD. After birth, affected newborns rapidly develop severe folate deficiency with megaloblastic anemia, diarrhea, oral mucositis, and recurrent infections (Table 1). Further symptoms include poor feeding, failure to thrive and neurologic manifestations including seizures and developmental delays [68]. If untreated, patients might develop ataxia and cognitive impairment. The disorder has been described in about 30 patients and mostly females are affected [52, 69].

The molecular bases of the hereditary folate malabsorption might be related to mutations in the *SLC46A1* gene, encoding the PCFT, which is present in the intestine and the choroid plexus [37, 39, 52, 70]. Patients deficient in PCFT develop severe folate deficiency suggesting that RFC (one folate carrier), that is expressed in the intestinal epithelium, does not significantly participate in folate absorption [71]. Treatment with large doses (oral or parenteral) of folinic acid (5-formylTHF) can normalize folate levels. CSF folate is far more difficult to normalize and patients require supraphysiological folate concentrations in blood to be able to increase CSF folate [72]. Monitoring the therapy is important and includes measurements of serum and CSF folate with the aim of achieving normal concentrations of folate in CSF.

Mutations in FR α , FR β , or RFC1 may be the cause of disturbed folate uptake. Folic acid treatment normalized serum folate and improved the clinical symptoms in one study, but red blood folate remained very low suggesting an impaired cellular uptake [73]. Patients show low uptake of labeled 5-methylTHF by stimulated lymphocytes and bone marrow cells or red cells [73]. The genetic background of these cases is unknown but might be variants of the hereditary folate malabsorption.

Cerebral folate deficiency

The disease is characterized by severe developmental disorders, mental retardation, epilepsy, and movement disorders [74]. Unlike children with hereditary folate malabsorption, those with CFD present with neurological disorders several years after birth. The disease is characterized by impaired folate transport across the choroid

Disorder	Gene Mendelian inheritance in man (MIM) code [45]	Involved enzyme/protein [EC number, cytogenetic location]	Function	References
5,10-Methylenetetrahydrofolate reductase deficiency	607093, #236250	5,10-Methylenetetrahydrofolate reductase (MTHFR) [EC 1.5.1.20, 1p36.3]	Conversion of 5,10-methyleneTHF to 5-methylTHF	[46–48]
Cerebral folate deficiency	136430, #613068	Folate receptor (FOLR1) [11q13.3-q14.1]	High affinity folate transporter, transport of 5-methylTHF across choroid plexus	[37, 49]
Glutamate formiminotransferase deficiency	606806, #229100	Formiminotransferase cyclodeaminase (FTCD) [EC 4.3.1.4, 21q22.3]	Channels 1-carbon units from formiminoglutamate to the folate pool	[50, 51]
Hereditary folate malabsorption	611672, #229050	Proton coupled folate transporter (PCFT or SLC46A1) [17q11.2]	High affinity folate transporter	[39, 52]
Cobalamin A deficiency	607481, #251100	Methylmalonic aciduria type A protein (MMAA) [4q31.21]	Involved in translocation of Cbl into the mitochondrion during adenosylcobalamin synthesis	[53, 54]
Cobalamin B deficiency	607568, #251110	Cobalamin adenosyltransferase (MMAB) [EC 2.5.1.17, 11q24.11]	Involved in adenosylcobalamin synthesis	[53, 54]
Cobalamin C deficiency	609831, #277400	Methylmalonic aciduria and homocystinuria type C protein (MMACHC) [1p34.1]	Involved in the binding and intracellular trafficking of Cbl	[55–57]
Cobalamin D deficiency	611935, #277410	Chromosome 2 open reading frame 25 (C2orf25) [2q23.2]	Protein involved in an early step of cobalamin metabolism	[58, 59]
Cobalamin E deficiency	602568, #236270	Methionine synthase reductase (MTRR) [EC 1.16.1.8, 5p15.2-p15.3]	Reductive regeneration of the Cbl cofactor	[48, 60]
Cobalamin F deficiency	612625, #277380	LMBR1 domain-containing protein 1 (LMBRD1) [6q13]	Probable lysosomal Cbl transporter	[61, 62]
Cobalamin G deficiency	156570, #250940	Methionine synthase (MTR) [EC 2.1.1.13, 1q43]	Transfer of methyl group from 5-methylTHF to homocysteine to form methionine and THF (Cbl-dependent)	[48, 63]
Methylmalonyl-CoA mutase deficiency	609058, #251000	Methylmalonyl-CoA mutase (MUT) [EC 5.4.99.2, 6p12.3]	Mitochondrial enzyme, catalyzes isomerization of methylmalonyl-CoA to succinyl-CoA	[64, 65]

Table 1 Inherited disorders of folate and cobalamin metabolism and transport.
Phenotype MIM number.

plexus, but not in the intestine. Therefore, when dietary folate intake is sufficient, patients show normal concentrations of serum folate, but very low CSF 5-methylTHF [71, 75]. Mutation in the *FOLR1* gene, which encodes the FR α , can cause severe 5-methylTHF deficiency in the CSF [37, 49]. Oral folinic acid can normalize the CSF folate status and lowers frequency and severity of epileptic seizures [37]. Cario et al. reported heterozygous mutations in both *FOLR1 C352T* and *FOLR1 C525A* genes, leading to nonsense mutations (FOLR1 Q118X and C175X) (Table 2) and the loss of FR-specific folate binding in two siblings [49]. Treatment with folinic acid led to improvement of the symptoms, whereas one of the siblings completely recovered. The homozygous missense mutation FOLR1 T313C was

also identified in one patient with CFD [76]. Additionally, 12% of children with any neurological disorder (71 of 584 patients) exhibited deficient CSF 5-methylTHF concentration [76]. FR α is the most abundant folate transporter in the choroid plexus, supporting its role as major folate transporter across the blood-brain barrier [37]. Normalizing CSF folate in children affected with CFD or those with hereditary folate malabsorption can be better achieved by administration of high doses of folinic acid or methylfolate [71]. Oral treatment is less effective than intramuscular injections. Using folic acid is not advisable, because it must be reduced first and the unmetabolized folic acid can bind to FR α with a high affinity thus preventing the binding of 5-methylTHF.

Gene	Genetic variant	Amino acid substitution	Reference SNP (rs) number
<i>MTHFR</i>	C677T	A222V	1801133
	A1298C	E429A	1801131
<i>MTR</i>	A2756G	D919G	1805087
<i>MTRR</i>	A66G	I22M	1801394
<i>SHMT1</i>	C1420T	L435F	1979277
<i>MTHFD1</i>	G1958A	R653Q	2236225
<i>DHFR</i>	A458T	D153V	121913223
<i>TYMS</i>	2R/3R (28 bp)	–	–
<i>RFC1</i>	G80A	H27R	1051266
<i>FOLR1</i>	C352T	Q118X	121918405
	C525A	C175X	121918406
	130_147 dup	P49PRKSQAP	121918843

Table 2 Common polymorphisms in folate-metabolizing enzymes.

CFD may be related to causes other than mutations in the *FOLR1* gene such as antibodies against the folate receptor [77, 78]. Since the formation of the antibodies can be downregulated with a milk-free diet, it has been suggested that soluble folate-binding proteins in cow’s milk [74] may induce antibody formation [79]. Furthermore, CFD has been found in patients with single nucleotide polymorphisms (SNPs) in the *DHFR* gene in two recent studies [44, 80]. *DHFR* deficiency caused megaloblastic anemia, normal total Hcy (tHcy), and low tetrahydrobiopterine in the CSF [44]. DNA sequencing revealed a homozygous *DHFR* C238T (Leu80Phe) missense mutation [44]. Treatment with folinic acid led to improvement of anemia and CSF 5-methylTHF levels, and seizure control. However, the neurodevelopmental improvement was less than that reported in patients with *FOLR1* mutations or FR autoantibodies [44]. CFD was also reported in patients with megaloblastic anemia who were heterozygous for the *DHFR* A458T polymorphism [80].

Other conditions with neurological manifestations that are associated with CFD are Rett syndrome [81], Kearns-Sayre syndrome [82], dihydropteridine reductase deficiency [83], Aicardi-Goutiere’s syndrome [84], schizophrenia [85], and hypomyelination with atrophy of the basal ganglion syndrome [86].

Glutamate formiminotransferase deficiency

Glutamate formiminotransferase deficiency is an autosomal recessive disorder caused by mutations in the *FTCD* gene [50]. Symptoms include elevated urinary levels of formiminoglutamic acid (FIGLU) after histidine load, megaloblastic anemia, mental retardation, and developmental delay. Heterozygous missense mutations (C457T

and G940C) in the *FTCD* gene were found in the mild form of the disease [51].

Severe 5,10-methylenetetrahydrofolate reductase deficiency

Severe *MTHFR* deficiency is an inborn disease which is associated with homocystinuria, developmental delay, decreased neurotransmitter levels, or seizures [87, 88]. One patient homozygous for the missense mutation in the *MTHFR* c.1129C>T was presented with severe psychomotor retardation, generalized cerebral atrophy, and hypomyelination on magnetic resonance imaging examination [89]. Folate (folic acid, 5-methylTHF, or folinic acid), vitamin B₆, Cbl, and methionine supplementations are the basic therapeutics. In addition, supplementation of large doses of betaine seemed to support the metabolic requirements of the brain by increasing SAM and SAM-dependent methyl transferases [87]. Improvement in the neurological signs has been reported, but it depends on the age of starting the betaine supplementation [87].

Common polymorphisms associated with neural tube defects

Common polymorphisms in the *MTHFR* gene have been described. The C>T substitution in exon 4 at bp 677 causes a substitution of valine for alanine and results in a thermolabile variant of the enzyme [90] that has a 50%–70% less activity when folate intake is limited [42, 90, 91] (Table 2). The TT genotype is found in 10%–20% of the European populations and it increases the risk of neural tube defects (NTDs). Another common polymorphism in the *MTHFR* gene is the A>C substitution at bp 1298 leading also to decreased enzyme activity but without marked effect on tHcy or folate plasma levels [92]. The prevalence of this polymorphism ranges from 6%–11% in Europe [93].

Folates are essential for brain development and function. Folate deficiency during pregnancy can cause NTDs in the offspring. NTDs are common severe congenital malformations that arise early in embryogenesis because of the failure of neural tube closure. Depending on the location of the lesion, NTDs are divided into spina bifida and anencephaly. Folate and Cbl deficiencies as well as elevated tHcy during pregnancy increase the risk of having a child with NTDs [94, 95]. Mutations in genes encoding enzymes involved in the folate/Hcy metabolism have been related to the risk of NTDs [96]. Several studies confirmed

the association between the MTHFR 677TT genotype and the risk of NTD [97, 98], but a number of studies failed to find an association [99, 100]. Shaw et al. found no association between the risk of spina bifida and 118 polymorphisms of folate-related genes [101]. However, the association seems to depend on folate and Cbl intake [46]. The effects of the polymorphisms seem to be counterbalanced by higher folate and Cbl intakes. This is the reason for the insignificant effect of these polymorphisms on NTD risk in countries applying mandatory fortification with folic acid.

In a study on fibroblasts from 41 NTD-affected fetuses, Ou et al. found a 7.2-fold higher risk for NTDs in samples with MTHFR 677TT genotype [102]. Christensen et al. reported an odds ratio (OR) (95% confidence interval) for NTD of 2.2 (0.82–5.99) for affected children, 2.0 (0.75–5.43) for mothers with an NTD child, and 6.0 (1.26–28.53) for mothers and child pairs with MTHFR 677TT [46]. Low red blood cell (RBC) folate further increased the risk of developing NTD, and RBC folate was lower in cases and case mothers compared to controls and control mothers, respectively. The MTHFR 677TT combined with low RBC folate resulted in an OR for NTD of 3.28 (0.84–12.85) for mothers and 13.43 (2.49–72.33) for cases [46].

The association between MTHFR A1298C polymorphism and NTD risk is not consistent. Few studies reported gene-gene interactions with the MTHFR C677T genotype [92, 103], other studies found a protective effect for the MTHFR 1298 C allele [104, 105]. In addition, the MTRR 66GG genotype in the mothers was associated with a 2.1-fold (OR 2.1, 95% CI 1.3–3.3) higher risk for having a child with NTD [106]. The MTR A2756G, the MTRR A66G [107], and the MTHFD1 1958AA polymorphisms [108] were also reported to enhance the maternal risk of having a child with spina bifida [28, 107]. Polymorphisms in RFC1 G80A [109], DHFR [110], and thymidylate synthase (TYMS; EC 2.1.1.45) 28-bp tandem repeat [111], and the presence of autoantibodies against the FR [112] might also be involved in maternal risk of having a child with NTD. Many NTDs can be prevented by improving maternal folate status before the conception. Therefore, it seems that the polymorphisms related to the folate cycle have no independent effect in case of high folate status.

Genetic defects affecting cobalamin-transport or dependent reactions

Cobalamin (Cbl, vitamin B₁₂) is a water soluble vitamin from the B-group. It is essential for cell growth and

division. Cbl deficiency can cause severe hematological and/or neurological manifestations. Low serum Cbl levels are associated with pregnancy loss [113, 114]. Inherited defects in Cbl metabolism are mostly associated with failure to thrive, irritation, feeding problems, and neurological or neurodevelopmental disorders. Hematological and neurological symptoms of folate and Cbl deficiency are similar which is consistent with the crosstalk between the folate and Cbl pathways.

Animal-based diet is the only source of Cbl for humans. The daily requirement for Cbl termed as recommended dietary allowance is 2.4 µg/day for adults [115]. Recent studies accessing blood concentrations of modern markers of cobalamin, suggested that the daily requirements for cobalamin should be set at >6 µg [116]. A relatively large amount of Cbl is stored in the body. The depletion of the vitamin takes several years to develop to deficiency when one stops to consume Cbl-containing diet.

Cbl is a cofactor for only two biochemical reactions in humans (Figure 2). Methylcobalamin (MeCbl) is a cofactor for the cytosolic enzyme MTR that transfers a methyl group from 5-methylTHF to Hcy converting it into methionine [117]. Adenosylcobalamin (AdoCbl) is a cofactor for the mitochondrial enzyme methylmalonyl-CoA mutase (MCM, EC 5.4.99.2) that converts succinyl-CoA into methylmalonyl-CoA. Classical nutritional Cbl deficiency is associated with low holotranscobalamin (holoTC) and elevated plasma concentration of tHcy and methylmalonic acid (MMA). Patients with defects in Cbl metabolism can show severely elevated MMA and/or tHcy without any evidence of low holoTC. This can lead the diagnoses and indicate that elevated MMA and/or tHcy are not related to transcobalamin (TC) deficiency, but to defects in the Cbl-dependent enzymes or trafficking proteins within the cells.

The absorption, transport, and dissimulation of Cbl are complex processes that require several proteins and cellular receptors. Food Cbl is first released from food proteins by means of salivary amylase and acidic conditions and is then bound to haptocorrin released in the saliva. Haptocorrin protects the vitamin from the acid surroundings in the stomach. In the alkaline environment of the intestine haptocorrin is degraded by pancreatic enzymes, and the vitamin liberated from food is recognized by intrinsic factor (IF). IF, another Cbl transporter, is a glycoprotein synthesized in the parietal cells of the stomach that binds only the forms of the vitamin which are active within the body. In the distal ileum, IF-bound-Cbl is taken up by a specific receptor called cubam. Cubam is composed of two proteins, cubilin (CUBN) and amnionless




Figure 2 Cobalamin homeostasis and inherited disorders.

AdoCbl, adenosylcobalamin; Cbl, cobalamin; Hcy, homocysteine; HOCbl, hydroxycobalamin; MCM, methylmalonyl-CoA mutase; MeCbl, methylcobalamin; MMA, methylmalonic acid; MTR, methionine synthase; SAH, S-adenosylhomocysteine; SAM, S-adenosylmethionine; TC, transcobalamin.

(AMN) [118]. Binding of IF-Cbl to cubam is mediated by CUB domains 5–8 in the CUBN protein. Amnionless is a small transmembrane protein that anchors cubilin to the cell membrane in the enterocytes [119]. In the enterocytes, Cbl is released and transferred to transcobalamin, forming the complex holoTC that is secreted into the blood by a yet unknown mechanism probably involving the multidrug resistance protein (MRP1) [120]. As the enterocytes have a high rate of synthesis of transcobalamin II (TC), it has been a common view that Cbl is secreted from the enterocytes in complex with TC. An alternative hypothesis is that free Cbl is transported from the cytosol across the basolateral cell surface into plasma, where it subsequently forms a complex with circulating TC [120]. In the blood, the major part of Cbl is bound to haptocorrin (70%–90%) which is called holohaptocorrin (metabolically inert fraction). Only 10%–30% of Cbl is bound to TC, which carries the metabolically active Cbl. The cellular uptake of Cbl is receptor mediated endocytosis via the TC-receptor or cubam. TC-bound Cbl is degraded in the lysosome and Cbl is released and directed into the two pathways requiring it as a cofactor. The lysosomal degradation and transport of Cbl within the cell are not fully understood.

Several inherited defects in Cbl absorption, transport, or assembly within the cell have been described (Table 1). Defects in any step involved in converting methylmalonyl-CoA to succinyl-CoA cause methylmalonylacidemia. This

can be due to either defective methylmalonyl-CoA mutase (MUT) or impaired synthesis or utilization of AdoCbl (CblA, B, D, and H). There are eight distinct complementation group defects of the intracellular Cbl metabolism (Table 1, Figure 2). The different types were identified by somatic complementation studies applied on fibroblasts isolated from the patients [121]. Moreover, the Cbl genetic defects have been identified on the molecular level [58, 122]. The CblF and CblC defects caused homocystinuria and methylmalonic aciduria [59, 123]. The CblD defect can cause either homocystinuria or methylmalonic aciduria or a combination of homocystinuria and methylmalonic aciduria [59]. The CblA, CblB, and MUT cause only methylmalonic aciduria, and CblE and G cause only homocystinuria. The molecular bases of these disorders have been partly elucidated [124]. The clinical features are similar and the start of the manifestations varies from a few weeks to the adulthood. Lifelong Cbl treatment is required but the prognoses may differ. Clinical symptoms include feeding difficulties, hypotonia, megaloblastic anemia, mental retardation [125], visual loss, or nystagmus. Neuroradiological studies on children with CblC/D defects have shown severe white matter abnormalities like edema, swelling [125], or hydrocephalus [126]. The exact mechanisms behind the neurological manifestations are not known, but may be related to elevated tHcy, lowered SAM causing hypomethylation, or accumulation of MMA that is neurotoxic.

Defects of cobalamin transport and absorption

CblA

The gene responsible for CblA was identified in 2002 by analyses of prokaryotic gene arrangements [53]. This disorder causes isolated methylmalonyluria.

Cobalamin C defect (CblC, MMACHC)

CblC is the most common severe disorder of intracellular Cbl metabolism. CblC is an autosomal recessive disorder of Cbl metabolism related to mutations in *MMACHC* gene [127]. The methylmalonic aciduria and homocystinuria type C protein (MMACHC) is responsible for processing the upper-axial ligands of dietary Cbl before AdoCbl and MeCbl can be synthesized [128, 129]. It causes impaired conversion of Cbl into its two metabolically active forms, MeCbl and AdoCbl. There are several known and recently identified mutations in the *MMACHC* gene [55, 56].

Severe neurological clinical symptoms are manifested in the early onset form during the first year of life. Symptoms include failure to thrive, microcephaly, feeding difficulties, hypotonia, vomiting, developmental delay, seizures, and speech delay. Symptoms are rarely responsive to treatment [130] in the infantile form of the disease. Retinal dysfunction has also been reported in two cases with CblC [131]. Several non-specific hematological symptoms can be seen like megaloblastic anemia and thrombocytopenia.

In 1984 [132] and later reports, a late onset form of CblC which is diagnosed in previously asymptomatic cases at older ages up to adulthood was identified [133, 134]. The first patient identified with CblC had acute onset of dementia, myelopathy, and motor neuron disease [132]. The late onset form has better outcome and response to treatment. A recent study on normal fibroblasts and fibroblasts from early onset CblC disorder identified several proteins that are downregulated in the mutant cells probably explaining some of the neurological manifestations of functional Cbl deficiency [135]. Hydroxycobalamin (HOCbl) did not cause any reduction in the excretion of tHcy from the mutant cells. Interestingly, because of the role of MTR in cellular folate retention, intracellular folate was lower in the CblC cells even after treatment with HOCbl compared to normal fibroblasts [135]. Moreover, protein markers related to brain function were found to be upregulated in fibroblasts from CblC patients and the level of expression was not restored to normal after incubation

with HOCbl [135]. This probably explains the neurocognitive manifestations of the disease or effects on brain development during the prenatal life.

There seems to be a genotype-phenotype association with some mutations: the c.271dupA and c.331C>T (R111X) mutations are more prevalent in the early onset disease. Some missense mutations c.394C>T and c.482G>A are associated with late-onset disease [55, 56]. Genotype-phenotype correlations were explained by variations in the levels of MMACHC mRNA being severely lower in the early-onset forms [55].

In several cases with late onset CblC defect (age at onset 16–41 years) thrombotic events were common. Moreover, six of 11 cases described had encephalopathy, two had seizures, six suffered from myelopathy, and six suffered from psychiatric disturbances [133]. Neurological symptoms, as shown by magnetic resonance imaging, dominated with abnormalities in the white matter area, cortical atrophy, or medullar lesions mostly reported [133]. None of the patients suffered from mental retardation and few remained free of neurological or psychiatric illnesses [133]. Death occurred in few cases despite treatment with Cbl. The mechanisms behind the neurological manifestations might be related to impaired methylation causing cerebral perivascular demyelination [136]. Interestingly, one case had recurrent thrombotic events that were prevented by HOCbl injections. Folinic acid and betaine were used as adjunct therapy to lower tHcy. Oral HOCbl was less effective than the injections in preventing the thrombosis [133].

CblD (MMADHC gene)

The disorder is caused by mutations in the methylmalonic aciduria and homocystinuria type D protein (*MMADHC*) gene that can result in isolated homocystinuria (variant 1), isolated methylmalonic aciduria (variant 2), or combined homocystinuria and methylmalonic aciduria. Patients unable to synthesize AdoCbl and those unable to produce MeCbl have methylmalonic aciduria or homocystinuria, respectively. The third complementation group are patients unable to synthesize both coenzyme forms and have therefore methylmalonic aciduria and homocystinuria [59]. Depending on the mutation, the clinical, cellular, and molecular phenotype of the CblD disorder is heterogeneous. Mutations affecting the N-terminus of *MMADHC* are thought to be associated with methylmalonic aciduria, and mutations affecting the C-terminus are associated with homocystinuria [137].

CblG

This group summarizes disorders in MTR that is related to disorders in the enzyme itself or in the co-factor, MeCbl. CblG is caused by defects in MeCbl synthesis that in turn cause MTR deficiency [138].

CblF (*LMBRD1* gene)

CblF is a disorder in Cbl lysosomal trafficking. This defect has been identified in 1986 as a distinct complementation group [121]. Fibroblasts from CblF patients are able to take Cbl via TC-receptor, but unable to release the vitamin from the lysosome [139]. This defect has been reported in only 13 patients so far [140].

Rutsch et al. recently identified the LMBD1 domain-containing protein (*LMBRD1*) gene on chromosome 6q13, that might be defect in CblF patients and might explain why Cbl is trapped in the lysosome [61]. The gene product is probably lysosomal cobalamin transporter (LMBD1), a lysosomal membrane protein with homology to lipocalin-interacting membrane receptor (LIMR). Five frame shift mutations in *LMBRD1* resulting in loss of LMBD1 function were identified. Fibroblasts of individuals with CblF showed improved synthesis of Cbl cofactor after transfection with wild-type *LMBD1*. Genetics defects on the molecular level have been recently identified in *LMBRD1* gene in one Turkish [140] and three Canadian [141] patients with CblF disorder. The LMBD1 protein is synthesized in the liver and is hypothesized to participate in Cbl uptake and transport in the lysosome [140].

The CblF disorder is manifested at an early age and patients have abnormal newborn screening. Patients are small for gestational age or growth retarded at birth. Beside homocystinuria and methylmalonic aciduria, cerebral seizures, intraventricular hemorrhage [140], failure to thrive, anemia, lethargy, feeding difficulties [61, 142], and developmental delay have been reported [61]. Other disorders have been reported in CblF patients like congenital heart failure, gastritis, ventricular hypertrophy, and hypotonia [61].

CblJ defect (mutations in *ABCD4* gene)

One previously identified peroxisomal ATP-binding cassette (ABC) transporter (*ABCD4*) has been recently shown to be involved in one inherited defect affecting Cbl metabolism. The *ABCD4* protein has been shown to colocalize with the lysosomal proteins LAMP1 and LMBD1 [143]. The

last protein is encoded by *LMBRD1* gene and is deficient in patients with CblF defect.

Imerslund-Gräsbeck syndrome (IGS) or juvenile megaloblastic anemia

This selective Cbl malabsorption is a rare autosomal recessive disorder characterized by Cbl deficiency that is responsive to treatment. Megaloblastic anemia is common and mild proteinuria occurs in approximately 50% of the patients [144, 145].

In 1960, the first description of the disease was by Gräsbeck as “selective cobalamin malabsorption with proteinuria” [146]. In 1963, Imerslund described a similar disorder called idiopathic chronic megaloblastic anemia in children [147]. The disease was first diagnosed in Finland and Norway where the estimated prevalence is 1:200,000. Although the exact prevalence worldwide is not known, many new cases have been reported from eastern Mediterranean countries. In contrast to the case of TC deficiency, symptoms do not appear directly after birth but from the age of a few months to 15 years [148]. Symptoms are rather unspecific and include fatigue, failure to grow or thrive, megaloblastic anemia, and mild neurological symptoms [149].

Investigations of the disorder identified several mutations in two different proteins that constitute the functional IF-receptor: CUBN and AMN [150]. CUBN and AMN form a complex called cubam that represents part of the IF receptor responsible for intestinal Cbl uptake and renal protein reabsorption. Imerslund-Gräsbeck syndrome (IGS) can lead to low expression of IF-receptor, increased degradation [151], or decreased affinity of IF-Cbl to the receptor [152]. A recent study identified genetic mutations in AMN causing a premature stop codon and a strong decrease in the luminal receptor activity [153]. Analyses of renal biopsy from a patient with AMN mutation showed no immunologic reaction for CUBN and an abnormal cytoplasmic, vesicular distribution of the receptor partner AMN suggesting that AMN depends on CUBN for correct localization in the human proximal tubule [154].

Defects in CUBN or AMN and those in IF cause similar symptoms and can be mistaken [155]. Life-long treatment with Cbl in IGS is necessary for preventing the symptoms.

The exact mechanism behind the transport of Cbl into the CNS is not clear. An active transport mechanism into the CNS has been proposed for Cbl. In one case of IGS, CSF Cbl was low and Cbl was required in short intervals for the remission of the psychiatric symptoms [156] suggesting

that AMN protein might be involved in Cbl transport into the brain [156].

Pediatric patients present with megaloblastic anemia, funicular myelosis, or benign proteinuria should be tested for IGS syndrome. The definitive test is based on genetic testing of the mutation, and Cbl should be administered lifelong.

Congenital gastric intrinsic factor deficiency

The discovery of new genes involved in Cbl assimilation and metabolism has improved our knowledge of the rare inborn errors of Cbl metabolism [119]. Juvenile Cbl deficiency leads to hematological and neurological disturbances. There is a lack of Cbl in congenital pernicious anemia due to gastrointestinal intrinsic factor (GIF) deficiency and megaloblastic anemia due to selective intestinal malabsorption of Cbl.

Overgaard et al. [157] recently described a case (15-year-old boy) of a compound heterozygous mutation in the *GIF* gene, with a previously described mutation (c.79+1G>A) and a novel mutation (c.290T>C; M97T) leading to a megaloblastic anemia in an adolescent. Serum Cbl was decreased and serum folate was normal. Cbl therapy together with oral iron supplement normalized blood parameters. Ament et al. [158] identified a specific GIF mutation [c.183_186delGAAT frame shift mutation (M61fs)] to be responsible for juvenile Cbl deficiency in cases of West-African origin. A child from Spain with a megaloblastic anemia due to GIF deficiency has been reported by Garcia Jimenez et al. [159]. The patient is a compound heterozygous in *GIF* gene for a splice site mutation inherited from his mother and a missense change inherited from his father. Leunbach et al. [160] reported a 28-month-old boy with pallor and weight stagnation, macrocytic anemia, and pancytopenia due to Cbl deficiency and a homozygous mutation in the intrinsic factor gene. His sister showed similar symptoms at the age of 15 months. All were given monthly cyanocobalamin injections which, however, caused leg cramps. Replacement with monthly HOCbl was successful.

Congenital transcobalamin deficiency

TC is a non-glycoprotein that has a half-life of 90 min [161]. The vascular endothelium is the major site of the synthesis of TC [162], but many other tissues can also synthesize it. TC deficiency might be caused by the absence of the

protein, error in RNA editing [163], or a protein that is not functional either because it can not bind to the TC-receptor or to Cbl [164, 165]. The molecular bases of several TC defects have been identified [163, 166, 167] after cloning the *TC* gene [168].

Newborns with TC deficiency are asymptomatic at birth. Symptoms are developed during the first few weeks of life and include megaloblastic anemia, failure to thrive, vomiting, infections, and neurological symptoms [163, 169]. Cbl treatment must be started as soon as possible and continue for life. Despite Cbl treatment many patients show continuous neurological symptoms like seizures, cerebral disturbances, and impaired visual abilities [163]. As approximately 80% of serum Cbl is bound to haptocorrin, total serum Cbl might be normal in patients with TC deficiency. This may give the wrong impression that Cbl status is normal and may thus delay the diagnosis of severe Cbl deficiency. Patients have severely increased concentrations of MMA and tHcy in blood and urine.

Haptocorrin (TC I) deficiency produces low serum Cbl levels similar to Cbl deficiency. Diagnosis is especially difficult when TC I deficiency is mild. The phenotype is asymptomatic. A prospective study found severe TC I deficiency with absence of TC I in 0.6% of 537 patients with low Cbl levels [170]. The low Cbl levels of TC I deficiency are usually misattributed to Cbl deficiency. Severe homozygous TC I deficiency features virtually undetectable TC I in plasma and secretions, and serum Cbl is usually <100 pmol/L but heterozygous have mild to moderate lowering of plasma TC I, and mildly lowered (100–150 pmol/L) or low-normal serum Cbl [170, 171].

The *TCN1* gene [Mendelian inheritance in man (MIM) 189905] is located on chromosome 11q11-q12.3 [172], has 9 exons of 59 to 191 bp and 8 introns of 160 bp to 3.2 kb, and encodes TC I, a protein of 433 amino acids [173]. Two mutations have been described, both are located in exon 2 of the 9-exon *TCN1* gene [171]; a 315C > T nonsense mutation and a G deletion at position 270 that causes a frame shift leading to a premature stop codon. These mutations lead to degradation of the transcripts via nonsense-mediated mRNA decay.

Conclusions and final remarks

Disorders in the folate and cobalamin transport or metabolism that affect folate and Cbl absorption, transport, or dependent enzymes, cause severe neurological symptoms that are in some instances reversible after supplementing

the co-enzyme form of the vitamin. Severe elevation in plasma or urine concentrations of tHcy and/or MMA can be considered as a screening test for these disorders. However, it is crucial to identify the defective pathway in order to supplement the proper form of the vitamin as early as possible. The neurological and neurodevelopmental complications in patients with defects in the folate or Cbl related reactions underline the importance of the vitamins in the pre- and post-natal development of the central nervous systems.

Conflict of interest statement

Authors' conflict of interest disclosure: The authors stated that there are no conflicts of interest regarding the publication of this article.

Research funding: None declared.

Employment or leadership: None declared.

Honorarium: None declared.

Received October 4, 2012; accepted October 7, 2012; previously published online November 20, 2012

References

1. Dierkes J, Bisse E, Nauck M, Orth M, Mayer H, Luley C, et al. The diagnostic value of serum homocysteine concentration as a risk factor for coronary artery disease. *Clin Chem Lab Med* 1998;36:453–7.
2. 2nd Conference on Hyperhomocysteinemia. Saarbrücken, Germany, 30–31 March 2001. *Clin Chem Lab Med* 2001;39:665–782.
3. 3rd Conference on Hyperhomocysteinemia. Saarbrücken, Germany, 11–12 April 2003. *Clin Chem Lab Med* 2003;41:1389–538.
4. 4th Conference on Hyperhomocysteinemia. Saarbrücken, Germany, 14–16 April 2005. *Clin Chem Lab Med* 2005;43:977–1170.
5. World Congress on Hyperhomocysteinemia. Saarbrücken, Germany, 05–09 June 2007. *Clin Chem Lab Med* 2007;45:1571–876.
6. Herrmann W. Homocysteine research: alive and kicking! *Clin Chem Lab Med* 2007;45:1571–4.
7. Herrmann W, Obeid R. Homocysteine: a biomarker in neurodegenerative diseases. *Clin Chem Lab Med* 2011;49:435–41.
8. Clarke R, Armitage J, Lewington S, Collins R. Homocysteine-lowering trials for prevention of vascular disease: protocol for a collaborative meta-analysis. *Clin Chem Lab Med* 2007;45:1575–81.
9. Huang T, Chen Y, Yang B, Yang J, Wahlqvist ML, Li D. Meta-analysis of B vitamin supplementation on plasma homocysteine, cardiovascular and all-cause mortality. *Clin Nutr* 2012;31:448–54.
10. Obeid R, McCaddon A, Herrmann W. The role of hyperhomocysteinemia and B-vitamin deficiency in neurological and neuropsychiatric diseases. *Clin Chem Lab Med* 2007;45:1590–606.
11. Quadri P, Fragiaco C, Pezzati R, Zanda E, Tettamanti M, Lucca U. Homocysteine and B vitamins in mild cognitive impairment and dementia. *Clin Chem Lab Med* 2005;43:1096–100.
12. Herrmann W, Obeid R. Biomarkers of neurodegenerative diseases. *Clin Chem Lab Med* 2011;49:343–4.
13. Mattsson N. CSF biomarkers in neurodegenerative diseases. *Clin Chem Lab Med* 2011;49:345–52.
14. van Harten AC, Kester MI, Visser PJ, Blankenstein MA, Pijnenburg YA, van der Flier WM, et al. Tau and p-tau as CSF biomarkers in dementia: a meta-analysis. *Clin Chem Lab Med* 2011;49:353–66.
15. Prvulovic D, Hampel H. Amyloid beta (A β) and phospho-tau (p-tau) as diagnostic biomarkers in Alzheimer's disease. *Clin Chem Lab Med* 2011;49:367–74.
16. Leoni V. The effect of apolipoprotein E (ApoE) genotype on biomarkers of amyloidogenesis, tau pathology and neurodegeneration in Alzheimer's disease. *Clin Chem Lab Med* 2011;49:375–83.
17. Rahmadi A, Steiner N, Munch G. Advanced glycation endproducts as gerontotoxins and biomarkers for carbonyl-based degenerative processes in Alzheimer's disease. *Clin Chem Lab Med* 2011;49:385–91.
18. Breen DP, Michell AW, Barker RA. Parkinson's disease – the continuing search for biomarkers. *Clin Chem Lab Med* 2011;49:393–401.
19. Eller M, Williams DR. Alpha-Synuclein in Parkinson disease and other neurodegenerative disorders. *Clin Chem Lab Med* 2011;49:403–8.
20. Steiner J, Bogerts B, Schroeter ML, Bernstein HG. S100B protein in neurodegenerative disorders. *Clin Chem Lab Med* 2011;49:409–24.
21. Kwon BK, Casha S, Hurlbert RJ, Yong VW. Inflammatory and structural biomarkers in acute traumatic spinal cord injury. *Clin Chem Lab Med* 2011;49:425–33.
22. Smith AD, Smith SM, de Jager CA, Whitbread P, Johnston C, Agacinski G, et al. Homocysteine-lowering by B vitamins slows the rate of accelerated brain atrophy in mild cognitive impairment: a randomized controlled trial. *PLoS ONE* 2010;5:e12244.
23. de Jager CA, Oulhaj A, Jacoby R, Refsum H, Smith AD. Cognitive and clinical outcomes of homocysteine-lowering B-vitamin treatment in mild cognitive impairment: a randomized controlled trial. *Int J Geriatr Psychiatry* 2012;27:592–600.
24. Obeid R, Kostopoulos P, Knapp JP, Kasoha M, Becker G, Fassbender K, et al. Biomarkers of folate and vitamin B12 are related in blood and cerebrospinal fluid. *Clin Chem* 2007;53:326–33.
25. Obeid R, Schlundt J, Umanskaya N, Herrmann W, Herrmann M. Folate is related to phosphorylated neurofilament-H and P-tau (Ser396) in rat brain. *J Neurochem* 2011;117:1047–54.
26. Geisel J, Zimbelmann I, Schorr H, Knapp JP, Bodis M, Hübner U, et al. Genetic defects as important factors for moderate hyperhomocysteinemia. *Clin Chem Lab Med* 2001;39:698–704.

27. Geisel J, Hubner U, Bodis M, Schorr H, Knapp JP, Obeid R, et al. The role of genetic factors in the development of hyperhomocysteinemia. *Clin Chem Lab Med* 2003;41:1427–34.
28. Gueant JL, Gueant-Rodriguez RM, Anello G, Bosco P, Brunaud L, Romano C, et al. Genetic determinants of folate and vitamin B12 metabolism: a common pathway in neural tube defect and Down syndrome? *Clin Chem Lab Med* 2003;41:1473–7.
29. Geisel J, Schorr H, Bodis M, Isber S, Hubner U, Knapp JP, et al. The vegetarian lifestyle and DNA methylation. *Clin Chem Lab Med* 2005;43:1164–9.
30. Geisel J, Schorr H, Heine GH, Bodis M, Hubner U, Knapp JP, et al. Decreased p66Shc promoter methylation in patients with end-stage renal disease. *Clin Chem Lab Med* 2007;45:1764–70.
31. Obeid R, Herrmann W. Homocysteine, folic acid and vitamin B12 in relation to pre- and postnatal health aspects. *Clin Chem Lab Med* 2005;43:1052–7.
32. Nelen WL. Hyperhomocysteinemia and human reproduction. *Clin Chem Lab Med* 2001;39:758–63.
33. Wang TT, Chandler CJ, Halsted CH. Intracellular pteroylpolyglutamate hydrolase from human jejunal mucosa. Isolation and characterization. *J Biol Chem* 1986;261:13551–5.
34. Hoffbrand AV, Tripp E, Houlihan CM, Scott JM. Studies on the uptake of synthetic conjugated folates by human marrow cells. *Blood* 1973;42:141–6.
35. Antony AC. Folate receptors. *Annu Rev Nutr* 1996;16:501–21.
36. Solanky N, Requena JA, D'Souza SW, Sibley CP, Glazier JD. Expression of folate transporters in human placenta and implications for homocysteine metabolism. *Placenta* 2010;31:134–43.
37. Steinfeld R, Grapp M, Kraetzner R, Dreha-Kulaczewski S, Helms G, Dechent P, et al. Folate receptor alpha defect causes cerebral folate transport deficiency: a treatable neurodegenerative disorder associated with disturbed myelin metabolism. *Am J Hum Genet* 2009;85:354–63.
38. Henderson GB, Suresh MR, Vitols KS, Huennekens FM. Transport of folate compounds in L1210 cells: kinetic evidence that folate influx proceeds via the high-affinity transport system for 5-methyltetrahydrofolate and methotrexate. *Cancer Res* 1986;46:1639–43.
39. Qiu A, Jansen M, Sakaris A, Min SH, Chattopadhyay S, Tsai E, et al. Identification of an intestinal folate transporter and the molecular basis for hereditary folate malabsorption. *Cell* 2006;127:917–28.
40. Zhao R, Min SH, Wang Y, Campanella E, Low PS, Goldman ID. A role for the proton-coupled folate transporter (PCFT-SLC46A1) in folate receptor-mediated endocytosis. *J Biol Chem* 2009;284:4267–74.
41. Smith DE, Smulders YM, Blom HJ, Popp J, Jessen F, Semmler A, et al. Determinants of the essential one-carbon metabolism metabolites, homocysteine, S-adenosylmethionine, S-adenosylhomocysteine and folate, in cerebrospinal fluid. *Clin Chem Lab Med* 2012;50:1641–7.
42. Stanger O. Physiology of folic acid in health and disease. *Curr Drug Metab* 2002;3:211–23.
43. Cook RJ. Folate metabolism. In: Carmel R, Jacobsen DW, editors. *Homocysteine in health and disease*, 1st ed. Cambridge, UK: Cambridge University Press, 2001:113–34.
44. Banka S, Blom HJ, Walter J, Aziz M, Urquhart J, Clouthier CM, et al. Identification and characterization of an inborn error of metabolism caused by dihydrofolate reductase deficiency. *Am J Hum Genet* 2011;88:216–25.
45. Online Mendelian inheritance in man. OMIM®. McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University :Baltimore, MD, 2011. Available from: <http://omim.org/> 2011. Accessed on 1 March 2012.
46. Christensen B, Arbour L, Tran P, Leclerc D, Sabbaghian N, Platt R, et al. Genetic polymorphisms in methylenetetrahydrofolate reductase and methionine synthase, folate levels in red blood cells, and risk of neural tube defects. *Am J Med Genet* 1999;84:151–7.
47. Watkins D, Ru M, Hwang HY, Kim CD, Murray A, Philip NS, et al. Hyperhomocysteinemia due to methionine synthase deficiency, cblG: structure of the MTR gene, genotype diversity, and recognition of a common mutation, P1173L. *Am J Hum Genet* 2002;71:143–53.
48. Bosco P, Gueant-Rodriguez RM, Anello G, Barone C, Namour F, Caraci F, et al. Methionine synthase (MTR) 2756 (A→G) polymorphism, double heterozygosity methionine synthase 2756 AG/methionine synthase reductase (MTRR) 66 AG, and elevated homocysteinemia are three risk factors for having a child with Down syndrome. *Am J Med Genet A* 2003;121A:219–24.
49. Cario H, Bode H, Debatin KM, Opladen T, Schwarz K. Congenital null mutations of the FOLR1 gene: a progressive neurologic disease and its treatment. *Neurology* 2009;73:2127–9.
50. Solans A, Estivill X, de la Luna S. Cloning and characterization of human FTCD on 21q22.3, a candidate gene for glutamate formiminotransferase deficiency. *Cytogenet Cell Genet* 2000;88:43–9.
51. Hilton JF, Christensen KE, Watkins D, Raby BA, Renaud Y, de la Luna S, et al. The molecular basis of glutamate formiminotransferase deficiency. *Hum Mutat* 2003;22:67–73.
52. Shin DS, Mahadeo K, Min SH, Diop-Bove N, Clayton P, Zhao R, et al. Identification of novel mutations in the proton-coupled folate transporter (PCFT-SLC46A1) associated with hereditary folate malabsorption. *Mol Genet Metab* 2011;103:33–7.
53. Dobson CM, Wai T, Leclerc D, Wilson A, Wu X, Dore C, et al. Identification of the gene responsible for the cblA complementation group of vitamin B12-responsive methylmalonic acidemia based on analysis of prokaryotic gene arrangements. *Proc Natl Acad Sci USA* 2002;99:15554–9.
54. Merinero B, Perez B, Perez-Cerda C, Rincon A, Desviat LR, Martinez MA, et al. Methylmalonic acidemia: examination of genotype and biochemical data in 32 patients belonging to mut, cblA or cblB complementation group. *J Inher Metab Dis* 2008;31:55–66.
55. Lerner-Ellis JP, Anastasio N, Liu J, Coelho D, Suormala T, Stucki M, et al. Spectrum of mutations in MMACHC, allelic expression, and evidence for genotype-phenotype correlations. *Hum Mutat* 2009;30:1072–81.
56. Lerner-Ellis JP, Tirone JC, Pawelek PD, Dore C, Atkinson JL, Watkins D, et al. Identification of the gene responsible for methylmalonic aciduria and homocystinuria, cblC type. *Nat Genet* 2006;38:93–100.
57. Liu MY, Yang YL, Chang YC, Chiang SH, Lin SP, Han LS, et al. Mutation spectrum of MMACHC in Chinese patients with combined methylmalonic aciduria and homocystinuria. *J Hum Genet* 2010;55:621–6.
58. Coelho D, Suormala T, Stucki M, Lerner-Ellis JP, Rosenblatt DS, Newbold RF, et al. Gene identification for the cblD defect of vitamin B12 metabolism. *N Engl J Med* 2008;358:1454–64.
59. Suormala T, Baumgartner MR, Coelho D, Zavadakova P, Kozich V, Koch HG, et al. The cblD defect causes either isolated or

- combined deficiency of methylcobalamin and adenosylcobalamin synthesis. *J Biol Chem* 2004;279:42742–9.
60. Wilson A, Leclerc D, Rosenblatt DS, Gravel RA. Molecular basis for methionine synthase reductase deficiency in patients belonging to the cblE complementation group of disorders in folate/cobalamin metabolism. *Hum Mol Genet* 1999;8:2009–16.
 61. Rutsch F, Gailus S, Miousse IR, Suormala T, Sagne C, Toliat MR, et al. Identification of a putative lysosomal cobalamin exporter altered in the cblF defect of vitamin B12 metabolism. *Nat Genet* 2009;41:234–9.
 62. Alfadhel M, Lillquist YP, Davis C, Junker AK, Stockler-Ipsiroglu S. Eighteen-year follow-up of a patient with cobalamin F disease (cblF): report and review. *Am J Med Genet A* 2011;155A:2571–7.
 63. Wilson A, Leclerc D, Saberi F, Campeau E, Hwang HY, Shane B, et al. Functionally null mutations in patients with the cblG-variant form of methionine synthase deficiency. *Am J Hum Genet* 1998;63:409–14.
 64. Froese DS, Gravel RA. Genetic disorders of vitamin B metabolism: eight complementation groups – eight genes. *Expert Rev Mol Med* 2010;12:e37.
 65. Worgan LC, Niles K, Tirone JC, Hofmann A, Verner A, Sammak A, et al. Spectrum of mutations in mut methylmalonic acidemia and identification of a common Hispanic mutation and haplotype. *Hum Mutat* 2006;27:31–43.
 66. Whitehead VM. Acquired and inherited disorders of cobalamin and folate in children. *Br J Haematol* 2006;134:125–36.
 67. Diop-Bove N, Kronn D, Goldman ID. Hereditary folate malabsorption. In: Pagon RA, et al. editors, *GeneReviews* [Internet]. 1993–2008 Jun 17 [updated 8 Dec 2011]. Seattle (WA): University of Washington, Seattle, 1993. Accessed 1 August 2012.
 68. Geller J, Kronn D, Jayabose S, Sandoval C. Hereditary folate malabsorption: family report and review of the literature. *Medicine (Baltimore)* 2002;81:51–68.
 69. Steinschneider M, Sherbany A, Pavlakis S, Emerson R, Lovelace R, De Vivo DC. Congenital folate malabsorption: reversible clinical and neurophysiologic abnormalities. *Neurology* 1990;40:1315.
 70. Shin DS, Zhao R, Yap EH, Fiser A, Goldman ID. A P425R mutation of the proton-coupled folate transporter causing hereditary folate malabsorption produces a highly selective alteration in folate binding. *Am J Physiol Cell Physiol* 2012;302:C1405–12.
 71. Zhao R, Diop-Bove N, Visentin M, Goldman ID. Mechanisms of membrane transport of folates into cells and across epithelia. *Annu Rev Nutr* 2011;31:177–201.
 72. Rosenblatt DS, Fenton WA. Inherited disorders of folate and cobalamin transport and metabolism. In: Scriver CR, et al., editors. *The metabolic and molecular bases of inherited disease*, 8th ed. New York: McGraw-Hill Medical Publishing Division, 2001:3897–933.
 73. Branda RF, Moldow CF, MacArthur JR, Wintrobe MM, Anthony BK, Jacob HS. Folate-induced remission in aplastic anemia with familial defect of cellular folate uptake. *N Engl J Med* 1978;298:469–75.
 74. Hyland K, Shoffner J, Heales SJ. Cerebral folate deficiency. *J Inherit Metab Dis* 2010;33:563–70.
 75. Ramaekers VT, Blau N. Cerebral folate deficiency. *Dev Med Child Neurol* 2004;46:843–51.
 76. Perez-Duenas B, Ormazabal A, Toma C, Torrico B, Cormand B, Serrano M, et al. Cerebral folate deficiency syndromes in childhood: clinical, analytical, and etiologic aspects. *Arch Neurol* 2011;68:615–21.
 77. Ramaekers VT, Rothenberg SP, Sequeira JM, Opladen T, Blau N, Quadros EV, et al. Autoantibodies to folate receptors in the cerebral folate deficiency syndrome. *N Engl J Med* 2005;352:1985–91.
 78. Frye RE, Sequeira JM, Quadros EV, James SJ, Rossignol DA. Cerebral folate receptor autoantibodies in autism spectrum disorder. *Mol Psychiatry* 2012. doi:10.1038/mp.2011.175.
 79. Ramaekers VT, Sequeira JM, Blau N, Quadros EV. A milk-free diet downregulates folate receptor autoimmunity in cerebral folate deficiency syndrome. *Dev Med Child Neurol* 2008;50:346–52.
 80. Cario H, Smith DE, Blom H, Blau N, Bode H, Holzmann K, et al. Dihydrofolate reductase deficiency due to a homozygous DHFR mutation causes megaloblastic anemia and cerebral folate deficiency leading to severe neurologic disease. *Am J Hum Genet* 2011;88:226–31.
 81. Ramaekers VT, Hansen SI, Holm J, Opladen T, Senderek J, Hausler M, et al. Reduced folate transport to the CNS in female Rett patients. *Neurology* 2003;61:506–15.
 82. Pineda M, Ormazabal A, Lopez-Gallardo E, Nascimento A, Solano A, Herrero MD, et al. Cerebral folate deficiency and leukoencephalopathy caused by a mitochondrial DNA deletion. *Ann Neurol* 2006;59:394–8.
 83. Kaufman S. Some metabolic relationships between bipterin and folate: implications for the “methyl trap hypothesis”. *Neurochem Res* 1991;16:1031–6.
 84. Blau N, Bonafe L, Krageloh-Mann I, Thony B, Kierat L, Hausler M, et al. Cerebrospinal fluid pterins and folates in Aicardi-Goutieres syndrome: a new phenotype. *Neurology* 2003;61:642–7.
 85. Ho A, Michelson D, Aaen G, Ashwal S. Cerebral folate deficiency presenting as adolescent catatonic schizophrenia: a case report. *J Child Neurol* 2010;25:898–900.
 86. Mercimek-Mahmutoglu S, van der Knaap MS, Baric I, Prayer D, Stoekler-Ipsiroglu S. Hypomyelination with atrophy of the basal ganglia and cerebellum (H-ABC). Report of a new case. *Neuropediatrics* 2005;36:223–6.
 87. Strauss KA, Morton DH, Puffenberger EG, Hendrickson C, Robinson DL, Wagner C, et al. Prevention of brain disease from severe 5,10-methylenetetrahydrofolate reductase deficiency. *Mol Genet Metab* 2007;91:165–75.
 88. Tsuji M, Takagi A, Sameshima K, Iai M, Yamashita S, Shinbo H, et al. 5,10-Methylenetetrahydrofolate reductase deficiency with progressive polyneuropathy in an infant. *Brain Dev* 2011;33:521–4.
 89. Goyette P, Christensen B, Rosenblatt DS, Rozen R. Severe and mild mutations in cis for the methylenetetrahydrofolate reductase (MTHFR) gene, and description of five novel mutations in MTHFR. *Am J Hum Genet* 1996;59:1268–75.
 90. Frosst P, Blom HJ, Milos R, Goyette P, Sheppard CA, Matthews RG, et al. A candidate genetic risk factor for vascular disease: a common mutation in methylenetetrahydrofolate reductase. *Nat Genet* 1995;10:111–3.
 91. Rosenblatt DS, Lue-Shing H, Arzoumanian A, Low-Nang L, Matiaszuk N. Methylenetetrahydrofolate reductase (MR) deficiency: thermolability of residual MR activity, methionine synthase activity, and methylcobalamin levels in cultured fibroblasts. *Biochem Med Metab Biol* 1992;47:221–5.
 92. Van Der Put NM, Gabreels F, Stevens EM, Smeitink JA, Trijbels FJ, Eskes TK, et al. A second common mutation in the methylenetetrahydrofolate reductase gene: an additional risk factor for neural-tube defects? *Am J Hum Genet* 1998;62:1044–51.


93. Christensen KE, Rozen R. Genetic variation: effect on folate metabolism and health. In: Bailey LB, editor. *Folate in health and disease*, 2nd ed. Boca Raton, London, New York: CRC Press, 2010:75–110.
94. MRC Vitamin Study Research Group. Prevention of neural tube defects: results of the Medical Research Council Vitamin Study. *Lancet* 1991;338:131–7.
95. Czeizel AE, Dudas I. Prevention of the first occurrence of neural-tube defects by periconceptional vitamin supplementation. *N Engl J Med* 1992;327:1832–5.
96. Van Der Put NM, Steegers-Theunissen RP, Frosst P, Trijbels FJ, Eskes TK, van den Heuvel LP, et al. Mutated methylenetetrahydrofolate reductase as a risk factor for spina bifida. *Lancet* 1995;346:1070–1.
97. Relton CL, Wilding CS, Pearce MS, Laffling AJ, Jonas PA, Lynch SA, et al. Gene-gene interaction in folate-related genes and risk of neural tube defects in a UK population. *J Med Genet* 2004;41:256–60.
98. Botto LD, Yang Q. 5,10-Methylenetetrahydrofolate reductase gene variants and congenital anomalies: a HuGE review. *Am J Epidemiol* 2000;151:862–77.
99. Speer MC, Worley G, Mackey JF, Melvin E, Oakes WJ, George TM. The thermolabile variant of methylenetetrahydrofolate reductase (MTHFR) is not a major risk factor for neural tube defect in American Caucasians. The NTD Collaborative Group. *Neurogenetics* 1997;1:149–50.
100. Shang Y, Zhao H, Niu B, Li W, Zhou R, Zhang T, et al. Correlation of polymorphism of MTHFRs and RFC-1 genes with neural tube defects in China. *Birth Defects Res A Clin Mol Teratol* 2008;82:3–7.
101. Shaw GM, Lu W, Zhu H, Yang W, Briggs FB, Carmichael SL, et al. 118 SNPs of folate-related genes and risks of spina bifida and conotruncal heart defects. *BMC Med Genet* 2009;10:49.
102. Ou CY, Stevenson RE, Brown VK, Schwartz CE, Allen WP, Khoury MJ, et al. 5,10 Methylenetetrahydrofolate reductase genetic polymorphism as a risk factor for neural tube defects. *Am J Med Genet* 1996;63:610–4.
103. Boduroglu K, Alanay Y, Alikasifoglu M, Aktas D, Tuncbilek E. Analysis of MTHFR 1298A>C in addition to MTHFR 677C>T polymorphism as a risk factor for neural tube defects in the Turkish population. *Turk J Pediatr* 2005;47:327–33.
104. Relton CL, Wilding CS, Laffling AJ, Jonas PA, Burgess T, Binks K, et al. Low erythrocyte folate status and polymorphic variation in folate-related genes are associated with risk of neural tube defect pregnancy. *Mol Genet Metab* 2004;81:273–81.
105. Candito M, Rivet R, Herbeth B, Boisson C, Rudigoz RC, Luton D, et al. Nutritional and genetic determinants of vitamin B and homocysteine metabolisms in neural tube defects: a multicenter case-control study. *Am J Med Genet A* 2008;146A:1128–33.
106. van der Linden I, den Heijer M, Afman LA, Gellekink H, Vermeulen SH, Kluijtmans LA, et al. The methionine synthase reductase 66A>G polymorphism is a maternal risk factor for spina bifida. *J Mol Med (Berl)* 2006;84:1047–54.
107. Doolin MT, Barbaux S, McDonnell M, Hoess K, Whitehead AS, Mitchell LE. Maternal genetic effects, exerted by genes involved in homocysteine remethylation, influence the risk of spina bifida. *Am J Hum Genet* 2002;71:1222–6.
108. Brody LC, Conley M, Cox C, Kirke PN, McKeever MP, Mills JL, et al. A polymorphism, R653Q, in the trifunctional enzyme methylenetetrahydrofolate dehydrogenase/methylenetetrahydrofolate cyclohydrolase/formyltetrahydrofolate synthetase is a maternal genetic risk factor for neural tube defects: report of the Birth Defects Research Group. *Am J Hum Genet* 2002;71:1207–15.
109. De Marco P, Calevo MG, Moroni A, Merello E, Raso A, Finnell RH, et al. Reduced folate carrier polymorphism (80A>G) and neural tube defects. *Eur J Hum Genet* 2003;11:245–52.
110. Johnson WG, Stenroos ES, Szychala JR, Chatkupt S, Ming SX, Buyske S. New 19 bp deletion polymorphism in intron-1 of dihydrofolate reductase (DHFR): a risk factor for spina bifida acting in mothers during pregnancy? *Am J Med Genet A* 2004;124A:339–45.
111. Volcik KA, Shaw GM, Zhu H, Lammer EJ, Laurent C, Finnell RH. Associations between polymorphisms within the thymidylate synthase gene and spina bifida. *Birth Defects Res A Clin Mol Teratol* 2003;67:924–8.
112. Taparia S, Gelineau-van WJ, Rosenquist TH, Finnell RH. Importance of folate-homocysteine homeostasis during early embryonic development. *Clin Chem Lab Med* 2007;45:1717–27.
113. Candito M, Magnaldo S, Bayle J, Dor JF, Gillet Y, Bongain A, et al. Clinical B12 deficiency in one case of recurrent spontaneous pregnancy loss. *Clin Chem Lab Med* 2003;41:1026–7.
114. Hubner U, Alwan A, Jouma M, Tabbaa M, Schorr H, Herrmann W. Low serum vitamin B12 is associated with recurrent pregnancy loss in Syrian women. *Clin Chem Lab Med* 2008;46:1265–9.
115. Institute of Medicine. Vitamin B12. Dietary reference intakes for thiamin, riboflavin, niacin, vitamin B6, folate, vitamin B12, pantothenic acid, biotin, and choline. USA, Washington, DC: National Academy Press, 2000:306–56.
116. Bor MV, Lydeking-Olsen E, Moller J, Nexø E. A daily intake of approximately 6 microg vitamin B-12 appears to saturate all the vitamin B-12-related variables in Danish postmenopausal women. *Am J Clin Nutr* 2006;83:52–8.
117. Banerjee RV, Matthews RG. Cobalamin-dependent methionine synthase. *FASEB J* 1990;4:1450–9.
118. Fyfe JC, Madsen M, Hojrup P, Christensen EI, Tanner SM, de la Chapelle A, et al. The functional cobalamin (vitamin B12)-intrinsic factor receptor is a novel complex of cubilin and amnionless. *Blood* 2004;103:1573–9.
119. Quadros EV. Advances in the understanding of cobalamin assimilation and metabolism. *Br J Haematol* 2010;148:195–204.
120. Beedholm-Ebsen R, van de Wetering K, Hardlei T, Nexø E, Borst P, Moestrup SK. Identification of multidrug resistance protein 1 (MRP1/ABCC1) as a molecular gate for cellular export of cobalamin. *Blood* 2010;115:1632–9.
121. Watkins D, Rosenblatt DS. Failure of lysosomal release of vitamin B12: a new complementation group causing methylmalonic aciduria (cblF). *Am J Hum Genet* 1986;39:404–8.
122. Leclerc D, Campeau E, Goyette P, Adjalla CE, Christensen B, Ross M, et al. Human methionine synthase: cDNA cloning and identification of mutations in patients of the cblG complementation group of folate/cobalamin disorders. *Hum Mol Genet* 1996;5:1867–74.
123. Nogueira C, Aiello C, Cerone R, Martins E, Caruso U, Moroni I, et al. Spectrum of MMACHC mutations in Italian and Portuguese patients with combined methylmalonic aciduria and homocystinuria, cblC type. *Mol Genet Metab* 2008;93:475–80.

124. Moras E, Hosack A, Watkins D, Rosenblatt DS. Mitochondrial vitamin B12-binding proteins in patients with inborn errors of cobalamin metabolism. *Mol Genet Metab* 2007;90:140–7.
125. Longo D, Fariello G, Dionisi-Vici C, Cannata V, Boenzi S, Genovese E, et al. MRI and 1H-MRS findings in early-onset cobalamin C/D defect. *Neuropediatrics* 2005;36:366–72.
126. Rossi A, Cerone R, Biancheri R, Gatti R, Schiaffino MC, Fonda C, et al. Early-onset combined methylmalonic aciduria and homocystinuria: neuroradiologic findings. *Am J Neuroradiol* 2001;22:554–63.
127. Martinelli D, Deodato F, Dionisi-Vici C. Cobalamin C defect: natural history, pathophysiology, and treatment. *J Inher Metab Dis* 2011;34:127–35.
128. Kim J, Gherasim C, Banerjee R. Decyanation of vitamin B12 by a trafficking chaperone. *Proc Natl Acad Sci USA* 2008;105:14551–4.
129. Hannibal L, Kim J, Brasch NE, Wang S, Rosenblatt DS, Banerjee R, et al. Processing of alkylcobalamins in mammalian cells: a role for the MMACHC (cblC) gene product. *Mol Genet Metab* 2009;97:260–6.
130. Rosenblatt DS, Aspler AL, Shevell MI, Pletcher BA, Fenton WA, Seashore MR. Clinical heterogeneity and prognosis in combined methylmalonic aciduria and homocystinuria (cblC). *J Inher Metab Dis* 1997;20:528–38.
131. Gaillard MC, Matthieu JM, Borruat FX. Retinal dysfunction in combined methylmalonic aciduria and homocystinuria (Cblc) disease: a spectrum of disorders. *Klin Monbl Augenheilkd* 2008;225:491–4.
132. Shinnar S, Singer HS. Cobalamin C mutation (methylmalonic aciduria and homocystinuria) in adolescence. A treatable cause of dementia and myelopathy. *N Engl J Med* 1984;311:451–4.
133. Thauvin-Robinet C, Roze E, Couvreur G, Horellou MH, Sedel F, Grabli D, et al. The adolescent and adult form of cobalamin C disease: clinical and molecular spectrum. *J Neurol Neurosurg Psychiatry* 2008;79:725–8.
134. Ben-Omran TI, Wong H, Blaser S, Feigenbaum A. Late-onset cobalamin-C disorder: a challenging diagnosis. *Am J Med Genet A* 2007;143A:979–84.
135. Hannibal L, DiBello PM, Yu M, Miller A, Wang S, Willard B, et al. The MMACHC proteome: hallmarks of functional cobalamin deficiency in humans. *Mol Genet Metab* 2011;103:226–39.
136. Powers JM, Rosenblatt DS, Schmidt RE, Cross AH, Black JT, Moser AB, et al. Neurological and neuropathologic heterogeneity in two brothers with cobalamin C deficiency. *Ann Neurol* 2001;49:396–400.
137. Miousse IR, Watkins D, Coelho D, Rupa T, Crombez EA, Vilain E, et al. Clinical and molecular heterogeneity in patients with the cblD inborn error of cobalamin metabolism. *J Pediatr* 2009;154:551–6.
138. Gulati S, Baker P, Li YN, Fowler B, Kruger W, Brody LC, et al. Defects in human methionine synthase in cblG patients. *Hum Mol Genet* 1996;5:1859–65.
139. Vassiliadis A, Rosenblatt DS, Cooper BA, Bergeron JJ. Lysosomal cobalamin accumulation in fibroblasts from a patient with an inborn error of cobalamin metabolism (cblF complementation group): visualization by electron microscope radioautography. *Exp Cell Res* 1991;195:295–302.
140. Gailus S, Hohne W, Gasnier B, Nurnberg P, Fowler B, Rutsch F. Insights into lysosomal cobalamin trafficking: lessons learned from cblF disease. *J Mol Med (Berl)* 2010;88:459–66.
141. Miousse IR, Watkins D, Rosenblatt DS. Novel splice site mutations and a large deletion in three patients with the cblF inborn error of vitamin B12 metabolism. *Mol Genet Metab* 2011;102:505–7.
142. Rosenblatt DS, Laframboise R, Pichette J, Langevin P, Cooper BA, Costa T. New disorder of vitamin B12 metabolism (cobalamin F) presenting as methylmalonic aciduria. *Pediatrics* 1986;78:51–4.
143. Coelho D, Kim JC, Miousse IR, Fung S, du MM, Buers I, et al. Mutations in ABCD4 cause a new inborn error of vitamin B(12) metabolism. *Nat Genet* 2012;44:1152–5.
144. Wahlstedt-Froberg V, Pettersson T, Aminoff M, Dugue B, Grasbeck R. Proteinuria in cubilin-deficient patients with selective vitamin B12 malabsorption. *Pediatr Nephrol* 2003;18:417–21.
145. Broch H, Imerslund O, Monn E, Hovig T, Seip M. Imerslund-Grasbeck anemia. A long-term follow-up study. *Acta Paediatr Scand* 1984;73:248–53.
146. Gräsbeck R, Gordin R, Kantero I, Kuhlback B. Selective vitamin B12 malabsorption and proteinuria in young people. A syndrome. *Acta Med Scand* 1960;167:289–96.
147. Imerslund O, Bjornstad P. Familial vitamin B12 malabsorption. *Acta Haematol* 1963;30:1–7.
148. Hauck FH, Tanner SM, Henker J, Laass MW. Imerslund-Grasbeck syndrome in a 15-year-old German girl caused by compound heterozygous mutations in CUBN. *Eur J Pediatr* 2008;167:671–5.
149. Gräsbeck R. Imerslund-Grasbeck syndrome [selective vitamin B(12) malabsorption with proteinuria]. *Orphanet J Rare Dis* 2006;1:17.
150. Tanner SM, Li Z, Bisson R, Acar C, Oner C, Oner R, et al. Genetically heterogeneous selective intestinal malabsorption of vitamin B12: founder effects, consanguinity, and high clinical awareness explain aggregations in Scandinavia and the Middle East. *Hum Mutat* 2004;23:327–33.
151. Gueant JL, Saunier M, Gastin I, Safi A, Lamireau T, Duclos B, et al. Decreased activity of intestinal and urinary intrinsic factor receptor in Grasbeck-Imerslund disease [corrected]. *Gastroenterology* 1995;108:1622–8.
152. Gueant JL, Chery C, Namour F, Aimone-Gastin I, Wustinger M. Decreased affinity of urinary intrinsic factor-cobalamin receptor in a case of Grasbeck-Imerslund syndrome. *Gastroenterology* 1999;116:1274–6.
153. Namour F, Dobrowoljski G, Chery C, Audonnet S, Feillet F, Sperl W, et al. Luminal expression of cubilin is impaired in Imerslund-Grasbeck syndrome with compound AMN mutations in intron 3 and exon 7. *Haematologica* 2011;96:1715–9.
154. Storm T, Emma F, Verroust PJ, Hertz JM, Nielsen R, Christensen EI. A patient with cubilin deficiency. *N Engl J Med* 2011;364:89–91.
155. Tanner SM, Li Z, Perko JD, Oner C, Cetin M, Altay C, et al. Hereditary juvenile cobalamin deficiency caused by mutations in the intrinsic factor gene. *Proc Natl Acad Sci USA* 2005;102:4130–3.
156. Luder AS, Tanner SM, de la Chapelle A, Walter JH. Amnionless (AMN) mutations in Imerslund-Grasbeck syndrome may be associated with disturbed vitamin B(12) transport into the CNS. *J Inher Metab Dis* 2008. doi:10.1007/s10545-007-0760-2.
157. Overgaard UM, Tanner SM, Birgens HS. Vitamin B12 deficiency in a 15-year old boy due to mutations in the intrinsic factor gene, GIF. *Br J Haematol* 2010;150:369–71.

158. Ament AE, Li Z, Sturm AC, Perko JD, Lawson S, Masterson M, et al. Juvenile cobalamin deficiency in individuals of African ancestry is caused by a founder mutation in the intrinsic factor gene GIF. *Br J Haematol* 2009;144:622–4.
159. Garcia Jimenez MC, Baldellou VA, Calvo Martin MT, Perez-Lungmus G, Lopez PJ. Hereditary juvenile cobalamin deficiency due to mutations in GIF gene. *An Pediatr (Barc)* 2008;69:56–8.
160. Leunbach TL, Johansen P, Tanner SM, Grasbeck R, Helgestad J. Homozygous mutation in the intrinsic factor gene in a child with severe vitamin B12 deficiency. *Ugeskr Laeger* 2011;173:2047–8.
161. Donaldson RM, Jr., Brand M, Serfilippi D. Changes in circulating transcobalamin II after injection of cyanocobalamin. *N Engl J Med* 1977;296:1427–30.
162. Quadros EV, Rothenberg SP, Jaffe EA. Endothelial cells from human umbilical vein secrete functional transcobalamin II. *Am J Physiol* 1989;256:C296–303.
163. Qian L, Quadros EV, Regec A, Zittoun J, Rothenberg SP. Congenital transcobalamin II deficiency due to errors in RNA editing. *Blood Cells Mol Dis* 2002;28:134–42.
164. Haurani FI, Hall CA, Rubin R. Megaloblastic anemia as a result of an abnormal transcobalamin II (Cardeza). *J Clin Invest* 1979;64:1253–9.
165. Seligman PA, Steiner LL, Allen RH. Studies of a patient with megaloblastic anemia and an abnormal transcobalamin II. *N Engl J Med* 1980;303:1209–12.
166. Li N, Rosenblatt DS, Seetharam B. Nonsense mutations in human transcobalamin II deficiency. *Biochem Biophys Res Commun* 1994;204:1111–8.
167. Li N, Rosenblatt DS, Kamen BA, Seetharam S, Seetharam B. Identification of two mutant alleles of transcobalamin II in an affected family. *Hum Mol Genet* 1994;3:1835–40.
168. Regec A, Quadros EV, Platika O, Rothenberg SP. The cloning and characterization of the human transcobalamin II gene. *Blood* 1995;85:2711–9.
169. Hall CA. The neurologic aspects of transcobalamin II deficiency. *Br J Haematol* 1992;80:117–20.
170. Carmel R. Mild transcobalamin I (haptocorrin) deficiency and low serum cobalamin concentrations. *Clin Chem* 2003;49:1367–74.
171. Carmel R, Parker J, Kelman Z. Genomic mutations associated with mild and severe deficiencies of transcobalamin I (haptocorrin) that cause mildly and severely low serum cobalamin levels. *Br J Haematol* 2009;147:386–91.
172. Shows TB, Alders M, Bennett S, Burbee D, Cartwright P, Chandrasekharappa S, et al. Report of the Fifth International Workshop on Human Chromosome 11 Mapping 1996. *Cytogenet Cell Genet* 1996;74:1–56.
173. Johnston J, Yang-Feng T, Berliner N. Genomic structure and mapping of the chromosomal gene for transcobalamin I (TCN1): comparison to human intrinsic factor. *Genomics* 1992;12:459–64.



Susanne H. Kirsch is at present employed at the Department of Clinical Chemistry and Laboratory Medicine of the Saarland University Hospital, Homburg, Germany. Recently, she finished working on her PhD in the field of folate. She studied Biology at the Saarland University (Saarbrücken, Germany) and received her diploma in 2004. From 2004 until 2008 she worked at the Department of Experimental Physics of the Saarland University. Her main research areas are folate, homocysteine, and vitamin B12 metabolism as well as method development using LC-MS/MS.



Prof. Dr. med. habil. Dr. rer. nat. Wolfgang Herrmann is as senior scientist at the Department of Clinical Chemistry and Laboratory Medicine, Medical School, Saarland University, 66421 Homburg, Germany. He is a member of the editorial board of the journal *“Clinical Chemistry and Laboratory Medicine”* and a member of the scientific advisory board of the journal *“Clinical Laboratory”*. Professor Herrmann is founding member and chairman of the *“Homocysteine Expert Panel”*, an organisation that supports research and spreads knowledge about homocysteine, vitamin deficiency and related diseases. His research focus has been in the areas of hyperhomocysteinemia, vitamin deficiency, lipid metabolism and degenerative diseases like cardiovascular, neurodegenerative or bone diseases. His Medline publication list comprises about 250 titles; approximately half of them involve research results related to the topic of homocysteine. Professor Herrmann editor of the textbook *“Vitamins in the Prevention of Human Diseases”* (DeGruyter, Berlin 2011), *Hypo- and Hypervitaminosis of B and D vitamins* (UNI-MED Verlag, Bremen 2013) and he is co-author of the textbook *“Laboratory and Diagnosis”* (Editor: L. Thomas, TH-Books, Frankfurt 2012).



Rima Obeid is Professor at the Department of Clinical Chemistry, Saarland University Hospital, Germany. She was assistant professor from 2006 to 2012 at the Department of Clinical Chemistry, University of the Saarland, Germany. In 2004, she received a two year research fellowship from the Alexander von Humboldt foundation. In 2003, she finished her PhD at the Department of Clinical Chemistry, University of the Saarland, Germany. In 1993, she graduated from the Department of Pharmacy, University of Damascus, Syria. She is expert in folate, choline and vitamin B12 metabolism in various physiological and pathological conditions.