

Włodzimierz M. Mikulski

## ON NATURALITY OF THE LEGENDRE OPERATOR

**Abstract.** We deduce that all natural operators of the type of the Legendre operator from the variational calculus in fibred manifolds are the constant multiples of the Legendre operator.

Given fibred manifolds  $Z_1 \rightarrow M$  and  $Z_2 \rightarrow M$  over the same base  $M$ , we denote by  $\mathcal{C}_M^\infty(Z_1, Z_2)$  the space of all base preserving fibred manifold morphisms of  $Z_1$  into  $Z_2$ . Let  $Y \rightarrow M$  be a fibred manifold,  $\dim(M) = m$ ,  $\dim(Y) = m + n$ . Let  $\lambda \in \mathcal{C}_M^\infty(J^s Y, \bigwedge^m T^* M)$  be an  $s$ -th order Lagrangian on  $Y \rightarrow M$ . Let  $\delta\lambda \in \mathcal{C}_{J^s Y}^\infty(J^s Y, V^* J^s Y \otimes \bigwedge^m T^* M)$  be the vertical differential of  $\lambda$  (the composition of the restriction  $\tilde{\delta}\lambda : V J^s Y \rightarrow V \bigwedge^m T^* M = \bigwedge^m T^* M \times_M \bigwedge^m T^* M$  of the differential  $d\lambda : T J^s Y \rightarrow T \bigwedge^m T^* M$  to the vertical sub-bundles with the second factor (essential) projection  $\bigwedge^m T^* M \times_M \bigwedge^m T^* M \rightarrow \bigwedge^m T^* M$ ). Let  $\Lambda(\lambda) : S^s T^* M \otimes V Y \rightarrow \bigwedge^m T^* M$  be the restriction of the vertical differential  $\delta\lambda : V J^s Y \rightarrow \bigwedge^m T^* M$  to the vector sub-bundle  $S^s T^* M \otimes V Y \subset V J^s Y$ , the kernel of  $V \pi_{s-1}^s : V J^s Y \rightarrow V J^{s-1} Y$ , where  $\pi_{s-1}^s : J^s Y \rightarrow J^{s-1} Y$  is the projection. The corresponding transformation  $\Lambda(\lambda) : J^s Y \rightarrow (\pi^{s-1})^* S^s T^* M \otimes V^* Y \otimes \bigwedge^m T^* M$  covering the identity of  $J^{s-1} Y$  (where the pull-back is given by the projection  $\pi^{s-1} : J^{s-1} Y \rightarrow Y$ ) is called the Legendre transformation determined by  $\lambda$ , [1]. It plays an important role in analytical mechanics, especially in the case of such  $\lambda$  (regular Lagrangians) for which  $\Lambda(\lambda) : J^s Y \rightarrow (\pi^{s-1})^* S^s T^* M \otimes V^* Y \otimes \bigwedge^m T^* M$  is a diffeomorphism (then it joints the Lagrange and Hamilton formalisms in fibred manifolds). Thus we have the Legendre operator

$$\Lambda : \mathcal{C}_M^\infty(J^s Y, \bigwedge^m T^* M) \rightarrow \mathcal{C}_Y^\infty(J^s Y, V^* Y \otimes S^s T^* M \otimes \bigwedge^m T^* M)$$

sending a Lagrangian  $\lambda \in \mathcal{C}_M^\infty(J^s Y, \bigwedge^m T^* M)$  into its Legendre transforma-

tion  $\Lambda(\lambda) \in \mathcal{C}_Y^\infty(J^s Y, V^* Y \otimes S^s T M \otimes \bigwedge^m T^* M)$ . In the present paper we prove the following theorem.

**THEOREM 1.** *Any local  $\mathcal{FM}_{m,n}$ -natural regular operator*

$$A : \mathcal{C}_M^\infty(J^s Y, \bigwedge^m T^* M) \rightarrow \mathcal{C}_Y^\infty(J^s Y, V^* Y \otimes S^s T M \otimes \bigwedge^m T^* M)$$

*is of the form  $A = c\Lambda$ ,  $c \in \mathbf{R}$ .*

**REMARK 1.** We recall that  $\mathcal{FM}_{m,n}$  is the category of fibred manifolds with  $m$ -dimensional bases and  $n$ -dimensional fibres and their fibred embeddings. The naturality means that for any  $\mathcal{FM}_{m,n}$ -morphism  $f : Y \rightarrow Y'$  and any  $s$ -th order Lagrangians  $\lambda \in \mathcal{C}_M^\infty(J^s Y, \bigwedge^m T^* M)$  and  $\lambda' \in \mathcal{C}_{M'}^\infty(J^s Y', \bigwedge^m T^* M')$  if  $\lambda$  and  $\lambda'$  are  $f$ -related then so are  $A(\lambda)$  and  $A(\lambda')$ . The regularity means that  $A$  transforms smoothly parametrized family of Lagrangians into smoothly parametrized family of respective type morphisms. The locality means that  $A(\lambda)_u$  depends on the germ of  $\lambda$  at  $u$ .

**Proof of Theorem 1.** From now on let  $(x^i, y^k)$ ,  $i = 1, \dots, m$ ,  $k = 1, \dots, n$  be the usual fibred coordinates on  $\mathbf{R}^{m,n}$ , the trivial bundle  $\mathbf{R}^m \times \mathbf{R}^n$  over  $\mathbf{R}^m$ . We will modify the proofs of the main results from the papers [3] and [4].

An  $\mathcal{FM}_{m,n}$ -morphism  $\varphi : \mathbf{R}^{m,n} \rightarrow \mathbf{R}^{m,n}$ ,  $\varphi((x^i), (y^k)) = ((x^i), (y^k - \sigma^k(x^i)))$  sends  $j_0^s((x), (\sigma^k))$  into

$$\Theta = j_0^s((x^i), (0)) \in (J^s \mathbf{R}^{m,n})_{(0,0)}.$$

Then (because of the invariance)  $A$  is uniquely determined by the evaluations

$$\langle A(\lambda)_\Theta, v \otimes \odot^s w \otimes u \rangle \in \mathbf{R}$$

for all  $\lambda \in \mathcal{C}_{\mathbf{R}^m}^\infty(J^s \mathbf{R}^{m,n}, \bigwedge^m T^* \mathbf{R}^m)$ , all  $v \in T_0 \mathbf{R}^n = V_{(0,0)} \mathbf{R}^{m,n}$ , all  $w \in T_0^* \mathbf{R}^m$  and all  $u \in \bigwedge^m T_0 \mathbf{R}^m$ .

Using the invariance of  $A$  with respect to  $\mathcal{FM}_{m,n}$ -morphism of the form  $id_{\mathbf{R}^m} \times \psi$  for linear isomorphisms  $\psi : \mathbf{R}^n \rightarrow \mathbf{R}^n$ , we get that  $A$  is uniquely determined by the evaluations

$$\langle A(\lambda)_\Theta, \frac{\partial}{\partial y^1} \otimes \odot^s w \otimes v \rangle \in \mathbf{R}$$

for all  $\lambda \in \mathcal{C}_{\mathbf{R}^m}^\infty(J^s \mathbf{R}^{m,n}, \bigwedge^m T^* \mathbf{R}^m)$ , all  $w \in T_0^* \mathbf{R}^m$  and all  $v \in \bigwedge^m T_0 \mathbf{R}^m$ .

Consider an arbitrary non-vanishing  $f : \mathbf{R}^m \rightarrow \mathbf{R}$ . There is a local diffeomorphism  $F : \mathbf{R}^m \rightarrow \mathbf{R}^m$  such that  $\frac{\partial F}{\partial x^1} = f$  and  $F(0) = 0$ . Then  $\mathcal{FM}_{m,n}$ -map  $(F, x^2, \dots, x^m, y^1, \dots, y^n)^{-1}$  preserves  $\Theta$ ,  $\frac{\partial}{\partial y^1} \otimes \odot^s w \otimes v$  and sends  $germ_0(d^m x)$  into  $germ_0(fd^m x)$ , where  $d^m x = dx^1 \wedge \dots \wedge dx^m$ . Then by the invariance,

regularity and density arguments,  $A$  is uniquely determined by the evaluations

$$\left\langle A(\lambda + bd^m x)_\Theta, \frac{\partial}{\partial y^1} \otimes \odot^s w \otimes v \right\rangle \in \mathbf{R}$$

for all  $\lambda \in \mathcal{C}_{\mathbf{R}^m}^\infty(J^s \mathbf{R}^{m,n}, \bigwedge^m T^* \mathbf{R}^m)$  with the condition  $\lambda(j_{x_o}^s((x^i), (0))) = 0$  for any  $x_o \in \mathbf{R}^m$ , all  $b \in \mathbf{R}$ , all  $w \in T_0^* \mathbf{R}^m$  and all  $v \in \bigwedge^m T_0 \mathbf{R}^m$ .

Then using the invariance of  $A$  with respect to  $\mathcal{FM}_{m,n}$ -maps of the form  $\varphi \times id$  with linear isomorphisms  $\varphi : \mathbf{R}^m \rightarrow \mathbf{R}^m$ , we see that  $A$  is uniquely determined by the evaluations

$$\left\langle A(\lambda + bd^m x)_\Theta, \frac{\partial}{\partial y^1} \otimes \odot^s d_0 x^m \otimes u^m \right\rangle \in \mathbf{R}$$

for all  $\lambda \in \mathcal{C}_{\mathbf{R}^m}^\infty(J^s \mathbf{R}^{m,n}, \bigwedge^m T^* \mathbf{R}^m)$  with the condition  $\lambda(j_{x_o}^s((x^i), (0))) = 0$  and all  $b \in \mathbf{R}$ , where  $u^m = \frac{\partial}{\partial x^1} \wedge \cdots \wedge \frac{\partial}{\partial x^m} \in \bigwedge^m T_0 \mathbf{R}^m$ .

Let  $\lambda$  and  $b$  be arbitrary as above. Using the invariance of  $A$  with respect to  $\mathcal{FM}_{m,n}$ -maps  $\psi_\tau = ((x^i), (\frac{1}{\tau^k} y^k))$  for  $\tau^k > 0$ , we get the homogeneity condition

$$\begin{aligned} A \left\langle (\psi_\tau)_*(\lambda + bd^m x)_\Theta, \frac{\partial}{\partial y^1} \otimes \odot^s d_0 x^m \otimes u^m \right\rangle &= \\ &= \tau^1 \left\langle A(\lambda + bd^m x)_\Theta, \frac{\partial}{\partial y^1} \otimes \odot^s d_0 x^m \otimes u^m \right\rangle. \end{aligned}$$

By Corollary 19.8 in [2] of the non-linear Peetre theorem we can assume that  $\lambda$  is a polynomial. It is easily seen that the coordinates of the polynomial  $(\psi_\tau)_* \lambda$  are the multiplication by monomials in  $\tau^k$ . The regularity of  $A$  implies that

$$\left\langle A(\lambda + bd^m x)_\Theta, \frac{\partial}{\partial y^1} \otimes \odot^s d_0 x^m \otimes u^m \right\rangle$$

is smooth in the coordinates of  $\lambda$  and  $b$ . Then by the homogeneity function theorem (and the above type of homogeneity) we deduce that

$$\left\langle A(\lambda + bd^m x)_\Theta, \frac{\partial}{\partial y^1} \otimes \odot^s d_0 x^m \otimes u^m \right\rangle$$

is a linear combination of the coordinates of  $\lambda$  on all  $x^{\tilde{\beta}} y_\beta^1 d^m x$  with coefficients being smooth functions in  $b$ , where  $(x^i, y_\beta^k)$  is the induced coordinate system on  $J^s \mathbf{R}^{m,n}$ . (Here and from now on  $\beta$  are  $m$ -tuples of non-negative integers with  $|\beta| \leq s$  and  $\tilde{\beta}$  are arbitrary  $m$ -tuples of non-negative integers.) In other words  $A$  is determined by the values

$$(*) \quad \left\langle A((ax^{\tilde{\beta}} y_\beta^1 + b)d^m x)_\Theta, \frac{\partial}{\partial y^1} \otimes \odot^s d_0 x^m \otimes u^m \right\rangle = a f_\beta^{\tilde{\beta}}(b)$$

for all  $a, b \in \mathbf{R}$ , all  $m$ -tuples  $\tilde{\beta}$  and all  $m$ -tuples  $\beta$  as above.

Let  $\tilde{\beta}_{i_o} \neq 0$  for some  $i_o = 1, \dots, m$ . We are going to use the invariance of  $A$  with respect to the locally defined  $\mathcal{FM}_{m,n}$ -map

$$\psi^{i_o} = (x^i, y^1 + x^{i_o}y^1, y^2, \dots, y^n)^{-1}$$

preserving  $x^i$ ,  $\Theta$ ,  $\frac{\partial}{\partial y^1}_0$ ,  $d_0 x^m$  and  $u^m$  and sending  $y_\beta^1$  into

$$y_\beta^1 + x^{i_o}y_\beta^1 + y_{\beta-1_{i_o}}^1 \quad (\text{if } \beta_{i_o} = 0 \text{ then the third term do not occur})$$

(because we have

$$\begin{aligned} y_\beta^1 \circ J^s((\psi^{i_o})^{-1})(j_{(x_o^i)}^s(x^i, \sigma^k)) &= \partial_\beta(\sigma^1 + x^{i_o}\sigma^1)(x_o^i) \\ &= \partial_\beta\sigma^1(x_o^i) + x^{i_o}\partial_\beta\sigma^1(x_o^i) + \partial_{\beta-1_{i_o}}\sigma^1(x_o^i) \\ &= (y_\beta^1 + x^{i_o}y_\beta^1 + y_{\beta-1_{i_o}}^1)(j_{(x_o^i)}^s(x^i, \sigma^k)), \end{aligned}$$

where  $\partial_\beta$  is the iterated partial derivative as indicated multiplied by  $\frac{1}{\beta!}$ . Then applying  $\psi^{i_o}$  to the left hand side of (\*) for  $\tilde{\beta} - 1_{i_o}$  instead of  $\tilde{\beta}$ , we see that the value (\*) is determined by the values (\*) for  $\beta - 1_{i_o}$  instead of  $\tilde{\beta}$ .

Continuing this procedure we see that the values (\*) are determined by the values (\*) for  $\tilde{\beta} = (0)$ . In other hand,  $A$  is determined by the values (\*) for  $\tilde{\beta} = (0)$  and all  $m$ -tuples  $\beta$  of non-negative integers with  $|\beta| \leq s$  and all reals  $a, b$ .

By the invariance of  $A$  with respect to  $\mathcal{FM}_{m,n}$ -maps  $((\tau^i x^i), (y^k))$  for  $\tau^i \neq 0$  we get the homogeneity conditions

$$\tau^{-\beta+(1, \dots, 1)} f_\beta^{(0)}(\tau^{(1, \dots, 1)} b) = \tau^{(1, \dots, 1, 1-s)} f_\beta^{(0)}(b).$$

If  $m \geq 2$ , then by the homogeneous function theorem this type of homogeneity gives that  $f_\beta^{(0)}$  are zero if it is not  $\beta_1 = \dots = \beta_{m-1} = \beta_m - s$ . Then if  $m \geq 2$ , we give that  $f_\beta^{(0)} = 0$  for all  $\beta \neq (0, \dots, 0, s)$ . Moreover, for  $\beta = (0, \dots, 0, s)$ ,  $f_\beta^{(0)}$  is constant. If  $m = 1$  this type of homogeneity gives that  $f_\beta^{(0)} = 0$  if  $\beta \neq (s)$ . Moreover  $f_{(s)}^{(0)}$  is constant. Then  $A$  is determined by the value

$$\left\langle A(y_{(0, \dots, 0, s)}^1 d^m x)_\Theta, \frac{\partial}{\partial y^1} \otimes \odot^s d_0 x^m \otimes u^m \right\rangle \in \mathbf{R}$$

if  $m \geq 2$ , and by the value

$$\left\langle A(y_{(s)}^1 d^1 x)_\Theta, \frac{\partial}{\partial y^1} \otimes \odot^s d_0 x^1 \otimes u^1 \right\rangle \in \mathbf{R}$$

if  $m = 1$ . Therefore the vector space of all  $A$  in question is 1-dimensional. This ends the proof of Theorem 1. ■

## References

- [1] I. Kolář, *A geometrical version of higher order Hamilton formalism in fibred manifolds*, J. Geom. Phys. **1** (2)(1984), 127–137.
- [2] I. Kolář, P. W. Michor, J. Slovák, *Natural Operations in Differential Geometry*, Springer, Berlin 1993.
- [3] W. M. Mikulski, *On naturality of the Helmholtz operator*, Archivum Math. Brno **41** (2) (2005), 145–149.
- [4] W. M. Mikulski, *On naturality of the formal Euler operator*, Demonstratio Math. **38** (1) (2005), 235–238.

INSTITUTE OF MATHEMATICS  
JAGIELLONIAN UNIVERSITY  
Reymonta 4  
KRAKÓW, POLAND  
e-mail: mikulski@im.uj.edu.pl

Received October 26, 2007.