

Thomas Mroz and Wolfgang Trutschnig*

A sharp inequality for Kendall's τ and Spearman's ρ of Extreme-Value Copulas

<https://doi.org/10.1515/demo-2018-0021>

Received November 14, 2018; accepted November 27, 2018

Abstract: We derive a new (lower) inequality between Kendall's τ and Spearman's ρ for two-dimensional Extreme-Value Copulas, show that this inequality is sharp in each point and conclude that the comonotonic and the product copula are the only Extreme-Value Copulas for which the well-known lower Hutchinson-Lai inequality is sharp.

Keywords: Concordance, Copula, Kendall tau, Spearman rho, Extreme Value Distribution

MSC: 60E05, 62E10, 28A12, 62H05

1 Introduction

It is well known that, on the one hand, Kendall's τ and Spearman's ρ are both measures of concordance, and that, on the other hand, they quantify different aspects of the underlying dependence structure (see [5]). Although a full characterization of the exact region Ω determined by all possible values of Kendall's τ and Spearman's ρ was only recently provided in [15], it has been well-known since the 1950s that for (continuous) random variables X, Y the value of $|\tau(X, Y) - \rho(X, Y)|$ can at most be $\frac{1}{2}$ (see [1, 2]). For standard subfamilies of copulas like Archimedean copulas and Extreme-Value copulas the values of Kendall's τ and Spearman's ρ may differ significantly less, determining the exact τ - ρ -region might, however, be even more difficult than determining Ω has been since in subfamilies handy dense subsets (like shuffles of the minimum copula M in case of Ω) may be hard to find or not even exist.

In 1990 Hutchinson and Lai conjectured that for continuous random variables X, Y such that Y is stochastically increasing in X and vice versa (see [7, 12] for a definition and equivalent formulations) the following inequalities hold:

$$-1 + \sqrt{1 + 3\tau(X, Y)} \leq \rho(X, Y) \leq \min \left\{ \frac{3\tau(X, Y)}{2}, 2\tau(X, Y) - \tau^2(X, Y) \right\} \quad (1)$$

In the sequel we will refer to the first inequality in (1) as lower Hutchinson-Lai inequality and to the second one as upper Hutchinson-Lai inequality. A counterexample to the upper Hutchinson-Lai inequality (concerning the $\frac{3\tau(X, Y)}{2}$ part) can be found in [12], and the lower Hutchinson-Lai inequality was disproved in [11]. On the other hand, Hürlimann (see [6]) provided a variational calculus based proof of the Hutchinson-Lai inequalities for an important family of stochastically increasing (continuous) random variables (X, Y) - those whose underlying copula C_A is an Extreme-Value copula. Extreme-Value copulas (EVCs, for short) form an important subclass of copulas that naturally arise in various fields of application like hydrology (see [14]) and finance (see [9, 10]), whenever maxima of i.i.d. sequences of random variables are considered. For more information on EVCs we refer to [3, 4, 13] and the references therein.

Thomas Mroz: Department for Mathematics, University of Salzburg, A-5020 Salzburg, Austria,

E-mails: ThomasMroz@a1.net

***Corresponding Author: Wolfgang Trutschnig:** Department for Mathematics, University of Salzburg, A-5020 Salzburg, Austria, E-mail: wolfgang@trutschnig.net

In the current paper we focus on EVCs and the lower Hutchinson-Lai inequality, show that it is only sharp for continuous random variables X, Y that are either comonotonic or independent, prove the validity of Conjecture (C2) in [15], i.e.

$$\frac{3\tau(X, Y)}{2 + \tau(X, Y)} \leq \rho(X, Y), \quad (2)$$

and then show that this inequality is sharp. As scale-invariant quantities, Kendall's τ and Spearman's ρ only depend on the underlying copula, we will therefore directly work with EVCs C_A and their corresponding Pickands dependence function A . Our proof is based on the sensitivity of τ and ρ with respect to certain modifications of piecewise linear Pickands dependence functions and does not use tools from variational calculus.

The rest of this paper is organized as follows: In Section 2 we introduce some notation and recall basic facts about two-dimensional Extreme-Value copulas that will be used in the sequel. Section 3 derives ineq. (2) with the help of two lemmata that are (in our opinion) interesting in themselves. An outlook to future work and the appendix containing a technical lemma needed in the proofs completes the paper.

2 Notation and preliminaries

In the sequel \mathcal{C} will denote the family of all two-dimensional *copulas*, $\mathcal{P}_{\mathcal{C}}$ the family of all *doubly stochastic measures*, i.e. the family of all probability measures on $[0, 1]^2$ whose marginals are uniformly distributed on $[0, 1]$; for background on copulas we refer to [4, 12] and [16]. For every $C \in \mathcal{C}$ the corresponding doubly stochastic measure will be denoted by μ_C . Letting d_{∞} denote the uniform metric on \mathcal{C} it is well known that $(\mathcal{C}, d_{\infty})$ is a compact metric space. $C \in \mathcal{C}$ is called *Extreme-Value copula* (EVC) if there exists a copula $B \in \mathcal{C}$ such that

$$C(x, y) = \lim_{n \rightarrow \infty} B^n(x^{\frac{1}{n}}, y^{\frac{1}{n}})$$

for all $x, y \in [0, 1]$. It is well known that the following three conditions are equivalent (see [3, 4, 12, 13]):

1. C is an EVC.
2. C is *max-stable*, i.e. $C(x, y) = C^n(x^{\frac{1}{n}}, y^{\frac{1}{n}})$ for all $n \in \mathbb{N}$ and all $x, y \in [0, 1]$.
3. There exists a *Pickands dependence function* A , i.e. a convex function $A : [0, 1] \rightarrow [0, 1]$ fulfilling $\max\{1-x, x\} \leq A(x) \leq 1$ for all $x \in [0, 1]$, such that

$$C(x, y) = (xy)^A\left(\frac{\ln(x)}{\ln(xy)}\right) \quad (3)$$

holds for all $x, y \in (0, 1)^2$.

In what follows we will only work with the convex set \mathcal{A} of all Pickands dependence functions and let \mathcal{C}_{EV} denote the family of all Extreme-Value copulas. For every $A \in \mathcal{A}$ we will write C_A for the copula induced by A according to (the right-hand side of) eq. (3). D^+A (D^-A) will denote the right-hand (left-hand) derivative of $A \in \mathcal{A}$. Setting $D^+A(1) := D^-A(1)$ it is straightforward to see that $D^+A : [0, 1] \rightarrow [-1, 1]$ is non-decreasing and right-continuous on $[0, 1]$. The Pickands function corresponding to the minimum copula M will be denoted by A_M , i.e. $A_M(x) = \max\{x, 1-x\}$. Furthermore $\|\cdot\|_{\infty}$ will denote the uniform norm on \mathcal{A} . For further properties of Pickands functions and their right-hand derivative we refer to [17] and the references therein.

It is well known that for every $A \in \mathcal{A}$ Spearman's ρ and Kendall's τ of the corresponding Extreme-Value Copula C_A can be calculated as

$$\begin{aligned} \tau(C_A) &= \int_0^1 \frac{t(1-t)}{A(t)} d(D^+A)(t), \\ \rho(C_A) &= 12 \int_0^1 \frac{1}{(1+A(t))^2} dt - 3. \end{aligned}$$

In the sequel we will simply write $\tau(A)$ and $\rho(A)$ instead of $\tau(C_A)$ and $\rho(C_A)$.

3 A sharp inequality between τ and ρ

We are now going to prove the inequality $\rho(A) \geq \frac{3\tau(A)}{2+\tau(A)}$ for all Pickands dependence functions $A \in \mathcal{A}$ and show that this inequality is sharp. Doing so we will first derive the result for the class of all piecewise linear Pickands dependence functions and then extend it to the full class \mathcal{A} via a standard denseness and continuity argument.

For every $\mathbf{x} = (x_1, x_2, \dots, x_n)$ with $x_0 := 0 < x_1 < x_2 < \dots < x_n < 1 =: x_{n+1}$ we will let $\mathcal{A}^{\mathbf{x}}$ denote the set of all $A \in \mathcal{A}$ which are linear on each of the intervals $[x_{i-1}, x_i], i \in \{1, \dots, n+1\}$. For $A \in \mathcal{A}^{\mathbf{x}}$, setting $y_i = A(x_i)$, the formulas for Kendall's τ and Spearman's ρ obviously become

$$\tau(A) = \sum_{i=1}^n \frac{x_i(1-x_i)}{y_i} \left(\frac{y_{i+1}-y_i}{x_{i+1}-x_i} - \frac{y_i-y_{i-1}}{x_i-x_{i-1}} \right), \tag{4}$$

$$\rho(A) = 12 \sum_{i=1}^{n+1} \frac{x_i-x_{i-1}}{(1+y_{i-1})(1+y_i)} - 3. \tag{5}$$

For $y_1 \in [A_M(x_1), 1]$, $\mathcal{A}_{y_1}^{\mathbf{x}}$ will denote the subclass of all $A \in \mathcal{A}^{\mathbf{x}}$ fulfilling $A(x_1) = y_1$. $T_{x_1, y_1} \in \mathcal{A}_{y_1}^{\mathbf{x}}$ denotes the triangular Pickands function induced by (x_1, y_1) , i.e. the only Pickands function which is linear on $[0, x_1]$ and on $[x_1, 1]$, and fulfills $T_{x_1, y_1}(x_1) = y_1$. For $A \in \mathcal{A}_{y_1}^{\mathbf{x}}$ and $x \in (0, x_1)$ we define the set I_x^A by

$$I_x^A = \left[\max \left\{ 1-x, A(x_1) - (x_1-x) \frac{A(x_2)-A(x_1)}{x_2-x_1} \right\}, 1 - \frac{1-A(x_1)}{x_1}x \right].$$

Obviously I_x^A coincides with the set of all $y \in [0, 1]$ such that there exists a (necessarily unique) Pickands function B which coincides with A on the interval $[x_1, 1]$, which is linear on $[0, x]$ and on $[x, x_1]$ and fulfills $B(x) = y$. In the sequel we will denote this function by $\varphi_{x,y}^{\mathbf{x}}(A)$ and refer to I_x^A as set of admissible y -values given A and x . Notice that for $y = 1 - \frac{1-A(x_1)}{x_1}x$ we have $\varphi_{x,y}^{\mathbf{x}}(A) = A$ and that for each $A \in \mathcal{A}_{y_1}^{\mathbf{x}}$ the admissible set I_x^A fulfills $I_x^A \subseteq I_x^{T_{x_1, y_1}}$.

Define $\Delta : \mathcal{A}^2 \rightarrow \mathbb{R}$ by

$$\Delta(A, B) = \rho(A) - \rho(B) - \left(\frac{3\tau(A)}{2+\tau(A)} - \frac{3\tau(B)}{2+\tau(B)} \right). \tag{6}$$

The subsequent two lemmata study properties of Δ which turn out to be key for deriving a sharp lower inequality for τ and ρ .

Lemma 3.1. *Suppose that $A \in \mathcal{A}_{y_1}^{\mathbf{x}}$, that $x \in (0, x_1)$ and that $y \in I_x^A$. Then the following inequality holds:*

$$\Delta(\varphi_{x,y}^{\mathbf{x}}(A), A) \geq \Delta(\varphi_{x,y}^{x_1}(T_{x_1, y_1}), T_{x_1, y_1}). \tag{7}$$

Moreover, in case of $y_1 < 1$ and $y < 1 - \frac{1-A(x_1)}{x_1}x$ we have equality in (7) if and only if $A = T_{x_1, y_1}$.




Figure 1: Example of a Pickands dependence function $A \in \mathcal{A}^x$ with $\mathbf{x} = (0.55, 0.85, 0.95)$ (black) and the Pickands function $\varphi_{x,y}^x(A)$ with $x = 0.3, y = 0.75$ (magenta).

Proof. Fix $A \in \mathcal{A}_{y_1}^x, x \in (0, x_1)$, and $y \in I_x^A$. Using equation (4) and (5) a straightforward calculation yields

$$\delta_\tau := \tau(\varphi_{x,y}^x(A)) - \tau(A) = \frac{(x_1 - x + xy_1 - x_1y)(y_1 - y - xy_1 + x_1y)}{(x_1 - x)yy_1}, \tag{8}$$

$$\delta_\rho := \rho(\varphi_{x,y}^x(A)) - \rho(A) = \frac{6(x_1(1 - y) - x(1 - y_1))}{(1 + y)(1 + y_1)}, \tag{9}$$

so both δ_τ and δ_ρ only depend on A in terms of x_1, y_1 and $y \in I_x^A$. Having this we directly get

$$\begin{aligned} \Delta(\varphi_{x,y}^x(A), A) &= \delta_\rho - \left(\frac{3\tau(\varphi_{x,y}^x(A))}{2 + \tau(\varphi_{x,y}^x(A))} - \frac{3\tau(A)}{2 + \tau(A)} \right) \\ &= \delta_\rho - \frac{6\delta_\tau}{(2 + \tau(A))(2 + \tau(A) + \delta_\tau)}, \end{aligned} \tag{10}$$

implying that $\Delta(\varphi_{x,y}^x(A), A)$ decreases if $\tau(A)$ decreases. In fact, in the case that x, y, x_1, y_1 are fixed, the quantities δ_ρ and δ_τ are fixed too, and A can only vary on the right hand side of x_1 , so equation (10) implies the stated monotonicity. Considering $y \in I_x^A \subseteq I_x^{T_{x_1, y_1}}$ as well as the fact that $A \leq T_{x_1, y_1}$ implies $\tau(A) \geq \tau(T_{x_1, y_1})$ inequality (7) now follows. The second assertion is a direct consequence of Lemma A.1 in the Appendix, which says that for arbitrary $A, B \in \mathcal{A}$ with $A \leq B$ and strict inequality in at least one point $t \in (0, 1)$ we have $\tau(B) < \tau(A)$. \square

Lemma 3.2. Suppose that $x_1 \in (0, 1)$ and that $y_1 \in [A_M(x_1), 1]$. Furthermore let $x \in (0, x_1)$ and $y \in I_x^{T_{x_1, y_1}}$ be arbitrary. Then

$$\Delta(\varphi_{x,y}^{x_1}(T_{x_1, y_1}), T_{x_1, y_1}) \geq 0 \tag{11}$$

holds. Moreover, we have equality in (11) if and only if $\varphi_{x,y}^{x_1}(T_{x_1,y_1})$ is a triangular Pickands dependence function too, i.e. if $\varphi_{x,y}^{x_1}(T_{x_1,y_1}) = T_{x,y}$.

Proof. Using $\tau(T_{x_1,y_1}) = \frac{1-y_1}{y_1}$ equation (10) simplifies to

$$\Delta(\varphi_{x,y}^{x_1}(T_{x_1,y_1}), T_{x_1,y_1}) = \frac{6N_1N_2N_3}{D_1D_2(D_3 + D_4 - D_5 + D_6)}$$

where

$$\begin{aligned} N_1 &= x(1 - y_1) - x_1(1 - y), & N_2 &= x(1 - y_1) - x_1(1 - y) + y_1 - y \\ N_3 &= y(x_1 + y_1) + y_1(1 - x) \\ D_1 &= 1 + y, & D_2 &= 1 + y_1, & D_3 &= x_1y(x_1 - x + y(1 - x_1)) \\ D_4 &= y_1(1 - x)(x_1 - x), & D_5 &= 2xyy_1(1 - x_1), & D_6 &= (1 - x)xy_1^2. \end{aligned}$$

To prove $\Delta(\varphi_{x,y}^{x_1}(T_{x_1,y_1}), T_{x_1,y_1}) \geq 0$ it suffices to show that both, the nominator and the denominator are non-negative for every $y \in I_x^{T_{x_1,y_1}}$, which can be done as follows:

(i) Obviously $N_3 > 0$ so the sign of the nominator is determined by N_1N_2 . Considering, firstly, that N_1N_2 is quadratic in y , secondly, that N_1N_2 is zero exactly for $y \in \{y_1 - (x_1 - x)\frac{1-y_1}{1-x_1}, 1 - \frac{1-y_1}{x_1}x\}$ (notice that the first point is not necessarily contained in $I_x^{T_{x_1,y_1}}$), and, thirdly, that the parabola opens downwards since the coefficient of the quadratic term is given by $-x_1(1 - x_1) < 0$ the nominator is non-negative.

(ii) Since D_1, D_2 are obviously positive the denominator is positive if and only if $D_3 + D_4 - D_5 + D_6 > 0$. Considering $y \in I_x^{T_{x_1,y_1}}$ there exists a unique point $b \in [0, 1 - \frac{1-y_1}{x_1}x - (y_1 - (x_1 - x)\frac{1-y_1}{1-x_1})]$ such that $y = y_1 - (x_1 - x)\frac{1-y_1}{1-x_1} + b$ holds. A straightforward calculation yields

$$\begin{aligned} D_3 + D_4 - D_5 + D_6 &= b^2x_1(1 - x_1) + b(x_1 - x)(2y_1 - x_1) \\ &\quad + (x_1 - x)y_1 \frac{x(1 - y_1) - 2x_1 + y_1 + 1}{1 - x_1}. \end{aligned}$$

The first summand is obviously non-negative, the same is true for the second one because of $y_1 \geq x_1$, and the third summand is positive since $x(1 - y_1) - 2x_1 + y_1 + 1 \geq x(1 - y_1) - x_1 + 1 > 0$. Considering that the second assertion is a direct consequence of the properties of the nominator mentioned in (i) the proof is complete. \square

Building upon the previous lemmata we can now proof the main result of this paper. It confirms the lower part of Conjecture (C2) in [15].

Theorem 3.3. *Every Pickands dependence function A fulfills $\rho(A) \geq \frac{3\tau(A)}{2+\tau(A)}$. The inequality is best possible in the sense that for every $u \in [0, 1]$ we can find a Pickands function A such that $\tau(A) = u$ and $\rho(A) = \frac{3u}{2+u}$.*

Proof. We first prove that the inequality holds for every piecewise linear Pickands dependence function by induction on the number of vertices: In the case of $n = 1$ and $x_1 \in (0, 1)$ each element A of \mathcal{A}^{x_1} is of triangular form, hence, using $\tau(T_{x_1,y_1}) = \frac{1-y_1}{y_1}$ and $\rho(T_{x_1,y_1}) = \frac{3(1-y_1)}{1+y_1}$, we get $\rho(A) - \frac{3\tau(A)}{2+\tau(A)} = 0$. In case of $n = 2$ vertices and $x_0 := 0 < x_1 < x_2 < 1$ each element A of $\mathcal{A}_{y_1}^{(x_1,x_2)}$ fulfills $A = \varphi_{x_1,y_1}^{x_2}(T_{x_2,y_2})$. Applying inequality (11) therefore yields

$$\rho(A) - \frac{3\tau(A)}{2 + \tau(A)} \geq \rho(T_{x_2,y_2}) - \frac{3\tau(T_{x_2,y_2})}{2 + \tau(T_{x_2,y_2})} = 0.$$

Assume now that $\rho(A) \geq \frac{3\tau(A)}{2+\tau(A)}$ holds for all piecewise linear Pickands functions with n vertices. Suppose that $\mathbf{x} = (x_1, x_2, \dots, x_n, x_{n+1})$ with $x_0 := 0 < x_1 < x_2 < \dots < x_n < x_{n+1} < 1 =: x_{n+2}$, fix $B \in \mathcal{A}^{\mathbf{x}}$ and set $y_1 = B(x_1)$. Since B can be represented as $B = \varphi_{x_1,y_1}^{(x_2,\dots,x_{n+1})}(A)$ for some piecewise linear A with at most n vertices, applying inequality (7) and inequality (11) yields

$$\rho(B) - \frac{3\tau(B)}{2 + \tau(B)} \geq \rho(A) - \frac{3\tau(A)}{2 + \tau(A)} \geq 0,$$

which completes the proof by induction.

The general result for not necessarily piecewise linear Pickands functions now follows via a standard denseness and continuity argument taking into account the following two facts: (i) The mapping $\Phi : (\mathcal{A}, \|\cdot\|_\infty) \rightarrow (\mathcal{C}_{EV}, d_\infty)$, defined by $\Phi(A) = C_A$ and the mapping $\Psi : \mathcal{C} \rightarrow [-1, 1]^2$, defined by $C \mapsto (\tau(C), \rho(C))$ are continuous (the latter is a direct consequence of the fact that τ and ρ are concordance measures, see [12]). (ii) The family of all piecewise linear Pickands functions is dense in $(\mathcal{A}, \|\cdot\|_\infty)$. The remaining assertion is a direct consequence of the identities $\tau(T_{x_1, y_1}) = \frac{1-y_1}{y_1}$ and $\rho(T_{x_1, y_1}) = \frac{3(1-y_1)}{1+y_1}$. \square

Remark 3.4. Theorem 3.3 implies that the lower Hutchinson-Lai inequality $\rho(C_A) \geq -1 + \sqrt{1 + 3\tau(C_A)}$ is only sharp for the Pickands functions A_M and $A_{II} = 1$, i.e. for the copulas M and II . Figure 2 depicts both Hutchinson-Lai inequalities together with the boundary of the τ - ρ -region of the full class \mathcal{C} as derived in [15] and the inequality derived in this paper.




Figure 2: The region determined by the Hutchinson-Lai inequalities (gray), the boundary of the full τ - ρ -region as recently established in [15] (green), and the sharp inequality derived in this paper (magenta).

Remark 3.5. Theorem 3.3 also implies that for every EVC C_A Kendall's τ can not exceed Spearman's ρ in general and that $\rho(C_A) \geq \frac{3\tau(C_A)}{2+\tau(C_A)} > \tau(C_A)$ holds for all $A \in \mathcal{C}_{EV} \setminus \{M, II\}$. Furthermore a straightforward calculation shows that the function $f : [0, 1] \rightarrow [0, 1]$, defined by $f(x) = \frac{3x}{2+x} - x$ attains its maximum at $x = \sqrt{6} - 2$, which means that for $\tau(C_A) = \sqrt{6} - 2$ we have $\rho(C_A) \geq \tau(C_A) + 5 - 2\sqrt{6} \approx \tau(C_A) + 0.101$.

4 Conclusion and outlook

We have shown that in the class of EVCs the lower Hutchinson-Lai inequality is only sharp for the comonotonic and the product copula, derived a new inequality, and proved that it is sharp in each point. We conjecture that in the class of EVCs the upper Hutchinson-Lai is not sharp either. Deriving the best-possible upper inequality will, however, be harder than deriving the sharp lower inequality has been.

A Appendix

For $A, B \in \mathcal{A}$ we will write $A \succ B$ if and only if $A(t) \geq B(t)$ for every $t \in [0, 1]$ with strict inequality in at least one point. The following Lemma and its proof is contained in [8], we include it for the sake of completeness.

Lemma A.1. *For $A, B \in \mathcal{A}$ with $A \succ B$ we have $\tau(C_A) < \tau(C_B)$.*

Proof. If $A \geq B$ then $C_A \leq C_B$ holds and we get

$$\begin{aligned} \tau(C_B) - \tau(C_A) &= -1 + 4 \int_{[0,1]^2} C_B d\mu_{C_B} + 1 - 4 \int_{[0,1]^2} C_A d\mu_{C_A} \\ &= 4 \left(\int_{[0,1]^2} (C_B - C_A) d\mu_{C_B} + \int_{[0,1]^2} C_A d\mu_{C_B} - \int_{[0,1]^2} C_A d\mu_{C_A} \right) \\ &= 4 \left(\int_{[0,1]^2} (C_B - C_A) d\mu_{C_B} + \int_{[0,1]^2} C_B d\mu_{C_A} - \int_{[0,1]^2} C_A d\mu_{C_A} \right) \\ &= 4 \left(\int_{[0,1]^2} (C_B - C_A) d\mu_{C_B} + \int_{[0,1]^2} (C_B - C_A) d\mu_{C_A} \right) \geq 0. \end{aligned} \tag{12}$$

Thereby the identity $\int_{[0,1]^2} C_1 d\mu_{C_2} = \int_{[0,1]^2} C_2 d\mu_{C_1}$, which holds for all two-dimensional copulas C_1, C_2 (see [12]) was used. According to [17], setting

$$L_D := \max\{x \in [0, 1] : D(x) = 1 - x\}, \quad R_D := \min\{x \in [0, 1] : D(x) = x\}$$

for every $D \in \mathcal{A}$ the support $\text{Supp}(\mu_{C_D})$ of μ_{C_D} coincides with the set $\{(x, y) \in [0, 1]^2 : f_{L_D}(x) \leq y \leq f_{R_D}(x)\}$, whereby $f_t : [0, 1] \rightarrow [0, 1]$ is defined as $f_t(x) = x^{\frac{1}{t}-1}$ for $t \in (0, 1)$, and as $f_0 = 0, f_1 = 1$ for $t \in \{0, 1\}$.

Suppose now that $A \succ B$ holds and that B does not coincide with $t \mapsto \max\{1 - t, t\}$ (in which case $\tau(C_A) < 1 = \tau(C_B)$ is trivial). Obviously $L_A \leq L_B < \frac{1}{2}$ as well as $R_A \geq R_B > \frac{1}{2}$ and we can find some $t_0 \in (L_B, R_B)$ fulfilling $A(t_0) > B(t_0)$. By continuity there exists some $\delta > 0$ such that $A(t) > B(t)$ holds for every $t \in [t, \bar{t}] := [t_0 - \delta, t_0 + \delta] \subseteq (L_B, R_B) \subseteq (L_A, R_A)$. For $x \in (0, 1)$ and $y \in (f_{\bar{t}}(x), f_t(x))$ we have $\frac{\ln(x)}{\ln(xy)} \in [t, \bar{t}]$, so equation (3) implies

$$\begin{aligned} \{(x, y) \in (0, 1)^2 : f_{\bar{t}}(x) < y < f_t(x)\} &\subset \{(x, y) \in (0, 1)^2 : C_B(x, y) > C_A(x, y)\} \\ &=: \{C_B > C_A\}. \end{aligned}$$

Considering

$$\{(x, y) \in (0, 1)^2 : f_{\bar{t}}(x) < y < f_t(x)\} \subset \text{Supp}(\mu_{C_B}) \subseteq \text{Supp}(\mu_{C_A})$$

shows $\mu_{C_B}(\{C_B > C_A\}) > 0$ and $\mu_{C_A}(\{C_B > C_A\}) = 0$, from which

$$\int_{[0,1]^2} (C_B - C_A) d\mu_{C_B} > 0 \quad \text{and} \quad \int_{[0,1]^2} (C_B - C_A) d\mu_{C_A} = 0$$

follows. Finally, applying eq. (12) yields $\tau(C_A) < \tau(C_B)$. □

References

- [1] Daniels, H. E. (1950). Rank correlation and population models. *J. R. Stat. Soc. Ser. B. Stat. Methodol.* 12(2), 171–181.
- [2] Durbin, J. and A. Stuart (1951). Inversions and rank correlation coefficients. *J. R. Stat. Soc. Ser. B. Stat. Methodol.* 13(2), 303–309.
- [3] de Haan, L. and S. I. Resnick (1977). Limit theory for multivariate sample extremes. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* 40(4), 317–337.
- [4] Durante, F. and C. Sempi (2016). *Principles of Copula Theory*. CRC Press, Boca Raton FL.
- [5] Fredricks, G. A. and R. B. Nelsen (2007). On the relationship between Spearman's rho and Kendall's tau for pairs of continuous random variables. *J. Statist. Plann. Inference* 137(7), 2143–2150.
- [6] Hürlimann, W. (2003). Hutchinson-Lai's conjecture for bivariate extreme value copulas. *Statist. Probab. Lett.* 61(2), 191–198.
- [7] Hutchinson, T. P. and C. D. Lai (1990). *Continuous Bivariate Distributions, Emphasising Applications*. Rumsby, Adelaide.
- [8] Kamnitsi, N., C. Genest, P. Jaworski, and W. Trutschnig (2018). On the size of the class of bivariate Extreme-Value copulas with a fixed value of Spearman's rho or Kendall's tau. *J. Math. Anal. Appl.*, to appear. Available at <https://doi.org/10.1016/j.jmaa.2018.11.057>.
- [9] Longin, F. and B. Solnik (2001). Extreme correlation of international equity markets. *J. Finance* 56(2), 649–676.
- [10] McNeil, A. J., R. Frey, and P. Embrechts (2015). *Quantitative Risk Management: Concepts, Techniques and Tools*. Revised Edition. Princeton University Press.
- [11] Munroe, P., T. Ransford, and C. Genest (2010). A counterexample to a conjecture of Hutchinson and Lai. *C. R. Math. Acad. Sci. Paris* 348(5-6), 305–310 (in french).
- [12] Nelsen, R. B. (2006). *An Introduction to Copulas*. Second edition. Springer, New York.
- [13] Pickands, J. (1981). Multivariate extreme value distributions. *Bull. Inst. Internat. Statist.* 49(2), 859–878. With a discussion (894–902).
- [14] Salvadori, G., C. De Michele, N. T. Kottegoda, and R. Rosso (2007). *Extremes in Nature - An Approach Using Copulas*. Springer, Dordrecht.
- [15] Schreyer, M., R. Paulin, and W. Trutschnig (2017). On the exact region determined by Kendall's τ and Spearman's ρ . *J. R. Stat. Soc. Ser. B. Stat. Methodol.* 79(2), 613–633.
- [16] Sempi, C. (2011). Copulae: Some mathematical aspects. *Appl. Stoch. Models Bus. Ind.* 27(1), 37–50.
- [17] Trutschnig, W., M. Schreyer, and J. Fernández-Sánchez (2016). Mass distributions of two-dimensional Extreme-Value copulas and related results. *Extremes* 19(3), 405–427.