

Research Article

Thu T. A. Le, Bao H. Dang, Thanh Q. C. Nguyen, Dam P. Nguyen, and Giao H. Dang*

Highly efficient removal of tetracycline and methyl violet 2B from aqueous solution using the bimetallic FeZn-ZIFs catalyst

Supplementary material




Figure S1: PXRD pattern of ZIF-8.




Figure S2: PXRD pattern of FeZn-ZIFs.

* **Corresponding author: Giao H. Dang**, College of Engineering, Can Tho University, Campus II, 3/2 Street, Ninh Kieu District, Can Tho City, Vietnam; Applied Chemical Engineering Lab, College of Engineering, Can Tho University, Campus II, 3/2 Street, Ninh Kieu District, Can Tho City, Vietnam, e-mail: dhgiao@ctu.edu.vn, tel: (+84) 292 3834267; fax: (+84) 292 383115

Thu T. A. Le, Bao H. Dang: College of Engineering, Can Tho University, Campus II, 3/2 Street, Ninh Kieu District, Can Tho City, Vietnam

Thanh Q. C. Nguyen: College of Natural Science, Can Tho University, Campus II, 3/2 Street, Ninh Kieu District, Can Tho City, Vietnam

Dam P. Nguyen: Department of Chemistry Education, Can Tho University, Campus II, 3/2 Street, Ninh Kieu District, Can Tho City, Vietnam




Figure S3: SEM images of FeZn-ZIFs (c: x10.000, d: x30.000).




Figure S4: EDX spectra of FeZn-ZIFs.




Figure S5: Nitrogen adsorption – desorption isotherm of FeZn-ZIFs.