

Hong-bo Tan, Xiu-qing Song and Hong Yan*

Photochemical oxidation of *N,N*-bis-(*tert*-butoxycarbonyl)-1,4-dihydropyrazine derivatives

Abstract: The photochemical oxidation of *N,N*-Bis-(*tert*-butoxycarbonyl)-1,4-dihydropyrazines was investigated by irradiation using a medium-pressure mercury lamp. The main products were isolated, and their structures were determined by spectral methods and single crystal X-ray diffraction analysis. Photooxygenation is suggested to be a [2+2] cycloaddition reaction of oxygen to the double bond.

Keywords: 1,4-dihydropyrazines; cycloaddition; photo-oxygenation; singlet oxygen.


DOI 10.1515/hc-2014-0202

Received December 4, 2014; accepted December 15, 2014

Introduction

The 1,4-dihydropyrazine moiety is a component of flavin coenzymes and several marine luciferins [1, 2]. The 1,4-dihydropyrazine system, with an 8π -electron structure and two enamine functions, has attracted much interest. Its antiaromatic character, electron-charge distribution in the molecule, properties of charge-transfer complexes, and role in redox-active biological molecules have been investigated [3–5]. Although the chemical properties of 1,4-dihydropyrazines have been studied for many years, their photochemical properties have received little attention. Here, a series of substituted 1,4-dihydropyrazines **1** were synthesized by the reported method [6, 7], and their photochemical oxidation properties were investigated. This work is a continuation of our studies of the photochemical properties of heterocyclic compounds, such as photodimerization and photochemical ring contraction

[8–11]. The photochemical reactions involving oxygen were found to result in the oxygenation of nitrogen heterocycles [12–14]. The photochemical oxidation properties of *N,N*-bis-(*tert*-butoxycarbonyl)-1,4-dihydropyrazines **1** were investigated by irradiation with a medium-pressure mercury lamp under an oxygen atmosphere. Control experiments were designed to study the mechanisms of the photooxygenation of compounds **1**.




Results and discussion

The photooxygenation of a solution of **1** in acetone in the presence of oxygen was investigated by irradiation with a 450-W medium-pressure mercury lamp. Acetone is a photosensitizer that enhances the photochemical reaction rate of the 1,4-dihydropyrazines [15–18]. After the reaction was completed, as judged by thin-layer chromatography (TLC) analysis, the solvent was removed under reduced pressure, and the residue was subjected to chromatography on silica gel using a mixed solvent of petroleum ether and ethyl acetate (20:3 v/v).


In the photooxygenation of **1a** for 6 h, product **2a** was obtained with a yield of 34% (Scheme 1). The formation of the methylated diol may involve a methyl radical that is generated by the irradiation of acetone via a Norrish type I reaction [19]. The initial diol structure is apparently the product derived from an intermediate dioxetane by reaction with H_2O as postulated in similar reports by Adam et al. [20, 21]. It can be suggested that H_2O and methyl radical in the system undergo a reaction with the dioxetane to yield **2a**. This suggestion was substantiated by control experiments, in which the formation of **2a** was accelerated with the addition of several drops of H_2O . When conventional solvents such as CH_2Cl_2 were used in the photooxygenation of **1a**, the diol was not found.

*Corresponding author: Hong Yan, College of Life Science and Bio-engineering, Beijing University of Technology, Beijing 100124, China, e-mail: hongyan@bjut.edu.cn

Hong-bo Tan and Xiu-qing Song: College of Life Science and Bio-engineering, Beijing University of Technology, Beijing 100124, China



Scheme 1




Scheme 2

In the photooxygenation of **1b** for 10 h, products **3b** and **4b** were obtained with yields of 22% and 24%, respectively (Scheme 2). The dioxetane **3b** is apparently the product of the reaction of **1b** with O_2 by [2+2] cycloaddition, as referred to by Adam et al. [22]. It can be suggested that the dioxetane is unstable and undergoes decomposition to the carbonyl compound **4b** by cleavage of the O-O and C-C bonds. Similar chemistry has been described [23–25]. In addition, it was observed that **3b** is formed first, and product **4b** is gradually formed as the reaction progresses. It was also observed that the isolated compound


3b is smoothly transformed to **4b** with a yield of 55% upon further irradiation.

In the photooxygenation of **1c** for 18 h, products **2c** and **5c** were obtained with the respective yields of 22% and 23% (Scheme 3). The formation of **2c** can be understood in terms of a similar process as the formation of **2a**, and the formation of **5c** is analogous to the formation of **4b** [26].

These findings form a basis for the suggested unified mechanism for the photooxygenation of **1** (Scheme 4). In the first step, substrate **1** undergoes a reaction with singlet oxygen by [2+2] cycloaddition that yields dioxetane **3**. Singlet oxygen is generated when triplet oxygen in air absorbs UV light (wavelengths <320 nm) [27–31]. The low energy gap between triplet and singlet oxygen allows for a sensitized excitation that can be achieved by UV light irradiation. The suggestion that singlet oxygen as a critical intermediate in the photooxygenation reaction of **1** was verified by flushing the reaction mixture with air or nitrogen. When air was continuously bubbled through the solution, the formation of **3** was accelerated, and the formation of **3** was abated under a nitrogen atmosphere. The role of singlet oxygen in this reaction was demonstrated by the addition of a quencher, which can capture the singlet oxygen, or a photosensitizer, which can convert the oxygen molecule to singlet oxygen. The quencher used



Scheme 3

Scheme 4 Proposed mechanism for the photooxygenation of **1**.




Figure 1: ORTEP diagrams of the crystal structure of **4b**.




Figure 2: ORTEP diagrams of the crystal structure of **5c**.

was 1,4-diazabicyclo[2.2.2]octane, and the formation of **3** was dramatically suppressed in its presence. By contrast, in the presence of rubrene, a known photosensitizer, the formation of **3** was considerably accelerated.

The formation of **2** from dioxetane **3** is proposed to be an intramolecular electron transfer-mediated cleavage of the peroxide. Because the O-O bond is weaker than the C-C bond, compound **3** is decomposed by a

stepwise process involving the homolysis of the peroxide bond to form a biradical intermediate, which then undergoes a reaction with H₂O to form a hydroxyl compound [32, 33]. The hydroxyl compound is methylated to give **2**. This methylation involves the methyl radical that is generated by irradiation of acetone (Norrish type I) [19]. The formation of **4** is proposed to originate from dioxetane **3** by a stepwise process that involves the

homolysis of the peroxide bond to form a biradical intermediate and a subsequent C-C bond cleavage [32, 34]. Compound **4** is directly converted to product **5** via a C-N bond cleavage, which is similar to the cleavage of *N*-boc oxamates, giving *N*-boc carbamates [26, 35]. Note that compound **4c** is unstable and is converted to the final product **5c** via a C-N bond cleavage. The three-dimensional structures of **4b** as *E*-isomer and **5c** as *Z*-isomer reflect the stability of **4** by the steric hindrance effect of the groups on the nitrogen atom. The increased steric interactions between the benzoyl and the phenyl groups lead to the *E*-isomer of **4b** being much more stable than the *Z*-isomer of **4b**.

Conclusions

Products of the photooxygenation of *N,N*-bis-(*tert*-butoxycarbonyl)-1,4-dihydropyrazines were isolated, and their structures were determined by ¹H NMR, ¹³C NMR, high-resolution mass spectral (HRMS), and single crystal X-ray diffraction analysis. The photooxygenation is suggested to proceed via the intermediary 1,2-dioxetane derived by [2+2] cycloaddition of singlet oxygen to the double bond.

Experimental

All chemicals were purchased from commercial sources and used without further purification. TLC was conducted on silica gel 60 F254 plates (Merck KGaA). Melting points were determined on a XT-5A digital melting point apparatus and are uncorrected. ¹H NMR spectra and ¹³C NMR spectra were recorded on a Bruker Avance 400 spectrometer at 400 and 100 MHz, respectively, in CDCl₃ or DMSO-*d*₆. HRMS analyses were conducted using a VG 70SE mass spectrometer operating in electrospray ionization mode. Irradiation for photochemical reactions was conducted using an Osram HBO 450W medium-pressure mercury lamp. The samples were irradiated in quartz cuvettes.

Preparation of *N,N*-bis-(*tert*-butoxycarbonyl)-1,4-dihydropyrazine derivatives 1

Compound **1a** was synthesized by the treatment of 1,4-dihydropyrazine bis(vinyl phosphate) with triethylammonium formate, palladium acetate, and triphenylphosphine in THF [6]. Compound **1b** was synthesized by a Pd-catalyzed Suzuki-Miyaura coupling reaction of 1,4-dihydropyrazine bis(vinyl phosphate) [6]. Compound **1c** was synthesized by the treatment of methyl ester of *N*-(4-toluenesulfonyl)-*N*-(*tert*-butoxycarbonyl)- α,β -didehydroalanine with DMAP and K₂CO₃ [7].

***N,N*-Bis-(*tert*-butoxycarbonyl)-1,4-dihydropyrazine (1a)** White solid isolated in 43% yield (lit. yield 45% [6]); mp 105–106°C (lit. mp 105–106°C [6]); ¹H NMR (CDCl₃): δ 1.45 (s, 18H), 5.77 (s, 2H), 5.89 (brs, 2H), 5.94 (br s, 2H), 6.04 (s, 2H).

***N,N*-Bis-(*tert*-butoxycarbonyl)-2,5-diphenyl-1,4-dihydropyrazine (1b)** White solid isolated in 69% yield (lit. yield 72% [6]); mp 188–189°C (lit. mp 188–189°C [6]); ¹H NMR (CDCl₃): δ 1.07 (s, 18H), 6.46 (s, 2H), 7.20–7.35 (m, 10H).

***N,N*-Bis-(*tert*-butoxycarbonyl)-2,5-bismethoxycarbonyl-1,4-dihydropyrazine (1c)** White solid isolated in 85% yield (lit. yield 89% [7]); mp 155–156°C (lit. mp 155–156°C [7]); ¹H NMR (CDCl₃): δ 1.50 (s, 18H), 3.80 (s, 6H), 7.09 (s, 2H).

General procedure for the photooxygenation of *N,N*-Bis-(*tert*-butoxycarbonyl)-1,4-dihydropyrazines 1

A solution of **1** (0.32 mmol) in 200 mL of acetone in a quartz vessel was irradiated using a 450-W medium-pressure mercury lamp. The progress of the reaction was monitored by TLC. After completion, the solvent was removed under reduced pressure, and the residue was subjected to chromatography on silica gel using a mixed solvent of petroleum ether and ethyl acetate (20:3 v/v) as an eluent to provide products **2–5**.

Di-*tert*-butyl 2-hydroxy-3-methoxy-2,3-dihydropyrazine-1,4-dicarboxylate (2a) Colorless needles; mp 147–148°C; ¹H NMR (DMSO-*d*₆): δ 6.31–6.46 (m, 1H, -OH), 5.94–6.14 (m, 2H, =CH), 5.17–5.55 (m, 2H, -CH), 3.20 (d, 3H, -OCH₃), 1.46 (s, 9H, -OC(CH₃)₃), 1.45 (s, 9H, -OC(CH₃)₃); ¹³C NMR (DMSO-*d*₆): δ 174.5, 152.5, 151.4, 73.0, 71.5, 55.1, 28.3, 28.2. HRMS (ESI). Calcd for C₁₅H₂₆N₂O₆ (M+Na⁺): *m/z* 353.36619. Found: *m/z* 353.36619.

1,4-Di-*tert*-butyl 2,5-dimethyl-2-hydroxy-3-methoxy-2,3-dihydropyrazine-1,2,4,5-tetracarboxylate (2c) Colorless needles; mp 155–156°C; ¹H NMR (CDCl₃): δ 7.41 (s, 1H, =CH), 5.42 (s, 1H, -CH), 4.48 (s, 1H, -OH), 3.84 (s, 3H, -OCH₃), 3.80 (s, 3H, -OCH₃), 3.37 (s, 3H, -OCH₃), 1.50 (s, 9H, -OC(CH₃)₃), 1.47 (s, 9H, -OC(CH₃)₃); ¹³C NMR (CDCl₃): δ 169.3, 164.3, 152.8, 150.7, 122.2, 109.5, 55.3, 53.7, 51.8, 27.9. HRMS (ESI). Calcd for C₁₉H₃₀N₂O₁₀ (M+Na⁺): *m/z* 469.43867. Found: *m/z* 469.43429.

Di-*tert*-butyl 1,4-diphenyl-7,8-dioxo-2,5-diazabicyclo[4.2.0]oct-3-ene-2,5-dicarboxylate (3b) White needles; mp 153–155°C; ¹H NMR (CDCl₃): δ 9.13 (s, 1H, =CH), 7.28–7.75 (m, 10H, Ar-H), 6.96 (s, 1H, -CH), 1.31 (s, 9H, -OC(CH₃)₃), 1.23 (s, 9H, -OC(CH₃)₃); ¹³C NMR (CDCl₃): δ 170.5, 161.0, 151.6, 151.2, 136.1, 135.5, 132.5, 129.1, 128.5, 128.4, 128.2, 125.3, 125.2, 123.7, 84.3, 27.5, 27.4. HRMS (ESI). Calcd for C₂₆H₃₀N₂O₆ (M+Na⁺): *m/z* 489.20016. Found: *m/z* 489.19871.

***tert*-Butyl (*E*)-benzoyl(2-(*N*-(*tert*-butoxycarbonyl)formamido)-2-phenylvinyl)carbamate (4b)** Colorless needles; mp 135–136°C; ¹H NMR (CDCl₃): δ 9.44 (s, 1H, -CHO), 7.24–7.53 (m, 10H, Ar-H), 6.52 (s, 1H, =CH), 1.36 (s, 9H, -OC(CH₃)₃), 1.10 (s, 9H, -OC(CH₃)₃); ¹³C NMR (CDCl₃): δ 171.3, 162.8, 152.1, 151.3, 135.4, 134.7, 131.8, 130.1, 128.7, 128.5, 128.3, 128.0, 127.9, 127.1, 27.7, 27.2. HRMS (ESI). Calcd for C₂₆H₃₀N₂O₆ (M+Na⁺): *m/z* 489.20039. Found: *m/z* 489.19871. Single crystal X-ray diffraction of **4b**: deposition CCDC-995232 (see next section).

Methyl (Z)-3-[(*tert*-butoxycarbonyl)amino]-2-[*N*-(*tert*-butoxycarbonyl)formamido]acrylate (5c) Colorless needles; mp 169–170°C; ^1H NMR (CDCl_3): δ 9.26 (s, 1H, -CHO), 8.01 (s, 1H, =CH), 6.61 (s, 1H, -NH), 3.77 (s, 3H, -OCH₃), 1.52 (s, 18H, -OC(CH₃)₃); ^{13}C NMR (CDCl_3): δ 164.0, 162.0, 151.5, 150.8, 85.0, 83.3, 52.0, 29.7, 28.0, 27.8. HRMS (ESI). Calcd for $\text{C}_{15}\text{H}_{24}\text{N}_2\text{O}_7$ ($\text{M}+\text{Na}^+$): m/z 367.35003. Found: m/z 367.34968. Single crystal X-ray diffraction of **5c**: deposition CCDC-995231 (see next section).

X-ray diffraction analysis of **4b** and **5c**

Crystals of **4b** and **5c** suitable for X-ray diffraction analysis were obtained by slow evaporation of an ethyl acetate solution of **4b** and **5c** at room temperature. The single crystal X-ray diffraction measurement was conducted on a Rigaku Saturn CCD area-detector diffractometer at 113(2) K using graphite monochromated Mo $K\alpha$ radiation ($\lambda=0.71070$ Å) in the ω and ϕ scanning mode (Figures 1 and 2). An empirical absorption correction was applied using the ABCOR program. All structures were solved by direct methods using the SHELXS-97 program and refined by full matrix least squares on F^2 using the SHELXL-97 program. All hydrogen atoms were geometrically fixed using the riding model. Files CCDC-995231 and CCDC-995232 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from www.ccdc.cam.ac.uk/conts/retrieving.html.

Acknowledgments: This work was financially supported by the Key Projects in the National Science and Technology Pillar Program (no. 2012ZX10001007-008-002).

References

- [1] Teranishi, K.; Goto, T. Synthesis and chemiluminescence of coelenterazine (Oplophorus luciferin) analogues. *Bull. Chem. Soc. Jpn.* **1990**, *63*, 3132–3140.
- [2] Goto, T.; Inoue, S.; Sugiura, S. Cypridina bioluminescence IV. Synthesis and chemiluminescence of 3, 7-dihydroimidazo [1,2-a] pyrazin-3-one and its 2-methyl derivative. *Tetrahedron Lett.* **1968**, *9*, 3873–3876.
- [3] Walsh, C. Flavin coenzymes: at the crossroads of biological redox chemistry. *Acc. Chem. Res.* **1980**, *13*, 148–155.
- [4] Brook, D. J.; Haltiwanger, R. C.; Koch, T. H. New type of charge-transfer complex from an antiaromatic electron donor. Possible radical cation stabilization by the captodative effect. *J. Am. Chem. Soc.* **1991**, *113*, 5910–5911.
- [5] Lichtblau, A.; Ehrend, A.; Hausen, H. D.; Kaim, W. *N,N'*-Disilylated 1,4-dihydropyrazines: organosilyl substitution reactions, structural effects of steric hindrance, and electron exchange with C60. *Chemische Berichte.* **1995**, *128*, 745–750.
- [6] Chaignaud, M.; Gillaizeau, I.; Ouhamou, N.; Coudert, G. New highlights in the synthesis and reactivity of 1,4-dihydropyrazine derivatives. *Tetrahedron* **2008**, *64*, 8059–8066.
- [7] Rodrigues, A.; Ferreira, P. M.; Monteiro, L. S. Synthesis and reactivity of a 1,4-dihydropyrazine derivative. *Tetrahedron* **2004**, *60*, 8489–8496.
- [8] Zhu, X.; Li, W.; Yan, H.; Zhong, R. Triplet phenacylimidazoliums-catalyzed photocycloaddition of 1,4-dihydropyridines: an experimental and theoretical study. *J. Photochem. Photobiol. A: Chem.* **2012**, *241*, 13–20.
- [9] Song, X.-Q.; Wang, H.-Q.; Yan, H.; Ni, C.-L.; Zhong, R.-G. Synthesis, NMR analysis and X-ray crystal structure of 6, 12-bis (4-fluorophenyl)-3, 9-dioxatetraasterane. *J. Mol. Struct.* **2011**, *1006*, 489–493.
- [10] Zhu, X.; Ni, C.; Yan, H.; Zhong, R. Synthesis and X-ray crystallographic analysis of 1,4-dihydropyridine photodimerization. *J. Photopolymer Sci. Technol.* **2009**, *22*, 379–384.
- [11] Xin, H.; Sun, W.; Yan, H.; Song, X. Photochemical ring contraction of 1-aryl-1,4-dihydropyrazine to 1-aryl-1H-imidazole. *J. Photochem. Photobiol. A: Chem.* **2013**, *267*, 49–54.
- [12] Boyer, J. H. Increasing the index of covalent oxygen bonding at nitrogen attached to carbon. *Chem. Rev.* **1980**, *80*, 495–561.
- [13] Adam, W.; Griesbeck, A. G.; Gollnick, K.; Knutzen-Mies, K. 1,2-Dioxetanes derived from 4,5-dimethyl-2,3-dihydrofuran and 4,5-dimethyl-2,3-dihydrothiophene. Synthesis via photooxygenation, activation parameters, and excitation properties. *J. Org. Chem.* **1988**, *53*, 1492–1495.
- [14] Castro, C.; Dixon, M.; Erden, I.; Ergonenc, P.; Keeffe, J. R.; Sukhovitsky, A. Dye-sensitized photooxygenation of the carbon-nitrogen double bond. *J. Org. Chem.* **1989**, *54*, 3732–3738.
- [15] Chu, W.; Tsui, S. Photoreductive model of disperse orange 11 in aqueous acetone and triethylamine. *J. Environ. Eng.* **2001**, *127*, 741–747.
- [16] Chu, W.; Tsui, S.-M. Photo-sensitization of diazo disperse dye in aqueous acetone. *Chemosphere* **1999**, *39*, 1667–1677.
- [17] Chu, W.; Choy, W. The mechanisms of rate enhancing and quenching of trichloroethene photodecay in the presence of sensitizer and hydrogen sources. *Water Res.* **2002**, *36*, 2525–2532.
- [18] Gruen, H.; Görner, H. Properties of 16,17-disubstituted dihydroviolanthrones formed by reaction of violanthrones with photogenerated radicals. *Photochem. Photobiol. Sci.* **2010**, *9*, 1088–1093.
- [19] Scaiano, J. C.; Stampeleskie, K. G.; Hallett-Tapley, G. L. Photochemical Norrish type I reaction as a tool for metal nanoparticle synthesis: importance of proton coupled electron transfer. *Chem. Commun.* **2012**, *48*, 4798–4808.
- [20] Murphy, S.; Adam, W. The elusive 1,4-dioxy biradical: revised mechanism for the formation of diol from 3,3-dimethyldioxetane in cyclohexadiene. *J. Am. Chem. Soc.* **1996**, *118*, 12916–12921.
- [21] Adam, W.; Bosio, S. G.; Turro, N. J. Highly diastereoselective dioxetane formation in the photooxygenation of enecarbamates with an oxazolidinone chiral auxiliary: steric control in the [2+2] cycloaddition of singlet oxygen through conformational alignment. *J. Am. Chem. Soc.* **2002**, *124*, 8814–8815.
- [22] Adam, W.; Peters, E. M.; Peters, K.; Platsch, H.; Schmidt, E.; Von Schnering, H. G.; Takayama, K. Synthesis, thermal stability, and chemiluminescence properties of the dioxetanes derived from 1,4-dioxins. *J. Org. Chem.* **1984**, *49*, 3920–3928.
- [23] Zhang, X.; Foote, C. S. 1,2-Dioxetane formation in photooxygenation of *N*-acylated indole derivatives. *J. Org. Chem.* **1993**, *58*, 5524–5527.
- [24] Singh, S.; Ullman, E. F. Reversible formation of excited states in intramolecular donor assisted chemiluminescence reactions of dioxetanes. *Chem. Commun.* **2003**, *14*, 1756–1757.

- [25] Kotani, H.; Ohkubo, K.; Fukuzumi, S. Photocatalytic oxygenation of anthracenes and olefins with dioxygen via selective radical coupling using 9-mesityl-10-methylacridinium ion as an effective electron-transfer photocatalyst. *J. Am. Chem. Soc.* **2004**, *126*, 15999–16006.
- [26] Berrée, F.; Michelot, G.; Le Corre, M. N-Boc ethyl oxamate: a new nitrogen nucleophile for use in Mitsunobu reactions. *Tetrahedron Lett.* **1998**, *39* (45), 8275–8276.
- [27] Zhang, X.; Foote, C. S.; Khan, S. I. Reactions of N-acylated indoles with singlet oxygen. *J. Org. Chem.* **1993**, *58*, 47–51.
- [28] Adam, W.; Ahrweiler, M.; Peters, K.; Schmiedeskamp, B. Oxidation of N-acylindoles by dimethyldioxirane and singlet oxygen: substituent effects on thermally persistent indole epoxides and dioxetanes. *J. Org. Chem.* **1994**, *59*, 2733–2739.
- [29] Clennan, E. L. New mechanistic and synthetic aspects of singlet oxygen chemistry. *Tetrahedron* **2000**, *56*, 9151–9179.
- [30] Clennan, E. L.; Pace, A. Advances in singlet oxygen chemistry. *Tetrahedron* **2005**, *61* (28), 6665–6691.
- [31] Larson, R. A.; Marley, K. A. Singlet oxygen in the environment. In *Environ. Photochem.* Springer: Berlin Heidelberg, 1999; pp 123–137.
- [32] Richardson, W. H.; Stiggal-Estberg, D. L.; Chen, Z.; Baker, J. C.; Burns, D. M.; Sherman David, G. Substituent effects upon efficiency of excited-state acetophenones produced on thermolysis of 3,4-diaryl-3,4-dimethyl-1,2-dioxetanes. *J. Org. Chem.* **1987**, *52*, 3143–3150.
- [33] Ando, W.; Kohmoto, S. Reactions of singlet oxygen. Formation of hydroxy ketones in the photo-oxygenation of fluorenylidene derivatives. *J. Chem. Soc. Chem. Commun.* **1978**, *3*, 120–121.
- [34] Schaap, A. P.; Gagnon, S. D.; Zaklika, K., Substituent effects on the decomposition of 1,2-dioxetanes: a Hammett correlation for substituted 1, 6-diaryl-2,5,7,8-tetraoxabicyclo [4.2. 0] octanes. *Tetrahedron Lett.* **1982**, *23*, 2943–2946.
- [35] Berrée, F.; Bazureau, J.-P.; Michelot, G.; Cone, M. L. N-Boc and N-CBz ethyl oxamates: new Gabriel reagents. *Synth. Commun.* **1999**, *29*, 2685–2693.