

Bing Zhao, Yue Fang, Ming-jie Ma, Qi-Gang Deng*, Yu Xu and Li-yan Wang

A new on-fluorescent sensor for Ag⁺ based on benzimidazole bearing bis(ethoxycarbonylmethyl) amino groups

DOI 10.1515/hc-2015-0002

Received January 19, 2015; accepted March 20, 2015; previously published online July 14, 2015

Abstract: A new fluorescent sensor based on a benzimidazole unit bearing bis(ethoxycarbonylmethyl) amino groups was designed and synthesized. The ligand exhibits strong sensitivity and selectivity for Ag⁺ by enhanced fluorescent intensity in the presence of a wide range of other tested metal ions in methanol. The colorimetric and fluorescent response to Ag⁺ can be conveniently detected even by the naked eye, which offers a facile method for visual detection of Ag⁺.

Keywords: Ag⁺ ion; benzimidazole; fluorescent sensor; naked eye.

Introduction

Some metal ions, such as Ca²⁺, Fe³⁺, Zn²⁺, K⁺, Mg²⁺, Ag⁺, and Na⁺, are playing a key biological role in human body [1], whereas other heavy metal ions, including Al³⁺, Cd²⁺, Cr³⁺, Hg²⁺, and Pb²⁺, can be harmful to organisms [2]. Accordingly, the development of fluorescent probes for the detection of various metal ions is becoming an important research area [3–6]. Over the last decades, the rational design and synthesis of efficient sensors to selectively recognize targets has been a hot topic in molecular recognition studies. Generally, a typical sensor is devised by a covalent linkage of three parts, namely, a receptor unit, a spacer unit, and a signaling unit, although there are some examples of spacer-free probes also. Usually, highly selective probes for transition or heavy elements that give a

positive response rather than fluorescent quenching upon the ion binding are preferred to promote the sensitivity factor [7–10]. The design of such turn-on silver ion (Ag⁺) sensors is an intriguing challenge because many transition elements often cause fluorescent quenching [11].

Most of the reported signaling units are mainly based on rhodamine [12] and coumarin fragments [13]. The imidazole unit [14, 15] is a newer fluorescent group that is becoming important in the field of chemosensors given its fluorescence off-on behavior that results from its particular structural properties [16]. Park and coworkers [17] have shown that a proper substitution of a tetraphenylimidazole scaffold can lead to white light-emitting compounds by the combination of excited-state intramolecular proton transfer and restricted energy transfer. In addition, a polymer containing thienimidazole systems can be used for both colorimetric and ratiometric detections of Hg²⁺ as well as fluorometric detection of Zn²⁺ via fluorescence turn-on response with augmented lifetime via the chelation of metal ions to both S and N heteroatoms [18]. Unfortunately, in the existing reports, the fluorescence enhancement in most cases is small and usually suffers from a high background interference. To our knowledge, few reports have been devoted to fluorescence enhancement of probes for Ag⁺.


Results and discussion

In continuation of our work on imidazole derivatives [19], a new silver ion probe **L** was synthesized and studied. The synthetic route to the ligand compound-bearing bis(ethoxycarbonylmethyl)amino and benzimidazole functionalities is outlined in Scheme 1.

The fluorescence properties of ligand **L** were investigated by UV-vis absorption and fluorescence emission spectra in methanol. The UV absorption spectrum of **L** in the presence of various cations is presented in Figure 1. The metal-free ligand shows no absorption higher than 350 nm. The addition of 10 equivalents of metal ion, including K⁺, Na⁺, Ca²⁺, Mg²⁺, Ba²⁺, Zn²⁺, Fe³⁺, Pb²⁺, Cu²⁺,

*Corresponding author: Qi-Gang Deng, Chemistry and Chemical Engineering Institute, Qiqihar University, Qiqihar 161006, China, e-mail: dqigang@163.com

Bing Zhao, Yue Fang, Ming-jie Ma, Yu Xu and Li-yan Wang:
Chemistry and Chemical Engineering Institute, Qiqihar University,
Qiqihar 161006, China



Scheme 1




Figure 1 Absorption spectra of **L** (10 μm) in the presence of 10 equivalents of different metal ions in methanol.

Co²⁺, Ni²⁺, Cd²⁺, and Hg²⁺, results in a little change in absorption (Figure 1), as analyzed by the UV absorption spectrum and the visible absorption observed by the naked eye. Upon the addition of 10 equivalents of Cr³⁺ or Al³⁺, the UV absorption peak exhibits a slightly red shift from 323 to 345 nm. By contrast, in the presence of Ag⁺ ion under otherwise similar conditions, the solution of **L** undergoes a significant color change from colorless to dark yellow (inset in Figure 1) with an emergence of a new absorption peak at approximately 424 nm. These results

show the selectivity of the interaction of compound **L** with silver ion in the presence of other ions.

During the spectrophotometric titration of the solution of **L** in methanol by solution of AgNO₃ in the same solvent, the optical density of the ligand absorption band ($\lambda_{\text{max}}=323$ nm) decreases, and the characteristic absorption band of the complex Ag-**L** with $\lambda_{\text{max}}=424$ nm appears and grows gradually with the increasing concentration of Ag⁺ (not shown). This process parallels the color change observed by the naked eye.

The insert shows color change to the naked eye upon the addition of Ag⁺ to the solution of **L** under otherwise similar conditions.

To further investigate the interaction of Ag⁺ and **L**, the fluorescence spectra were analyzed (Figure 2). Compound **L** shows a very weak fluorescence at 424 nm in the absence of metal ions with the fluorescence intensity of approximately 26 a.u. After the addition of metal ion, including Ba²⁺, Ca²⁺, Cd²⁺, Co²⁺, Cu²⁺, Fe³⁺, Hg²⁺, K⁺, Mg²⁺, Na²⁺, Ni²⁺, Pb²⁺, or Zn²⁺, changes in the fluorescence emission are minimal. Even in the presence of Al³⁺ or Cr³⁺ when a slight red shift in UV absorption is observed, the fluorescence spectra are still very similar to that of the free ligand **L**. By contrast, the addition of Ag⁺ (10 equivalents) causes a remarkable enhancement of fluorescence intensity at




Figure 2 Selective fluorescent spectrum of the interaction of Ag ion and **L**. (A) Fluorescence spectra of **L** (10.0 μm) in the absence and presence of various cations (100.0 μm) in methanol solution ($\lambda_{\text{ex}}=323$ nm). (B) Fluorescence intensity ratios F/F_0 of **L** (5.0×10^{-7} M, constant) at 424 nm ($\lambda_{\text{ex}}=323$ nm) in methanol in the presence of Ag⁺ only (F_0) and in the presence of Ag⁺ and 10 equivalents of additional metal ion (F).

424 nm by a factor of approximately 5.5 (Figure 2A). To validate the selectivity of **L** toward Ag⁺, competition experiments in the presence of Ag⁺ only (fluorescence intensity F_0) and in the presence of Ag⁺ and other ions (fluorescence intensity F) were conducted. As shown in Figure 2B, the maximum fluorescence intensity ratio F/F_0 is approximately 1.28 for K⁺ and the minimum value of 0.86 is observed in the presence of Pb²⁺. These results demonstrate that the enhancement in fluorescence intensity resulting from the addition of Ag⁺ is not influenced significantly by the addition of the background metal ions. Additional experiments showed that the fluorescence quantum yield of the ligand **L** at 424 nm is increased from 0.21 to 0.72 in the presence of Ag⁺ in methanol solution.

The fluorescence titration experiments of **L** with Ag⁺ in a methanol solution were also investigated in the range of molar ratios [Ag⁺]/[**L**]=0–60. The fluorescence intensity increases with the increasing concentration of Ag⁺ and reaches a plateau with a ratio of [Ag⁺]/[**L**] >20 (not shown).

To determine the stoichiometry of the Ag-**L** complex, Job's method was applied using fluorescence titration experiments (Figure 3). As can be seen, the fluorescence intensity reaches a maximum for the ratio of [Ag⁺]/{[Ag⁺]+[**L**]} of 0.5, which indicates a 1:1 stoichiometry of Ag⁺ to **L** in the complex.

Conclusions

A new fluorescence probe **L** for Ag⁺ was synthesized and analyzed. The enhancement in fluorescence intensity of **L** upon interaction with Ag⁺ may be explained on the basis of the thermodynamically favorable photoinduced electron transfer (PET) mechanism between **L** and Ag⁺. The binding of Ag⁺ to **L** through the lone pairs of N and

O atoms hinders the PET process leading to fluorescence intensity enhancement.

Experimental

¹H NMR (600 MHz) and ¹³C NMR (150 MHz) spectra were measured on the Bruker instrument. UV-vis absorption spectra were recorded on a TU-1901 ultraviolet and visible spectrophotometer (1 cm quartz cell) at 25°C. Fluorescence measurements were conducted on a Perkin Elmer LS55 fluorescence spectrometer using a 1-cm quartz cell at 25°C, with excitation and emission slit widths of 10 and 4 nm, respectively, and excitation wavelength at 323 nm. Starting materials **2** and **3** were synthesized according to the reported procedures [20].

Synthesis of ethyl *N*-{[4-(1*H*-benzimidazol-2-yl)]phenyl]}-*N*-(ethoxycarbonyl)methyl]-aminoacetate (**L**)

A mixture of *o*-phenylenediamine (5.30 g, 5 mmol) and compound **3** (1.46 g, 5 mmol) in methanol was heated under reflux at 65°C for approximately 6 h until the reaction was completed, as judged by TLC analysis, and then concentrated under reduced pressure. The crude product **L** was subjected to silica gel column chromatography eluting with mixtures of hexanes and ethyl acetate in the ratios from 10:1 to 7:3. The free base **L** was obtained in 77% yield as a gray solid; mp 170.4–171.4°C; IR (KBr, cm⁻¹): 3241, 3109, 2947, 2865, 1678, 1603, 1559, 1511, 1432, 1285, 1103; ¹H NMR (600 MHz, DMSO-*d*₆): δ 12.55 (s, 1H), 7.96 (d, 2H, *J* = 8.4 Hz), 7.57 (d, 1H, *J* = 7.2 Hz), 7.45 (d, 1H, *J* = 7.2 Hz), 7.13 (m, 2H), 6.71 (d, 2H, *J* = 8.4 Hz), 4.29 (s, 4H), 4.15 (q, 4H, *J* = 13.8 Hz), 1.22 (t, 6H, *J* = 7.2 Hz); ¹³C NMR (150 MHz, DMSO-*d*₆): δ 170.7, 152.3, 149.5, 128.0, 119.4, 112.4, 61.0, 53.0, 14.6. Anal. Calcd for C₂₁H₂₃N₃O₄: C, 66.13; H, 6.08; N, 11.02. Found: C, 66.29; H, 6.17; N, 10.99.

Acknowledgments: This work was supported by the Program for Education Department of Heilongjiang Province (no. 12541862) and Qiqihar University Graduate Innovation Fund Grants (No. YJSCX2014-028X).

References

- [1] Dang F. F.; Lei, K. W.; Liu, W. S. A new highly selective fluorescent silver probe. *J. Fluoresc.* **2008**, *18*, 149–153.
- [2] Liu, D. L.; Pang, T.; Ma, K. F.; Jiang, W.; Bao, X. F. A new highly sensitive and selective fluorescence chemosensor for Cr³⁺ based on rhodamine B and a 4,13-diaza-18-crown 6-ether conjugate. *RSC Adv.* **2014**, *4*, 2563–2567.
- [3] Aragoni, M. C.; Arca, M.; Bencini, A.; Blake, A. J.; Caltagirone, C.; Decortes, A.; Demartin, F. Coordination chemistry of *N*-amino-propyl pendant arm derivatives of mixed *N*/*S*-, and *N*/*S*/*O*-donor macrocycles, and construction of selective fluorimetric chemosensors for heavy metal ions. *J. Chem. Soc., Dalton Trans.*, **2005**, *21*, 2994–3004.




Figure 3 A 1:1 stoichiometry of the host-guest relationship realized from the Job's plot between the probe **L** and the Ag⁺.

- [4] Prodi, L. Luminescent chemosensors: from molecules to nanoparticles. *New J. Chem.* **2005**, *29*, 20–31.
- [5] Fernandez, Y. D.; Gramatges, A. P.; Amendola, V.; Foti, F.; Mangano, C.; Pallavicini, P.; Patroni, S. Using micelles for a new approach to fluorescent sensors for metal cations. *Chem. Commun.* **2004**, *14*, 1650–1651.
- [6] Beer, P. D.; Gale, P. A. Anion recognition and sensing: the state of the art and future perspectives. *Angew. Chem. Int. Ed.* **2001**, *40*, 486–516.
- [7] Jose, D. A.; Kumar, D. K.; Kar, P.; Verma, S.; Ghosh, A.; Ganguly, B.; Ghosh, H. N.; Das, A. Role of positional isomers on receptor-anion binding and evidence for resonance energy transfer. *Tetrahedron* **2007**, *63*, 12007–12014.
- [8] Linder, M. C.; Hazegh-Azam, M. Copper biochemistry and molecular biology. *Tetrahedron* **1996**, *63*, 797s–811s.
- [9] Moon, K. S.; Singh, N.; Lee, G. W.; Jang, D. O. Colorimetric anion chemosensor based on 2-aminobenzimidazole: naked-eye detection of biologically important anions. *Tetrahedron* **2007**, *63*, 9106–9111.
- [10] Gale, P. A. Anion receptor chemistry. *Chem. Commun.* **2011**, *47*, 82–86.
- [11] Duan, Y. W.; Tang, H. Y.; Guo, Y.; Song, Z. K.; Peng, M. J.; Yan, Y. The synthesis and study of the fluorescent probe for sensing Cu²⁺ based on a novel coumarin Schiff-base. *Chin. Chem. Lett.* **2014**, *25*, 1082–1086.
- [12] Tang, L. J.; Guo, J. J.; Cao, Y. H.; Zhao, N. New application of a known molecule: rhodamine B 8-hydroxy-2-quinolinecarboxaldehyde Schiff base as a colorimetric and fluorescent “off-on” probe for copper (II). *J. Fluoresc.* **2012**, *22*, 1603–1608.
- [13] Yuan, L.; Lin, W. Y.; Yang, Y. T.; Song, J. Z.; Wang, J. L. Rational design of a highly reactive ratiometric fluorescent probe for cyanide. *Org. Lett.* **2011**, *13*, 3730–3733.
- [14] Padalkar, V. S.; Tathe, A.; V.D. Gupta, V. D.; Patil, V. S.; Phatangare, K.; Sekar, N. Synthesis and photo-physical characteristics of ESIPT inspired 2-Substituted benzimidazole, benzoxazole and benzothiazole fluorescent derivatives. *J. Fluoresc.* **2012**, *22*, 311–322.
- [15] Joo, T. Y.; Singh, N.; Lee, G. W.; Jang, D. O. Benzimidazole-based ratiometric fluorescent receptor for selective recognition of acetate. *Tetrahedron Lett.* **2007**, *48*, 8846–8850.
- [16] Zhang, T. T.; Chen, X. P.; Liu, J. T.; Zhang, L. Z.; Chu, J. M.; Su, L.; Zhao, B. X. A high sensitive fluorescence turn-on probe for imaging Zn²⁺ in aqueous solution and living cells. *RSC Adv.* **2014**, *4*, 16973–16978.
- [17] Park, S.; Kwon, J. E.; Kim, S. H.; Seo, J.; Chung, K.; Park, S. Y.; Jang, D. J.; Medina, B. M.; Gierschner, J.; Park, S. Y. A white-light-emitting molecule: frustrated energy transfer between constituent emitting centers. *J. Am. Chem. Soc.* **2009**, *131*, 14043–14049.
- [18] Satapathy, R.; Wu, Y. H.; H.C. Lin, H. C. Novel thieno-imidazole based probe for colorimetric detection of Hg²⁺ and fluorescence turn-on response of Zn²⁺. *Org. Lett.* **2012**, *14*, 2564–2567.
- [19] Zhao, B.; Y.Y. Ruan, Y. Y.; Ma, M. J.; Deng, Q. G.; Wang, L. Y.; Feng, Y. Q.; Gao, Y. One-pot synthesis of 11,23-bis(imidazol-1-yl)calyx[4]arene derivatives based on the upper rim of bisaminocalix[4]arene. *Heterocycles* **2013**, *87*, 1917–1924.
- [20] Meng, X. M.; Zhu, M. Z.; Liu, L.; Guo, Q. X. Novel highly selective fluorescent chemosensors for Zn(II). *Tetrahedron Lett.* **2006**, *47*, 1559–1562.