

Vidya S. Dofe*, Aniket P. Sarkate, Zarina M. Shaikh and Charansingh H. Gill

Ultrasound-assisted synthesis and antimicrobial activity of tetrazole-based pyrazole and pyrimidine derivatives

<https://doi.org/10.1515/hc-2017-0067>

Received April 16, 2017; accepted September 18, 2017; previously published online December 21, 2017

Abstract: New tetrazole-based pyrazole and pyrimidine derivatives were synthesized by an ultrasound irradiation method. All compounds were characterized by infrared spectroscopy (IR), ^1H nuclear magnetic resonance (NMR), ^{13}C NMR, mass spectrometry (MS) and elemental analysis and assessed *in vitro* for their efficacy as antimicrobial agents against four bacteria (*Staphylococcus aureus*, *Bacillus subtilis*, *Escherichia coli*, *Pseudomonas aeruginosa*) and two fungi (*Candida albicans*, *Aspergillus niger*). Compounds **8a**, **8e**, **9a**, **9b** and **9e** show potent activity against the tested strains compared to the reference drugs chloramphenicol and clotrimazole.

Keywords: antimicrobial; pyrazole; pyrimidine; tetrazole; ultrasound.

Introduction

Ultrasound-accelerated chemical reactions proceed via the formation and adiabatic collapse of transient cavitation bubbles. Ultrasound irradiation has become an alternative energy source for organic reactions ordinarily accomplished by heating. Many reactions can be conducted smoothly by sonication to provide improved yields and increased selectivity [1].

Life-threatening invasive infections caused by microorganisms have increased to alarming levels all over the world [2]. In the past decades, antibiotics have been used as antimicrobial drugs. However, many human pathogens no longer

respond to these antibiotics. Therefore, it is necessary to find new and efficacious drugs to treat these problems. The substitution of fluorine in potential drug molecules can improve their therapeutic efficacy through hydrogen bonding interactions at the active sites of the enzyme [3]. Thus, fluorine substitution remains an important aspect in the development of more active and selective drug candidates.

The synthetic versatility of tetrazole is due to its widespread applications in the field of medicinal chemistry. Tetrazole is considered a carboxylic acid analog because of their similar $\text{p}K_{\text{a}}$ values and planar delocalized systems. An advantage of tetrazole derivatives over carboxylic acids is that they are resistant to various biological degradation processes contributing to longer bioavailability of drugs [4, 5]. These factors, prime in the potential applications of tetrazole derivatives, are responsible for anticancer [6], antifungal [7], antitubercular [8], anti-HIV [9], antioxidant [10] and hormonal properties of the bioactive agents [11]. Pyrazole derivatives have showed significant biological activities, such as antimicrobial [12], analgesic [13], anti-inflammatory [14] and anticancer activities [15]. Pyrimidines also occupy a distinct and unique place in medicinal chemistry [16]. Pyrimidine derivatives are of interest due to their pharmacological properties such as antitumor [17], antimalarial [18], antiviral [19], antifungal [20], antibacterial [21], anti-inflammatory [22], analgesic [23] and antiprotozoal [24] activities. In continuation of our earlier work on design and synthesis of pharmacologically significant heterocycles [25], herein we report an efficient green synthesis and antimicrobial activity of new tetrazole containing pyrazole and pyrimidine derivatives.

Results and discussion

Chemistry


New tetrazole-based pyrazole derivatives **8a–f** and tetrazole-based pyrimidine derivatives **9a–f** were synthesized using conventional heating and ultrasound irradiation methods (Scheme 1). The precursors **4a–f** were prepared in good yield according to the literature

*Corresponding author: Vidya S. Dofe, Department of Chemistry, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad 431 004, India, e-mail: vidya.dofe84@gmail.com

Aniket P. Sarkate: Department of Chemical Technology, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad 431 004, India

Zarina M. Shaikh: Department of Microbiology, Maulana Azad College of Arts, Science and Commerce, Aurangabad 431 001, India

Charansingh H. Gill: Department of Chemistry, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad 431 004, India



Scheme 1 Reagents and conditions: (i) acetic anhydride, pyridine, 100°C, 3–4 h; (ii) AlCl₃, 150°C, 3–4 h; (iii) 4-fluorobenzaldehyde, KOH, EtOH, room temperature (rt), 2–3 h; (iv) H₂O₂, NaOH, 0°C to rt, 2–3 h; (v) 2-chloroacetonitrile, K₂CO₃, DMF, rt, 3–4 h; (vi) sodium azide, zinc bromide, H₂O, reflux, 100°C, 4–5 h; (vii) hydrazine hydrate, ethanol, ()), 65°C, 45–55 min; (viii) thiourea, KOH, ethanol, ()), 65°C, 15–25 min.

procedure [26, 27]. Compounds **4a–f** were converted into corresponding substituted 3-hydroxychromones **5a–f** by oxidative cyclization using hydrogen peroxide in ethanol. The hydroxychromones **5a–f** were alkylated with 2-chloroacetonitrile in the presence of K₂CO₃ in *N,N'*-dimethylformamide (DMF) at room temperature to afford the substituted 2-(2-(4-fluorophenyl)-4-oxo-4*H*-chromen-3-yloxy)acetonitriles **6a–f**. The reaction of **6a–f** with sodium azide and zinc bromide in water at 100°C furnished 3-[(1*H*-tetrazol-5-yl)methoxy]-2-(4-fluorophenyl)-4*H*-chromen-4-one derivatives **7a–f**. Treatment of **7a–f** with hydrazine hydrate in ethanol under ultrasound irradiation yielded the desired pyrazoles **8a–f**. In addition, treatment of compounds **7a–f** with thiourea in ethanolic KOH under ultrasound irradiation furnished the desired pyrimidines **9a–f**. All synthesized compounds were characterized by ¹H nuclear magnetic resonance (NMR), ¹³C NMR, mass spectrometry (MS) and elemental analysis.

Antimicrobial activity

All compounds were evaluated for antibacterial activity against Gram-positive bacteria *Staphylococcus aureus* (NCIM 2079) and *Bacillus subtilis* (NCIM 2920) and Gram-negative bacteria *Escherichia coli* (NCIM 2065) and *Pseudomonas aeruginosa* (NCIM 2200) and for antifungal

activity against *Candida albicans* (NCIM 3471) and *Aspergillus niger* (NCIM 596). The minimum inhibitory concentration (MIC) values were determined by the micro-broth dilution method. The antibacterial drug chloramphenicol and antifungal drug clotrimazole were used as reference antibiotics. The results are shown in Table 1. Of interest are high activities of compounds **9a** and **9e** against *A. niger*. Compound **9e** is also active against *C. albicans*. The activities of **8e**, **9a** and **9b** against *S. aureus* exceed those of the reference drug chloramphenicol.

Table 1 Antimicrobial screening results of bioactive compounds **8** and **9** using the micro-broth dilution method.

No.	Minimum inhibitory concentration (µg/mL)					
	Bacteria				Fungi	
	<i>Sa</i>	<i>Bs</i>	<i>Ec</i>	<i>Pa</i>	<i>Ca</i>	<i>An</i>
8e	25	200	50	200	75	50
9a	25	200	175	150	75	25
9b	25	75	125	100	100	50
9e	50	50	75	75	50	12.5
Chloramphenicol	50	25	50	50	–	–
Clotrimazole	–	–	–	–	50	25

Sa, *Staphylococcus aureus*; *Bs*, *Bacillus subtilis*; *Ec*, *Escherichia coli*; *Pa*, *Pseudomonas aeruginosa*; *Ca*, *Candida albicans*; *An*, *Aspergillus niger*.

Conclusion

Newly synthesized compounds **8e**, **9a**, **9b** and **9e** are highly potent antimicrobial agents.

Experimental

The progress of each reaction was monitored by thin-layer chromatography (TLC) using Merck silica gel 60 F₂₅₄ plates and visualized with ultraviolet (UV) light and iodine. Melting points were determined in open capillaries and are uncorrected. Infrared spectra were recorded on a Carry 600 Series Fourier-transform infrared spectroscopy (FT-IR) spectrophotometer using KBr pellets. ¹H NMR (400 MHz) and ¹³C NMR (100 MHz) spectra were recorded on a Bruker AVANCE II 400 NMR spectrometer in CDCl₃ or DMSO-*d*₆. Mass spectra were recorded on a Waters quadrupole time-of-flight (Q-TOF) micro instrument equipped with an electrospray ionization (ESI) source. Elemental analysis was performed on a Perkin-Elmer EAL-240 elemental analyzer.

General procedure for synthesis of compounds 5a–f

A mixture of (*E*)-3-(4-fluorophenyl)-1-(2-hydroxyphenyl)prop-2-en-1-one (**4a–f**, 1 mmol), ethanol (15 mL), NaOH (10%, 5 mL) and hydrogen peroxide (30%, 1.5 mL) was stirred vigorously for 30 min and then kept for 2–3 h in ice. The progress of the reaction was monitored by TLC using ethyl acetate/hexane as an eluent. After completion of the reaction, the mixture was poured into ice water and acidified with 1 M hydrochloric acid (HCl). The precipitate was collected by filtration, washed with water and crystallized from chloroform/ethanol to afford a pure product.

2-(4-Fluorophenyl)-3-hydroxy-4H-chromen-4-one (5a) This compound was synthesized from (*E*)-3-(4-fluorophenyl)-1-(2-hydroxyphenyl)prop-2-en-1-one (**4a**); a yellow solid; yield 74%; mp 152–154°C; IR: 3297 (OH), 2925 (ArH), 1618 (C=O, pyrone), 1572 cm⁻¹ (C=C); ¹H NMR (CDCl₃): δ 7.11 (br s, 1H, ArH), 7.22 (t, 2H, *J* = 8.5 Hz, ArH), 7.42 (m 1H, ArH), 7.58 (d, 1H, *J* = 8.5 Hz, ArH), 7.71 (m, 1H, ArH), 8.24 (d, 1H, *J* = 1.8 Hz, ArH), 8.26 (s, 1H, OH), 8.27–8.30 (m, 1H, ArH); ¹³C NMR (CDCl₃): δ 115.7, 115.9, 118.2, 120.7, 124.6, 125.5, 127.3, 129.9, 133.7, 138.2, 144.1, 155.3, 162.3, 164.8, 173.4. ESI-MS. Calcd for C₁₅H₉FO₃ (M+H)⁺: *m/z* 257.0. Found: *m/z* 257.0.

6-Chloro-2-(4-fluorophenyl)-3-hydroxy-4H-chromen-4-one (5b) This compound was synthesized from (*E*)-1-(5-chloro-2-hydroxyphenyl)-3-(4-fluorophenyl)prop-2-en-1-one (**4b**); a yellow solid; yield 77%; mp 213–215°C; ¹H NMR (DMSO-*d*₆): δ 7.08–7.25 (m, 2H, ArH), 7.68–7.82 (m, 2H, ArH), 8.12–8.23 (m, 2H, ArH), 8.51 (s, 1H, ArH), 9.08 (br s, 1H, OH); ¹³C NMR (DMSO-*d*₆): δ 114.1, 117.5, 120.6, 125.4, 127.4, 128.4, 129.7, 130.2, 133.6, 148.5, 155.4, 157.6, 160.8, 162.5, 172.9. ESI-MS. Calcd for C₁₅H₈ClFO₃ (M+H)⁺: *m/z* 291.0224. Found: *m/z* 291.2223.

2-(4-Fluorophenyl)-3-hydroxy-8-methyl-4H-chromen-4-one (5c) This compound was synthesized from (*E*)-3-(4-fluorophenyl)-1-(2-hydroxy-3-methylphenyl)prop-2-en-1-one (**4c**); a yellow solid, yield 69%; mp 160–162°C; ¹H NMR (DMSO-*d*₆): δ 2.54 (s, 3H, CH₃),

7.24–7.29 (m, 3H, ArH), 7.53 (d, 1H, *J* = 7 Hz, ArH), 7.93 (d, 1H, *J* = 7.7 Hz, ArH), 8.26 (m, 2H, ArH), 9.38 (s, 1H, OH); ¹³C NMR (DMSO-*d*₆): δ 15.3, 115.1, 115.3, 121, 122.3, 123.6, 127.1, 127.9, 129.6, 133.7, 138.7, 143.5, 152.8, 161.3, 163.7, 173. ESI-MS. Calcd for C₁₆H₁₁FO₃ (M+H)⁺: *m/z* 271.077. Found: *m/z* 271.2154.

2-(4-Fluorophenyl)-3-hydroxy-6-methyl-4H-chromen-4-one (5d) This compound was synthesized from (*E*)-3-(4-fluorophenyl)-1-(2-hydroxy-5-methylphenyl)prop-2-en-1-one (**4d**); a yellow solid, yield 73%; mp 171–173°C; ¹H NMR (DMSO-*d*₆): δ 2.46 (s, 3H, CH₃), 7.27 (t, 2H, *J* = 8.6 Hz, ArH), 7.54 (s, 2H, ArH), 7.91 (s, 1H, ArH), 8.27–8.30 (m, 2H, ArH), 9.31 (s, 1H, OH); ¹³C NMR (DMSO-*d*₆): δ 20.5, 115, 115.2, 117.7, 120.9, 123.8, 127.7, 129.8, 130.6, 134.5, 138.7, 143.9, 152.8, 161.3, 163.8, 172.8. ESI-MS. Calcd for C₁₆H₁₁FO₃ (M+H)⁺: *m/z* 271.0770. Found: *m/z* 271.0234.

6,8-Dichloro-2-(4-fluorophenyl)-3-hydroxy-4H-chromen-4-one (5e) This compound was synthesized from (*E*)-1-(3,5-dichloro-2-hydroxyphenyl)-3-(4-fluorophenyl)prop-2-en-1-one (**4e**); a yellow solid; yield 75%; mp 203–205°C; ¹H NMR (DMSO-*d*₆): δ 7.25 (t, 2H, *J* = 8.8 Hz, ArH), 7.82 (d, 1H, *J* = 2.6 Hz, ArH), 7.96 (d, 1H, *J* = 2.2 Hz, ArH), 8.29 (m, 2H, ArH), 9.81 (s, 1H, OH); ¹³C NMR (DMSO-*d*₆): δ 115.2, 115.4, 122.8, 123, 123.7, 127.1, 128.8, 129.8, 132.6, 139.2, 144.5, 148.4, 161.5, 164, 171.3. ESI-MS. Calcd for C₁₅H₇Cl₂FO₃ (M+H)⁺: *m/z* 324.9835. Found: *m/z* 325.0267.

6-Chloro-2-(4-fluorophenyl)-3-hydroxy-7-methyl-4H-chromen-4-one (5f) This compound was synthesized from (*E*)-1-(5-chloro-2-hydroxy-4-methylphenyl)-3-(4-fluorophenyl)prop-2-en-1-one (**4f**); a yellow solid, yield 79%; mp 222–224°C; ¹H NMR (DMSO-*d*₆): δ 2.5 (s, 3H, CH₃), 7.28 (m, 2H, ArH), 7.64 (br s, 1H, ArH), 8.03 (s, 1H, ArH), 8.27 (br s, 2H, ArH), 9.54 (s, 1H, OH); ¹³C NMR (DMSO-*d*₆): δ 20.3, 115.1, 115.3, 120.1, 120.5, 124, 127.5, 129.9, 130.1, 138.8, 141.8, 144.3, 152.8, 161.4, 163.9, 171.8. ESI-MS. Calcd for C₁₆H₁₀ClFO₃ (M+H)⁺: *m/z* 305.0381. Found: *m/z* 305.0251.

General procedure for synthesis of compounds 6a–f

To a stirred solution of **5a–f** (1 mmol) in DMF was added potassium carbonate (2 mmol) and portion-wise 2-chloroacetonitrile (1 mmol) at room temperature for about 3–4 h. The mixture was quenched with crushed ice and the resultant precipitate was filtered and crystallized from ethanol.

2-(2-(4-Fluorophenyl)-4-oxo-4H-chromen-3-yloxy)acetonitrile (6a) This compound was prepared from **5a**; a white solid; yield 78%; mp 140–142°C; IR: ν 3012 (Ar-H), 2965 (C-H), 2346 (C≡N), 1634 (C=O), 1604 (C=C), 1238 (C-F) cm⁻¹; ¹H NMR (CDCl₃): δ 5.11 (s, 2H, OCH₂), 7.20–2.27 (m, 2H, ArH), 7.45 (m, 1H, ArH), 7.57 (dd, 1H, *J* = 1.1, 8.6 Hz, ArH), 7.74 (ddt, 1H, *J* = 1.5, 7.8, 8.0 Hz, ArH), 8.08 (m, 2H, ArH), 8.25 (dd, 1H, *J* = 1.5, 7.8 Hz, ArH); ¹³C NMR (CDCl₃): δ 56.1, 115.1, 115.9, 116.1, 118.1, 123.8, 125.4, 125.8, 126.1, 131.3, 134.2, 137.5, 155.3, 156.4, 163.1, 165.6, 174.1. Anal. Calcd for C₁₇H₁₀FNO₃: C, 69.15; H, 3.41; N, 4.74. Found: C, 69.07; H, 3.35; N, 4.68.

2-(6-Chloro-2-(4-fluorophenyl)-4-oxo-4H-chromen-3-yloxy)acetonitrile (6b) This compound was prepared from **5b**; a white solid, yield 86%; mp 168–170°C; IR: ν 3069 (Ar-H), 2952 (C-H), 2360 (C≡N), 1628 (C=O), 1600 (C=C), 1144 (C-F) 755 (C-Cl) cm⁻¹; ¹H NMR (DMSO-*d*₆):

δ 5.14 (s, 2H, OCH₂), 7.34 (t, 2H, $J=8.0$ Hz, ArH), 7.73 (d, 1H, $J=8.0$ Hz, ArH), 7.79 (dd, 1H, $J=2.6, 9.2$ Hz, ArH), 8.07 (d, 1H, $J=2.6$ Hz, ArH), 8.13 (m, 2H, ArH); ¹³C NMR (DMSO-*d*₆): δ 56.4, 115.5, 115.7, 120.4, 124, 124.3, 125.7, 130.3, 131.1, 131.2, 134.1, 137.6, 153.1, 155.4, 162.4, 165, 172.1. Anal. Calcd for C₁₇H₉ClFNO₃: C, 61.93; H, 2.75; N, 4.25. Found: C, 61.98; H, 2.82; N, 4.33.

2-(2-(4-Fluorophenyl)-8-methyl-4-oxo-4H-chromen-3-yloxy) acetonitrile (6c) This compound was prepared from **5c**; a white solid; yield 71%; mp 176–178°C; IR: ν 3032 (Ar-H), 2915 (C-H), 2360 (C≡N), 1633 (C=O), 1604 (C=C), 1193 (C-F) cm⁻¹; ¹H NMR (CDCl₃): δ 2.58 (s, 3H, CH₃), 5.12 (s, 2H, OCH₂), 7.23–7.27 (m, 2H, ArH), 7.34 (t, 1H, $J=7.0$ Hz, ArH), 7.57 (d, 1H, $J=7.0$ Hz, ArH), 8.07–8.13 (m, 3H, ArH); ¹³C NMR (CDCl₃): δ 15.8, 56, 115.1, 115.9, 116.2, 123.4, 123.8, 125.1, 126.5, 127.6, 131.3, 135, 137.4, 153.8, 155.8, 163.1, 165.6, 174.4. Anal. Calcd for C₁₈H₁₂FNO₃: C, 69.90; H, 3.91; N, 4.53. Found: C, 69.96; H, 3.94; N, 4.56.

2-(2-(4-Fluorophenyl)-6-methyl-4-oxo-4H-chromen-3-yloxy) acetonitrile (6d) This compound was prepared from **5d**; a white solid; yield 74%; mp 162–164°C; IR: ν 3087 (Ar-H), 2974 (C-H), 2369 (C≡N), 1632 (C=O), 1603 (C=C), 1187 (C-F) cm⁻¹; ¹H NMR (DMSO-*d*₆): δ 2.53 (s, 3H, CH₃), 5.15 (s, 2H, OCH₂), 7.36 (s, 2H, ArH), 7.61 (s, 2H, ArH), 7.90 (s, 1H, ArH), 8.12 (s, 2H, ArH); ¹³C NMR (DMSO-*d*₆): δ 20.5, 56.4, 115.5, 115.7, 118, 122.9, 124.1, 126.2, 131.1, 131.1, 134.8, 135.4, 136.3, 137.5, 153, 154.9, 162.3, 173.1. Anal. Calcd for C₁₈H₁₂FNO₃: C, 69.90; H, 3.91; N, 4.53. Found: C, 69.82; H, 3.95; N, 4.66.

2-(6,8-Dichloro-2-(4-fluorophenyl)-4-oxo-4H-chromen-3-yloxy) acetonitrile (6e) This compound was prepared from **5e**; a white solid; yield 77%; mp 177–179°C; IR: ν 3064 (Ar-H), 2916 (C-H), 2361 (C≡N), 1625 (C=O), 1601 (C=C), 1180 (C-F) 717 (C-Cl) cm⁻¹; ¹H NMR (CDCl₃): δ 5.12 (s, 2H, OCH₂), 7.26 (t, 2H, $J=8.6$ Hz, ArH), 7.77 (d, 1H, $J=2.2$ Hz, ArH), 8.10 (d, 1H, $J=2.2$ Hz, ArH), 8.18 (m, 2H, ArH); ¹³C NMR (CDCl₃): δ 56.1, 114.8, 116.2, 116.4, 123.8, 124.6, 125.5, 131.2, 131.5, 131.6, 134.2, 137.6, 149.4, 156.3, 163.5, 166, 172.5. Anal. Calcd for C₁₇H₈Cl₂FNO₃: C, 56.07; H, 2.21; N, 3.85. Found: C, 56.18; H, 2.24; N, 3.79.

2-(6-Chloro-2-(4-fluorophenyl)-7-methyl-4-oxo-4H-chromen-3-yloxy)acetonitrile (6f) This compound was prepared from **5f**; a white solid, yield 82%; mp 170–172°C; IR: ν 3013 (Ar-H), 2915 (C-H), 2360 (C≡N), 1599 (C=O), 1552 (C=C), 1203 (C-F) 750 (C-Cl) cm⁻¹; ¹H NMR (DMSO-*d*₆): 2.54 (s, 3H, CH₃), 5.14 (s, 2H, OCH₂), 7.35 (s, 2H, ArH), 7.70 (s, 1H, ArH), 8.04 (s, 1H, ArH), 8.12 (s, 2H, ArH); ¹³C NMR (DMSO-*d*₆): δ 20.3, 56.4, 115.5, 115.6, 115.7, 120.3, 122.4, 124.2, 125.9, 125.9, 131.1, 131.1, 137.5, 142.9, 153.1, 155.1, 167.4, 172.1. Anal. Calcd for C₁₈H₁₁ClFNO₃: C, 62.89; H, 3.23; N, 4.07. Found: C, 62.93; H, 3.28; N, 4.11.

General procedure for synthesis of compounds 7a–f

To a mixture of sodium azide (1.5 mmol) and zinc bromide (1.5 mmol) in water (20 mL) was added compound **6a–f** (1 mmol). The mixture was then heated at reflux for 4–5 h with vigorous stirring. After completion of the reaction, as monitored by TLC using chloroform/ methanol as an eluent, the mixture was quenched with crushed ice, and the resultant solid was filtered and crystallized from ethanol.

3-[(1H-Tetrazol-5-yl)methoxy]-2-(4-fluorophenyl)-4H-chromen-4-one (7a) This compound was prepared from **6a**; a white solid, yield: 81%; mp 200–202°C; IR: ν 3444 (N-H), 3030 (Ar-H), 2917 (C-H),

1606 (C=O), 1554 (C=C), 1238 (C-F) cm⁻¹; ¹H NMR (DMSO-*d*₆): δ 5.47 (s, 2H, OCH₂), 7.26 (t, 2H, $J=9.0$ Hz, ArH), 7.48–7.52 (m, 1H, ArH), 7.70 (d, 1H, $J=8.0$ Hz, ArH), 7.82 (ddd, 1H, $J=8.5, 7.1, 1.7$ Hz, ArH), 8.01–8.05 (m, 2H, ArH), 8.16 (dd, 1H, $J=8.0, 1.7$ Hz, ArH); ¹³C NMR (DMSO-*d*₆): δ 61.8 (O-CH₂), 115.3, 115.6, 118.3, 123.4, 124.9, 125.1, 126.2, 130.9, 131, 134.1, 138.2, 154.7, 154.9, 162 (tetrazole C), 164.5 (C-F), 173.5 (C=O). Anal. Calcd for C₁₇H₁₁FN₄O₃: C, 60.36; H, 3.28; N, 16.56. Found: C, 60.29; H, 3.25; N, 16.51.

3-[(1H-Tetrazol-5-yl)methoxy]-6-chloro-2-(4-fluorophenyl)-4H-chromen-4-one (7b) This compound was prepared from **6b**; a white solid; yield 85%; mp 203–205°C; IR: ν 3435 (N-H), 3035 (Ar-H), 2992 (C-H), 1622 (C=O), 1605 (C=C), 1197 (C-F), 718 (C-Cl) cm⁻¹; ¹H NMR (DMSO-*d*₆): δ 5.42 (s, 2H, OCH₂), 7.17 (t, 2H, $J=9.0$ Hz, ArH), 7.64 (d, 1H, $J=9.0$ Hz, ArH), 7.72 (dd, 1H, $J=9.0, 2.6$ Hz, ArH), 7.98 (m, 2H, ArH), 8.04 (d, 1H, $J=2.6$ Hz, ArH); ¹³C NMR (DMSO-*d*₆): δ 61.7 (O-CH₂), 115.4, 115.6, 120.7, 123.8, 124.5, 125.9, 125.9, 129.9, 131, 134, 138.2, 153.2, 155.2, 162.1 (tetrazole C), 164.6 (C-F), 172.5 (C=O). Anal. Calcd for C₁₇H₁₀ClFN₄O₃: C, 54.78; H, 2.70; N, 15.03. Found: C, 54.84; H, 2.72; N, 15.23.

3-[(1H-Tetrazol-5-yl)methoxy]-2-(4-fluorophenyl)-8-methyl-4H-chromen-4-one (7c) This compound was prepared from **6c**; a white solid; yield 73%; mp 198–200°C; IR: ν 3435 (N-H), 3035 (Ar-H), 2992 (C-H), 1622 (C=O), 1605 (C=C), 1197 cm⁻¹ (C-F); ¹H NMR (DMSO-*d*₆): δ 2.33 (s, 3H, CH₃), 5.61 (s, 2H, OCH₂), 6.97 (s, 1H, ArH), 7.10–7.22 (m, 2H, ArH), 7.53 (s, 1H, ArH), 7.69–7.80 (m, 3H, ArH), 8.41 (s, 1H, NH); ¹³C NMR (DMSO-*d*₆): δ 15.1 (CH₃), 63.5 (O-CH₂), 114.9, 115.2, 122.3, 123, 124.3, 126.6, 127.1, 128.2, 130.4, 134.2, 139, 149.5, 152.6, 161.5 (tetrazole C), 164 (C-F), 173.8 (C=O). Anal. Calcd for C₁₈H₁₃FN₄O₃: C, 61.36; H, 3.72; N, 15.90. Found: C, 61.45; H, 3.69; N, 15.98.

3-[(1H-Tetrazol-5-yl)methoxy]-2-(4-fluorophenyl)-6-methyl-4H-chromen-4-one (7d) This compound was prepared from **6d**; a white solid; yield 75%; mp 202–204°C; IR: ν 3423 (N-H), 3021 (Ar-H), 2918 (C-H), 1602 (C=O), 1554 (C=C), 1174 cm⁻¹ (C-F); ¹H NMR (DMSO-*d*₆): δ 2.46 (s, 3H, CH₃), 5.47 (s, 2H, OCH₂), 7.27 (t, 2H, $J=9.0$ Hz, ArH), 7.57–7.63 (m, 2H, ArH), 7.90 (s, 1H, ArH), 8.00 (dd, 2H, $J=9.0, 5.3$ Hz, ArH); ¹³C NMR (DMSO-*d*₆): δ 20.4 (CH₃), 61.8 (O-CH₂), 115.3, 115.5, 118, 123.1, 124.1, 126.3, 126.3, 130.9, 134.7, 135.2, 138.2, 153, 154.7, 162 (tetrazole C), 164.5 (C-F), 173.4 (C=O). Anal. Calcd for C₁₈H₁₃FN₄O₃: C, 61.36; H, 3.72; N, 15.90. Found: C, 61.34; H, 3.65; N, 15.81.

3-[(1H-Tetrazol-5-yl)methoxy]-6,8-dichloro-2-(4-fluorophenyl)-4H-chromen-4-one (7e) This compound was prepared from **6e**; a white solid; yield 80%; mp 194–196°C; IR: ν 3423 (N-H), 3040 (Ar-H), 2917 (C-H), 1660 (C=O), 1603 (C=C), 1158 (C-F), 760 cm⁻¹ (C-Cl); ¹H NMR (DMSO-*d*₆): δ 5.50 (s, 2H, OCH₂), 7.24 (t, 2H, $J=8.4$ Hz, ArH), 7.33–7.37 (m, 1H, ArH), 8.00 (d, 2H, $J=5.5$ Hz, ArH), 8.08 (dd, 1H, $J=8.4, 5.5$ Hz, ArH); ¹³C NMR (DMSO-*d*₆): δ 62 (O-CH₂), 115.6, 115.8, 123, 123.8, 125.2, 129.6, 131, 133.5, 138.6, 139.4, 148.9, 154.6, 162.2 (tetrazole C), 164.7 (C-F), 172 (C=O). Anal. Calcd for C₁₇H₉Cl₂FN₄O₃: C, 50.15; H, 2.23; N, 13.76. Found: C, 50.33; H, 2.32; N, 13.87.

3-[(1H-Tetrazol-5-yl)methoxy]-6-chloro-2-(4-fluorophenyl)-7-methyl-4H-chromen-4-one (7f) This compound was prepared from **6f**; a white solid; yield 78%; mp 178–180°C; IR: ν 3495 (N-H), 3077 (Ar-H), 2924 (C-H), 1638 (C=O), 1618 (C=C), 1166 (C-F) 771 cm⁻¹ (C-Cl); ¹H NMR (DMSO-*d*₆): δ 2.44 (s, 3H, CH₃), 5.45 (s, 2H, OCH₂), 7.30 (t, 2H, $J=9.0$ Hz, ArH), 7.73–7.76 (m, 1H, ArH), 7.92–7.96 (m, 3H, ArH);

^{13}C NMR (DMSO- d_6): δ 20.1 (CH_3), 61.8 ($\text{O}-\text{CH}_2$), 115.4, 115.7, 120.6, 122.5, 123.9, 126.1, 130.6, 130.9, 131, 138.1, 142.7, 153, 154.9, 162 (tetrazole C), 164.5 (C-F), 172.4 (C=O). Anal. Calcd for $\text{C}_{18}\text{H}_{12}\text{ClFN}_4\text{O}_3$: C, 55.90; H, 3.13; N, 14.49. Found: C, 55.97; H, 3.21; N, 14.58.

General procedure for synthesis of compounds 8a–f

Conventional method Hydrazine hydrate (1 mmol) was added to a solution of **7a–f** (1 mmol) in ethanol (10 mL) and the mixture was heated for 5–6 h. After completion of the reaction, as monitored by TLC, 10 mL of water was added, and the resultant precipitate was filtered, dried under reduced pressure and crystallized from ethanol.

Ultrasound-assisted method Hydrazine hydrate (1 mmol) was added to a solution of **7a–f** (1 mmol) in ethanol (10 mL) and the mixture was ultrasonicated for 45–55 min at 65°C. After completion of the reaction, as monitored by TLC, 10 mL of water was added, and the resultant precipitate was filtered, dried and crystallized from ethanol.

2-[4-((1H-Tetrazol-5-yl)methoxy)-5-(4-fluorophenyl)-1H-pyrazol-3-yl]phenol (8a) This compound was prepared from **7a**; a yellow solid; conventional method time 340 min, yield 70%; ultrasound-assisted method time 51 min, yield 88%; mp 269–271°C; IR: ν 3943 (OH), 3352 (NH), 1884 (C=N), 1246 (C-F) cm^{-1} ; ^1H NMR (DMSO- d_6): δ 5.08 (s, 2H, OCH_2), 6.89 (t, 1H, $J=7.3$ Hz, ArH), 6.96 (d, 1H, $J=8.1$ Hz, ArH), 7.19–7.23 (m, 3H, ArH), 7.63–7.88 (m, 3H, ArH), 10.80 (br s, 1H, OH), 12.69 (br s, 1H, NH), 13.54 (br s, 1H, NH); ^{13}C NMR (DMSO- d_6): δ 64.0, 99.5, 115.4, 115.6, 116.1, 119.2, 119.3, 127.7, 127.8, 129.0, 129.1, 136.5, 153.0, 153.1, 154.9, 160.5, 162.9. Anal. Calcd for $\text{C}_{17}\text{H}_{13}\text{FN}_6\text{O}_2$: C, 57.95; H, 3.72; N, 23.85. Found: C, 57.92; H, 3.69; N, 23.76.

2-[4-((1H-Tetrazol-5-yl)methoxy)-5-(4-fluorophenyl)-1H-pyrazol-3-yl]-4-chlorophenol (8b) This compound was prepared from **7b**; a yellow solid; conventional method time 315 min, yield 75%; ultrasound-assisted method time 45 min, yield 93%; mp 228–230°C; IR: ν 3905 (OH), 3387 (NH), 1852 (C=N), 1216 (C-F) cm^{-1} ; ^1H NMR (DMSO- d_6): δ 5.08 (s, 2H, OCH_2), 6.97 (d, 1H, $J=8.4$ Hz, ArH), 7.24–7.30 (m, 3H, ArH), 7.58–7.89 (m, 3H, ArH), 10.72 (br s, 1H, OH), 12.86 (s, 1H, NH), 13.68 (s, 1H, NH); ^{13}C NMR (DMSO- d_6): δ 64.3, 99.5, 115.7, 117.3, 117.6, 123.0, 126.5, 126.6, 127.7, 127.8, 128.5, 136.7, 153.4, 153.7, 153.8, 160.6, 163.0. Anal. Calcd for $\text{C}_{17}\text{H}_{12}\text{ClFN}_6\text{O}_2$: C, 52.79; H, 3.13; N, 21.73. Found: C, 52.63; H, 3.10; N, 21.68.

2-[4-((1H-Tetrazol-5-yl)methoxy)-5-(4-fluorophenyl)-1H-pyrazol-3-yl]-6-methylphenol (8c) This compound was prepared from **7c**; a yellow solid; conventional method time 350 min, yield 77%; ultrasound-assisted method time 54 min, yield 84%; mp 250–252°C; IR: ν 3918 (OH), 3346 (NH), 1815 (C=N), 1245 (C-F) cm^{-1} ; ^1H NMR (CDCl_3): δ 2.57 (s, 3H, CH_3), 5.14 (s, 2H, OCH_2), 6.91–6.98 (m, 2H, ArH), 7.36 (t, 1H, $J=7.7$ Hz, ArH), 7.59 (d, 1H, $J=7.7$ Hz, ArH), 7.65–7.69 (m, 3H, ArH), 10.79 (br s, 1H, OH), 12.80 (s, 1H, NH), 13.71 (s, 1H, NH); ^{13}C NMR (CDCl_3): δ 15.8, 63.9, 99.8, 115.1, 115.9, 116.2, 119.4, 119.5, 127.6, 127.7, 129.5, 129.7, 137.0, 153.4, 153.6, 155.3, 160.7, 163.1. Anal. Calcd for $\text{C}_{18}\text{H}_{15}\text{FN}_6\text{O}_2$: C, 59.01; H, 4.13; N, 22.94. Found: C, 59.13; H, 3.98; N, 22.81.

2-[4-((1H-Tetrazol-5-yl)methoxy)-5-(4-fluorophenyl)-1H-pyrazol-3-yl]-4-methylphenol (8d) This compound was prepared from **7d**; a yellow solid; conventional method time 320 min, yield 79%; ultrasound-assisted method time 48 min, yield 91%; mp 240–242°C; IR: ν

3955 (OH), 3346 (NH), 1865 (C=N), 1258 (C-F) cm^{-1} ; ^1H NMR (DMSO- d_6): δ 2.52 (s, 3H, CH_3), 5.19 (s, 2H, OCH_2), 6.35–6.42 (m, 2H, ArH), 6.92–6.94 (m, 2H, ArH), 7.64–7.89 (m, 3H, ArH), 10.84 (br s, 1H, OH), 12.78 (br s, 1H, NH), 13.73 (br s, 1H, NH); ^{13}C NMR (DMSO- d_6): δ 16.0, 64.2, 99.6, 115.5, 115.7, 116.5, 119.8, 119.9, 127.6, 127.7, 129.7, 129.8, 136.6, 154.2, 154.9, 155.8, 161.1, 163.6. Anal. Calcd for $\text{C}_{18}\text{H}_{15}\text{FN}_6\text{O}_2$: C, 59.01; H, 4.13; N, 22.94. Found: C, 58.96; H, 4.27; N, 22.98.

2-[4-((1H-Tetrazol-5-yl)methoxy)-5-(4-fluorophenyl)-1H-pyrazol-3-yl]-4,6-dichlorophenol (8e) This compound was prepared from **7e**; a yellow solid; conventional method time 330 min, yield 71%; ultrasound-assisted method time 45 min, yield 95%; mp 225–227°C; IR: ν 3936 (OH), 3386 (NH), 1855 (C=N), 1282 (C-F) cm^{-1} ; ^1H NMR (CDCl_3): δ 5.15 (s, 2H, OCH_2), 6.42 (t, 2H, $J=8.8$ Hz, ArH), 6.91 (d, 1H, $J=2.6$ Hz, ArH), 7.69 (d, 1H, $J=2.6$ Hz, ArH), 7.90 (dd, 2H, $J=8.8$, 5.2 Hz, ArH), 10.81 (br s, 1H, OH), 12.73 (br s, 1H, NH), 13.82 (br s, 1H, NH); ^{13}C NMR (CDCl_3): δ 64.6, 99.9, 115.3, 115.4, 116.8, 120.1, 120.2, 127.9, 128.0, 129.4, 129.6, 137.3, 154.7, 155.0, 156.2, 161.3, 163.8. Anal. Calcd for $\text{C}_{17}\text{H}_{11}\text{Cl}_2\text{FN}_6\text{O}_2$: C, 48.47; H, 2.63; N, 19.95. Found: C, 48.55; H, 2.61; N, 19.98.

2-[4-((1H-Tetrazol-5-yl)methoxy)-5-(4-fluorophenyl)-1H-pyrazol-3-yl]-4-chloro-5-methylphenol (8f) This compound was prepared from **7f**; a yellow solid; conventional method time 330 min, yield 72%; ultrasound-assisted method time 45 min, yield 92%; mp 278–280°C; IR: ν 3945 (OH), 3335 (NH), 1852 (C=N), 1273 (C-F) cm^{-1} ; ^1H NMR (DMSO- d_6): δ 2.52 (s, 3H, CH_3), 5.19 (s, 2H, OCH_2), 6.52 (s, 1H, ArH), 6.89 (s, 2H, ArH), 7.72–7.75 (m, 2H, ArH), 7.93 (s, 1H, ArH), 10.86 (br s, 1H, OH), 12.71 (br s, 1H, NH), 13.88 (br s, 1H, NH); ^{13}C NMR (DMSO- d_6): δ 16.4, 64.8, 99.4, 115.9, 116.0, 116.8, 119.6, 119.7, 127.9, 128.0, 129.5, 129.7, 136.8, 154.4, 155.1, 155.9, 160.8, 163.5. Anal. Calcd for $\text{C}_{18}\text{H}_{14}\text{ClFN}_6\text{O}_2$: C, 53.94; H, 3.52; N, 20.97. Found: C, 53.96; H, 3.50; N, 21.01.

General procedure for synthesis of compounds 9a–f

Conventional method To a mixture of compound **7a–f** (1 mmol) and KOH (1.5 mmol) in ethanol (10 mL), thiourea (1.5 mmol) was added and the mixture was heated for 3–4 h. After completion of the reaction, as monitored by TLC, the reaction mass was cooled to room temperature, poured on crushed ice and acidified with concentrated HCl. The resultant solid was filtered and crystallized from ethanol.

Ultrasound-assisted method To a mixture of compound **7a–f** (1 mmol) and KOH (1.5 mmol) in ethanol (10 mL), thiourea (1.5 mmol) was added and the reaction mass was ultrasonicated for 15–25 min at 65°C. After completion of the reaction, as monitored by TLC, the mixture was cooled to room temperature, poured on crushed ice and acidified with concentrated HCl. The resultant precipitate was filtered and crystallized from ethanol.

5-[(1H-Tetrazol-5-yl)methoxy]-6-(4-fluorophenyl)-4-(2-hydroxyphenyl)pyrimidine-2(1H)-thione (9a) This compound was prepared from **7a**; a yellow solid; conventional method time 230 min, yield 73%; ultrasound-assisted method time 21 min, yield 87%; mp 163–165°C; IR ν : 3919 (OH), 3423 (NH), 3313 (NH), 3040 (CH) cm^{-1} ; ^1H NMR (DMSO- d_6): δ 4.79 (s, 2H, OCH_2), 7.19 (t, 2H, $J=8.8$ Hz, ArH), 7.26–7.30 (m, 2H, ArH), 7.43–7.46 (m, 1H, ArH), 7.76 (m, 1H, ArH), 7.97 (dd, 2H, $J=5.5$, 8.8 Hz, ArH), 10.10 (br s, 1H, OH), 10.81 (br s, 1H, NH), 11.19 (br s, 1H, NH); ^{13}C NMR (DMSO- d_6): δ 74.5, 115.2, 115.4, 116.5, 117.4,

119.1, 120.0, 120.9, 129.5, 130.1, 130.5, 131.7, 135.5, 155.8, 157.9, 159.6, 161.4, 198.7. Anal. Calcd for $C_{18}H_{11}FN_6O_2S$: C, 54.54; H, 3.31; N, 21.20; S, 8.09. Found: C, 54.47; H, 3.45; N, 21.06; S, 8.22.

5-[(1H-Tetrazol-5-yl)methoxy]-4-(5-chloro-2-hydroxyphenyl)-6-(4-fluorophenyl)pyrimidine-2(1H)-thione (9b) This compound was prepared from **7b**; a yellow solid; conventional method time 220 min, yield 80%; ultrasound-assisted method time 15 min, yield 95%; mp 148–150°C; IR: ν 3975 (OH), 3424 (NH), 3383 (NH), 3067 (CH) cm^{-1} ; 1H NMR (DMSO- d_6): δ 4.81 (s, 2H, OCH₂), 6.90 (d, 1H, $J=8.8$ Hz, ArH), 7.19 (t, 2H, $J=8.8$ Hz, ArH), 7.25 (dd, 1H, $J=8.8$, 2.6 Hz, ArH), 7.32 (d, 1H, $J=2.6$ Hz, ArH), 8.00 (m, 2H, ArH), 10.10 (br s, 1H, OH), 10.81 (br s, 1H, NH), 11.19 (br s, 1H, NH); ^{13}C NMR (DMSO- d_6): δ 63.5, 115.4, 115.6, 117.9, 122.5, 122.8, 129.4, 130.4, 130.9, 131.6, 131.7, 146.4, 157.9, 159.6, 161.7, 162.2, 164.7, 190.6. Anal. Calcd for $C_{18}H_{12}ClFN_6O_2S$: C, 50.18; H, 2.81; N, 19.51; S, 7.44. Found: C, 50.25; H, 2.87; N, 19.49; S, 7.48.

5-[(1H-Tetrazol-5-yl)methoxy]-6-(4-fluorophenyl)-4-(2-hydroxy-3-methylphenyl)pyrimidine-2(1H)-thione (9c) This compound was prepared from **7c**; a yellow solid; conventional method time 220 min, yield 74%; ultrasound-assisted method time 24 min, yield 82%; mp 215–217°C; IR: ν 3935 (OH), 3475 (NH), 3364 (NH), 3057 (CH) cm^{-1} ; 1H NMR (DMSO- d_6): δ 2.49 (s, 3H, CH₃), 4.85 (s, 2H, OCH₂), 6.95 (s, 1H, ArH), 7.18–7.20 (m, 2H, ArH), 7.69 (s, 1H, ArH), 7.92–7.96 (m, 3H, ArH), 10.16 (br s, 1H, OH), 10.87 (br s, 1H, NH), 11.21 (br s, 1H, NH); ^{13}C NMR (DMSO- d_6): δ 15.8, 68.7, 115.7, 115.8, 118.1, 122.8, 122.9, 129.0, 130.2, 131.3, 131.8, 132.2, 146.9, 158.2, 159.3, 161.5, 161.9, 165.0, 195.2. Anal. Calcd for $C_{19}H_{15}FN_6O_2S$: C, 55.60; H, 3.68; N, 20.48; S, 7.81. Found: C, 55.57; H, 3.72; N, 20.39; S, 7.76.

5-[(1H-Tetrazol-5-yl)methoxy]-6-(4-fluorophenyl)-4-(2-hydroxy-5-methylphenyl)pyrimidine-2(1H)-thione (9d) This compound was prepared from **7d**; a yellow solid; conventional method time 235 min, yield 75%; ultrasound-assisted method time 24 min, yield 94%; mp 314–316°C; IR: ν 3954 (OH), 3424 (NH), 3364 (NH), 3095 (CH) cm^{-1} ; 1H NMR (DMSO- d_6): δ 2.44 (s, 3H, CH₃), 4.87 (s, 2H, OCH₂), 6.97 (t, 2H, $J=8.8$ Hz, ArH), 7.17–7.23 (m, 2H, ArH), 7.74 (s, 1H, ArH), 7.90 (dd, 2H, $J=8.8$, 5.5 Hz, ArH), 10.11 (br s, 1H, OH), 10.82 (br s, 1H, NH), 11.19 (br s, 1H, NH); ^{13}C NMR (DMSO- d_6): δ 16.3, 68.4, 115.9, 116.0, 118.5, 123.1, 123.4, 128.6, 129.8, 131.2, 131.5, 132.6, 147.7, 158.4, 160.2, 161.6, 161.8, 165.3, 194.9. Anal. Calcd for $C_{19}H_{15}FN_6O_2S$: C, 55.60; H, 3.68; N, 20.48; S, 7.81. Found: C, 55.65; H, 3.61; N, 20.55; S, 7.74.

5-[(1H-Tetrazol-5-yl)methoxy]-4-(3,5-dichloro-2-hydroxyphenyl)-6-(4-fluorophenyl)pyrimidine-2(1H)-thione (9e) This compound was prepared from **7e**; a yellow solid; conventional method time 195 min, yield 78%; ultrasound-assisted method time 18 min, yield 91%; mp 183–185°C; IR: ν 3935 (OH), 3453 (NH), 3390 (NH), 3054 (CH) cm^{-1} ; 1H NMR (DMSO- d_6): δ 4.83 (s, 2H, OCH₂), 6.24 (t, 2H, $J=8.5$ Hz, ArH), 7.37–7.39 (m, 1H, ArH), 7.78 (d, 2H, $J=5.5$ Hz, ArH), 7.92 (dd, 1H, $J=8.5$, 5.5 Hz, ArH), 10.09 (br s, 1H, OH), 10.87 (br s, 1H, NH), 11.24 (br s, 1H, NH); ^{13}C NMR (DMSO- d_6): δ 67.1, 115.6, 115.9, 116.8, 117.7, 119.5, 120.2, 121.0, 129.8, 130.2, 130.7, 131.2, 136.1, 156.4, 157.5, 159.9, 162.0, 193.6. Anal. Calcd for $C_{18}H_{11}Cl_2FN_6O_2S$: C, 46.46; H, 2.38; N, 18.06; S, 6.89. Found: C, 46.40; H, 2.48; N, 18.01; S, 6.97.

5-[(1H-Tetrazol-5-yl)methoxy]-4-(5-chloro-2-hydroxy-4-methylphenyl)-6-(4-fluorophenyl)pyrimidine-2(1H)-thione (9f) This compound was prepared from **7f**; a yellow solid;

conventional method time 210 min, yield 76%; ultrasound-assisted method time 21 min, yield 92%; mp 277–279°C; IR: ν 3954 (OH), 3473 (NH), 3323 (NH), 3036 (CH) cm^{-1} ; 1H NMR (DMSO- d_6): δ 2.45 (s, 3H, CH₃), 4.86 (s, 2H, OCH₂), 6.18 (t, 2H, $J=8.8$ Hz, ArH), 7.67–7.70 (m, 2H, ArH), 7.98–8.01 (m, 2H, ArH), 10.15 (br s, 1H, OH), 10.91 (br s, 1H, NH), 11.32 (br s, 1H, NH); ^{13}C NMR (DMSO- d_6): δ 17.3, 67.5, 115.9, 116.0, 116.9, 117.3, 119.6, 120.6, 121.4, 130.3, 130.7, 131.5, 131.9, 137.4, 155.6, 157.2, 160.4, 162.6, 194.2. Anal. Calcd for $C_{19}H_{14}ClFN_6O_2S$: C, 51.30; H, 3.17; N, 18.89; S, 7.21. Found: C, 51.38; H, 3.22; N, 18.91; S, 7.13.

Antimicrobial activity

Antibacterial activity of the synthesized compounds was tested *in vitro* against Gram-positive bacteria *S. aureus* (NCIM 2079) and *B. subtilis* (NCIM 2920) and Gram-negative bacteria *E. coli* (NCIM 2065) and *P. aeruginosa* (NCIM 2200). The compounds were also screened for antifungal activity against *C. albicans* (NCIM 3471) and *A. niger* (NCIM 596). Compounds were diluted in dimethyl sulfoxide (DMSO) with 1 $\mu g/mL$ concentrations for bioassay. Micro-broth dilution method [28] was used to determine the MICs of compounds in 96-well micro-titer plates. Test compounds were serially diluted in growth media. Plates were incubated at 30°C for fungi and 37°C for bacteria for 24 h. All experiments were carried out in triplicates.

Acknowledgments: V.S.D. is grateful to the Council of Scientific and Industrial Research (CSIR), New Delhi, India, for providing Senior Research Fellowship and Sophisticated Analytical Instrumentation Facility (SAIF). Panjab University provided spectral data. We are thankful to the Head, Department of Chemistry, Dr. Babasaheb Ambedkar Marathwada University, for providing laboratory facility. We also thank the Department of Microbiology, Maulana Azad College of Arts, Science and Commerce, Aurangabad, for providing biological assay results.

References

- [1] Jin, H.; Zhao, H.; Zhao, F.; Li, S.; Liu, W.; Zhou, G.; Tao, K.; Hou, T. Efficient epoxidation of chalcones with urea-hydrogen peroxide under ultrasound irradiation. *Ultrason. Sonochem.* **2009**, *16*, 304–307.
- [2] Grare, M.; Mourer, M.; Fontanay, S.; Regnouf-de-Vains, J. B.; Finance, C.; Duval, R. E. In vitro activity of para-guanidinoethyl-calix[4]arene against susceptible and antibiotic-resistant Gram-negative and Gram-positive bacteria. *J. Antimicrob. Chemother.* **2007**, *60*, 575–581.
- [3] Anzahae, M. Y.; Watts, J. K.; Alla, N. R.; Nicholson, A. W.; Damha, M. J. Energetically important C-H...F-C pseudohydrogen bonding in water: evidence and application to rational design of oligonucleotides with high binding affinity. *J. Am. Chem. Soc.* **2011**, *133*, 728–731.
- [4] Young, A. M.; Audus, K. L.; Proudfoot, J.; Yazdani, M. Tetrazole compounds: the effect of structure and pH on Caco-2 cell permeability. *J. Pharm. Sci.* **2006**, *95*, 717–725.

- [5] Myznikov, L. V.; Hrabalek, A.; Koldobskii, G. I. Drugs in the tetrazole series. *Chem. Heterocycl. Compd.* **2007**, *43*, 1–9.
- [6] Alam, M.; Nami, S. A.; Husain, A.; Lee, D.; Park, S. Synthesis, characterization, X-ray diffraction, antimicrobial and in vitro cytotoxicity studies of 7a-Aza-B-homostigmast-5-eno [7a,7-d] tetrazole. *C. R. Chim.* **2013**, *16*, 201–206.
- [7] Bondaryk, M.; Lukowska-Chojnacka, E.; Staniszewska, M. Tetrazole activity against *Candida albicans*. The role of KEX2 mutations in the sensitivity to (\pm)-1-[5-(2-chlorophenyl)-2H-tetrazol-2-yl]propan-2-yl acetate. *Bioorg. Med. Chem. Lett.* **2015**, *25*, 2657–2663.
- [8] Karabanovich, G.; Roh, J.; Smutny, T.; Nemecek, J.; Vicherek, P.; Stolarikova, J.; Vejsova, M.; Dufkova, I.; Vavrova, K.; Pavek, P.; et al. 1-Substituted-5-[(3,5-dinitrobenzyl)sulfanyl]-1H-tetrazoles and their isosteric analogs: a new class of selective antitubercular agents active against drug-susceptible and multidrug-resistant mycobacteria. *Eur. J. Med. Chem.* **2014**, *82*, 324–340.
- [9] Muraglia, E.; Kinzel, O. D.; Laufer, R.; Miller, M. D.; Moyer, G.; Munshi, V.; Orvieto, F.; Palumbi, M. C.; Pescatore, G.; Rowley, M.; et al. Tetrazolethioacetanilides: potent non-nucleoside inhibitors of WT HIV reverse transcriptase and its K103N mutant. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 2748–2752.
- [10] Pegklidou, K.; Koukoulitsa, C.; Nicolaou, I.; Demopoulos, V. J. Design and synthesis of novel series of pyrrole based chemotypes and their evaluation as selective aldose reductase inhibitors. A case of bioisosterism between a carboxylic acid moiety and that of a tetrazole. *Bioorg. Med. Chem.* **2010**, *18*, 2107–2114.
- [11] Li, J.; Chen, S. Y.; Tao, S.; Wang, H.; Li, J. J.; Swartz, S.; Musial, C.; Hernandez, A. A.; Flynn, N.; Murphy, B. J.; et al. Design and synthesis of tetrazole-based growth hormone secretagogue: the SAR studies of the O-benzyl serine side chain. *Bioorg. Med. Chem. Lett.* **2008**, *18*, 1825–1829.
- [12] Isloor, A. M.; Kalluraya, B.; Shetty, P. Regioselective reaction: synthesis, characterization and pharmacological studies of some new Mannich bases derived from 1,2,4-triazoles. *Eur. J. Med. Chem.* **2009**, *44*, 3784–3787.
- [13] Isloor, A. M.; Kalluraya, B.; Rao, M. Sydnone derivatives: part IV; synthesis of 3-aryl-4-(substituted pyrazolidene hydrazine-4-thiazolyl) sydnones as possible analgesic and anticonvulsant agents. *J. Saudi Chem. Soc.* **2000**, *4*, 265–270.
- [14] Bekhita, A. A.; Abdel-Aziem, T. Design, synthesis and biological evaluation of some pyrazole derivatives as anti-inflammatory-antimicrobial agents. *Bioorg. Med. Chem.* **2004**, *12*, 1935–1945.
- [15] Bertuzzi, G.; Locatelli, E.; Colecchia, D.; Calandro, P.; Bonini, B. F.; Chandanshive, J. Z.; Mazzanti, A.; Zani, P.; Chiariello, M.; Franchini, M. C. Straightforward synthesis of a novel ring-fused pyrazole-lactam and in vitro cytotoxic activity on cancer cell lines. *Eur. J. Med. Chem.* **2016**, *117*, 1–7.
- [16] Mohamed, M. S.; Awad, S. M.; Sayed, A. I. Synthesis of certain pyrimidine derivatives as antimicrobial agents and anti-inflammatory agents. *Molecules* **2010**, *15*, 1882–1890.
- [17] Zhang, N.; Ayril-Kaloustian, S.; Nguyen, T.; Hernandezb, R.; Beyerb, C. 2-Cyanoaminopyrimidines as a class of antitumor agents that promote tubulin polymerization. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 3003–3005.
- [18] Katiyar, S. B.; Bansal, I.; Saxena, J. K.; Chauhan, P. M. S. Syntheses of 2,4,6-trisubstituted pyrimidine derivatives as a new class of antifilarial topoisomerase II inhibitors. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 47–50.
- [19] Lu, X.; Chen, Y.; Guo, Y.; Liu, Z.; Shi, Y.; Xu, Y.; Wang, X.; Zhang, Z.; Liu, J. The design and synthesis of N-1-alkylated-5-aminoaryalkyl-substituted-6-methyluracils as potential non-nucleoside HIV-1 RT inhibitors. *Bioorg. Med. Chem.* **2007**, *15*, 7399–7407.
- [20] Morgan, A.; Cofer, C.; Stevens, D. L. Iclaprim: a novel dihydrofolate reductase inhibitor for skin and soft tissue infections. *Future Microbiol.* **2009**, *4*, 131–144.
- [21] Prakash, O.; Kumar, R.; Kumar, R.; Tyagi, P.; Kuhad, R. C. Organiodine(III) mediated synthesis of 3,9-diaryl- and 3,9-difuryl-bis-1,2,4-triazolo[4,3-a][4,3-c]pyrimidines as antibacterial agents. *Eur. J. Med. Chem.* **2007**, *42*, 868–872.
- [22] Sondhi, S. M.; Jain, S.; Dinodia, M.; Shuklab, R.; Raghbir, R. One pot synthesis of pyrimidine and bispyrimidine derivatives and their evaluation for anti-inflammatory and analgesic activities. *Bioorg. Med. Chem.* **2007**, *15*, 3334–3344.
- [23] Bruno, O.; Brullo, C.; Schenone, S.; Ranise, A.; Bondavalli, F.; Barocelli, E.; Tognolini, M.; Magnanini, F.; Ballabeni, V. Progress in 5H[1]benzopyrano[4,3-d]pyrimidin-5-amine series: 2-Methoxy derivatives effective as antiplatelet agents with analgesic activity. *Farmaco.* **2002**, *57*, 753–758.
- [24] McCarthy, O.; Musso-Buendia, A.; Kaiser, M.; Brun, R.; Ruiz-Perez, L. M.; Johansson, N. G.; Pacanowska, D. G.; Gilbert, I. H. Design, synthesis and evaluation of novel uracil acetamide derivatives as potential inhibitors of plasmodium falciparum dUTP nucleotide hydrolase. *Eur. J. Med. Chem.* **2009**, *44*, 678–688.
- [25] Diwakar, S. D.; Bhagwat, S. S.; Shingare, M. S.; Gill, C. H. Substituted 3-((Z)-2-(4-nitrophenyl)-2-(1H-tetrazol-5-yl) vinyl)-4H-chromen-4-ones as novel anti-MRSA agents: synthesis, SAR, and in-vitro assessment. *Bioorg. Med. Chem. Lett.* **2008**, *18*, 4678–4681.
- [26] Gharpure, M.; Choudhary, R.; Ingle, V.; Juneja, H. Synthesis of new series of 3-hydroxy/acetoxo-2-phenyl-4H-chromen-4-ones and their biological importance. *J. Chem. Sci.* **2013**, *125*, 575–582.
- [27] Dofe, V. S.; Sarkate, A. P.; Lokwani, D. K.; Kathwate, S. H.; Gill, C. H. Synthesis, antimicrobial evaluation, and molecular docking studies of novel chromone based 1,2,3-triazoles. *Res. Chem. Int.* **2017**, *43*, 15–28.
- [28] Wiegand, I.; Hilpert, K.; Hancock, R. E. Agar and broth dilution methods to determine the minimal inhibitory concentration (MIC) of antimicrobial substances. *Nat. Protoc.* **2008**, *3*, 163–175.