

Prescribe

BROMFED® - BROMFED-PD®

Timed-release Capsules
(antihistamine/decongestant action)

BROMFED® CAPSULES Timed-release capsules
(antihistamine/decongestant action)

DESCRIPTION: A light green and clear capsule containing white beads.

Each capsule contains:

Brompheniramine maleate	12 mg
Pseudoephedrine hydrochloride (d-isoephedrine HCl)	120 mg

BROMFED-PD® CAPSULES Timed-release capsules
(antihistamine/decongestant action)

DESCRIPTION: A dark green and clear capsule containing white beads.

Each capsule contains:

Brompheniramine maleate	6 mg
Pseudoephedrine hydrochloride (d-isoephedrine HCl)	60 mg

Both products are in a specially prepared base to provide prolonged action.

BROMFED® TABLETS

DESCRIPTION: A white scored tablet. Each tablet contains:

Brompheniramine maleate	4 mg
Pseudoephedrine hydrochloride	60 mg

These products contain ingredients of the following therapeutic classes:
antihistamine and decongestant.

CLINICAL PHARMACOLOGY: Brompheniramine maleate is an alkylamine type antihistamine. This group of antihistamines are among the most active histamine antagonists and are generally effective in relatively low doses. The drugs are not so prone to produce drowsiness and are among the most suitable agents for day time use; but again, a significant proportion of patients do experience this effect. Pseudoephedrine hydrochloride is a sympathomimetic which acts predominantly on alpha receptors and has little action on beta receptors. It therefore functions as an oral nasal decongestant with minimal CNS stimulation.

INDICATIONS: For the treatment of the symptoms of seasonal and perennial allergic rhinitis, and vasomotor rhinitis, including nasal obstruction (congestion).

CONTRAINDICATIONS: Hypersensitivity to any of the ingredients. Also contraindicated in patients with severe hypertension, severe coronary artery disease, patients on MAO inhibitor therapy, patients with narrow-angle glaucoma, urinary retention, peptic ulcer and during an asthmatic attack.

BROMFED®/BROMFED-PD®

WARNINGS: Considerable caution should be exercised in patients with hypertension, diabetes mellitus, ischemic heart disease, hyperthyroidism, increased intraocular pressure and prostatic hypertrophy. The elderly (60 years or older) are more likely to exhibit adverse reactions.

Antihistamines may cause excitability, especially in children. At doses higher than the recommended dose, nervousness, dizziness or sleeplessness may occur.

PRECAUTIONS: General: Caution should be exercised in patients with high blood pressure, heart disease, diabetes or thyroid disease. The antihistamine in this product may exhibit additive effects with other CNS depressants, including alcohol.

INFORMATION FOR PATIENTS: Antihistamine may cause drowsiness and ambulatory patients who operate machinery or motor vehicles should be cautioned accordingly.

DRUG INTERACTIONS: MAO inhibitors and beta adrenergic blockers increase the effects of sympathomimetics. Sympathomimetics may reduce the antihypertensive effects of methyldopa, mecamylamine, reserpine and veratrum alkaloids. Concomitant use of antihistamines with alcohol and other CNS depressants may have an additive effect.

PREGNANCY: The safety of use of this product in pregnancy has not been established.

ADVERSE REACTIONS: Adverse reactions include drowsiness, lassitude, nausea, giddiness, dryness of mouth, blurred vision, cardiac palpitations, flushing, increased irritability or excitement (especially in children).

BROMFED® CAPSULES

DOSAGE AND ADMINISTRATION: Adults and children over 12 years of age—1 capsule orally every 12 hours.

HOW SUPPLIED: Bottles of 100—500

BROMFED-PD® CAPSULES

DOSAGE AND ADMINISTRATION: Children 6 to 12 years of age—1 capsule every 12 hours. Adults—2 capsules every 12 hours.

HOW SUPPLIED: Bottles of 100—500

BROMFED® TABLETS

DOSAGE AND ADMINISTRATION: Adults and children 12 and over: One tablet every 4 hours not to exceed 6 doses in 24 hours. Children 6 to 12 years: One-half tablet every 4 hours not to exceed 6 doses in 24 hours. Do not give to children under 6 years except under the advice and supervision of a physician.

HOW SUPPLIED: Bottles of 100

BROMFED® SYRUP

DOSAGE AND ADMINISTRATION: Adults and children 12 years of age and over: 2 teaspoonfuls orally every 4-6 hours. Children 6 to under 12 years of age: 1 teaspoon orally every 4-6 hours. Do not exceed 4 doses in 24 hours. Children under 6 years of age, consult a physician.

HOW SUPPLIED: Bromfed Syrup is available in 16 fl. oz. bottles.

CAUTION: FEDERAL (U.S.A.) LAW PROHIBITS DISPENSING WITHOUT A PRESCRIPTION.

Muro


Pharmaceutical Inc.

Tewksbury, MA 01876-9987

New members

Bertrand, V. Paul, PCOM '68; Chicago, IL
Bonefeld, Robert, MSUCOM '80; Treasure Island, FL
Boyd, Seward, Jr., UOMHS '81; Washington, DC
Burns, Patrick, PCOM '80; Washington, DC
Ellis, Richard L., Lt., KCOM '84; Port Hueneme, LA
Gangewere, Keith D., PCOM '80; Richmond Heights, OH
Huddleston, Don M., TCOM '76; Rolla, MO
Kaplan, Daniel B., PCOM '73; Philadelphia, PA
Latella, Joseph X., UOMHS '70; Webster City, IA
Manley, Louis J., UOMHS '56; Orlando, FL
Olson, Allen L., KCOM '73; Marquette, MI
Pardoe, Theresa, M., UOMHS '86; Milwaukee, WI
Pratt, Allan, T., UOMHS '76; Kissimmee, FL
Ramone, Michael D., KCOM '84; Drexel Hill, PA
Robb, Thomas M., UOMHS '76; Youngtown, OH
Sebastian, Peter, R., PCOM '78; Salisbury, MD
Shanks, James R., KCOM '73; Clearwater, FL
Smith, Leon, PCOM '71; Muskegon, MI

Coming soon...



An ideal new
companion for

SELDANE[®]

(terfenadine)

Merrell Dow