

The purpose of this quiz is to provide a convenient means of self-assessment of your reading of the scientific content of this issue of JAOA. Enter your answers to the questions in the spaces provided so that you can easily check them with the answers that will be published next month.

To apply for CME credit, transfer your answers to the mail-in card on page 101 and return it to the CME office. So that you may complete this self-assessment in privacy, use only your member number to apply for CME credit. The CME office will record only the fact that you have completed the self-assessment test. Any grading will be done by the Editorial Department only for the purpose of planning areas of study which may be helpful to cover in future issues of JAOA.

1. The study of the use of hyalurate sodium *v* air in extracapsular cataract extraction demonstrated a statistically significant advantage in utilizing healon over air in protecting against endothelial cell loss in cataract surgery.
☐ (a) True
☐ (b) False
2. Which of the following is not considered a part of the joint system of the shoulder girdle?
☐ (a) First rib
☐ (b) Glenohumeral joint
☐ (c) Second rib
☐ (d) Suprahumeral articulation
☐ (e) Scapulothoracic articulation
3. Which muscle group is considered to be the initiator of humeral abduction?
☐ (a) Supraspinatus
☐ (b) Rhomboid major
☐ (c) Teres major
☐ (d) Latissimus dorsi
☐ (e) All of the above are equally important in initiating humeral abduction.
4. The initial treatment of phlegmasia cerulea dolens is
☐ (a) venous thrombectomy.
☐ (b) thrombolytic therapy.
☐ (c) heparinization.
5. The majority of iliofemoral thromboses occur on which side (because of anatomical considerations)?
☐ (a) Right
☐ (b) Left
6. The classical retinal lesions found in systemic lupus erythematosus (SLE) is cotton wool spots.
☐ (a) True
☐ (b) False
7. The dreaded complication of antimalarial treatment in SLE is irreversible retinal toxicity leading to blindness.
☐ (a) True
☐ (b) False
8. All the following are true EXCEPT:
☐ (a) Takayasu's arteritis has been associated with elevated antistreptolysin-O titers in some cases.
☐ (b) Purified protein derivative (PPD) positivity has been found in many of Takayasu's patients.
☐ (c) *Mycobacterium tuberculosis* organisms have been identified in arterial histologic sections.
☐ (d) There may be a genetic predisposition to Takayasu's arteritis with HLA association.
9. All the following are false EXCEPT:
☐ (a) Takayasu's arteritis is more common in men.
☐ (b) Most patients affected are elderly females.
☐ (c) The primary blood vessels involved are large and medium arteries.
☐ (d) Early symptoms include prominent joint pain and swelling.
10. According to recent reports, the number of AIDS cases currently reported in the United States is about
☐ (a) 1.5 million.
☐ (b) 150,000.
☐ (c) 61,000.
☐ (d) 25,000.
11. Bisexual and homosexual men, male and female IV drug users, and male homosexual IV drug users are at highest risk for contracting AIDS. This combined population accounts for what percentage of the AIDS cases reported in the United States?
☐ (a) 50%
☐ (b) 66%
☐ (c) 75%
☐ (d) 91%
12. Recent surveys suggest that osteopathic medical schools and their allied health care centers are currently prepared to deal with the AIDS crisis.
☐ (a) True
☐ (b) False