

Abdul Razak*, Omar Ibrahim Alhaidari and Javed Ahmed

Interventions for reducing late-onset sepsis in neonates: an umbrella review

<https://doi.org/10.1515/jpm-2022-0131>

Received March 11, 2022; accepted August 17, 2022;

published online October 31, 2022

Abstract

Objectives: Neonatal sepsis is one of the leading causes of neonatal deaths in neonatal intensive care units. Hence, it is essential to review the evidence from systematic reviews on interventions for reducing late-onset sepsis (LOS) in neonates.

Methods: PubMed and the Cochrane Central were searched from inception through August 2020 without any language restriction. Cochrane reviews of randomized clinical trials (RCTs) assessing any intervention in the neonatal period and including one or more RCTs reporting LOS. Two authors independently performed screening, data extraction, assessed the quality of evidence using Cochrane Grading of Recommendations Assessment, Development and Evaluation, and assessed the quality of reviews using a measurement tool to assess of multiple systematic reviews 2 tool.

Results: A total of 101 high-quality Cochrane reviews involving 612 RCTs and 193,713 neonates, evaluating 141 interventions were included. High-quality evidence showed a reduction in any or culture-proven LOS using antibiotic lock therapy for neonates with central venous catheters (CVC). Moderate-quality evidence showed a decrease in any LOS with antibiotic prophylaxis or vancomycin prophylaxis for neonates with CVC, chlorhexidine for skin or cord care, and kangaroo care for low birth weight babies. Similarly, moderate-quality evidence showed reduced culture-proven

LOS with intravenous immunoglobulin prophylaxis for preterm infants and probiotic supplementation for very low birth weight (VLBW) infants. Lastly, moderate-quality evidence showed a reduction in fungal LOS with the use of systemic antifungal prophylaxis in VLBW infants.

Conclusions: The overview summarizes the evidence from the Cochrane reviews assessing interventions for reducing LOS in neonates, and can be utilized by clinicians, researchers, policymakers, and consumers for decision-making and translating evidence into clinical practice.

Keywords: infection; late-onset sepsis; neonatal sepsis; neonate; overview; sepsis.

Introduction

Sepsis is traditionally defined as the isolation of the pathogen from sterile body fluid such as blood or cerebrospinal fluid. In adults, the consensus definition includes life-threatening organ dysfunction caused by a dysregulated response to infection [1]. However, in neonates, no such consensus definition is available [2]. Sepsis manifesting in neonates within the first 72 h of life is classified as early-onset, whereas it is considered late-onset if the manifestations appear beyond 72 h of life. Advances in obstetric care and prophylactic intrapartum antibiotics have reduced the risk of early-onset sepsis [3, 4]. However, the incidence of late-onset sepsis (LOS) has increased in parallel with the improved survival of extremely premature (gestational age <28 weeks) and very low birth weight (VLBW, birth weight <1,500 g) neonates [3, 4]. It affects 0.61–14.2 percent of hospitalized newborn infants and is inversely related to the degree of prematurity [5].

Sepsis remains the major contributor to global mortality, and the World Health Organization has declared it a global health priority [6]. Despite effective treatment strategies, including appropriate antimicrobial treatment, it remains a leading cause of neonatal death and a significant contributor to neonatal morbidities in neonatal intensive care units and the community. It is therefore imperative to identify the preventive measures that reduce the risk of sepsis in the neonatal period. Randomized clinical trials (RCTs) and their systematic

*Corresponding author: Dr. Abdul Razak, MD, Monash Newborn, Monash Children's Hospital, Department of Paediatrics, Monash University, 246 Clayton Road, Clayton, VIC 3168, Australia; and Division of Neonatology, Department of Pediatrics, King Abdullah Bin Abdulaziz University Hospital, Princess Nourah Bint Abdulrahman University, Riyadh, Saudi Arabia, Phone: +966560636849, E-mail: abdul.razak@monash.edu. <https://orcid.org/0000-0002-6185-3694>

Omar Ibrahim Alhaidari, Division of Neonatology, Department of Pediatrics, King Abdullah Bin Abdulaziz University Hospital, Princess Nourah Bint Abdulrahman University, Riyadh, Saudi Arabia; and Department of Pediatrics, McMaster Children's Hospital, McMaster University, ON, Canada

Javed Ahmed, Department of Pediatrics, McMaster Children's Hospital, McMaster University, ON, Canada

reviews investigating a broad range of interventions realize the potential for interventions of interest to reduce the risk of neonatal sepsis. Given that several risk factors impact the risk of neonatal sepsis, there is a need to systematically examine all potentially relevant interventions that can contribute to the prevention of neonatal sepsis. To our knowledge, no published overview has compiled and summarized the evidence from systematic reviews on interventions for preventing neonatal sepsis in one coherent document. This overview will help clinicians, researchers, policymakers, funding bodies, and consumers assist decision-making and evidence translation.

Materials and methods

We conducted this systematic review as per Preferred Reporting Items for Overviews of Systematic Reviews including Harms (PRIO-harms) guideline [7] and the approach outlined in the Cochrane Handbook for Systematic Reviews of Interventions [8]. We registered the protocol for the systematic review with PROSPERO (registration number CRD42020192513), the international prospective register for systematic reviews (<https://www.crd.york.ac.uk/PROSPERO>).

Search strategy

We conducted a comprehensive search of the literature using appropriate pre-specified search terms within the following databases: PubMed and the Cochrane Central Register of Controlled Trials (CENTRAL) from inception (Supplemental Information) through August 6, 2020. We did not apply language restrictions. We also read the reference lists of included systematic reviews to identify eligible studies. Finally, we searched the “related articles” feature in PubMed and hand searched in Cochrane Neonatal (<https://neonatal.cochrane.org>) for missing reviews.

Study selection (eligibility criteria)

Type of studies: In this overview, we included Cochrane reviews RCTs or quasi-RCTs assessing neonatal interventions. The Cochrane review must consist of one or more RCTs reporting at least one of the pre-specified outcomes of this review. We excluded non-Cochrane reviews, Cochrane reviews of non-randomized studies, and Cochrane reviews with no RCTs reporting the outcome.

Population: We included Cochrane reviews assessing interventions in term and preterm neonates. We also included reviews reporting data on mixed infant (age less than one year) populations as long as separate data was provided for neonates.

Interventions and comparisons: We considered all types of interventions following the birth of the newborn infant (delivery room interventions) or interventions in the neonatal period (neonatal interventions within 28 days from the birth) in a neonatal intensive care unit or community, compared with placebo, no treatment, or an

alternative treatment. We excluded Cochrane reviews assessing non-neonatal interventions (interventions in pregnancy or before childbirth) and interventions beyond the neonatal period. However, interventions initiated in the neonatal period and continued beyond the neonatal period were also included.

Outcomes

Primary outcomes:

1. Any LOS: Clinically suspected or microbiologically confirmed LOS, as defined by the Cochrane review authors.
2. Culture-proven LOS: Microbiologically confirmed LOS, as defined by the Cochrane review authors.

Secondary outcomes:

1. Bacterial LOS: LOS caused by a bacterial organism, as defined by the Cochrane review authors.
2. Fungal LOS: LOS caused by a fungal organism, as defined by the Cochrane review authors.

We excluded Cochrane reviews evaluating early-onset sepsis as our overview focuses on neonatal interventions and not childbirth or pregnancy interventions. However, we included Cochrane reviews evaluating both early-onset and late-onset sepsis as long as reviews reporting separate data on LOS were available.

Study selection and data extraction

The author (A.R.) performed the literature search across the databases using pre-specified terms (Supplementary Information). We managed the citations retrieved through the search using Covidence. Authors, J.A. and O.A., independently performed the titles and abstracts screening, independently evaluated the full-text articles of the short-listed reports, and independently extracted the information, including review title and authors, date last assessed as up-to-date, number of trials, number of participants and their characteristics, interventions and comparisons, outcomes relevant to this overview, and quality of RCTs assessed by review authors, and summary intervention effects (relative risks (RR), odds risk or absolute risk differences and their 95% confidence intervals (CI), model used for meta-analysis) for all neonates and subgroup of neonates (preterm and term neonates).

Assessment of methodological quality of included reviews

Quality of included reviews: Two authors (J.A. and A.R.) independently assessed the methodological quality of each Cochrane systematic review using the AMSTAR 2 (A Measurement Tool to Assess Systematic Reviews 2) instrument [9]. AMSTAR 2 evaluates the review methodology against 16 distinct domains, of which 7 are critical domains. Eleven domains on AMSTAR 2 are rated as yes or no, and five domains are rated as yes, partially yes, and no. The overall confidence in the results of the review was rated as high (no or one non-critical weakness), moderate (more than one non-critical weakness), low (one critical flaw with or without non-critical weaknesses), and critically low (more than one critical flaw with or without non-critical weaknesses).

Quality of included studies within reviews: We reported, instead of reassessing, the quality of included RCTs within reviews according to the review authors' judgment.

Quality of evidence in the included reviews: We reported the quality of evidence using the Cochrane Grading of Recommendations Assessment, Development, and Evaluation (GRADE) approach according to the review authors' judgment as high certainty, moderate certainty, low certainty, or very low certainty [10]. Two authors (J.A. and O.A.) independently assessed the evidence certainty using the Cochrane GRADE if the Cochrane review authors did not provide the assessment of the evidence certainty. Discussions and consensus with the third author (A.R.) resolved discrepancies in the review process.

Data synthesis

We summarized the results of included reviews in the summary of the findings table. We categorized the interventions into 5 categories: effective interventions (high certainty in the evidence of effectiveness), possible effective interventions (moderate certainty in the

evidence of effectiveness), ineffective interventions (high certainty in the evidence of lack of effectiveness (or harm)), possible ineffective interventions (moderate certainty in the evidence of lack of effectiveness (or harm)), and no conclusions possible (low or very low certainty in the evidence) [8].

Results

The database search results and study selection log are shown in Figure 1. The search yielded 1,047 from databases and 1 article from other sources. After removing the duplicates and unrelated articles by title and abstract screening, we included 286 reviews for full-text screening. Finally, 101 Cochrane reviews were included after excluding 185 reviews [11–111]. All the 101 included Cochrane reviews were of high quality, and the detailed evaluation as per AMSTAR 2 tool is provided in Table 1 of the Supplementary Information.

Table 1 provides the details of the included reviews, including the summary assessment of the primary and




Figure 1: Study flow diagram outlining stages of search results and filtering process.

Table 1: Summary of Cochrane reviews assessing intervention on late-onset neonatal sepsis.

Review ID	Population	Intervention	Comparison	Sepsis	No. of RCTs	No. of patients included in the RCTs	Summary of measure	Estimate (95% CI)	Certainty of evidence, GRADE
Abramamatha 2019	<37 w	Routine monitoring of gastric residuals	No monitoring	CP	2	141	RR, fixed	1.46 [0.85, 2.52]	Low ^{1,3,a}
		Criteria 1 gastric residue monitoring	Criteria 2 gastric residue monitoring	CP	1	87	RR, fixed	5.35 [0.26, 108.27]	Low ^{1,3,a}
Aher 2020	<37 w or LBW Neonates requiring PN	Late ESAs	P/NI	CP	5	551	RR, fixed	0.75 [0.52, 1.09]	Low ^{1,3,a}
Alnsworth 2015		Percutaneous CVC	Peripheral cannula	CP	6	549	RR, fixed	0.95 [0.72, 1.25]	Moderate ^{1,a}
Ardell 2015	<37 w	Animal derived surfactant	Protein free synthetic surfactant	Bacterial	10	5,219	RR, fixed	1.00 [0.91, 1.10]	Moderate ^{1,a}
Ardell 2018	<37 w	0.2 mg IV vit K	0.2 mg IM	Any	1	52	RR, fixed	1.0 [0.28, 3.58]	Low ^{1,3}
			0.5 mg IM	Any	1	54	RR, fixed	0.86 [0.26, 2.86]	Low ^{1,3}
			Any IM	Any	1	80	RR, fixed	0.92 [0.31, 2.72]	Low ^{1,3,a}
Austin 2015	<32 w or VLBW	Oral/topical antifungal	P/NI	Fungal	4	1800	RR, fixed	0.20 [0.14, 0.27]	Low ^{1,2,a}
			Systemic	Fungal	3	326	RR, fixed	1.89 [0.66, 5.39]	Low ^{1,3,a}
			Delayed	Bacterial	1	75	RR, fixed	1.14 [0.81, 1.60]	Low ^{1,3,a}
Bahadue 2012	RDS	Early surfactant (animal derived surfactant)	Non-impregnated CVC	CP	1	86	RR, fixed	0.11 [0.01, 0.87]	Low ^{1,3,a}
Balain 2015	Neonates with CVC	Antimicrobial-impregnated CVC	No glucose reduction	Bacterial	1	23	RR, fixed	0.61 [0.29, 1.25]	Low ^{1,3,a}
Bottino 2011	<32 w or VLBW	Insulin infusion	Glucose reduction	Bacterial	1	23	RR, fixed	0.46 [0.10, 2.03]	Low ^{1,3,a}
				Fungal	1	23	RR, fixed	0.18 [0.01, 3.47]	Low ^{1,3,a}
Brion 2003	<37 w	Vitamin E	P/NI	Any	4	1,009	RR, fixed	1.52 [1.13, 2.04]	Moderate ^{1,a}
			Another vitamin E	Any	1	44	RR, fixed	0.33 [0.04, 2.96]	Low ^{1,3,a}
Brown 2014	Neonates with severe GI disease	Glutamine	P/NI	CP	3	263	RR, fixed	1.37 [0.89, 2.11]	Moderate ^{1,a}
Carr 2003	Neonates requiring NICU	G-CSF or GM-CSF prophylaxis	P/NI	CP	1	75	RR, fixed	0.66 [0.36, 1.20]	Low ^{1,3,a}
				Any	2	284	RR, fixed	1.02 [0.76, 1.37]	Low ^{1,3,a}
Cleminson 2015	<32 w or VLBW	Systemic antifungal	P/NI	Fungal	10	1,371	RR, fixed	0.43 [0.31, 0.59]	Moderate ^{1,a}
			Oral/Topical	Fungal	3	326	RD, fixed	-0.03 [-0.07, 0.02]	Low ^{1,3,a}
			Routine care	CP	8	2086	RR, fixed	1.13 [0.97, 1.31]	Low ^{1,3}
Cleminson 2016	<37 w	Topical ointment or cream (sunflower seed oil, aquaphor, petroleum jelly, beiersdorf inc., bepanthen, 70% lanolin, 30% olive oil, eucerin crème)		CoNS	6	1839	RR, fixed	1.30 [1.03, 1.65]	Moderate ^{1,a}
				Other bacteria	6	1839	RR, fixed	0.84 [0.63, 1.12]	Moderate ^{1,a}
				Fungal	6	1839	RD, fixed	0.01 [-0.01, 0.03]	Moderate ^{1,a}
				CP	6	844	RR, fixed	0.71 [0.51, 1.01]	Low ^{1,3}
				CoNS	5	775	RR, fixed	0.15 [0.02, 1.16]	Low ^{1,3,a}
				Other bacteria	5	775	RR, fixed	0.70 [0.47, 1.05]	Very low ^{1,2,3,a}
				Fungal	5	775	RR, fixed	1.93 [0.42, 8.78]	Very low ^{1,2,3,a}

Table 1: (continued)

Review ID	Population	Intervention	Comparison	Sepsis	No. of RCTs	No. of patients included in the RCTs	Summary of measure	Estimate (95% CI)	Certainty of evidence, GRADE
Collins 2015	<37 w	Topical ointment or cream (sunflower seed oil, aquaphor)	Topical oil (sunflower seed oil, aquaphor)	CP	1	316	RR, fixed	0.91 [0.57, 1.46]	Low ^{1,3,a}
				Other bacteria	1	316	RR, fixed	0.90 [0.53, 1.50]	Low ^{1,3,a}
				Fungal	1	316	RR, fixed	1.35 [0.31, 5.94]	Low ^{1,3,a}
				Any	1	88	RR, fixed	0.35 [0.17, 0.69]	Low ^{1,a}
Collins 2016	<37 w	Breastfeeding, with supplements without bottle feeding	Late discharge (once full suckling is established)	Infection (any)	3	500	RR, fixed	0.70 [0.35, 1.42]	Moderate ^{1,4}
Conde 2016	LBW infant	Kangaroo care	Conventional care	Any (latest follow up)	8	1,463	RR, fixed	0.50 [0.36, 0.69]	Moderate ¹
				Any (at discharge or 40–41 w)	5	1,239	RR, fixed	0.35 [0.22, 0.54]	Moderate ^{1,a}
Craft 2000	Neonates <1,500 g or with CVC requiring PN	Early kangaroo care	Late kangaroo care	Any	1	73	RR, fixed	0.42 [0.12, 1.49]	Low ^{1,3,a}
				Any	4	290	RR, fixed	0.11 [0.05, 0.24]	Moderate ^{1, a}
				CoNS	4	221	RR, fixed	0.33 [0.19, 0.59]	Low ^{1,2,a}
				Bacterial	3	583	RR, fixed	0.73 [0.57, 0.93]	Low ^{1,3,a}
Darlow 2003	<37 w or <2000 g	Selenium supplementation	P/NI	Any	3	947	RR, fixed	0.89 [0.76, 1.04]	Low ^{1,3,a}
				Any	1	80	RR, fixed	No difference b/w groups	Low ^{1,3,a}
De paoli 2008	<37 w requiring CPAP	Once a week vit A	Standard dose	Any	1	80	RR, fixed	No difference b/w groups	Low ^{1,3,a}
				CP	1	87	RR, fixed	1.02 [0.66, 1.58]	Low ^{1,3,a}
				Suspected	1	87	RR, fixed	0.93 [0.60, 1.44]	Low ^{1,3,a}
				CP	1	100	RR, fixed	1.09 [0.53, 2.24]	Low ^{1,3,a}
Doyle 2017 (early PNCS)	Preterm at risk of BPD	Infant flow driver	Medicorp prong	Infection (LOS, any)	25	4,101	RR, fixed	1.05 [0.96, 1.15]	Moderate ^{1,1,a}
Doyle 2017 (late PNCS)	Preterm at risk of BPD	Early corticosteroids (dexamethasone and hydrocortisone)	P/NI	Infection (LOS, any)	18	1,349	RR, fixed	1.14 [0.97, 1.34]	High
Fenton 2020	LBW	Late corticosteroids (dexamethasone or hydrocortisone)	P/NI	CP	1	30	RR, fixed	0.44 [0.04, 4.32]	Low ^{1,3,a}
Flenady 2003	Neonates	Radiant warmers	Incubators	Any	2	90	RR, fixed	0.93 [0.66, 1.30]	Low ^{1,3,a}
				CP	1	60	RR, fixed	0.6 [0.16, 2.29]	Low ^{1,3, a}
Foster 2015	Neonates with IV	IV filter	P/NI	CP	2	530	RR, fixed	0.86 [0.59, 1.27]	Moderate ¹
Foster 2017	Term or preterm	Routine oro-or-nasopharyngeal suction at birth	No	Infection (LOS (<7 days), any)	1	509	RR, fixed	0.76 [0.42, 1.36]	Low ¹

Table 1: (continued)

Review ID	Population	Intervention	Comparison	Sepsis	No. of RCTs	No. of patients included in the RCTs	Summary of measure	Estimate (95% CI)	Certainty of evidence, GRADE
Gordon 2017	Neonates with UVC	Early planned UVC removal	Later planned UVC removal	CP	1	210	RR, fixed	0.65 [0.35, 1.22]	Low ^{1,3}
				CoNS	1	210	RR, fixed	0.49 [0.19, 1.26]	Low ^{1,3}
				Other bacteria	1	210	RR, fixed	0.61 [0.21, 1.81]	Low ^{1,3}
Gray 2011	<37 w	Cot-nursing	Incubator	Fungi	1	210	RR, fixed	1.96 [0.18, 21.31]	Low ^{1,3}
Howlett 2019	<37 w or LBW	Inositol (IV followed by PO)	P/NI	Any	2	108	RR, fixed	0.27 [0.01, 6.41]	Low ^{1,3}
		Inositol (repeat dose)	P/NI	Bacterial	2	701	RR, fixed	1.33 [1.00, 1.75]	High
		Inositol (single dose)	P/NI	Bacterial	4	1,067	RR, fixed	1.21 [0.95, 1.54]	Moderate ³
Ibrahim 2011	<37 w	Steroids	P/NI	Bacterial	1	74	RR, fixed	1.46 [0.71, 2.97]	Moderate ^{3,a}
	(primary treatment for hypotension)		P/NI	Bacterial	1	18	RR, fixed	0.33 [0.09, 1.23]	Moderate ^{3,a}
	<37 w (treatment for refractory hypotension)	Steroids	P/NI	Bacterial	1	40	RR, fixed	0.60 [0.20, 1.82]	Low ^{1,3,a}
			Other drug (e.g. inotrope)						
			P/NI	Bacterial	2	65	RR, fixed	1.09 [0.29, 4.10]	Low ^{1,3,a}
Imdad 2013	Neonates	Antiseptics (chlorhexidine)	Antiseptics (salicylic acid powder)	Any	1	213	RR, random	1.11 [0.07, 17.50]	Low ^{1,3,a}
Inglis 2005 (antibiotics for UVC)	Neonates with UVC	Any prophylactic antibiotic	P/NI	Any	1	29	N/A	N/A (no difference between groups/no events)	Low ^{1,3,a}
Inglis 2007 (antibiotics for UAC)	Neonates with UAC	Any prophylactic antibiotic	P/NI	Any	2	212	N/A	Not done (one trial reported no significant difference whereas another trial reported significant difference between groups)	Low ^{1,3,a}
Inglis 2007 (antibiotics for ventilated neonates)	Ventilated neonates	Antibiotic	P/NI	Any	2	N/A	N/A	Meta-analysis for sepsis is N/A (one trial reported no difference in LOS)	Low ^{1,3,a}
Jacobs 2013	>35 w	Therapeutic hypothermia	P/NI	CP	8	1,222	RR, fixed	0.87 [0.60, 1.26]	Moderate ^{1,a}
Jardine 2008	Neonates with CVC	Antibiotics (vancomycin, amoxicillin)	P/NI	Bacterial	2	201	RR, fixed	0.38 [0.18, 0.82]	Moderate ^{1,a}
				Any	2	201	RR, fixed	0.40 [0.20, 0.78]	Moderate ^{1,a}
Kabra 2005	Neonates with UVC	ML-UVCs	SL-UVCs	Suspected	2	56	RR, fixed	0.89 [0.29, 2.78]	Low ^{1,3,a}
Kapoor 2019 (lipid for term and late preterm)	>37 w with surgical problems	Fish oil LE	Non-fish oil LE	CP	2	51	RR, fixed	1.05 [0.47, 2.34]	Very low ^{1,3}
	>37 w cholestasis	Fish oil LE	Non-fish oil LE	Any	3	93	RR, fixed	1.04 [0.53, 2.02]	Low ^{1,3,a}
				Any	2	40	RR, fixed	1.21 [0.50, 2.92]	Very low ^{1,3}

Table 1: (continued)

Review ID	Population	Intervention	Comparison	Sepsis	No. of RCTs	No. of patients included in the RCTs	Summary of measure	Estimate (95% CI)	Certainty of evidence, GRADE
Kapoor 2019 (lipid for preterm)	<37 w	Fish oil LE	Non-fish oil LE	CP	7	774	RR, fixed	1.16 [0.91, 1.48]	Low ^{1,3}
			Another fish oil LE	Any	1	55	RR, fixed	1.69 [0.56, 5.11]	Low ^{1,3}
		Alternative LE	Another alternative-LE	Any	1	59	RR, fixed	1.93 [0.65, 5.73]	Low ^{1,3}
		Alternative LE	Soybean LE	CP	2	164	RR, fixed	1.22 [0.54, 2.78]	Low ^{1,3}
Kelly 2017	Asymptomatic newborns with MSAF	Antibiotics	P/NI	CP	1	250	RD, fixed	-0.01 [-0.07, 0.04]	Low ^{1,3}
				Suspected	1	250	RD, fixed	-0.03 [-0.10, 0.05]	Low ^{1,3}
	Symptomatic newborns with MSAF	Antibiotics	P/NI	CP	3	445	RD, fixed	0.00 [-0.02, 0.03]	Low ^{1,3}
Kuti 2021	Pregnant women, mothers, other caregivers, and HCWs	2% chlorhexidine gluconate	Alcohol hand sanitizer	CP	1	2,932	RR, fixed	2.26 [1.73, 2.93]	Very low ^{1,3}
		4% chlorhexidine gluconate	Triclosan 1%	Any	1	2,932	RR, fixed	2.19 [1.79, 2.69]	Very low ^{1,3}
				Any	1	1916	RR, fixed	6.01 [3.56, 10.14]	Low ^{1,3,a}
Lai 2016 (antimicrobial dressing for CVC)	Neonates with CVC	Chlorhexidine dressing	Polyurethane dressing	CP	1	655	RR, fixed	1.18 [0.53, 2.65]	Moderate ³
				Suspected	1	705	RR, fixed	1.06 [0.75, 1.52]	Moderate ³
Lai 2016 (co bedding)	<37 w, twins	Co-bedding	P/NI	Any	2	59	RR, fixed	0.45 [0.07, 2.86]	Very low ^{1,3,a}
				CP	3	65	RR, fixed	0.84 [0.30, 2.31]	Very low ^{1,3}
Lassi 2019	Neonates	Maternal and neonates care education	P/NI	Infection, including sepsis (any)	2	42,043	RR, random	0.88 [0.72, 1.08]	Moderate ^{1,a}
Lemyre 2016	<37 w	NIPPV for RDS	NCPAP for RDS	Any	2	136	RR, fixed	0.78 [0.36, 1.70]	Moderate ¹
Matviya 2013	<37 w or LBW with symptomatic PDA	Surgical ligation	Medical treatment	Any	1	154	RR, fixed	1.14 [0.62, 2.09]	Low ^{1,3,a}
McCall 2018	<37 w or LBW	Plastic wrap or bag	Routine care	Bacterial	2	830	RR, fixed	0.88 [0.70, 1.10]	Low ^{1,3,a}
		Thermal mattress	Routine care	Bacterial	1	102	RR, fixed	0.92 [0.48, 1.79]	Low ^{1,3,a}
McMullan 2018	Neonates under-going CVC removal	Cephazolin	P/NI	CP	1	88	RR, fixed	0.09 [0.01, 1.60]	Low ^{1,3}
				Suspected	1	88	RR, fixed	0.33 [0.01, 7.97]	Low ^{1,3}
Meyer 2018	<37 w	Respiratory support before cord clamping	P/N	CP	1	150	RR, fixed	1.67 [0.41, 6.73]	Low ^{1,3,a}
Moe-Byrne 2016	<37 w	Glutamine	P/NI	CP	11	2,815	RR, fixed	0.94 [0.86, 1.04]	Moderate ^{2,4}
Moon K 2020	>34 w	Early PN	Late PN	CP	1	209	RR, fixed	0.22 [0.05, 1.01]	Low ^{1,3}
				Any	1	209	RR, fixed	0.53 [0.32, 0.89]	Low ^{1,3}
Morgan 2013	VLBW or <32 w	Early trophic feeding	No trophic feeds	CP	3	237	RR, fixed	1.06 [0.72, 1.56]	Low ^{1,3,a}
Morgan 2014	VLBW or <32 w	Delayed feeds (>4 days after birth)	Early feeds	CP	2	457	RR, fixed	1.27 [0.95, 1.70]	Low ^{1,3,a}
Muelbert 2019	<37 w	Smell and taste of milk	P/NI	CP	1	51	RR, fixed	2.46 [0.27, 22.13]	Low ^{1,3}
Nasuf 2018	<37 w	OPC	P/NI	CP	6	335	RR, fixed	0.86 [0.56, 1.33]	Very low ^{1,3}

Table 1: (continued)

Review ID	Population	Intervention	Comparison	Sepsis	No. of RCTs	No. of patients included in the RCTs	Summary of measure	Estimate (95% CI)	Certainty of evidence, GRADE
Ng 2016	<37 w at risk of CLD	Salbutamol	P/NI	CP	1	173	RR, fixed	1.06 [0.54, 2.06]	Low ^{1,3,a}
Ng 2017	<37 w	Cromolyn sodium	P/NI	Any	2	64	RR, fixed	0.82 [0.41, 1.63]	Low ^{1,3,a}
Ng 2019	<37 w	Hydrolyzed formula	Standard formula	CP	1	60	RR, fixed	1.50 [0.27, 8.34]	Low ^{1,3,a}
Oddie 2017	<32 w or VLBW	Slow feeding advancement	Faster feed advancement	CP	8	3,392	RR, fixed	1.15 [1.00, 1.32]	Low ^{1,3}
Ohlsson 2020 (ibuprofen for prevention of PDA)	<37 w and LBW EOS	Ibuprofen	P/NI	Bacterial	2	201	RR, fixed	2.70 [1.10, 6.59]	Low ^{1,3,a}
Ohlsson 2020 (ibuprofen for treatment of PDA)	<37 w or LBW	Ibuprofen	Indomethacin	Bacterial	7	735	RR, fixed	1.22 [0.84, 1.76]	Low ^{1,3,a}
		Ibuprofen, PO	Indomethacin, IV/PO	Bacterial	2	53	RD, fixed	0.03 [-0.22, 0.28]	Low ^{1,3,a}
		Ibuprofen, PO	Ibuprofen, IV	Bacterial	3	236	RR, fixed	0.82 [0.54, 1.25]	Low ^{1,3,a}
		High dose ibuprofen	Standard dose ibuprofen	Bacterial	1	70	RR, fixed	0.93 [0.51, 1.68]	Low ^{1,3,a}
Ohlsson 2020 (IVIG for prevention of infection)	Preterm and LBW	Early ibuprofen IVIG	Expectant P/NI	Bacterial CP CP 'serious infection'	1 10 16	105 3,975 4,986	RR, fixed RR, fixed RR, fixed	0.90 [0.58, 1.41] 0.85 [0.74, 0.98] 0.82 [0.74, 0.92]	Moderate ^{3,a} Moderate ² Moderate ²
Ohlsson 2020 (paracetamol for PDA)	PDA <37 w or LBW	Paracetamol	Ibuprofen P/NI	Bacterial	4	472	RR, fixed	0.88 [0.64, 1.21]	Low ^{1,3,a}
			Ibuprofen P/NI	Bacterial	1	48	RR, fixed	1.45 [0.36, 5.79]	Moderate ^{3,a}
			Indomethacin P/NI	Bacterial	2	277	RR, fixed	1.14 [0.59, 2.19]	Low ^{1,3,a}
Ohlsson 2020 (early EPO)	<37 w and/or LBW	Erythropoietin	P/NI	CP	12	2,180	RR, fixed	0.87 [0.74, 1.02]	Moderate ^{1,a}
Onland 2017 (late inhaled steroid for prevention of BPD)	<36 w	Darbepoetin alfa Inhaled corticosteroids (budesonide, fluticasone, beclomethasone)	P/NI P/NI	CP Any	1 5	62 107	RR, fixed RR, fixed	1.13 [0.38, 3.30] 0.90 [0.50, 1.64]	Low ^{1,3,a} Low ^{1,3,a}
Onland 2017 (systemic steroid for prevention of BPD)	Preterm at risk of BPD	Lower cumulative dexamethasone dose	Higher cumulative dexamethasone dose	CP Suspected	6 2	230 72	RR, fixed RR, fixed	0.91 [0.60, 1.38] 1.03 [0.62, 1.70]	Low ^{1,3,a} Low ^{1,3,a}
Osborn 2007	<37 w	Thyroid hormone therapy	P/NI	Any	3	278	RR, fixed	0.78 [0.53, 1.16]	Moderate ^{3,a}
Osborn 2018	Neonates requiring PN	High amino acid High amino acid (at the start) High amino acid (at max dose)	Low Low Low P/NI	Bacterial Bacterial Bacterial Any CP Bacterial Fungal Any	15 5 1 12 12 8 6 3	1,255 319 127 5,425 5,425 3,565 3,266 564	RR, fixed RR, fixed RR, fixed RR, fixed RR, fixed RR, fixed RR, fixed RR, fixed	0.96 [0.79, 1.18] 0.94 [0.65, 1.38] 0.94 [0.63, 1.41] 0.80 [0.72, 0.89] 0.83 [0.72, 0.94] 0.86 [0.74, 1.00] 0.23 [0.10, 0.54] 0.25 [0.14, 0.46]	Low ^{1,4,a} Low ^{1,3,a} Low ^{1,3,a} Moderate ^{3,a} Low ^{1,4} Low ^{1,2} Low ^{1,2} Low ^{1,2} Low ^{1,3}
Pammi 2020	<37 w	Lactoferrin Lactoferrin + probiotics							

Table 1: (continued)

Review ID	Population	Intervention	Comparison	Sepsis	No. of RCTs	No. of patients included in the RCTs	Summary of measure	Estimate (95% CI)	Certainty of evidence, GRADE
Pfister 2009	<37 w or LBW	Protein with surfactant	Protein-free surfactant	Bacterial	1	319	RR, fixed	0.28 [0.11, 0.72]	Low ^{1,3}
Premkumar 2019	<37 w	Human milk fortifier		Fungal	2	494	RR, fixed	0.24 [0.08, 0.71]	Low ^{1,3}
Quigley 2019	<37 w or LBW	Formula	Bovine milk fortifier	CP	1	1,036	RR, fixed	1.00 [0.87, 1.14]	Moderate ^{1,a}
Rojas-Reyes 2012	<32 w	Prophylactic surfactant	Donor human milk	CP	1	125	RR, fixed	0.54 [0.25, 1.21]	Low ^{1,3}
Schulzke 2014	<32 w or VLBW	Nebulized pentoxifylline	Rescue surfactant	Bacterial	5	1,025	RR, fixed	0.94 [0.79, 1.12]	Moderate ^{1,a}
(pentoxifylline for BPD)			P/NI	CP	6	2,438	RR, fixed	0.83 [0.64, 1.08]	Low ^{1,2,a}
Schulzke 2014	<37 w	Physical activity			1	100	RR, fixed	1.22 [0.63, 2.36]	Low ^{1,3,a}
(physical activity for bone mineralization)									
Seger 2009	<37 w	Animal surfactant	P/NI	CP	1	16	N/A	The trial reported no difference in sepsis between the groups	Low ^{1,3,a}
Shah 2008	Term or preterm requiring CVC	Heparin infusion	P/NI	Any	4	1,012	RR, fixed	1.14 [0.87, 1.48]	Moderate ^{1,a}
Shah 2009	<32 w or VLBW	Anti-staphylococcal immunoglobulin A-21	P/NI	Any	3	477	RR, fixed	0.82 [0.43, 1.57]	High ^a
				CoNS	2	2,488	RR, fixed	1.07 [0.94, 1.22]	Moderate ^{1,a}
				Other bacterial	2	2,488	RR, fixed	0.87 [0.72, 1.06]	Moderate ^{1,a}
				Any	2	2,488	RR, fixed	1.00 [0.91, 1.09]	Moderate ^{1,a}
				CoNS	1	206	RR, fixed	0.86 [0.32, 2.28]	Moderate ^{3,a}
				Other bacterial	1	206	RR, fixed	0.93 [0.53, 1.64]	Moderate ^{3,a}
				Any	1	206	RR, fixed	0.93 [0.54, 1.62]	Moderate ^{3,a}
				CP	6	1,121	RD, fixed	1.17 [0.99, 1.38]	Moderate ^{1,a}
Shah 2017 (prevention of BPD)	VLBW neonates	Early inhaled steroids	P/NI	Any	1	278	RR, fixed	1.04 [0.73, 1.49]	Low ^{1,3,a}
Shah 2017 (prevention of BPD)	VLBW or <32 w	Inhaled steroid	Systemic corticosteroids	CP	2	368	RR, fixed	1.07 [0.79, 1.45]	Low ^{1,3,a}
Shah 2017 (treatment of BPD)	VLBW or <32 w	Inhaled steroid	Systemic corticosteroids	Suspected	1	52	RR, random	1.25 [1.00, 1.56]	Low ^{1,3,a}
Shah 2018	Neonates requiring RBC transfusion	Short transfusion	Longer transfusion						
Sharif 2020	<32 w	Probiotics	P/NI	CP	47	9,762	RR, fixed	0.89 [0.82, 0.97]	Moderate ¹
Sinclair 2011	<32 w or VLBW	Early parenteral lipid	Delayed	CP	1	29	RR, fixed	0.81 [0.13, 5.01]	Low ^{1,3,a}
		Insulin infusion for hyperglycemia	Standard care	CP	1	386	RR, fixed	0.92 [0.63, 1.34]	Low ^{1,3,a}

Table 1: (continued)

Review ID	Population	Intervention	Comparison	Sepsis	No. of RCTs	No. of patients included in the RCTs	Summary of measure	Estimate (95% CI)	Certainty of evidence, GRADE
Singh 2015	<32 w	Bovine lung lavage surfactant extract (prophylaxis)	Modified bovine minced lung surfactant extract	Bacterial	2	1,123	RR, fixed	1.08 [0.91, 1.28]	Moderate ^{1,a}
	<37 w	Bovine lung lavage surfactant extract (rescue)	Modified bovine minced lung surfactant extract	Bacterial	6	2,228	RR, fixed	1.00 [0.87, 1.15]	Moderate ^{1,a}
Sinha 2015	Term or late preterm	Modified bovine minced lung (rescue) Chlorhexidine for skin or cord care, in hospital Chlorhexidine for skin or cord care in the community	Porcine minced lung P/NI P/NI	Bacterial Any Any	6 2 1	526 13,033 203	RR, fixed RR, fixed RR, fixed	1.13 [0.87, 1.46] 0.98 [0.82, 1.16] 0.69 [0.49, 0.95]	Low ^{1,3,a} Moderate ^{1,a} Moderate ^{3,a}
Soll 1997	<30 w	Animal surfactant	P/NI	CP	4	914	RR, fixed	1.06 [0.81, 1.38]	Low ^{1,2,a}
Soll 2009	<30 w <1,250 g with RDS or at risk	Multiple doses of surfactant (synthetic surfactant, curosurf)	Single dose	Bacterial	2	1,169	RR, fixed	0.85 [0.70, 1.04]	Moderate ^{1,a}
Spence 1999	Neonates who required endotracheal intubation	Nasal intubation	Oral	CP Suspected	1 1	86 91	RR, fixed RR, fixed	1.0 [0.15, 6.78] 2.1 [0.89, 4.91]	Low ^{1,3,a} Low ^{1,3,a}
Subramaniam 2016	<32 w or VLBW	Prophylactic CPAP	Supportive care	Any	3	568	RR, fixed	1.04 [0.64, 1.69]	Moderate ^{1,a}
Symington 2006		Assisted ventilation	Assisted ventilation	Any	1	425	RR, random	0.59 [0.33, 1.04]	Low ^{1,3,a}
Taylor 2015	<37 w Neonates with CVC	NIDCAP Antibiotic lock	P/NI P/NI	Any CP	1 3	21 271	RR, fixed RR, fixed	0.92 [0.71, 1.18] 0.15 [0.06, 0.40]	Low ^{1,3,a} High
Ullman 2013	Neonate with any IV/arterial catheter	Less frequent for intravenous administration set replacement	Frequent	Any CP	3 1	271 148	RR, fixed RR, fixed	0.25 [0.12, 0.49] 0.58 [0.25, 1.35]	High Low ^{1,3,a}
Ungerer 2004	Term born to mothers with risk of infection (GBS)	Prophylactic antibiotic (intramuscular penicillin G)	Selective antibiotics use	CP	1	67	RD, fixed	0.00 [-0.06, 0.06]	Low ^{1,3,a}
	Term with other risk factors, PROM, fever)	Prophylactic antibiotics (intramuscular penicillin and kanamycin)	Selective antibiotics use	CP	1	49	RR, fixed	0.12 [0.01, 2.04]	Low ^{1,3,a}
Walsh 2019	<37 w	Iodine supplementation	P/NI	Any	1	1,259	RR, fixed	1.09 [0.96, 1.24]	High ^a
Webster 2003	Visitors	Over gowns	No gowns	Any	4	3,979	RR, random	0.95 [0.40, 2.23]	Low ^{1,2,a}
Whitelaw 2001	Infants <3 months with severe IVH	Acetazolamide + furosemide	Standard therapy	CNS infection	1	117	RR, fixed	1.20 [0.57, 2.52]	Low ^{1,3,a}
Whitelaw 2017	Infants with IVH	Repeated lumbar/Ventricular tapping	Conservative	CNS infection	2	195	RR, fixed	1.73 [0.53, 5.67]	Low ^{1,3}

Table 1: (continued)

Review ID	Population	Intervention	Comparison	Sepsis	No. of RCTs	No. of patients included in the RCTs	Summary of measure	Estimate (95% CI)	Certainty of evidence, GRADE
Wilkinson 2016	<37 w	HFNC	CPAP (primary respiratory support after birth)	CP	4	439	RR, fixed	1.29 [0.66, 2.54]	Low ^{1,3,a}
			CPAP (post-extubation)	CP	2	529	RR, fixed	0.92 [0.59, 1.43]	Low ^{1,3,a}
			NIPPV (primary respiratory support after birth)	CP	1	76	RR, fixed	1.33 [0.32, 5.56]	Low ^{1,3,a}
Zupan 2004	Neonates	Topical cord antiseptics	P/NI	Any	7	N/A	RR, fixed	No meta-analysis was conducted; No sepsis was noted in either arm in all the trials	Low ^{1,a}

^aCochrane reviews, where evidence's certainty was unavailable, were evaluated by two independent authors based on Grading of Recommendations, Assessment, Development and Evaluation (1, Risk of bias; 2, heterogeneity; 3, imprecision/wide confidence interval; 4, publication bias (asymmetry in funnel plot)/other reasons; 5, indirectness), with conflicts resolved by consensus and discussion with the third author; The effect estimates highlighted in bold represent statistically significant differences between the groups; BPD, bronchopulmonary dysplasia; CC, catheter colonization; CLD, chronic lung disease; CoNS, coagulase-negative staphylococcus; CP, culture-proven; CPAP, continuous positive airway pressure; CVC, central venous catheter; DBM, donor breast milk; ESAs, erythropoiesis-stimulating agents; GBS, group B *streptococcus*; G-CSF, granulocyte colony stimulating factor; GI, gastrointestinal; GM-CSF, granulocyte-macrophage colony stimulating factor; GRADE, grading of recommendations, assessment development and evaluation; HFNC, high flow nasal cannula; IVH, intraventricular hemorrhage; IVIG, intravenous immunoglobulin; IV, intravenous infusions; LBW, low birth weight; LE, lipid emulsion; ML-UVCs, multiple lumen umbilical venous catheters; MSAF, meconium-stained amniotic fluid; N/A: not available; NCPAP, nasal continuous positive airway pressure; NICU, neonatal intensive care unit; NIDCAP, neonates individualized developmental care and assessment program; NIPPV, nasal intermittent positive pressure ventilation; OPC, oropharyngeal colostrum; PCVC, percutaneous central venous catheters; PDA, patent ductus arteriosus; PO, per oral; PN, parenteral nutrition; P/NI: placebo/no intervention; PNCS, postnatal corticosteroids; PSBI, possible serious bacterial infection; RBCs, red blood cells; RDS, respiratory distress syndrome; RCTs, randomised clinical trials; RR, risk ratio; RD, risk difference; SL-UVCs, single lumen umbilical venous catheters; UAC, umbilical artery catheters; UVC, umbilical venous catheters; VLBW, very low birth weight.

secondary outcomes of the study. The subgroup and sensitivity analysis details are provided in Table 2 of the Supplementary Information. Most reviews evaluated one intervention, but few evaluated two or more interventions. Overall, 101 included reviews evaluated 141 interventions with LOS as one of the primary or secondary outcomes. Few reviews assessed various types of LOS (any, bacterial or fungal), and the majority of them evaluated LOS as culture-positive sepsis ($n=65$), followed by any sepsis ($n=55$), bacterial sepsis ($n=34$), and fungal sepsis ($n=10$).

The number of RCTs in 101 reviews ranged from one to 47. The number of neonatal participants in each trial ranged from 16 to 42,043. In total, there were 612 RCTs involving 193,713 term and preterm neonates. Of the 101 included reviews, 28 provided GRADE assessments for one or more LOS outcomes. For the remainder of the reviews ($n=73$), two study authors performed the GRADE assessment independently for all the primary and secondary outcomes. The details are provided in Table 1. The interventions on LOS were categorized into five categories based on the effectiveness and certainty of the evidence, as summarized in Table 2.

Effective interventions

High-quality evidence showed reduced LOS (any sepsis, RR (95% CI) 0.15 (0.06, 0.40) [103] or culture-proven sepsis, RR (95% CI) 0.25 (0.12, 0.49)) [103] when antibiotic lock therapy, compared with placebo, was used for neonates with central venous catheters (CVC). We found no high-quality, effective intervention for reducing bacterial or fungal sepsis.

Possible effective interventions

Any sepsis

Moderate-quality evidence showed antibiotic prophylaxis for neonates with CVC (RR (95% CI) 0.38 (0.18, 0.82)) [47], vancomycin prophylaxis for neonates requiring parenteral nutrition or neonates with CVC (RR (95% CI) 0.11 (0.05, 0.24)) [28], chlorhexidine for skin or cord care (RR (95% CI) 0.69 (0.49, 0.95) [97], and kangaroo care for low birth weight babies (RR (95% CI) 0.50 (0.36, 0.69)) [27] reduced any LOS.

Culture-proven sepsis

Moderate-quality evidence showed intravenous immunoglobulin prophylaxis (RR (95% CI) 0.85 (0.74, 0.98)) [72] for preterm infants and probiotic supplementation for VLBW

infants (RR (95% CI) 0.89 (0.82, 0.97) [94]) reduced culture-proven sepsis.

Bacterial sepsis

Moderate-quality evidence showed antibiotic prophylaxis for neonates with CVC (RR (95% CI) 0.40 (0.20, 0.78)) [47] reduced bacterial LOS.

Fungal sepsis

Moderate-quality evidence showed systemic antifungal prophylaxis VLBW infants (RR (95% CI) 0.43 (0.31, 0.59)) [23] reduced fungal LOS.

Ineffective interventions

High-quality evidence showed no difference in LOS with use of late postnatal corticosteroids for bronchopulmonary dysplasia prevention (RR (95% CI) 1.14 (0.97, 1.34)) [32], heparin infusion for neonates with CVC (RR (95% CI) 0.82 (0.43, 1.57)) [90], iodine supplementation (RR (95% CI) 1.09 (0.96, 1.24)) [106], and inositol supplementation (RR (95% CI) 1.33 (1.00, 1.75)) [40].

Possible ineffective interventions and no conclusions possible

Many interventions where the evidence certainty was moderate-quality with lack of effectiveness (or harm) and where the evidence certainty was low-to very low-quality were categorized in these groups as listed in Table 2.

Discussion

Summary of main results

This overview included 101 high-quality Cochrane reviews involving 612 RCTs and 193,713 neonates that evaluated 141 interventions.

Overall completeness and applicability of evidence

Only 101 reviews (around 20% of 496 reviews in the Cochrane Neonatal) were eligible for inclusion in the overview. Although there were 101 reviews involving 612 RCTs

Table 2: Summary of effectiveness of interventions on late-onset neonatal sepsis.

	LOS, any (suspected or confirmed)	LOS, culture-proven	LOS, bacterial	LOS, fungal
Effective interventions	– Antibiotic lock for CVC	– Antibiotic lock for CVC	– None	– None
Possible effective interventions	– Kangaroo care for LBW babies	–	–	–
	– Vancomycin prophylaxis for VLBW babies requiring PN or babies with CVC requiring PN	– IVIG prophylaxis (<37 w or LBW)	– Antibiotic prophylaxis for neonates with CVC	– Systemic antifungal prophylaxis
	– Antibiotic prophylaxis for neonates with CVC	– Probiotics supplementation (<32 w)		
	– Chlorhexidine for skin or cord care (in community setting) ^a			
Ineffective interventions	– Late postnatal corticosteroids for BPD prevention	– None	– Inositol supplementation (>1 dose) (<37 w)	– None
	– Heparin infusion for CVC			
	– Iodine supplementation (<37 w)			
Possible ineffective interventions	– Vitamin E supplementation			
	– Supplemental feeds with bottle vs. not (<37 w)	– PCVC vs. PVC for parenteral nutrition	– Animal-derived surfactant vs. protein-free synthetic surfactant	– Topical ointment/cream
	– Early postnatal corticosteroids for BPD prevention	– Glutamine supplementation for severe GI disease or preterm infants	– Topical ointment/cream	– Inositol supplementation (1 dose) (<37 w)
	– Chlorhexidine vs. Polyurethane dressing for CVC	– Intravenous inline filter	– Steroid as primary treatment for hypotension (<37 w)	– Early vs. expectant ibuprofen for PDA (<37 w)
	– Maternal education bundle	– Chlorhexidine vs. Polyurethane dressing for CVC	– Early vs. expectant ibuprofen for PDA (<37 w)	
	– NIPPV vs. NCPAP for RDS (<37 w)	– Early erythropoietin prophylaxis (<37 w)	– Paracetamol prophylaxis for PDA (<37 w)	
	– Thyroid hormone prophylaxis	– Protein vs. Protein-free surfactant	– Anti-staphylococcal immunoglobulin prophylaxis	
	– Animal-derived surfactant for RDS (<37 w)	– Formula vs. donor human milk (<37 w)	– Bovine lung lavage vs. modified bovine minced surfactant for RDS (rescue or prophylaxis)	
	– Anti-staphylococcal immunoglobulin prophylaxis	– Early inhaled steroids for CLD prevention	– Multiple vs. single dose surfactant (<30 w)	
	– Chlorhexidine for skin or cord care (in hospital setting) ^a			
No conclusions possible	– Prophylactic NCPAP (<32 w)	– Routine monitoring of gastric residuals	– Early vs. delayed animal-derived surfactant	– Prophylactic oral/topical antifungal
	– Prophylactic intravenous vs. intramuscular vitamin K	– Criterion-based monitoring of gastric residuals	– Insulin infusion for hyperglycemia	– Prophylactic vs. systemic antifungal
	– Aqueous vs. oil-based vitamin E	– Late ESAs	– Topical oil	– Insulin infusion for hyperglycemia
	– G-CSF or GM-CSF prophylaxis	– Antimicrobial-impregnated CVC	– Vancomycin prophylaxis for VLBW babies requiring PN via CVC	– Topical oil
	– Early discharge on gavage feeds	– G-CSF or GM-CSF prophylaxis	– Ibuprofen prophylaxis for PDA	– Early vs. late UVC removal
	– Vitamin a (any dose, high dose) supplementation (<32 w)	– Topical ointment/cream or oil	– Selenium supplementation (<37 w)	

Table 2: (continued)

LOS, any (suspected or confirmed)	LOS, culture-proven	LOS, bacterial	LOS, fungal
- Radiant warmer vs. incubator care	- High vs. low protein intake	- Early vs. late UVC removal	- Lactoferrin or lactoferrin with
- Routine oro-or-nasopharyngeal suction at birth	- Radiant warmer vs. incubator care	- Steroid vs. inotrope primary treatment for hypotension (<37 w)	probiotics supplementation (<37 w)
- Cot-nursing vs. incubator care	- Early vs. late UVC removal	- Steroid for refractory hypotension (<37 w)	
- Cord antiseptics care (chlorhexidine vs. salicylic acid)	- Fish- vs. non-fish-oil LE (<37 w, <37 w with surgical problems)	- Plastic wrap or thermal mattress during delivery (<37 w)	
- Antibiotics prophylaxis for UAC/UVC/Ventilated neonates	- Antibiotics for babies born through MSAF	- Ibuprofen vs. Indomethacin for PDA treatment (<37 w)	
- ML-UVCs vs. SL-UVCs	- Co-bedding preterm twin babies	- Ibuprofen: PO vs. IV and high dose vs. standard dose (<37 w)	
- Fish- vs. non-fish-oil LE (<37 w with surgical problems or cholestasis)	- Cephalosol for CVC removal	- Paracetamol vs. Ibuprofen or indomethacin for PDA (<37 w)	
- Antibiotics for asymptomatic babies born through MSAF	- Respiratory support before cord clamping	- High vs. low amino acid PN (overall, at initiation, and at maximum dose)	
- Co-bedding preterm twin babies	- Early vs. Late PN (>34 w)	- Lactoferrin or lactoferrin with probiotics supplementation (<37 w)	
- Surgical ligation vs. medical treatment for PDA (<37 w)	- Trophic feeds (<32 w)	- Prophylactic vs. rescue surfactant (<32 w)	
- Cephalosol for CVC removal	- Delayed feeding (4/> days after birth) in <32 w	- Bovine vs. porcine surfactant for RDS (rescue) (<37 w)	
- Early vs. Late PN (>34 w)	- Smell and taste of milk (<37 w)		
- Cromolyn sodium nebulization (<37 w, CLD)	- Oropharyngeal colostrum (<37 w)		
- Late inhaled corticosteroids (<37 w, BPD)	- Salbutamol (<37 w, CLD)		
- High vs. low dexamethasone (<37 w, BPD)	- Hydrolyzed formula feeding (<37 w)		
- Lactoferrin or lactoferrin with probiotics supplementation (<37 w)	- Slow vs. faster feed advancement (<32 w)		
- Short binasal prongs vs. single nasal prongs for NCPAP support	- Darbepoetin prophylaxis (<37 w)		
- Inhaled vs. systemic steroids for CLD prevention	- High vs. low dexamethasone (<37 w, BPD)		
- Short vs. long RBC transfusion	- Lactoferrin supplementation (<37 w)		
- Nasal vs. Oral intubation	- Human milk vs. bovine milk fortifier (<37 w)		
- Prophylactic NCPAP vs. Assisted ventilation (<32 w)	- Short binasal prongs vs. single nasal prongs for NCPAP support		
- NIDCAP (<37 w)	- Nebulized pentoxifylline for CLD (<32 w)		
- Gowns for attendants of the neonates	- Physical activity for bone mineralization (<37 w)		
- Acetazolamide and fructose for babies with severe IVH	- Inhaled vs. systemic steroids for CLD prevention or treatment		
	- Early vs. late parenteral lipids (<32 w)		
	- Insulin infusion for hyperglycemia (<32w)		

Table 2: (continued)

LOS, any (suspected or confirmed)	LOS, culture-proven	LOS, bacterial	LOS, fungal
– Repeated lumbar/Ventricular tapping for infants with IVH – Alcohol vs. chlorhexidine for hand hygiene	– Animal-derived surfactant for all <30 w preterm infants – Nasal vs. Oral intubation – Less frequent vs. frequent intravenous administration set replacement – Antibiotic prophylaxis for neonates with risk factors – HFNC vs. CPAP (primary support or post-extubation) OR NIPPV (primary support) – Alcohol vs. chlorhexidine for hand hygiene		

^aMothers of infants were also participants and received vaginal chlorhexidine washes. The interventions are categorized into 5 categories based on the effectiveness of the intervention and the certainty of the evidence as (1) effective interventions (high certainty in the evidence of effectiveness) (2) possible effective interventions (moderate certainty in the evidence of effectiveness) (3) ineffective interventions (high certainty in the evidence of lack of effectiveness or harm) (4) possible ineffective interventions (moderate certainty in the evidence of lack of effectiveness or harm), and (5) no conclusions possible (low or very low certainty in evidence). CLD, chronic lung disease; CVC, central venous catheter; ESA, erythropoietin stimulating agents; G-CSF, granulocyte colony-stimulating factor; GI: gastrointestinal; GM-CSF, granulocyte monocyte colony-stimulating factor; HIE: hypoxic-ischemic encephalopathy; HFNC, high flow nasal cannula; IV, intravenous; IVH, intraventricular hemorrhage; LBW, low birth weight; LE, lipid emulsion; LOS, late-onset sepsis; ML-UVCs, multiple lumen umbilical venous catheters; MSAF, meconium-stained amniotic fluid; NCPAP, nasal continuous positive airway pressure; NIDCAP, neonates individualized developmental care and assessment program; NIPPV, nasal intermittent positive pressure ventilation; PCVC, percutaneous central venous catheter; PDA, patent ductus arteriosus; PN, parenteral nutrition; PO, per oral; PVC, peripheral venous cannula; RDS, respiratory distress syndrome; SL-UVCs, single lumen umbilical venous catheters; UAC, umbilical artery catheters; UVC, umbilical venous catheters.

and 193,713 neonates, the body of evidence was reduced as not all the included reviews reported the primary outcome of interest of the overview. Further, the data on secondary outcomes was limited as only one-third of reviews reported bacterial sepsis, and 10% reported fungal sepsis. All reviews explicitly defined various types of LOS; however, there was some variation in the definition between and within the trials included in the reviews.

The overview's scope was limited to studying the effectiveness of interventions on LOS; hence, we did not examine the effects of interventions on other outcomes. This is important as some interventions found ineffective in reducing sepsis may have considerable influence on other important outcomes. For example, in this overview, we found high-quality evidence that late postnatal corticosteroids for preventing bronchopulmonary dysplasia [32] are ineffective in reducing sepsis; however, they are beneficial in lowering extubation failure, death, or bronchopulmonary dysplasia, which are other important outcomes that were not studied [32]. Furthermore, clinicians must consider many other factors, such as effect size, harms, cost-effectiveness, and generalizability, by referring to the original studies or individual Cochrane reviews for considering any therapy in neonatal practice [112–114]. For example, though the overview found moderate-quality evidence of effectiveness that probiotics supplementation reduces LOS in very preterm infants, clinicians must consider other factors, such as harms, including probiotics-associated sepsis, anticipated benefit, quality assurance, cost-effectiveness, and so on [112–115].

Quality of evidence

All the 101 included Cochrane reviews were high-quality and rated a low risk of bias based on the AMSTAR 2 tool [9]. The risk of bias of included RCTs in those reviews, assessed using the approach outlined in the Cochrane Handbook for Systematic Reviews of Interventions, was variable; however, it was incorporated in the GRADE approach for evaluating the certainty of the evidence. The most common reason for downgrading the quality was study limitations, followed by imprecision and heterogeneity.

Strengths and limitations

This is a first overview providing a comprehensive summary of Cochrane reviews evaluating interventions to reduce LOS and shall be helpful for clinicians and researchers to aid in choosing interventions for reducing LOS. Also, the overview

examines the certainty of the evidence for sepsis-related outcomes that were not evaluated in nearly three-fourths of the Cochrane reviews. Further, the overview provides a rigorous assessment of the evidence depending on the effectiveness of interventions and the certainty of the evidence. The search was comprehensive, and the reporting was transparent, based on the PRIO-harms guideline.

The overview focuses on interventions to reduce LOS and only evaluates sepsis-related outcomes. It did not assess other important outcomes, so clinicians should be careful while inferring this data and consider various other factors, as mentioned above, while applying the evidence to their clinical practice. In addition, the overview did not assess early-onset sepsis as the overview focuses on neonatal interventions and not childbirth or pregnancy interventions. Finally, the overview included only Cochrane reviews, but we realize that most interventions were likely assessed in the Cochrane reviews.

To conclude, the overview summarizes the evidence from the Cochrane reviews assessing interventions in the neonatal period for reducing LOS. Clinicians, researchers, policymakers, and consumers can utilize it for evidence translation and decision-making. Clinicians, however, are recommended to assess the effects of the interventions on other outcomes, including harms, and consider other aspects, such as feasibility, generalizability, anticipated benefit based on the effect size, and others, while translating evidence into clinical practice.

Acknowledgments: Authors would like to acknowledge Ahmed Hamed for initial screening of the articles for the systematic review.

Research funding: No specific funding was sought for this project. Dr. Razak's research is supported by Monash University, Australia.

Author contributions: Abdul Razak conceptualized and designed the study, oversaw the conduct of the review, assessed the quality of systematic reviews, provided intellectual content, drafted the manuscript and approved the final version. Javed Ahmed co-conceptualized and designed the study, performed the initial screening of the articles, abstracted the data, assessed the quality of systematic reviews, revised the manuscript and approved the final version. Omar Ibrahim Alhaidari co-conceptualized and designed the study, performed the screening of the articles, abstracted the data, revised the manuscript and approved the final version. All authors have accepted responsibility for the entire content of this manuscript and approved its submission.

Conflict of interest: The authors have indicated they have no potential conflict of interest to disclose.

Informed consent: Not applicable.

Ethical approval: Not applicable.

Protocol registration: The protocol for the systematic review was registered with PROSPERO (registration number CRD42020192513), the international prospective register for systematic reviews.

References

1. Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, et al. The third international consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA* 2016; 315:801–10.
2. Wynn JL, Wong HR, Shanley TP, Bizzarro MJ, Saiman L, Polin RA. Time for a neonatal-specific consensus definition for sepsis. *Pediatr Crit Care Med* 2014;15:523–8.
3. Bizzarro MJ, Raskind C, Baltimore RS, Gallagher PG. Seventy-five years of neonatal sepsis at Yale: 1928-2003. *Pediatrics* 2005; 116:595–602.
4. Shim GH, Kim SD, Kim HS, Kim ES, Lee HJ, Lee JA, et al. Trends in epidemiology of neonatal sepsis in a tertiary center in Korea: a 26-year longitudinal analysis, 1980-2005. *J Kor Med Sci* 2011;26: 284–9.
5. Dong Y, Speer CP. Late-onset neonatal sepsis: recent developments. *Arch Dis Child Fetal Neonatal Ed* 2015;100: F257–63.
6. Reinhart K, Daniels R, Kissoon N, Machado FR, Schachter RD, Finfer S. Recognizing sepsis as a global health priority — a WHO resolution. *N Engl J Med* 2017;377:414–7.
7. Bougioukas KI, Liakos A, Tsapas A, Ntzani E, Haidich AB. Preferred reporting items for overviews of systematic reviews including harms checklist: a pilot tool to be used for balanced reporting of benefits and harms. *J Clin Epidemiol* 2018;93: 9–24.
8. Pollock M, Fernandes RM, Becker LA, Pieper DLH. Chapter V: overviews of reviews. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al, editors. *Cochrane handbook for systematic reviews of interventions* version 6.2 (updated February 2021). Cochrane; 2021.
9. Shea BJ, Reeves BC, Wells G, Thuku M, Hamel C, Moran J, et al. Amstar 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ* 2017;358:1–9.
10. Schunemann HJ, Guyatt G, Oxman A, editors. *GRADE handbook for grading quality of evidence and strength of recommendations*. West Sussex, United Kingdom: Cochrane; 2015.
11. Abiramalatha T, Thanigainathan S, Ninan B. Routine monitoring of gastric residual for prevention of necrotising enterocolitis in preterm infants. *Cochrane Database Syst Rev* 2019;7:Cd012937.
12. Aher SM, Ohlsson A. Late erythropoiesis-stimulating agents to prevent red blood cell transfusion in preterm or low birth weight infants. *Cochrane Database Syst Rev* 2020;1:1–87.
13. Ainsworth S, McGuire W. Percutaneous central venous catheters versus peripheral cannulae for delivery of parenteral nutrition in neonates. *Cochrane Database Syst Rev* 2015;10:1–25.
14. Ardell S, Offringa M, Ovelman C, Soll R. Prophylactic vitamin K for the prevention of vitamin K deficiency bleeding in preterm neonates. *Cochrane Database Syst Rev* 2018;2:1–41.
15. Ardell S, Pfister RH, Soll R. Animal derived surfactant extract versus protein free synthetic surfactant for the prevention and treatment of respiratory distress syndrome. *Cochrane Database Syst Rev* 2015;8:1–49.
16. Austin N, Cleminson J, Darlow BA, McGuire W. Prophylactic oral/topical non-absorbed antifungal agents to prevent invasive fungal infection in very low birth weight infants. *Cochrane Database Syst Rev* 2015;2015:Cd003478.
17. Bahadue FL, Soll R. Early versus delayed selective surfactant treatment for neonatal respiratory distress syndrome. *Cochrane Database Syst Rev* 2012;11:1–53.
18. Balain M, Oddie SJ, McGuire W. Antimicrobial-impregnated central venous catheters for prevention of catheter-related bloodstream infection in newborn infants. *Cochrane Database Syst Rev* 2015;9:1–21.
19. Bottino M, Cowett RM, Sinclair JC. Interventions for treatment of neonatal hyperglycemia in very low birth weight infants. *Cochrane Database Syst Rev* 2011;1:1–35.
20. Brion LP, Bell EF, Raghuveer TS. Vitamin E supplementation for prevention of morbidity and mortality in preterm infants. *Cochrane Database Syst Rev* 2003;4:1–258.
21. Brown J, Moe-Byrne T, McGuire W. Glutamine supplementation for young infants with severe gastrointestinal disease. *Cochrane Database Syst Rev* 2014:Cd005947.
22. Carr R, Modi N, Doré C. G-CSF and GM-CSF for treating or preventing neonatal infections. *Cochrane Database Syst Rev* 2003;2003:Cd003066.
23. Cleminson J, Austin N, McGuire W. Prophylactic systemic antifungal agents to prevent mortality and morbidity in very low birth weight infants. *Cochrane Database Syst Rev* 2015;2015: Cd003850.
24. Cleminson J, McGuire W. Topical emollient for preventing infection in preterm infants. *Cochrane Database Syst Rev* 2016; 2016:Cd001150.
25. Collins CT, Gillis J, McPhee AJ, Sukanuma H, Makrides M. Avoidance of bottles during the establishment of breast feeds in preterm infants. *Cochrane Database Syst Rev* 2016;10:1–51.
26. Collins CT, Makrides M, McPhee AJ. Early discharge with home support of gavage feeding for stable preterm infants who have not established full oral feeds. *Cochrane Database Syst Rev* 2015;2015:Cd003743.
27. Conde-Agudelo A, Díaz-Rossello JL. Kangaroo mother care to reduce morbidity and mortality in low birthweight infants. *Cochrane Database Syst Rev* 2016;2016:Cd002771.
28. Craft AP, Finer NN, Barrington KJ. Vancomycin for prophylaxis against sepsis in preterm neonates. *Cochrane Database Syst Rev* 2000;2000:Cd001971.
29. Darlow BA, Austin NC. Selenium supplementation to prevent short-term morbidity in preterm neonates. *Cochrane Database Syst Rev* 2003;4:1–18.

30. Darlow BA, Graham PJ, Rojas-Reyes MX. Vitamin A supplementation to prevent mortality and short- and long-term morbidity in very low birth weight infants. *Cochrane Database Syst Rev* 2016;8:1–60.
31. De Paoli AG, Davis PG, Faber B, Morley CJ. Devices and pressure sources for administration of nasal continuous positive airway pressure (NCPAP) in preterm neonates. *Cochrane Database Syst Rev* 2008;1:1–44.
32. Doyle LW, Cheong JL, Ehrenkranz RA, Halliday HL. Late (> 7 days) systemic postnatal corticosteroids for prevention of bronchopulmonary dysplasia in preterm infants. *Cochrane Database Syst Rev* 2017;10:CD001145.
33. Doyle LW, Cheong JL, Ehrenkranz RA, Halliday HL. Early (< 8 days) systemic postnatal corticosteroids for prevention of bronchopulmonary dysplasia in preterm infants. *Cochrane Database Syst Rev* 2017;10:1–117.
34. Fenton TR, Premji SS, Al-Wassia H, Sauve RS. Higher versus lower protein intake in formula-fed low birth weight infants. *Cochrane Database Syst Rev* 2014;2014:CD003959.
35. Flenady V, Woodgate PG. Radiant warmers versus incubators for regulating body temperature in newborn infants. *Cochrane Database Syst Rev* 2003;4:1–51.
36. Foster JP, Dawson JA, Davis PG, Dahlen HG. Routine oro/nasopharyngeal suction versus no suction at birth. *Cochrane Database Syst Rev* 2017;4:CD010332.
37. Foster JP, Richards R, Showell MG, Jones LJ. Intravenous in-line filters for preventing morbidity and mortality in neonates. *Cochrane Database Syst Rev* 2015;8:1–25.
38. Gordon A, Greenhalgh M, McGuire W. Early planned removal of umbilical venous catheters to prevent infection in newborn infants. *Cochrane Database Syst Rev* 2017;10:CD012142.
39. Gray PH, Flenady V. Cot-nursing versus incubator care for preterm infants. *Cochrane Database Syst Rev* 2011;8:1–35.
40. Howlett A, Ohlsson A, Plakkal N. Inositol in preterm infants at risk for or having respiratory distress syndrome. *Cochrane Database Syst Rev* 2019;7:CD000366.
41. Ibrahim H, Sinha IP, Subhedar NV. Corticosteroids for treating hypotension in preterm infants. *Cochrane Database Syst Rev* 2011. <https://doi.org/10.1002/14651858.cd003662.pub4>.
42. Imdad A, Bautista RM, Senen KA, Uy ME, Mantaring JB 3rd, Bhutta ZA. Umbilical cord antiseptics for preventing sepsis and death among newborns. *Cochrane Database Syst Rev* 2013;5:1–93.
43. Inglis GD, Davies MW. Prophylactic antibiotics to reduce morbidity and mortality in neonates with umbilical venous catheters. *Cochrane Database Syst Rev* 2005;4:1–13.
44. Inglis GD, Jardine LA, Davies MW. Prophylactic antibiotics to reduce morbidity and mortality in ventilated newborn infants. *Cochrane Database Syst Rev* 2007;1:1–17.
45. Inglis GD, Jardine LA, Davies MW. Prophylactic antibiotics to reduce morbidity and mortality in neonates with umbilical artery catheters. *Cochrane Database Syst Rev* 2007:CD004697.
46. Jacobs SE, Berg M, Hunt R, Tarnow-Mordi WO, Inder TE, Davis PG. Cooling for newborns with hypoxic ischaemic encephalopathy. *Cochrane Database Syst Rev* 2013;1:1–91.
47. Jardine LA, Inglis GD, Davies MW. Prophylactic systemic antibiotics to reduce morbidity and mortality in neonates with central venous catheters. *Cochrane Database Syst Rev* 2008;1:1–18.
48. Kabra NS, Kumar M, Shah SS. Multiple versus single lumen umbilical venous catheters for newborn infants. *Cochrane Database Syst Rev* 2005;3:1–20.
49. Kapoor V, Malviya MN, Soll R. Lipid emulsions for parenterally fed term and late preterm infants. *Cochrane Database Syst Rev* 2019;6:CD013171.
50. Kapoor V, Malviya MN, Soll R. Lipid emulsions for parenterally fed preterm infants. *Cochrane Database Syst Rev* 2019;6:CD013163.
51. Kelly LE, Shivananda S, Murthy P, Srinivasjois R, Shah PS. Antibiotics for neonates born through meconium-stained amniotic fluid. *Cochrane Database Syst Rev* 2017;6:CD006183.
52. Kuti BP, Ogunlesi TA, Oduwole O, Oranganje C, Udoh EE, Meremikwu MM. Hand hygiene for the prevention of infections in neonates. *Cochrane Database Syst Rev* 2021;1:CD013326.
53. Lai NM, Foong SC, Foong WC, Tan K. Co-bedding in neonatal nursery for promoting growth and neurodevelopment in stable preterm twins. *Cochrane Database Syst Rev* 2016;4:CD008313.
54. Lai NM, Taylor JE, Tan K, Choo YM, Ahmad Kamar A, Muhamad NA. Antimicrobial dressings for the prevention of catheter-related infections in newborn infants with central venous catheters. *Cochrane Database Syst Rev* 2016;3:CD011082.
55. Lassi ZS, Kedzior SGE, Bhutta ZA. Community-based maternal and newborn educational care packages for improving neonatal health and survival in low- and middle-income countries. *Cochrane Database Syst Rev* 2019;11:1–51.
56. Lemyre B, Laughon M, Bose C, Davis PG. Early nasal intermittent positive pressure ventilation (NIPPV) versus early nasal continuous positive airway pressure (NCPAP) for preterm infants. *Cochrane Database Syst Rev* 2016;12:1–64.
57. Malviya MN, Ohlsson A, Shah SS. Surgical versus medical treatment with cyclooxygenase inhibitors for symptomatic patent ductus arteriosus in preterm infants. *Cochrane Database Syst Rev* 2013;2013:CD003951.
58. McCall EM, Alderdice F, Halliday HL, Vohra S, Johnston L. Interventions to prevent hypothermia at birth in preterm and/or low birth weight infants. *Cochrane Database Syst Rev* 2018;2:CD004210.
59. McMullan RL, Gordon A. Antibiotics at the time of removal of central venous catheter to reduce morbidity and mortality in newborn infants. *Cochrane Database Syst Rev* 2018;3:CD012181.
60. Meyer MP, Nevill E, Wong MM. Provision of respiratory support compared to no respiratory support before cord clamping for preterm infants. *Cochrane Database Syst Rev* 2018;3:1–43.
61. Moe-Byrne T, Brown JV, McGuire W. Glutamine supplementation to prevent morbidity and mortality in preterm infants. *Cochrane Database Syst Rev* 2016;3:1–49.
62. Moon K, Athalye-Jape GK, Rao U, Rao SC. Early versus late parenteral nutrition for critically ill term and late preterm infants. *Cochrane Database Syst Rev* 2020;4:CD013141.
63. Morgan J, Bombell S, McGuire W. Early trophic feeding versus enteral fasting for very preterm or very low birth weight infants. *Cochrane Database Syst Rev* 2013:CD000504.
64. Morgan J, Young L, McGuire W. Delayed introduction of progressive enteral feeds to prevent necrotising enterocolitis in very low birth weight infants. *Cochrane Database Syst Rev* 2014;12:1–28.

65. Muelbert M, Lin L, Bloomfield FH, Harding JE. Exposure to the smell and taste of milk to accelerate feeding in preterm infants. *Cochrane Database Syst Rev* 2019;7:Cd013038.
66. Nasuf AWA, Ojha S, Dorling J. Oropharyngeal colostrum in preventing mortality and morbidity in preterm infants. *Cochrane Database Syst Rev* 2018;9:Cd011921.
67. Ng DHC, Klassen JRL, Embleton ND, McGuire W. Protein hydrolysate versus standard formula for preterm infants. *Cochrane Database Syst Rev* 2019;7:1–44.
68. Ng G, da Silva O, Ohlsson A. Bronchodilators for the prevention and treatment of chronic lung disease in preterm infants. *Cochrane Database Syst Rev* 2016;12:Cd003214.
69. Ng G, Ohlsson A. Cromolyn sodium for the prevention of chronic lung disease in preterm infants. *Cochrane Database Syst Rev* 2017;1:Cd003059.
70. Oddie SJ, Young L, McGuire W. Slow advancement of enteral feed volumes to prevent necrotising enterocolitis in very low birth weight infants. *Cochrane Database Syst Rev* 2017;8:Cd001241.
71. Ohlsson A, Aher SM. Early erythropoiesis-stimulating agents in preterm or low birth weight infants. *Cochrane Database Syst Rev* 2020;2:1–136.
72. Ohlsson A, Lacy JB. Intravenous immunoglobulin for preventing infection in preterm and/or low birth weight infants. *Cochrane Database Syst Rev* 2020;1:Cd000361.
73. Ohlsson A, Shah PS. Paracetamol (acetaminophen) for patent ductus arteriosus in preterm or low birth weight infants. *Cochrane Database Syst Rev* 2020;4:1–121.
74. Ohlsson A, Shah SS. Ibuprofen for the prevention of patent ductus arteriosus in preterm and/or low birth weight infants. *Cochrane Database Syst Rev* 2020;1:1–65.
75. Ohlsson A, Walia R, Shah SS. Ibuprofen for the treatment of patent ductus arteriosus in preterm or low birth weight (or both) infants. *Cochrane Database Syst Rev* 2020;9:1–253.
76. Onland W, De Jaegere AP, Offringa M, van Kaam A. Systemic corticosteroid regimens for prevention of bronchopulmonary dysplasia in preterm infants. *Cochrane Database Syst Rev* 2017;1:Cd010941.
77. Onland W, Offringa M, van Kaam A. Late (≥ 7 days) inhalation corticosteroids to reduce bronchopulmonary dysplasia in preterm infants. *Cochrane Database Syst Rev* 2017;8:Cd002311.
78. Osborn DA, Hunt RW. Prophylactic postnatal thyroid hormones for prevention of morbidity and mortality in preterm infants. *Cochrane Database Syst Rev* 2007;1:1–78.
79. Osborn DA, Schindler T, Jones LJ, Sinn JK, Bolisetty S. Higher versus lower amino acid intake in parenteral nutrition for newborn infants. *Cochrane Database Syst Rev* 2018;3:Cd005949.
80. Pammi M, Suresh G. Enteral lactoferrin supplementation for prevention of sepsis and necrotizing enterocolitis in preterm infants. *Cochrane Database Syst Rev* 2020;3:Cd007137.
81. Pfister RH, Soll R, Wiswell TE. Protein-containing synthetic surfactant versus protein-free synthetic surfactant for the prevention and treatment of respiratory distress syndrome. *Cochrane Database Syst Rev* 2009;4:1–32.
82. Premkumar MH, Pammi M, Suresh G. Human milk-derived fortifier versus bovine milk-derived fortifier for prevention of mortality and morbidity in preterm neonates. *Cochrane Database Syst Rev* 2019;2019:1–31.
83. Quigley M, Embleton ND, McGuire W. Formula versus donor breast milk for feeding preterm or low birth weight infants. *Cochrane Database Syst Rev* 2019. <https://doi.org/10.1002/14651858.cd002971.pub3>.
84. Rojas-Reyes MX, Morley CJ, Soll R. Prophylactic versus selective use of surfactant in preventing morbidity and mortality in preterm infants. *Cochrane Database Syst Rev* 2012;2:1–72.
85. Schulzke SM, Kaempfen S, Patole SK. Pentoxifylline for the prevention of bronchopulmonary dysplasia in preterm infants. *Cochrane Database Syst Rev* 2014:Cd010018.
86. Schulzke SM, Kaempfen S, Trachsel D, Patole SK. Physical activity programs for promoting bone mineralization and growth in preterm infants. *Cochrane Database Syst Rev* 2014;4:1–44.
87. Seger N, Soll R. Animal derived surfactant extract for treatment of respiratory distress syndrome. *Cochrane Database Syst Rev* 2009. <https://doi.org/10.1002/14651858.cd007836>.
88. Shah A, Brunskill SJ, Desborough MJ, Doree C, Trivella M, Stanworth SJ. Transfusion of red blood cells stored for shorter versus longer duration for all conditions. *Cochrane Database Syst Rev* 2018;12:Cd010801.
89. Shah PS, Kaufman DA. Antistaphylococcal immunoglobulins to prevent staphylococcal infection in very low birth weight infants. *Cochrane Database Syst Rev* 2009;2:1–36.
90. Shah PS, Shah VS. Continuous heparin infusion to prevent thrombosis and catheter occlusion in neonates with peripherally placed percutaneous central venous catheters. *Cochrane Database Syst Rev* 2008;2:1–24.
91. Shah SS, Ohlsson A, Halliday HL, Shah VS. Inhaled versus systemic corticosteroids for the treatment of bronchopulmonary dysplasia in ventilated very low birth weight preterm infants. *Cochrane Database Syst Rev* 2017;10:1–53.
92. Shah SS, Ohlsson A, Halliday HL, Shah VS. Inhaled versus systemic corticosteroids for preventing bronchopulmonary dysplasia in ventilated very low birth weight preterm neonates. *Cochrane Database Syst Rev* 2017;10:1–50.
93. Shah VS, Ohlsson A, Halliday HL, Dunn M. Early administration of inhaled corticosteroids for preventing chronic lung disease in very low birth weight preterm neonates. *Cochrane Database Syst Rev* 2017;1:1–61.
94. Sharif S, Meader N, Oddie SJ, Rojas-Reyes MX, McGuire W. Probiotics to prevent necrotising enterocolitis in very preterm or very low birth weight infants. *Cochrane Database Syst Rev* 2020;10:1–133.
95. Sinclair JC, Bottino M, Cowett RM. Interventions for prevention of neonatal hyperglycemia in very low birth weight infants. *Cochrane Database Syst Rev* 2011;10:1–79.
96. Singh N, Halliday HL, Stevens TP, Suresh G, Soll R, Rojas-Reyes MX. Comparison of animal-derived surfactants for the prevention and treatment of respiratory distress syndrome in preterm infants. *Cochrane Database Syst Rev* 2015;12:1–83.
97. Sinha A, Sazawal S, Pradhan A, Ramji S, Opiyo N. Chlorhexidine skin or cord care for prevention of mortality and infections in neonates. *Cochrane Database Syst Rev* 2015;3:1–59.
98. Soll R, Özek E. Prophylactic animal derived surfactant extract for preventing morbidity and mortality in preterm infants. *Cochrane Database Syst Rev* 1997;4:1–36.
99. Soll R, Özek E. Multiple versus single doses of exogenous surfactant for the prevention or treatment of neonatal respiratory distress syndrome. *Cochrane Database Syst Rev* 2009;1:1–25.

100. Spence K, Barr P. Nasal versus oral intubation for mechanical ventilation of newborn infants. *Cochrane Database Syst Rev* 2000;1999: Cd000948.
101. Subramaniam P, Ho JJ, Davis PG. Prophylactic nasal continuous positive airway pressure for preventing morbidity and mortality in very preterm infants. *Cochrane Database Syst Rev* 2021;10: 1–72.
102. Symington AJ, Pinelli J. Developmental care for promoting development and preventing morbidity in preterm infants. *Cochrane Database Syst Rev* 2006;2:1–61.
103. Taylor JE, Tan K, Lai NM, McDonald SJ. Antibiotic lock for the prevention of catheter-related infection in neonates. *Cochrane Database Syst Rev* 2015;6:1–48.
104. Ullman AJ, Cooke ML, Gillies D, Marsh NM, Daud A, McGrail MR, et al. Optimal timing for intravascular administration set replacement. *Cochrane Database Syst Rev* 2013;2013: Cd003588.
105. Ungerer RL, Lincetto O, McGuire W, Saloojee H, Gulmezoglu AM. Prophylactic versus selective antibiotics for term newborn infants of mothers with risk factors for neonatal infection. *Cochrane Database Syst Rev* 2004;4:1–16.
106. Walsh V, Brown JVE, McGuire W. Iodine supplementation for the prevention of mortality and adverse neurodevelopmental outcomes in preterm infants. *Cochrane Database Syst Rev* 2019; 2:1–37.
107. Webster J, Pritchard MA. Gowning by attendants and visitors in newborn nurseries for prevention of neonatal morbidity and mortality. *Cochrane Database Syst Rev* 2003;2003: Cd003670.
108. Whitelaw A, Brion LP, Kennedy CR, Odd D. Diuretic therapy for newborn infants with posthemorrhagic ventricular dilatation. *Cochrane Database Syst Rev* 2001;2:1–18.
109. Whitelaw A, Lee-Kelland R. Repeated lumbar or ventricular punctures in newborns with intraventricular hemorrhage. *Cochrane Database Syst Rev* 2017;4:1–28.
110. Wilkinson D, Andersen C, O'Donnell CPF, De Paoli AG, Manley BJ. High flow nasal cannula for respiratory support in preterm infants. *Cochrane Database Syst Rev* 2016;2:1–76.
111. Zupan J, Garner P, Omari AA. Topical umbilical cord care at birth. *Cochrane Database Syst Rev* 2004;2004: Cd001057.
112. Razak A, Alshehri N. Azithromycin for preventing bronchopulmonary dysplasia in preterm infants: a systematic review and meta-analysis. *Pediatr Pulmonol* 2021;56:957–66.
113. Razak A, Hussain A. Lactoferrin supplementation to prevent late-onset sepsis in preterm infants: a meta-analysis. *Am J Perinatol* 2021;38:283–90.
114. Razak A, Patel RM, Gautham KS. Use of probiotics to prevent necrotizing enterocolitis: evidence to clinical practice. *JAMA Pediatr* 2021;175:773–4.
115. Razak A, Alshehri N. Azithromycin for preventing bronchopulmonary dysplasia in preterm infants: A systematic review and meta-analysis. *Pediatr Pulmonol* 2021;56:957–66.

Supplementary Material: The online version of this article offers supplementary material (<https://doi.org/10.1515/jpm-2022-0131>).