



Allan Jenkinson, Sarah Zaidi, Ravindra Bhat, Anne Greenough and Theodore Dassios*

Carboxyhaemoglobin levels in infants with hypoxic ischaemic encephalopathy

<https://doi.org/10.1515/jpm-2023-0174>

Received April 25, 2023; accepted August 3, 2023;
published online August 29, 2023

Abstract

Objectives: Hypoxic ischaemic encephalopathy (HIE) is associated with oxidative stress. A potential marker of oxidative damage is carboxyhaemoglobin (COHb) which is the product of the reaction between carbon monoxide and haemoglobin and is routinely assessed on blood gas analysis. Our objective was to test the hypothesis that higher COHb levels would be associated with worse outcomes in infants treated for HIE.

Methods: A retrospective, observational study was performed of all infants who received whole body hypothermia for HIE at a tertiary neonatal intensive care unit between January 2018 and August 2021. For each participating infant, the highest COHb level per day was recorded for days one, three and five after birth.

Results: During the study period, 67 infants with a median (IQR) gestational age of 40 (38–41) weeks underwent therapeutic hypothermia for HIE. The median (IQR) COHb level on day three was higher in infants without electroencephalographic seizures (1.4 [1.1–1.4] %) compared with infants with seizures (1.1 [0.9–1.3] %, $p=0.024$). The median (IQR) COHb on day five was higher in infants without MRI brain abnormalities (1.4 [1.2–1.7] %) compared with infants with MRI abnormalities (1.2 [1.0–1.4] %, $p=0.032$). The COHb level was not significantly different between the nine infants who died compared to the infants who survived.

Conclusions: COHb levels were higher in infants with HIE without seizures and in those with normal MRI brain

examinations. We suggest that carbon monoxide has a potential protective role in HIE.

Keywords: oxidation; neonates; seizures; MRI brain

Introduction

Hypoxic ischaemic encephalopathy (HIE) is a serious birth complication resulting from impaired cerebral blood flow and oxygen delivery in term infants [1]. Treatment response and outcome prediction are measured using a combination of the results of clinical examination, amplitude integrated electroencephalography (aEEG) and magnetic resonance imaging (MRI) of the brain [2]. The severity of the clinical picture, however, can be subjective depending on the assessor and the interpretation of aEEG can be affected by hypothermia treatment or anti-epileptic medication, while some infants are too unstable for transport to an MRI scanner [3]. As a result, the early predictive abilities of those modalities can be hindered.

The pathophysiology of HIE involves oxidative stress which is most marked during the secondary phase of injury occurring 6–15 h after the initial event, potentially as a result of reperfusion injury [4]. By-products of the oxidative processes may act as useful biomarkers in these infants, providing an additional non-invasive means of risk stratifying neonates with HIE [5]. A potential marker of oxidative stress in the neonatal population is carboxyhaemoglobin (COHb) [6] which is the product of the reaction between carbon monoxide and haemoglobin. COHb levels are available as part of routine blood gas analysis. Carboxyhaemoglobin levels have previously been shown to correlate with the length of invasive ventilation in paediatric intensive care [7] and the development of bronchopulmonary dysplasia and intraventricular haemorrhage in preterm infants [8]. We hypothesised that infants with severe HIE, that is those with abnormal MRI brain findings and/or who had clinical or electrical seizures, would have elevated COHb levels during the first days after birth compared to those with less severe HIE.

*Corresponding author: Dr. Theodore Dassios, Department of Women and Children's Health, School of Life Course Sciences, Faculty of Life Sciences and Medicine, King's College London, London, UK; and Neonatal Intensive Care Centre, King's College Hospital NHS Foundation Trust, Denmark Hill, London, SE5 9RS, UK, E-mail: theodore.dassios@kcl.ac.uk

Allan Jenkinson and Anne Greenough, Department of Women and Children's Health, School of Life Course Sciences, Faculty of Life Sciences and Medicine, King's College London, London, UK. <https://orcid.org/0000-0002-8672-5349> (A. Greenough)

Sarah Zaidi, GKT School of Medical Education, London, UK

Ravindra Bhat, Neonatal Intensive Care Centre, King's College Hospital NHS Foundation Trust, London, UK

Subjects and methods

Study design and subjects

A retrospective, observational study of all infants who received whole body hypothermia for HIE at a tertiary neonatal intensive care unit (NICU) between January 2018 and August 2021 was undertaken. Infants were identified via the BadgerNet Neonatal Electronic Patient Record (Clevermed, Edinburgh, UK). Infants with haemolytic jaundice and a positive direct antiglobulin test were excluded from the study as carbon monoxide is a by-product of heme degradation and is associated with increased COHb [9]. Infants were included if they had at least five arterial blood gases in the first 3 days after birth. The study was registered with the Clinical Governance Department. The Health Research Authority Toolkit of the National Health System, United Kingdom confirmed that the study would not need regulatory approval by a research ethics committee.

Information from the medical notes

The following information was collected from the medical notes: meconium stained amniotic fluid, emergency caesarean section, sex, gestational age (weeks), birth weight (kg), Apgar score at 1, 5 and 10 min, arterial cord blood pH, base deficit and lactate. Additionally, the highest blood aspartate aminotransferase (AST) and creatinine levels in the first 3 days, the duration of invasive ventilation (days), clinical or echocardiographic evidence of pulmonary hypertension [10], use of inotropic agents [11], electrical or clinical seizures [12], MRI abnormalities consistent with HIE [13] and survival to discharge from neonatal care were recorded. All infants that received whole body hypothermia for HIE had a MRI brain before discharge from neonatal care, beyond the pseudo-normalisation window at the end of the first week of life [14].

Blood gas analysis

Arterial or capillary blood gas analysis was performed by the medical or nursing team as part of routine clinical care. Samples were analysed using the ABL90 FLEX PLUS analyser (Radiometer UK Ltd) and results were stored in the archived database on the blood gas analyser hard disk. All samples stored in the designated study period were extracted from the archived database for analysis of the COHb levels. COHb results were expressed as a fraction (%) of the total haemoglobin concentration. For each participating infant, the highest COHb level per day was recorded for days one, three and five after birth. This time period was selected as we have previously demonstrated that in the first month of life, the COHb levels were highest during the first 3 days and that during that period there was the greatest difference in levels between infants who developed oxidative diseases and those that did not [8]. Furthermore, the majority of infants did not have regular blood gas monitoring after day six.

Clinical management

Active whole body hypothermia was initiated within 6 h from birth either at the hospital or by the dedicated neonatal transport team attending the referring hospital and maintained for 72 h. Assisted

ventilation was initiated due to poor condition at birth and maintained if there was an absence of adequate respiratory drive and the administration of sedative and anticonvulsive agents. The fraction of inspired oxygen was manually adjusted to achieve oxygen saturation levels measured by pulse oximetry (SpO_2) of 95–100 % [15] and ventilation settings were adjusted to achieve a partial pressure of carbon dioxide of 4.5–6.5 kPa [16].

Statistical analysis

Continuous data were tested for normality with the Kolmogorov-Smirnov test and found to be non-normally distributed and thus presented as median and interquartile range (IQR) and the Mann Whitney U test used to determine if differences were statistically significant. The strength of relationships between continuous variables was examined using Spearman's Rho non parametric bivariate correlation analysis. Statistical analysis was performed using SPSS software, version 27.0 (SPSS Inc, Chicago, IL, USA).

Results

During the study period, 67 infants (44 male) underwent therapeutic hypothermia for HIE. Their median (IQR) gestational age was 40 (38–41) weeks and birth weight was 3.39 (2.08–3.66) kg (Table 1).

Seizures (electrographic or clinical) were reported in 33/66 (50 %). MRI features consistent with HIE were present in 20/62 (30 %). Nine infants (13 %) died from HIE before discharge from neonatal care.

The median (IQR) COHb level on day three was higher in infants without EEG confirmed seizures compared to in infants with seizures ($p=0.024$, Table 2). The median (IQR) COHb on day five was higher in infants without MRI abnormalities compared to in infants with MRI abnormalities

Table 1: Demographics and clinical characteristics. Data presented as median (IQR).

Gestational age, weeks	40 (38–41)
Birth weight, kg	3.39 (2.08–3.66)
Male	44/67 (65 %)
Inborn	42/67 (62.7 %)
Emergency LSCS	29/66 (43 %)
Apgar at 1 min	2 (0–3)
Apgar at 5 min	4 (2–6)
Apgar at 10 min	6 (4–8)
pH cord gas	7.0 (6.9–7.2)
BE cord gas	-11.9 (-16.7 to -7.1)
Lactate cord gas, mmol/L	10 (6–14)
Meconium stained amniotic fluid	23 (34 %)
Pulmonary hypertension	9 (13 %)
Need for inotropic support	27/67 (40 %)

LSCS, lower segment caesarean section; BE, base excess.

Table 2: COHb levels according to presence of seizures and MRI brain results. Data are presented as median [IQR].

	No seizures	Seizures	p-Value	Normal MRI	Abnormal MRI	p-Value
COHb day 1	1.4 [1.1–1.6]	1.3 [1.0–1.5]	0.534	1.4 [1.1–1.5]	1.45 [1.1–1.6]	0.360
COHb day 3	1.4 [1.1–1.4]	1.1 [0.9–1.3]	0.024	1.3 [1.0–1.4]	1.2 [1–1.5]	0.946
COHb day 5	1.4 [1.2–1.7]	1.4 [1.1–1.5]	0.391	1.4 [1.2–1.7]	1.2 [1.0–1.5]	0.032

($p=0.032$, Table 2). The COHb levels were not significantly different between the nine infants who died compared to those who survived.

The COHb levels on any study day were not significantly different between infants with or without meconium stained amniotic fluid or pulmonary hypertension or who need inotrope support (Supplementary Table 1). The COHb levels on any of the study days did not significantly correlate with arterial cord blood pH, base deficit, lactate, AST or creatinine levels (Supplementary Table 2).

Discussion

We have demonstrated that COHb levels were higher on day three in infants without seizures compared to infants with seizures and higher on day five in infants with normal MRI brain examinations compared to infants with abnormal MRI scans.

These findings do not support our hypothesis that infants with severe HIE would have higher COHb levels during the first days after birth compared to those with less severe HIE. It was interesting that on all three study days better outcomes (survival, no seizures, normal MRI examination) were associated with higher COHb levels compared to the unfavourable outcomes (death, seizures, abnormal MRI). We postulate that the consistency of these finding suggests that elevated COHb levels have a different mechanism in HIE than in our initial hypothesis.

Whilst carbon monoxide is known to be a toxic agent, recent evidence has suggested that when given at low doses it might have neuroprotective properties [17–19]. Elevated carboxyhaemoglobin levels may thus represent an increased endogenous neuroprotective production of carbon monoxide in response to the hypoxic-ischemic insult. This effect might be mediated by the up-regulation of enzymes responsible for the heme catabolism which produce carbon monoxide as a by-product. Indeed, animal studies have highlighted the protective effect of carbon monoxide supplementation in ameliorating hepatobiliary dysfunction during catabolism of heme molecules in endotoxemic livers [20]. Furthermore, exogenous carbon monoxide has also

been shown to have anti-inflammatory, antiapoptotic and cytoprotective effects in animal studies [18, 21]. In an animal study where mice were exposed to carbon monoxide at a concentration of either 200 or 250 ppm for a period of 1 h after an hypoxic insult, it was demonstrated that 250 ppm of carbon monoxide were effective in preserving cortical volumes following a mild hypoxic-ischemic injury [3]. In further animal studies, Liu et al. also showed that carbon monoxide reperfusion relieved oxidative injury and inhibited apoptosis and autophagy [22]. Our results then are interesting in pointing towards carbon monoxide not as an index of oxidative stress, but as a potential neuroprotective pathway that could be further explored as highlighted in the aforementioned animal studies. It is interesting to note that there were no significant correlations between cord gas variables (pH, BE, lactate), AST, platelets and creatinine levels and COHb levels on any of the study days. Those observations also highlight that COHb may be of limited clinical utility as a marker of oxidative stress in HIE.

This is the first human study pointing to a possible protective role of carbon monoxide in hypoxic ischemic injury. A limitation to our study is the lack of control samples, however, COHb levels would not be measured serially in infants without any pathology. In our cohort we also report a relatively low mortality rate so we are underpowered to report significant differences relating to this outcome. Our mortality rate, however, is not dissimilar to those in previously published national reports [23].

Conclusions

In conclusion higher carboxyhaemoglobin levels were associated with normal MRI brain findings and the absence of electrical seizures in infants with HIE. Those findings are consistent with a possible cytoprotective effect of carbon monoxide. Trials are warranted to investigate the potential therapeutic effects of carbon monoxide in the context of HIE.

Acknowledgments: We thank Mrs Deirdre Gibbons for secretarial assistance.

Research ethics: The study was registered with the Clinical Governance Department of KCH. The Health Research Authority Toolkit of the National Health System, United Kingdom confirmed that the study would not need regulatory approval by a Research Ethics Committee.

Informed consent: Not applicable.

Author contributions: AJ: data collection, analysis and writing the first draft of the manuscript. SZ: data collection, revision of the manuscript. RB: data analysis and interpretation, revision of the manuscript. AG: conceptualisation, supervision, critical revision. TD: conceptualisation, supervision, analysis, critical revision. All authors have accepted responsibility for the entire content of this manuscript and approved its submission.

Competing interests: The authors state no conflict of interest.

Research funding: None declared.

Data availability: Data will be made available on reasonable request.

References

- Allen KA, Brandon DH. Hypoxic ischemic encephalopathy: pathophysiology and experimental treatments. *Newborn Infant Nurs Rev* 2011;11:125–33.
- van Laerhoven H, de Haan TR, Offringa M, Post B, van der Lee JH. Prognostic tests in term neonates with hypoxic-ischemic encephalopathy: a systematic review. *Pediatrics* 2013;131:88–98.
- Douglas-Escobar M, Weiss MD. Hypoxic-ischemic encephalopathy: a review for the clinician. *JAMA Pediatr* 2015;169:397–403.
- Wassink G, Gunn ER, Drury PP, Bennet L, Gunn AJ. The mechanisms and treatment of asphyxial encephalopathy. *Front Neurosci* 2014;8:40.
- Douglas-Escobar M, Weiss MD. Biomarkers of hypoxic-ischemic encephalopathy in newborns. *Front Neurol* 2012;3:144.
- Tokuriki S, Okuno T, Ohta G, Ohshima Y. Carboxyhemoglobin formation in preterm infants is related to the subsequent development of bronchopulmonary dysplasia. *Dis Markers* 2015;2015:620921.
- Chawla A, Ray S, Matettore A, Peters MJ. Arterial carboxyhaemoglobin levels in children admitted to PICU: a retrospective observational study. *PLoS One* 2019;14:e0209452.
- Bednarczuk N, Williams EE, Greenough A, Dassios T. Carboxyhaemoglobin levels and free-radical-related diseases in prematurely born infants. *Early Hum Dev* 2022;164:105523.
- Bailey DGN, Fuchs H, Hentschel R. Carboxyhemoglobin – the forgotten parameter of neonatal hyperbilirubinemia. *J Perinat Med* 2017;45: 613–17.
- Lakshminrusimha S, Shankaran S, Laptook A, McDonald S, Keszler M, Van Meurs K, et al. Pulmonary hypertension associated with hypoxic-ischemic encephalopathy-antecedent characteristics and comorbidities. *J Pediatr* 2018;196:45–51.e3.
- Ovali F. Hemodynamic changes and evaluation during hypoxic-ischemic encephalopathy and therapeutic hypothermia. *Early Hum Dev* 2022;167:105563.
- Kharoshankaya L, Stevenson NJ, Livingstone V, Murray DM, Murphy BP, Ahearne CE, et al. Seizure burden and neurodevelopmental outcome in neonates with hypoxic-ischemic encephalopathy. *Dev Med Child Neurol* 2016;58:1242–8.
- Rutherford M, Malamateniou C, McGuinness A, Allsop J, Biarge MM, Counsell S. Magnetic resonance imaging in hypoxic-ischaemic encephalopathy. *Early Hum Dev* 2010;86:351–60.
- Parmentier CEJ, de Vries LS, Groenendaal F. Magnetic resonance imaging in (near)-term infants with hypoxic-ischemic encephalopathy. *Diagnostics* 2022;12. <https://doi.org/10.3390/diagnostics12030645>.
- Morgan MC, Maina B, Waiyego M, Mutinda C, Aluvaala J, Maina M, et al. Oxygen saturation ranges for healthy newborns within 24 hours at 1800 m. *Arch Dis Child Fetal Neonatal Ed* 2017;102:F266–8.
- Wong SK, Chim M, Allen J, Butler A, Tyrrell J, Hurley T, et al. Carbon dioxide levels in neonates: what are safe parameters? *Pediatr Res* 2022; 91:1049–56.
- Kohmoto J, Nakao A, Kaizu T, Tsung A, Ikeda A, Tomiyama K, et al. Low-dose carbon monoxide inhalation prevents ischemia/reperfusion injury of transplanted rat lung grafts. *Surgery* 2006;140:179–85.
- Leffler CW, Parfenova H, Jaggar JH. Carbon monoxide as an endogenous vascular modulator. *Am J Physiol Heart Circ Physiol* 2011;301:H1–11.
- Parfenova H, Tcheranova D, Basuroy S, Fedinec AL, Liu J, Leffler CW. Functional role of astrocyte glutamate receptors and carbon monoxide in cerebral vasodilation response to glutamate. *Am J Physiol Heart Circ Physiol* 2012;302:H2257–66.
- Kyokane T, Norimizu S, Taniai H, Yamaguchi T, Takeoka S, Tsushima E, et al. Carbon monoxide from heme catabolism protects against hepatobiliary dysfunction in endotoxin-treated rat liver. *Gastroenterology* 2001;120:1227–40.
- Wolff DG, Bidlack WR. The formation of carbon monoxide during peroxidation of microsomal lipids. *Biochem Biophys Res Commun* 1976; 73:850–7.
- Liu H, Yu S, Peng Y, Chang X, Yu X. The protective effects of carboxyhemoglobin during the resuscitation from hemorrhagic shock in rats. *Oncotarget* 2017;8:83619–25.
- Shipley L, Gale C, Sharkey D. Trends in the incidence and management of hypoxic-ischaemic encephalopathy in the therapeutic hypothermia era: a national population study. *Arch Dis Child Fetal Neonatal Ed* 2021; 106:529–34.

Supplementary Material: This article contains supplementary material (<https://doi.org/10.1515/jpm-2023-0174>).