Open Mathematics

Research Article

Shigeru Furuichi* and Hamid Reza Moradi

On further refinements for Young inequalities

https://doi.org/10.1515/math-2018-0115 Received July 21, 2018; accepted October 18, 2018.

Abstract: In this paper sharp results on operator Young's inequality are obtained. We first obtain sharp multiplicative refinements and reverses for the operator Young's inequality. Secondly, we give an additive result, which improves a well-known inequality due to Tominaga. We also provide some estimates for the difference $A^{1/2} \left(A^{-1/2}BA^{-1/2}\right)^{\nu} A^{1/2} - \{(1-\nu)A + \nu B\}$ for $\nu \notin [0,1]$.

Keywords: Operator inequality, Young inequality, Weighted arithmetic and geometric mean, Positive operator

MSC: 47A63, 26D07, 47A60

1 Introduction

This note lies in the scope of operator inequalities. We assume that the reader is familiar with the continuous functional calculus and the Kubo-Ando theory [1].

It is to be understood throughout the paper that the capital letters present bounded linear operators acting on a Hilbert space \mathcal{H} . A is positive (written $A \ge 0$) in case $\langle Ax, x \rangle \ge 0$ for all $x \in \mathcal{H}$ also an operator A is said to be strictly positive (denoted by A > 0) if A is positive and invertible. If A and B are self-adjoint, we write $B \ge A$ in case $B - A \ge 0$. As usual, by I we denote the identity operator.

The weighted arithmetic mean ∇_{ν} , geometric mean \sharp_{ν} , and harmonic mean $!_{\nu}$, for $\nu \in [0, 1]$ and a, b > 0, are defined as follows:

$$a\nabla_{\nu}b = (1-\nu)a + \nu b, \quad a_{\parallel\nu}b = a^{1-\nu}b^{\nu}, \quad a!_{\nu}b = \left\{(1-\nu)a^{-1} + \nu b^{-1}\right\}^{-1}.$$

If $v = \frac{1}{2}$, we denote the arithmetic, geometric, and harmonic means, respectively, by ∇ , \sharp and !, for the simplicity. Like the scalar cases, the operator arithmetic mean, the operator geometric mean, and the operator harmonic mean for A, B > 0 can be stated in the following form:

$$A\nabla_{\nu}B=\left(1-\nu\right)A+\nu B,\quad A\sharp_{\nu}B=A^{\frac{1}{2}}\left(A^{-\frac{1}{2}}BA^{-\frac{1}{2}}\right)^{\nu}A^{\frac{1}{2}},\quad A!_{\nu}B=\left\{(1-\nu)A^{-1}+\nu B^{-1}\right\}^{-1}.$$

The celebrated arithmetic-geometric-harmonic-mean inequalities for scalars assert that if a, b > 0, then

$$a!_{V}b \leq a!_{V}b \leq a\nabla_{V}b.$$
 (1)

Generalization of the inequalities (1) to operators can be seen as follows: If A, B > 0, then

$$A!_{\nu}B \leq A\sharp_{\nu}B \leq A\nabla_{\nu}B.$$

The last inequality above is called the operator Young inequality. During the past years, several refinements and reverses were given for Young's inequality, see for example [2–4].

^{*}Corresponding Author: Shigeru Furuichi: Department of Information Science, College of Humanities and Sciences, Nihon University, 3-25-40, Sakurajyousui, Setagaya-ku, Tokyo, 156-8550, Japan, E-mail: furuichi@chs.nihon-u.ac.jp Hamid Reza Moradi: Young Researchers and Elite Club, Mashhad Branch, Islamic Azad University, Mashhad, Iran, E-mail: hrmoradi@mshdiau.ac.ir

Zuo et al. showed in [5, Theorem 7] that the following inequality holds:

$$K(h,2)^{r}A_{\parallel\nu}B \leq A\nabla_{\nu}B, \quad r = \min\{\nu, 1-\nu\}, K(h,2) = \frac{(h+1)^{2}}{4h}, h = \frac{M}{m}$$
 (2)

whenever $0 < m'I \le B \le mI < MI \le A \le M'I$ or $0 < m'I \le A \le mI < MI \le B \le M'I$. As the authors mentioned in [5], the inequality (2) improves the following refinement of Young's inequality involving Specht's ratio $S(t) = \frac{t^{\frac{1}{l-1}}}{e \log t^{\frac{1}{l-1}}} (t > 0, t \ne 1)$ (see [6, Theorem 2]),

$$S(h^r)A\sharp_{\nu}B\leq A\nabla_{\nu}B.$$

Under the above assumptions, Dragomir proved in [7, Corollary 1] that

$$A\nabla_{\nu}B \leq \exp\left[\frac{\nu(1-\nu)}{2}(h-1)^2\right]A\sharp_{\nu}B.$$

We remark that there is no relationship between two constants $K(h,2)^r$ and $\exp\left[\frac{v(1-v)}{2}(h-1)^2\right]$ in general. In [3, 8] we proved some sharp multiplicative reverses of Young's inequality. In this brief note, as the continuation of our previous works, we establish sharp bounds for the arithmetic, geometric and harmonic mean inequalities. Moreover, we shall show some additive-type refinements and reverses of Young's inequality. We will formulate our new results in a more general setting, namely the sandwich assumption $sA \le B \le tA$ $(0 < s \le t)$. Additionally, we present some Young type inequalities for the wider range of v; i.e., $v \notin [0, 1]$.

2 Main results

In our previous work [8], we gave new sharp inequalities for reverse Young inequalities. In this section we firstly give new sharp inequalities for Young inequalities, as limited cases in the first inequalities both (i) and (ii) of the following theorem.

Theorem 2.1. Let A, B > 0 such that $sA \le B \le tA$ for some scalars $0 < s \le t$ and let $f_v(x) \equiv \frac{(1-v)+vx}{x^v}$ for x > 0, and $v \in [0, 1]$.

- (i) If $t \le 1$, then $f_v(t)A \not\vdash_v B \le A \nabla_v B \le f_v(s)A \not\vdash_v B$.
- (ii) If $s \ge 1$, then $f_v(s)A \not\parallel_v B \le A \nabla_v B \le f_v(t)A \not\parallel_v B$.

Proof. Since $f'_{\nu}(x) = \nu(1-\nu)(x-1)x^{-\nu-1}$, $f_{\nu}(x)$ is monotone decreasing for $0 < x \le 1$ and monotone increasing for $x \ge 1$.

- (i) For the case $0 < s \le x \le t \le 1$, we have $f_v(t) \le f_v(x) \le f_v(s)$, which implies $f_v(t)A \not\parallel_V B \le A \nabla_v B \le f_v(s)A \not\parallel_V B$ by the standard functional calculus.
- (ii) For the case $1 \le s \le x \le t$, we have $f_{\nu}(s) \le f_{\nu}(x) \le f_{\nu}(t)$ which implies $f_{\nu}(s)A \not\parallel_{\nu} B \le A \nabla_{\nu} B \le f_{\nu}(t)A \not\parallel_{\nu} B$ by the standard functional calculus.

Remark 2.2. It is worth emphasizing that each assertion in Theorem 2.1 implies the other one. For instance, assume that the assertion (ii) holds, i.e.,

$$f_{V}(s) \le f_{V}(x) \le f_{V}(t), \quad 1 \le s \le x \le t. \tag{3}$$

Let $t \le 1$, then $1 \le \frac{1}{t} \le \frac{1}{x} \le \frac{1}{x}$. Hence (3) ensures that

$$f_{V}\left(\frac{1}{t}\right) \leq f_{V}\left(\frac{1}{x}\right) \leq f_{V}\left(\frac{1}{s}\right).$$

So

$$\frac{\left(1-\nu\right)t+\nu}{t^{1-\nu}}\leq\frac{\left(1-\nu\right)x+\nu}{x^{1-\nu}}\leq\frac{\left(1-\nu\right)s+\nu}{s^{1-\nu}}.$$

Now, by replacing v by 1 - v we get

$$\frac{\left(1-v\right)+vt}{t^{v}}\leq\frac{\left(1-v\right)+vx}{x^{v}}\leq\frac{\left(1-v\right)+vs}{s^{v}}$$

which means

$$f_{v}(t) \leq f_{v}(x) \leq f_{v}(s), \quad 0 < s \leq x \leq t \leq 1.$$

In the same spirit, we can derive (ii) from (i).

Corollary 2.3. Let $A, B > 0, m, m', M, M' > 0, and v \in [0, 1].$

(i) If 0 < m'I < A < mI < MI < B < M'I, then

$$\frac{m\nabla_{V}M}{m \#_{V} M} A \# B \leq A \nabla_{V} B \leq \frac{m'\nabla_{V}M'}{m' \#_{V} M'} A \# B. \tag{4}$$

(ii) If $0 < m'I \le B \le mI < MI \le A \le M'I$, then

Proof. We use again the function $f_{\nu}(x) = \frac{(1-\nu)+\nu x}{x^{\nu}}$ in this proof. The condition (i) is equivalent to $I \leq \frac{M}{m}I \leq A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \leq \frac{M'}{m'}I$, so that we get $f_{\nu}(\frac{M}{m})A \not\parallel_{\nu} B \leq A \nabla_{\nu} B \leq f_{\nu}(\frac{M'}{m'})A \not\parallel_{\nu} B$ by putting $s = \frac{M}{m}$ and $t = \frac{M'}{m'}$ in (ii) of Theorem 2.1.

Similarly, the condition (ii) is equivalent to $\frac{m'}{M'}I \leq A^{-\frac{1}{2}}BA^{-\frac{1}{2}} \leq \frac{m}{M}I \leq I$, so that we get $f_{\nu}(\frac{m}{M})A \not\parallel_{\nu} B \leq I$ $A \nabla_{\nu} B \leq f_{\nu}(\frac{m'}{M'}) A \not\parallel_{\nu} B$ by putting $s = \frac{m'}{M'}$ and $t = \frac{m}{M}$ in (i) of Theorem 2.1.

Note that the second inequalities in both (i) and (ii) of Theorem 2.1 and Corollary 2.3 are special cases of [8, Theorem A].

Remark 2.4. It is remarkable that the inequalities $f_v(t) \le f_v(x) \le f_v(s)$ (0 < $s \le x \le t \le 1$) given in the proof of Theorem 2.1 are sharp, since the function $f_v(x)$ for $s \le x \le t$ is continuous. So, all results given from Theorem 2.1 are similarly sharp. As a matter of fact, let A = MI and B = mI, then from LHS of (5), we infer

$$A \nabla_{V} B = (M \nabla_{V} m) I$$
 and $A \sharp_{V} B = (M \sharp_{V} m) I$.

Consequently,

$$\frac{M\nabla_{\nu}m}{M\sharp_{\nu}m}A\sharp_{\nu}B=A\nabla_{\nu}B.$$

To see that the constant $\frac{m\nabla_v M}{m_{\theta} M}$ in the LHS of (4) can not be improved, we consider A = mI and B = MI, then

$$\frac{m\nabla_{\nu}M}{m\#_{\nu}M}A\#_{\nu}B=A\nabla_{\nu}B.$$

By replacing A, B by A^{-1} , B^{-1} , respectively, the refinement and reverse of non-commutative geometricharmonic mean inequality can be obtained as follows:

Corollary 2.5. Let $A, B > 0, m, m', M, M' > 0, and v \in [0, 1].$

(i) If $0 < m'I \le A \le mI < MI \le B \le M'I$, then

$$\frac{m'!_{\nu}M'}{m'!_{\nu}M'}A\sharp_{\nu}B\leq A!_{\nu}B\leq \frac{m!_{\nu}M}{m\sharp_{\nu}M}A\sharp_{\nu}B.$$

(ii) If $0 < m'I \le B \le mI < MI \le A \le M'I$, then

$$\frac{M'!_{\nu}m'}{M'!_{\nu}m'}A_{\sharp\nu}B \leq A!_{\nu}B \leq \frac{M!_{\nu}m}{M!_{\nu}m}A_{\sharp\nu}B.$$

Now, we give a new sharp reverse inequality for Young's inequality as an additive-type in the following.

Theorem 2.6. Let A, B > 0 such that $sA \le B \le tA$ for some scalars $0 < s \le t$, and $v \in [0, 1]$. Then

$$A \nabla_{\nu} B - A \sharp_{\nu} B \le \max \left\{ g_{\nu}(s), g_{\nu}(t) \right\} A \tag{6}$$

where $g_v(x) \equiv (1 - v) + vx - x^v$ for $s \le x \le t$.

Proof. Straightforward differentiation shows that $g_{\nu}''(x) = \nu(1-\nu)x^{\nu-2} \ge 0$ and $g_{\nu}(x)$ is continuous on the interval [s,t], so

$$g_{V}(x) \leq \max \{g_{V}(s), g_{V}(t)\}.$$

Therefore, by applying similar arguments as in the proof of Theorem 2.1, we reach the desired inequality (6). This completes the proof of theorem. \Box

Corollary 2.7. Let A, B > 0 such that $mI \le A, B \le MI$ for some scalars 0 < m < M. Then for $v \in [0, 1]$,

$$A \nabla_{V} B - A \sharp_{V} B \leq \xi A$$

where $\xi = \max \left\{ \frac{1}{M} \left(M \nabla_{\nu} m - M \sharp_{\nu} m \right), \frac{1}{m} \left(m \nabla_{\nu} M - m \sharp_{\nu} M \right) \right\}.$

Remark 2.8. We claim that if A, B > 0 such that $mI \le A$, $B \le MI$ for some scalars 0 < m < M with $h = \frac{M}{m}$ and $v \in [0, 1]$, then

$$A \nabla_{\nu} B - A \sharp_{\nu} B \leq \max \left\{ g_{\nu}(h), g_{\nu}\left(\frac{1}{h}\right) \right\} A \leq L(1, h) \log S(h) A$$

holds, where $L(x, y) = \frac{y - x}{\log y - \log x} (x \neq y)$ is the logarithmic mean and the term S(h) refers to the Specht's ratio. Indeed, we have the inequalities

$$(1-v)+vh-h^{v} \leq L(1,h)\log S(h), \quad (1-v)+v\frac{1}{h}-h^{-v} \leq L(1,h)\log S(h),$$

which were originally proved in [9, Lemma 3.2], thanks to $S(h) = S(\frac{1}{h})$ and $L(1,h) = L(1,\frac{1}{h})$. Therefore, our result, Theorem 2.6, improves the well-known result by Tominaga [9, Theorem 3.1],

$$A \nabla_{\nu} B - A \mu_{\nu} B \leq L(1, h) \log S(h) A$$
.

Since $g_{\nu}(x)$ is convex, we can not obtain a general result on the lower bound for $A \nabla_{\nu} B - A \sharp_{\nu} B$. However, if we impose the conditions, we can obtain new sharp inequalities for Young inequalities as an additive-type in the first inequalities both (i) and (ii) in the following proposition. (At the same time, of course, we also obtain the upper bounds straightforwardly.)

Proposition 2.9. Let A, B > 0 such that $sA \le B \le tA$ for some scalars $0 < s \le t$, $v \in [0, 1]$, and g_v is defined as in Theorem 2.6.

- (i) If $t \le 1$, then $g_v(t) A \le A \nabla_v B A \mu_v B \le g_v(s) A$.
- (ii) If $s \ge 1$, then $g_v(s) A \le A \nabla_v B A \sharp_v B \le g_v(t) A$.

Proof. It follows from the fact that $g_v(x)$ is monotone decreasing for $0 < x \le 1$ and monotone increasing for $x \ge 1$.

Corollary 2.10. *Let* A, B > 0, m, m', M, M' > 0, and $v \in [0, 1]$.

(i) If $0 < m'I \le A \le mI < MI \le B \le M'I$, then

$$\frac{1}{m}\left(m\nabla_{\nu}M-m_{\sharp\nu}M\right)A\leq A\nabla_{\nu}B-A_{\sharp\nu}B\leq \frac{1}{m'}\left(m'\nabla_{\nu}M'-m'_{\sharp\nu}M'\right)A.$$

(ii) If $0 < m'I \le B \le mI < MI \le A \le M'I$, then

$$\frac{1}{M}\left(M\nabla_{\nu}m-M_{\sharp\nu}m\right)A\leq A\nabla_{\nu}B-A_{\sharp\nu}B\leq \frac{1}{M'}\left(M'\nabla_{\nu}m'-M'_{\sharp\nu}m'\right)A.$$

In the following we use the notations ∇_V and \downarrow_V to distinguish from the operator means ∇_V and \downarrow_V :

$$A \nabla_{\nu} B = (1 - \nu) A + \nu B$$
, $A \mid_{\nu} B = A^{\frac{1}{2}} (A^{-\frac{1}{2}} B A^{-\frac{1}{2}})^{\nu} A^{\frac{1}{2}}$

for $v \notin [0, 1]$. Notice that, since A, B > 0, the expressions $A \nabla_v B$ and $A \mid_{V} B$ are also well-defined.

Remark 2.11. It is known (and easy to show) that for any A, B > 0,

$$A \nabla_{v} B \leq A |_{v} B$$
, for $v \notin [0, 1]$.

Assume $g_{v}(x)$ is defined as in Theorem 2.6. By an elementary computation we have

$$\begin{cases} g'_{v}(x) > 0 \text{ for } v \notin [0, 1] \text{ and } 0 < x \le 1 \\ g'_{v}(x) < 0 \text{ for } v \notin [0, 1] \text{ and } x > 1 \end{cases}.$$

Now, in the same way as above we have also for any $v \notin [0, 1]$ *:*

(i) If $0 < m'I \le A \le mI \le MI \le B \le M'I$, then

$$\frac{1}{m'}\left(m'_{\natural \nu}M'-m'_{\triangledown \nu}M'\right)A\leq A_{\natural \nu}B-A_{\triangledown \nu}B\leq \frac{1}{m}\left(m_{\natural \nu}M-m_{\triangledown \nu}M\right)A.$$

On account of assumptions, we also infer

$$(m' \downarrow_{\mathcal{V}} M' - m' \bigtriangledown_{\mathcal{V}} M') I \leq A \downarrow_{\mathcal{V}} B - A \bigtriangledown_{\mathcal{V}} B \leq (m \downarrow_{\mathcal{V}} M - m \bigtriangledown_{\mathcal{V}} M) I.$$

(ii) If $0 < m'I \le B \le mI \le MI \le A \le M'I$, then

$$\frac{1}{M}\left(M_{\natural\nu}m-M_{\bigtriangledown\nu}m\right)A\leq A_{\natural\nu}B-A_{\bigtriangledown\nu}B\leq \frac{1}{M'}\left(M'_{\natural\nu}m'-M'_{\bigtriangledown\nu}m'\right)A.$$

On account of assumptions, we also infer

$$(M_{\natural_{V}}m - M_{\bigtriangledown_{V}}m)I \leq A_{\natural_{V}}B - A_{\bigtriangledown_{V}}B \leq (M'_{\natural_{V}}m' - M'_{\bigtriangledown_{V}}m')I.$$

In addition, with the same assumption to Theorem 2.6 except for $v \notin [0, 1]$ *, we have*

$$\min\{g_{\nu}(s),g_{\nu}(t)\}A\leq A \bigtriangledown_{\nu}B-A_{\frac{1}{\nu}}B,$$

since we have $\min\{g_{\nu}(s), g_{\nu}(t)\} \leq g_{\nu}(x)$ by $g_{\nu}''(x) \leq 0$, for $\nu \notin [0, 1]$.

Acknowledgement: The authors thank anonymous referees for giving valuable comments and suggestions to improve our manuscript. The author (S.F.) was partially supported by JSPS KAKENHI Grant Number 16K05257.

References

- Kubo F., Ando T., Means of positive linear operators, Math. Ann., 1980, 246(3), 205-224.
- Furuichi S., Moradi H.R., On some refinements for mean inequalities, To appear in Rocky Mountain J. Math., https://projecteuclid.org/euclid.rmjm/1524880819
- Gümüş I.H., Moradi H.R., Sababheh M., More accurate operator means inequalities, J. Math. Anal. Appl., 2018, 465(1), 267-280.
- [4] Liao W., Wu J., Zhao J., New versions of reverse Young and Heinz mean inequalities with the Kantorovich constant, Taiwanese J. Math., 2015, 19(2), 467-479.
- Zuo H., Shi G., Fujii M., Refined Young inequality with Kantorovich constant, J. Math. Inequal., 2011, 5(4), 551-556. [5]
- [6] Furuichi S., Refined Young inequalities with Specht's ratio, J. Egyptian Math. Soc., 2012, 20(1), 46-49.
- Dragomir S.S., On new refinements and reverse of Young's operator inequality, arXiv:1510.01314v1.
- Furuichi S., Moradi H.R., Sababheh M., New sharp inequalities for operator means, Linear Multilinear Algebra., https://doi.org/10.1080/03081087.2018.1461189
- Tominaga M., Specht's ratio in the Young inequality, Sci. Math. Jpn., 2002, 55(3), 583-588.