

Open Mathematics

Research Article

Young Bae Jun and Eun Hwan Roh*

MBJ-neutrosophic ideals of BCK/BCI -algebras

<https://doi.org/10.1515/math-2019-0106>

Received September 6, 2018; accepted February 19, 2019

Abstract: The notion of MBJ-neutrosophic ideal is introduced, and its properties are investigated. Conditions for an MBJ-neutrosophic set to be an MBJ-neutrosophic ideal are provided. In a BCK/BCI -algebra, a condition for an MBJ-neutrosophic set to be an MBJ-neutrosophic ideal is given. In a BCK -algebra, a condition for an MBJ-neutrosophic subalgebra to be an MBJ-neutrosophic ideal is given. In a BCI -algebra, conditions for an MBJ-neutrosophic ideal to be an MBJ-neutrosophic subalgebra are considered. In an (S) - BCK -algebra, we show that every MBJ-neutrosophic ideal is an MBJ-neutrosophic \circ -subalgebra, and a characterization of an MBJ-neutrosophic ideal is established.

Keywords: MBJ-neutrosophic set, MBJ-neutrosophic subalgebra, MBJ-neutrosophic ideal, MBJ-neutrosophic \circ -subalgebra

MSC: 06F35, 03G25, 03E72

1 Introduction

Different types of uncertainties are encountered in many complex systems and/or in many practical situations like behavioral, biological and chemical etc. In order to handle uncertainties in many real applications, the fuzzy set was introduced by L.A. Zadeh [1] in 1965. The intuitionistic fuzzy set on a universe X was introduced by K. Atanassov in 1983 as a generalization of fuzzy set. As a more general platform that extends the notions of classic set, (intuitionistic) fuzzy set and interval valued (intuitionistic) fuzzy set, the notion of neutrosophic set is developed by Smarandache [2–4]. Neutrosophic algebraic structures in BCK/BCI -algebras are discussed in the papers [5–14] and [15]. In [16], the notion of MBJ-neutrosophic sets is introduced as another generalization of neutrosophic set, it is applied to BCK/BCI -algebras. Mohseni et al. [16] introduced the concept of MBJ-neutrosophic subalgebras in BCK/BCI -algebras, and investigated related properties. They gave a characterization of MBJ-neutrosophic subalgebra, and established a new MBJ-neutrosophic subalgebra by using an MBJ-neutrosophic subalgebra of a BCI -algebra. They considered the homomorphic inverse image of MBJ-neutrosophic subalgebra, and discussed translation of MBJ-neutrosophic subalgebra.

In this paper, we apply the notion of MBJ-neutrosophic sets to ideals of BCK/BI -algebras. We introduce the concept of MBJ-neutrosophic ideals in BCK/BCI -algebras, and investigate several properties. We provide a condition for an MBJ-neutrosophic subalgebra to be an MBJ-neutrosophic ideal in a BCK -algebra. We provide conditions for an MBJ-neutrosophic set to be an MBJ-neutrosophic ideal in a BCK/BCI -algebra.

Young Bae Jun: Department of Mathematics Education, Gyeongsang National University, Jinju 52828, Korea; E-mail: skywine@gmail.com

***Corresponding Author: Eun Hwan Roh:** Department of Mathematics Education, Chinju National University of Education, Jinju 52673, Korea; E-mail: ehroh9988@gmail.com

We discuss relations between MBJ-neutrosophic subalgebras, MBJ-neutrosophic \circ -subalgebras and MBJ-neutrosophic ideals. In a BCI-algebra, we provide conditions for an MBJ-neutrosophic ideal to be an MBJ-neutrosophic subalgebra. In an (S)-BCK-algebra, we consider a characterization of an MBJ-neutrosophic ideal.

2 Preliminaries

By a BCI-algebra, we mean a set X with a binary operation $*$ and a special element 0 that satisfies the following conditions:

- (I) $((x * y) * (x * z)) * (z * y) = 0$,
- (II) $(x * (x * y)) * y = 0$,
- (III) $x * x = 0$,
- (IV) $x * y = 0, y * x = 0 \Rightarrow x = y$

for all $x, y, z \in X$. If a BCI-algebra X satisfies the following identity:

$$(V) (\forall x \in X) (0 * x = 0),$$

then X is called a BCK-algebra.

By a weakly BCK-algebra (see [17]), we mean a BCI-algebra X satisfying $0 * x \leq x$ for all $x \in X$.

Every BCK/BCI-algebra X satisfies the following conditions:

$$(\forall x \in X) (x * 0 = x), \quad (2.1)$$

$$(\forall x, y, z \in X) (x \leq y \Rightarrow x * z \leq y * z, z * y \leq z * x), \quad (2.2)$$

$$(\forall x, y, z \in X) ((x * y) * z = (x * z) * y), \quad (2.3)$$

$$(\forall x, y, z \in X) ((x * z) * (y * z) \leq x * y), \quad (2.4)$$

where $x \leq y$ if and only if $x * y = 0$. Any BCI-algebra X satisfies the following conditions (see [17]):

$$(\forall x, y \in X) (x * (x * (x * y)) = x * y), \quad (2.5)$$

$$(\forall x, y \in X) (0 * (x * y) = (0 * x) * (0 * y)). \quad (2.6)$$

A BCI-algebra X is said to be *p-semisimple* (see [17]) if

$$(\forall x \in X) (0 * (0 * x) = x). \quad (2.7)$$

In a *p-semisimple* BCI-algebra X , the following holds:

$$(\forall x, y \in X) (0 * (x * y) = y * x, x * (x * y) = y). \quad (2.8)$$

A BCI-algebra X is said to be *associative* (see [17]) if

$$(\forall x, y, z \in X) ((x * y) * z = x * (y * z)). \quad (2.9)$$

By an (S)-BCK-algebra, we mean a BCK-algebra X such that, for any $x, y \in X$, the set

$$\{z \in X \mid z * x \leq y\}$$

has the greatest element, written by $x \circ y$ (see [18]).

A nonempty subset S of a BCK/BCI-algebra X is called a *subalgebra* of X if $x * y \in S$ for all $x, y \in S$. A subset I of a BCK/BCI-algebra X is called an *ideal* of X if it satisfies:

$$0 \in I, \quad (2.10)$$

$$(\forall x \in X) (\forall y \in I) (x * y \in I \Rightarrow x \in I). \quad (2.11)$$

A subset I of a BCI -algebra X is called a *closed ideal* of X (see [17]) if it is an ideal of X which satisfies:

$$(\forall x \in X)(x \in I \Rightarrow 0 * x \in I). \quad (2.12)$$

By an *interval number* we mean a closed subinterval $\tilde{a} = [a^-, a^+]$ of I , where $0 \leq a^- \leq a^+ \leq 1$. Denote by $[I]$ the set of all interval numbers. Let us define what is known as *refined minimum* (briefly, rmin) and *refined maximum* (briefly, rmax) of two elements in $[I]$. We also define the symbols “ \succeq ”, “ \preceq ”, “ $=$ ” in case of two elements in $[I]$. Consider two interval numbers $\tilde{a}_1 := [a_1^-, a_1^+]$ and $\tilde{a}_2 := [a_2^-, a_2^+]$. Then

$$\begin{aligned} \text{rmin} \{ \tilde{a}_1, \tilde{a}_2 \} &= [\min \{ a_1^-, a_2^- \}, \min \{ a_1^+, a_2^+ \}], \\ \text{rmax} \{ \tilde{a}_1, \tilde{a}_2 \} &= [\max \{ a_1^-, a_2^- \}, \max \{ a_1^+, a_2^+ \}], \\ \tilde{a}_1 \succeq \tilde{a}_2 &\Leftrightarrow a_1^- \geq a_2^-, a_1^+ \geq a_2^+, \end{aligned}$$

and similarly we may have $\tilde{a}_1 \preceq \tilde{a}_2$ and $\tilde{a}_1 = \tilde{a}_2$. To say $\tilde{a}_1 \succ \tilde{a}_2$ (resp. $\tilde{a}_1 \prec \tilde{a}_2$) we mean $\tilde{a}_1 \succeq \tilde{a}_2$ and $\tilde{a}_1 \neq \tilde{a}_2$ (resp. $\tilde{a}_1 \preceq \tilde{a}_2$ and $\tilde{a}_1 \neq \tilde{a}_2$). Let $\tilde{a}_i \in [I]$ where $i \in \Lambda$. We define

$$\text{rinf } \tilde{a}_i = \left[\inf_{i \in \Lambda} a_i^-, \inf_{i \in \Lambda} a_i^+ \right] \quad \text{and} \quad \text{rsup } \tilde{a}_i = \left[\sup_{i \in \Lambda} a_i^-, \sup_{i \in \Lambda} a_i^+ \right].$$

Let X be a nonempty set. A function $A : X \rightarrow [I]$ is called an *interval-valued fuzzy set* (briefly, an *IVF set*) in X . Let $[I]^X$ stand for the set of all IVF sets in X . For every $A \in [I]^X$ and $x \in X$, $A(x) = [A^-(x), A^+(x)]$ is called the *degree of membership* of an element x to A , where $A^- : X \rightarrow I$ and $A^+ : X \rightarrow I$ are fuzzy sets in X which are called a *lower fuzzy set* and an *upper fuzzy set* in X , respectively. For simplicity, we denote $A = [A^-, A^+]$.

Let X be a non-empty set. A *neutrosophic set* (NS) in X (see [3]) is a structure of the form:

$$A := \{ \langle x; A_T(x), A_I(x), A_F(x) \rangle \mid x \in X \},$$

where $A_T : X \rightarrow [0, 1]$ is a truth membership function, $A_I : X \rightarrow [0, 1]$ is an indeterminate membership function, and $A_F : X \rightarrow [0, 1]$ is a false membership function. For the sake of simplicity, we shall use the symbol $A = (A_T, A_I, A_F)$ for the neutrosophic set

$$A := \{ \langle x; A_T(x), A_I(x), A_F(x) \rangle \mid x \in X \}.$$

We refer the reader to the books [17, 18] for further information regarding BCK/BCI -algebras, and to the site “<http://fs.gallup.unm.edu/neutrosophy.htm>” for further information regarding neutrosophic set theory.

Let X be a non-empty set. By an *MBJ-neutrosophic set* in X (see [16]), we mean a structure of the form:

$$\mathcal{A} := \{ \langle x; M_A(x), \tilde{B}_A(x), J_A(x) \rangle \mid x \in X \},$$

where M_A and J_A are fuzzy sets in X , which are called a truth membership function and a false membership function, respectively, and \tilde{B}_A is an IVF set in X which is called an indeterminate interval-valued membership function.

For the sake of simplicity, we shall use the symbol $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ for the MBJ-neutrosophic set

$$\mathcal{A} := \{ \langle x; M_A(x), \tilde{B}_A(x), J_A(x) \rangle \mid x \in X \}.$$

Let X be a BCK/BCI -algebra. An MBJ-neutrosophic set $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ in X is called an *MBJ-neutrosophic subalgebra* of X (see [16]) if it satisfies:

$$(\forall x, y \in X) \left(\begin{aligned} &M_A(x * y) \geq \min \{ M_A(x), M_A(y) \}, \\ &\tilde{B}_A(x * y) \succeq \text{rmin} \{ \tilde{B}_A(x), \tilde{B}_A(y) \}, \\ &J_A(x * y) \leq \max \{ J_A(x), J_A(y) \}. \end{aligned} \right) \quad (2.13)$$

3 MBJ-neutrosophic ideals of BCK/BCI -algebras

Definition 3.1. Let X be a BCK/BCI -algebra. An MBJ-neutrosophic set $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ in X is called an *MBJ-neutrosophic ideal* of X if it satisfies:

$$(\forall x \in X) \begin{pmatrix} M_A(0) \geq M_A(x) \\ \tilde{B}_A(0) \succeq \tilde{B}_A(x) \\ J_A(0) \leq J_A(x) \end{pmatrix} \quad (3.1)$$

and

$$(\forall x, y \in X) \begin{pmatrix} M_A(x) \geq \min\{M_A(x * y), M_A(y)\} \\ \tilde{B}_A(x) \succeq \text{rmin}\{\tilde{B}_A(x * y), \tilde{B}_A(y)\} \\ J_A(x) \leq \max\{J_A(x * y), J_A(y)\} \end{pmatrix}. \quad (3.2)$$

An MBJ-neutrosophic ideal $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ of a BCI -algebra X is said to be *closed* if

$$(\forall x \in X) \begin{pmatrix} M_A(0 * x) \geq M_A(x) \\ \tilde{B}_A(0 * x) \succeq \tilde{B}_A(x) \\ J_A(0 * x) \leq J_A(x) \end{pmatrix}. \quad (3.3)$$

Example 3.2. Consider a set $X = \{0, 1, 2, a\}$ with the binary operation $*$ which is given in Table 1. Then $(X; *, 0)$ is a BCI -algebra (see [17]). Let $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ be an MBJ-neutrosophic set in X defined by Table 2.

Table 1: Cayley table for the binary operation “ $*$ ”.

$*$	0	1	2	a
0	0	0	0	a
1	1	0	0	a
2	2	2	0	a
a	a	a	a	0

Table 2: MBJ-neutrosophic set $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$.

X	$M_A(x)$	$\tilde{B}_A(x)$	$J_A(x)$
0	0.7	[1.0, 1.0]	0.2
1	0.5	[0.2, 0.6]	0.2
2	0.4	[0.2, 0.6]	0.7
a	0.3	[0.2, 0.6]	0.7

It is routine to verify that $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is a closed MBJ-neutrosophic ideal of X .

Proposition 3.3. Let X be a BCK/BCI -algebra. Then every MBJ-neutrosophic ideal $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ of X satisfies the following assertion.

$$x * y \leq z \Rightarrow \begin{cases} M_A(x) \geq \min\{M_A(y), M_A(z)\}, \\ \tilde{B}_A(x) \succeq \text{rmin}\{\tilde{B}_A(y), \tilde{B}_A(z)\}, \\ J_A(x) \leq \max\{J_A(y), J_A(z)\} \end{cases} \quad (3.4)$$

for all $x, y, z \in X$.

Proof. Let $x, y, z \in X$ be such that $x * y \leq z$. Then

$$M_A(x * y) \geq \min\{M_A((x * y) * z), M_A(z)\} = \min\{M_A(0), M_A(z)\} = M_A(z),$$

$$\tilde{B}_A(x * y) \geq \text{rmin}\{\tilde{B}_A((x * y) * z), \tilde{B}_A(z)\} = \text{rmin}\{\tilde{B}_A(0), \tilde{B}_A(z)\} = \tilde{B}_A(z),$$

and

$$J_A(x * y) \leq \max\{J_A((x * y) * z), J_A(z)\} = \max\{J_A(0), J_A(z)\} = J_A(z).$$

It follows that

$$M_A(x) \geq \min\{M_A(x * y), M_A(y)\} = \min\{M_A(y), M_A(z)\},$$

$$\tilde{B}_A(x) \geq \text{rmin}\{\tilde{B}_A(x * y), \tilde{B}_A(y)\} = \text{rmin}\{\tilde{B}_A(y), \tilde{B}_A(z)\},$$

and

$$J_A(x) \leq \max\{J_A(x * y), J_A(y)\} = \max\{J_A(y), J_A(z)\}.$$

This completes the proof. \square

Theorem 3.4. Every MBJ-neutrosophic set in a BCK/BCI-algebra X satisfying (3.1) and (3.4) is an MBJ-neutrosophic ideal of X .

Proof. Let $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ be an MBJ-neutrosophic set in X satisfying (3.1) and (3.4). Note that $x * (x * y) \leq y$ for all $x, y \in X$. It follows from (3.4) that

$$M_A(x) \geq \min\{M_A(x * y), M_A(y)\},$$

$$\tilde{B}_A(x) \geq \text{rmin}\{\tilde{B}_A(x * y), \tilde{B}_A(y)\},$$

and

$$J_A(x) \leq \max\{J_A(x * y), J_A(y)\}.$$

Therefore $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic ideal of X . \square

Theorem 3.5. Given an MBJ-neutrosophic set $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ in a BCK/BCI-algebra X , if (M_A, J_A) is an intuitionistic fuzzy ideal of X , and B_A^- and B_A^+ are fuzzy ideals of X , then $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic ideal of X .

Proof. It is sufficient to show that \tilde{B}_A satisfies the condition

$$(\forall x \in X)(\tilde{B}_A(0) \geq \tilde{B}_A(x)) \quad (3.5)$$

and

$$(\forall x, y \in X)(\tilde{B}_A(x) \geq \text{rmin}\{\tilde{B}_A(x * y), \tilde{B}_A(y)\}). \quad (3.6)$$

For any $x, y \in X$, we get

$$\tilde{B}_A(0) = [B_A^-(0), B_A^+(0)] \geq [B_A^-(x), B_A^+(x)] = \tilde{B}_A(x)$$

and

$$\begin{aligned} \tilde{B}_A(x) &= [B_A^-(x), B_A^+(x)] \\ &\geq [\min\{B_A^-(x * y), B_A^-(y)\}, \min\{B_A^+(x * y), B_A^+(y)\}] \\ &= \text{rmin}\{[B_A^-(x * y), B_A^+(x * y)], [B_A^-(y), B_A^+(y)]\} \\ &= \text{rmin}\{\tilde{B}_A(x * y), \tilde{B}_A(y)\}. \end{aligned}$$

Therefore $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic ideal of X . \square

If $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic ideal of a BCK/BCI-algebra X , then

$$\begin{aligned} [B_A^-(x), B_A^+(x)] &= \tilde{B}_A(x) \succeq \text{rmin}\{\tilde{B}_A(x * y), \tilde{B}_A(y)\} \\ &= \text{rmin}\{[B_A^-(x * y), B_A^+(x * y)], [B_A^-(y), B_A^+(y)]\} \\ &= [\min\{B_A^-(x * y), B_A^-(y)\}, \min\{B_A^+(x * y), B_A^+(y)\}] \end{aligned}$$

for all $x, y \in X$. It follows that $B_A^-(x) \geq \min\{B_A^-(x * y), B_A^-(y)\}$ and $B_A^+(x) \geq \min\{B_A^+(x * y), B_A^+(y)\}$. Thus B_A^- and B_A^+ are fuzzy ideals of X . But (M_A, J_A) is not an intuitionistic fuzzy ideal of X as seen in Example 3.2. This shows that the converse of Theorem 3.5 is not true.

Given an MBJ-neutrosophic set $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ in a BCK/BCI-algebra X , we consider the following sets.

$$\begin{aligned} U(M_A; t) &:= \{x \in X \mid M_A(x) \geq t\}, \\ U(\tilde{B}_A; [\delta_1, \delta_2]) &:= \{x \in X \mid \tilde{B}_A(x) \succeq [\delta_1, \delta_2]\}, \\ L(J_A; s) &:= \{x \in X \mid J_A(x) \leq s\}, \end{aligned}$$

where $t, s \in [0, 1]$ and $[\delta_1, \delta_2] \in [I]$.

Theorem 3.6. *An MBJ-neutrosophic set $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ in a BCK/BCI-algebra X is an MBJ-neutrosophic ideal of X if and only if the non-empty sets $U(M_A; t)$, $U(\tilde{B}_A; [\delta_1, \delta_2])$ and $L(J_A; s)$ are ideals of X for all $t, s \in [0, 1]$ and $[\delta_1, \delta_2] \in [I]$.*

Proof. Suppose that $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic ideal of X . Let $t, s \in [0, 1]$ and $[\delta_1, \delta_2] \in [I]$ be such that $U(M_A; t)$, $U(\tilde{B}_A; [\delta_1, \delta_2])$ and $L(J_A; s)$ are non-empty. Obviously, $0 \in U(M_A; t) \cap U(\tilde{B}_A; [\delta_1, \delta_2]) \cap L(J_A; s)$. For any $x, y, a, b, u, v \in X$, if $x * y \in U(M_A; t)$, $y \in U(M_A; t)$, $a * b \in U(\tilde{B}_A; [\delta_1, \delta_2])$, $b \in U(\tilde{B}_A; [\delta_1, \delta_2])$, $u * v \in L(J_A; s)$ and $v \in L(J_A; s)$, then

$$\begin{aligned} M_A(x) &\geq \min\{M_A(x * y), M_A(y)\} \geq \min\{t, t\} = t, \\ \tilde{B}_A(a) &\succeq \text{rmin}\{\tilde{B}_A(a * b), \tilde{B}_A(b)\} \succeq \text{rmin}\{[\delta_1, \delta_2], [\delta_1, \delta_2]\} = [\delta_1, \delta_2], \\ J_A(u) &\leq \max\{J_A(u * v), J_A(v)\} \leq \min\{s, s\} = s, \end{aligned}$$

and so $x \in U(M_A; t)$, $a \in U(\tilde{B}_A; [\delta_1, \delta_2])$ and $u \in L(J_A; s)$. Therefore $U(M_A; t)$, $U(\tilde{B}_A; [\delta_1, \delta_2])$ and $L(J_A; s)$ are ideals of X .

Conversely, assume that the non-empty sets $U(M_A; t)$, $U(\tilde{B}_A; [\delta_1, \delta_2])$ and $L(J_A; s)$ are ideals of X for all $t, s \in [0, 1]$ and $[\delta_1, \delta_2] \in [I]$. Assume that $M_A(0) < M_A(a)$, $\tilde{B}_A(0) \prec \tilde{B}_A(a)$ and $J_A(0) > J_A(a)$ for some $a \in X$. Then $0 \notin U(M_A; M_A(a)) \cap U(\tilde{B}_A; \tilde{B}_A(a)) \cap L(J_A; J_A(a))$, which is a contradiction. Hence $M_A(0) \geq M_A(x)$, $\tilde{B}_A(0) \succeq \tilde{B}_A(x)$ and $J_A(0) \leq J_A(x)$ for all $x \in X$. If

$$M_A(a_0) < \min\{M_A(a_0 * b_0), M_A(b_0)\}$$

for some $a_0, b_0 \in X$, then $a_0 * b_0 \in U(M_A; t_0)$ and $b_0 \in U(M_A; t_0)$ but $a_0 \notin U(M_A; t_0)$ for $t_0 := \min\{M_A(a_0 * b_0), M_A(b_0)\}$. This is a contradiction, and thus $M_A(a) \geq \min\{M_A(a * b), M_A(b)\}$ for all $a, b \in X$. Similarly, we can show that $J_A(a) \leq \max\{J_A(a * b), J_A(b)\}$ for all $a, b \in X$. Suppose that $\tilde{B}_A(a_0) \prec \text{rmin}\{\tilde{B}_A(a_0 * b_0), \tilde{B}_A(b_0)\}$ for some $a_0, b_0 \in X$. Let $\tilde{B}_A(a_0 * b_0) = [\lambda_1, \lambda_2]$, $\tilde{B}_A(b_0) = [\lambda_3, \lambda_4]$ and $\tilde{B}_A(a_0) = [\delta_1, \delta_2]$. Then

$$[\delta_1, \delta_2] \prec \text{rmin}\{[\lambda_1, \lambda_2], [\lambda_3, \lambda_4]\} = [\min\{\lambda_1, \lambda_3\}, \min\{\lambda_2, \lambda_4\}],$$

and so $\delta_1 < \min\{\lambda_1, \lambda_3\}$ and $\delta_2 < \min\{\lambda_2, \lambda_4\}$. Taking

$$[\gamma_1, \gamma_2] := \frac{1}{2} (\tilde{B}_A(a_0) + \text{rmin}\{\tilde{B}_A(a_0 * b_0), \tilde{B}_A(b_0)\})$$

implies that

$$\begin{aligned} [\gamma_1, \gamma_2] &= \frac{1}{2} ([\delta_1, \delta_2] + [\min\{\lambda_1, \lambda_3\}, \min\{\lambda_2, \lambda_4\}]) \\ &= [\frac{1}{2}(\delta_1 + \min\{\lambda_1, \lambda_3\}), \frac{1}{2}(\delta_2 + \min\{\lambda_2, \lambda_4\})]. \end{aligned}$$

It follows that

$$\min\{\lambda_1, \lambda_3\} > \gamma_1 = \frac{1}{2}(\delta_1 + \min\{\lambda_1, \lambda_3\}) > \delta_1$$

and

$$\min\{\lambda_2, \lambda_4\} > \gamma_2 = \frac{1}{2}(\delta_2 + \min\{\lambda_2, \lambda_4\}) > \delta_2.$$

Hence $[\min\{\lambda_1, \lambda_3\}, \min\{\lambda_2, \lambda_4\}] \succ [\gamma_1, \gamma_2] \succ [\delta_1, \delta_2] = \tilde{B}_A(a_0)$, and therefore $a_0 \notin U(\tilde{B}_A; [\gamma_1, \gamma_2])$. On the other hand,

$$\tilde{B}_A(a_0 * b_0) = [\lambda_1, \lambda_2] \succeq [\min\{\lambda_1, \lambda_3\}, \min\{\lambda_2, \lambda_4\}] \succ [\gamma_1, \gamma_2]$$

and

$$\tilde{B}_A(b_0) = [\lambda_3, \lambda_4] \succeq [\min\{\lambda_1, \lambda_3\}, \min\{\lambda_2, \lambda_4\}] \succ [\gamma_1, \gamma_2],$$

that is, $a_0 * b_0, b_0 \in U(\tilde{B}_A; [\gamma_1, \gamma_2])$. This is a contradiction, and therefore $\tilde{B}_A(x) \succeq \text{rmin}\{\tilde{B}_A(x * y), \tilde{B}_A(y)\}$ for all $x, y \in X$. Consequently $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic ideal of X . \square

Theorem 3.7. Given an ideal I of a BCK/BCI-algebra X , let $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ be an MBJ-neutrosophic set in X defined by

$$M_A(x) = \begin{cases} t & \text{if } x \in I, \\ 0 & \text{otherwise,} \end{cases} \quad \tilde{B}_A(x) = \begin{cases} [\gamma_1, \gamma_2] & \text{if } x \in I, \\ [0, 0] & \text{otherwise,} \end{cases} \quad J_A(x) = \begin{cases} s & \text{if } x \in I, \\ 1 & \text{otherwise,} \end{cases} \quad (3.7)$$

where $t \in (0, 1]$, $s \in [0, 1]$ and $\gamma_1, \gamma_2 \in (0, 1]$ with $\gamma_1 < \gamma_2$. Then $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic ideal of X such that $U(M_A; t) = U(\tilde{B}_A; [\gamma_1, \gamma_2]) = L(J_A; s) = I$.

Proof. Let $x, y \in X$. If $x * y \in I$ and $y \in I$, then $x \in I$ and so

$$\begin{aligned} M_A(x) &= t = \min\{M_A(x * y), M_A(y)\}, \\ \tilde{B}_A(x) &= [\gamma_1, \gamma_2] = \text{rmin}\{[\gamma_1, \gamma_2], [\gamma_1, \gamma_2]\} = \text{rmin}\{\tilde{B}_A(x * y), \tilde{B}_A(y)\}, \\ J_A(x) &= s = \max\{J_A(x * y), J_A(y)\}. \end{aligned}$$

If any one of $x * y$ and y is contained in I , say $x * y \in I$, then $M_A(x * y) = t$, $\tilde{B}_A(x * y) = [\gamma_1, \gamma_2]$, $J_A(x * y) = s$, $M_A(y) = 0$, $\tilde{B}_A(y) = [0, 0]$ and $J_A(y) = 1$. Hence

$$\begin{aligned} M_A(x) &\geq 0 = \min\{t, 0\} = \min\{M_A(x * y), M_A(y)\}, \\ \tilde{B}_A(x) &\succeq [0, 0] = \text{rmin}\{[\gamma_1, \gamma_2], [0, 0]\} = \text{rmin}\{\tilde{B}_A(x * y), \tilde{B}_A(y)\}, \\ J_A(x) &\leq 1 = \max\{s, 1\} = \max\{J_A(x * y), J_A(y)\}. \end{aligned}$$

If $x * y, y \notin I$, then $M_A(x * y) = 0 = M_A(y)$, $\tilde{B}_A(x * y) = [0, 0] = \tilde{B}_A(y)$ and $J_A(x * y) = 1 = J_A(y)$. It follows that

$$\begin{aligned} M_A(x) &\geq 0 = \min\{0, 0\} = \min\{M_A(x * y), M_A(y)\}, \\ \tilde{B}_A(x) &\succeq [0, 0] = \text{rmin}\{[0, 0], [0, 0]\} = \text{rmin}\{\tilde{B}_A(x * y), \tilde{B}_A(y)\}, \\ J_A(x) &\leq 1 = \max\{1, 1\} = \max\{J_A(x * y), J_A(y)\}. \end{aligned}$$

It is obvious that $M_A(0) \geq M_A(x)$, $\tilde{B}_A(0) \succeq \tilde{B}_A(x)$ and $J_A(0) \leq J_A(x)$ for all $x \in X$. Therefore $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic ideal of X . Obviously, we have $U(M_A; t) = U(\tilde{B}_A; [\gamma_1, \gamma_2]) = L(J_A; s) = I$. \square

Theorem 3.8. For any non-empty subset I of X , let $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ be an MBJ-neutrosophic set in X which is given in (3.7). If $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic ideal of X , then I is an ideal of X .

Proof. Obviously, $0 \in I$. Let $x, y \in X$ be such that $x * y \in I$ and $y \in I$. Then $M_A(x * y) = t = M_A(y)$, $\tilde{B}_A(x * y) = [\gamma_1, \gamma_2] = \tilde{B}_A(y)$ and $J_A(x * y) = s = J_A(y)$. Thus

$$\begin{aligned} M_A(x) &\geq \min\{M_A(x * y), M_A(y)\} = t, \\ \tilde{B}_A(x) &\succeq \text{rmin}\{\tilde{B}_A(x * y), \tilde{B}_A(y)\} = [\gamma_1, \gamma_2], \\ J_A(x) &\leq \max\{J_A(x * y), J_A(y)\} = s, \end{aligned}$$

and hence $x \in I$. Therefore I is an ideal of X . \square

Theorem 3.9. *In a BCK-algebra, every MBI-neutrosophic ideal is an MBI-neutrosophic subalgebra.*

Proof. Let $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ be an MBI-neutrosophic ideal of a BCK-algebra X . Since $(x * y) * x \leq y$ for all $x, y \in X$, it follows from Proposition 3.3 that

$$\begin{aligned} M_A(x * y) &\geq \min\{M_A(x), M_A(y)\}, \\ \tilde{B}_A(x * y) &\succeq \text{rmin}\{\tilde{B}_A(x), \tilde{B}_A(y)\}, \\ J_A(x * y) &\leq \max\{J_A(x), J_A(y)\} \end{aligned}$$

for all $x, y \in X$. Hence $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBI-neutrosophic subalgebra of a BCK-algebra X . \square

The converse of Theorem 3.9 may not be true as seen in the following example.

Example 3.10. Consider a BCK-algebra $X = \{0, 1, 2, 3\}$ with the binary operation $*$ which is given in Table 3. Let $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ be an MBI-neutrosophic set in X defined by Table 4. Then $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBI-neutrosophic subalgebra of X , but it is not an MBI-neutrosophic ideal of X since

$$\tilde{B}_A(1) \not\succeq \text{rmin}\{\tilde{B}_A(1 * 2), \tilde{B}_A(2)\}.$$

Table 3: Cayley table for the binary operation “ $*$ ”.

$*$	0	1	2	3
0	0	0	0	0
1	1	0	0	1
2	2	1	0	2
3	3	3	3	0

Table 4: MBI-neutrosophic set $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$.

X	$M_A(x)$	$\tilde{B}_A(x)$	$J_A(x)$
0	0.7	[0.3, 0.8]	0.2
1	0.4	[0.2, 0.6]	0.3
2	0.4	[0.3, 0.8]	0.4
3	0.6	[0.2, 0.6]	0.5

We provide a condition for an MBJ-neutrosophic subalgebra to be an MBJ-neutrosophic ideal in a BCK-algebra.

Theorem 3.11. Let $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ be an MBJ-neutrosophic subalgebra of a BCK-algebra X satisfying the condition (3.4). Then $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic ideal of X .

Proof. For any $x \in X$, we get

$$M_A(0) = M_A(x * x) \geq \min\{M_A(x), M_A(x)\} = M_A(x),$$

$$\begin{aligned}\tilde{B}_A(0) &= \tilde{B}_A(x * x) \succeq \text{rmin}\{\tilde{B}_A(x), \tilde{B}_A(x)\} \\ &= \text{rmin}\{[B_A^-(x), B_A^+(x)], [B_A^-(x), B_A^+(x)]\} \\ &= [B_A^-(x), B_A^+(x)] = \tilde{B}_A(x),\end{aligned}$$

and

$$J_A(0) = J_A(x * x) \leq \max\{J_A(x), J_A(x)\} = J_A(x).$$

Since $x * (x * y) \leq y$ for all $x, y \in X$, it follows from (3.4) that

$$\begin{aligned}M_A(x) &\geq \min\{M_A(x * y), M_A(y)\}, \\ \tilde{B}_A(x) &\succeq \text{rmin}\{\tilde{B}_A(x * y), \tilde{B}_A(y)\}, \\ J_A(x) &\leq \max\{J_A(x * y), J_A(y)\}\end{aligned}$$

for all $x, y \in X$. Therefore $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic ideal of X . \square

Theorem 3.9 is not true in a BCI-algebra as seen in the following example.

Example 3.12. Let $(Y, *, 0)$ be a BCI-algebra and let $(\mathbb{Z}, -, 0)$ be an adjoint BCI-algebra of the additive group $(\mathbb{Z}, +, 0)$ of integers. Then $X = Y \times \mathbb{Z}$ is a BCI-algebra and $I = Y \times \mathbb{N}$ is an ideal of X where \mathbb{N} is the set of all non-negative integers (see [17]). Let $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ be an MBJ-neutrosophic set in X defined by

$$M_A(x) = \begin{cases} t & \text{if } x \in I, \\ 0 & \text{otherwise,} \end{cases} \quad \tilde{B}_A(x) = \begin{cases} [\gamma_1, \gamma_2] & \text{if } x \in I, \\ [0, 0] & \text{otherwise,} \end{cases} \quad J_A(x) = \begin{cases} s & \text{if } x \in I, \\ 1 & \text{otherwise,} \end{cases} \quad (3.8)$$

where $t \in (0, 1]$, $s \in [0, 1)$ and $\gamma_1, \gamma_2 \in (0, 1]$ with $\gamma_1 < \gamma_2$. Then $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic ideal of X by Theorem 3.7. But it is not an MBJ-neutrosophic subalgebra of X since

$$M_A((0, 0) * (0, 1)) = M_A((0, -1)) = 0 < t = \min\{M_A((0, 0)), M_A(0, 1)\},$$

$$\tilde{B}_A((0, 0) * (0, 2)) = \tilde{B}_A((0, -2)) = [0, 0] \prec [\gamma_1, \gamma_2] = \text{rmin}\{\tilde{B}_A((0, 0)), \tilde{B}_A(0, 2)\},$$

and/or

$$J_A((0, 0) * (0, 3)) = J_A((0, -3)) = 1 > s = \max\{J_A((0, 0)), J_A(0, 3)\}.$$

Definition 3.13. An MBJ-neutrosophic ideal $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ of a BCI-algebra X is said to be closed if

$$(\forall x \in X)(M_A(0 * x) \geq M_A(x), \tilde{B}_A(0 * x) \succeq \tilde{B}_A(x), J_A(0 * x) \leq J_A(x)). \quad (3.9)$$

Theorem 3.14. In a BCI-algebra, every closed MBJ-neutrosophic ideal is an MBJ-neutrosophic subalgebra.

Proof. Let $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ be a closed MBJ-neutrosophic ideal of a BCI-algebra X . Using (3.2), (2.3), (III) and (3.3), we have

$$M_A(x * y) \geq \min\{M_A((x * y) * x), M_A(x)\} = \min\{M_A(0 * y), M_A(x)\} \geq \min\{M_A(y), M_A(x)\},$$

$$\tilde{B}_A(x * y) \succeq \text{rmin}\{\tilde{B}_A((x * y) * x), \tilde{B}_A(x)\} = \text{rmin}\{\tilde{B}_A(0 * y), \tilde{B}_A(x)\} \succeq \text{rmin}\{\tilde{B}_A(y), \tilde{B}_A(x)\},$$

and

$$J_A(x * y) \leq \max\{J_A((x * y) * x), J_A(x)\} = \max\{J_A(0 * y), J_A(x)\} \leq \max\{J_A(y), J_A(x)\}$$

for all $x, y \in X$. Hence $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic subalgebra of X . \square

Theorem 3.15. *In a weakly BCK-algebra, every MBJ-neutrosophic ideal is closed.*

Proof. Let $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ be an MBJ-neutrosophic ideal of a weakly BCK-algebra X . For any $x \in X$, we obtain

$$M_A(0 * x) \geq \min\{M_A((0 * x) * x), M_A(x)\} = \min\{M_A(0), M_A(x)\} = M_A(x),$$

$$\tilde{B}_A(0 * x) \succeq \text{rmin}\{\tilde{B}_A((0 * x) * x), \tilde{B}_A(x)\} = \text{rmin}\{\tilde{B}_A(0), \tilde{B}_A(x)\} = \tilde{B}_A(x),$$

and

$$J_A(0 * x) \leq \max\{J_A((0 * x) * x), J_A(x)\} = \max\{J_A(0), J_A(x)\} = J_A(x).$$

Therefore $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is a closed MBJ-neutrosophic ideal of X . \square

Corollary 3.16. *In a weakly BCK-algebra, every MBJ-neutrosophic ideal is an MBJ-neutrosophic subalgebra.*

The following example shows that any MBJ-neutrosophic subalgebra is not an MBJ-neutrosophic ideal in a BCI-algebra.

Example 3.17. Consider a BCI-algebra $X = \{0, a, b, c, d, e\}$ with the $*$ -operation in Table 5. Let $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ be an MBJ-neutrosophic set in X defined by Table 6. It is routine to verify that $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic subalgebra of X . But it is not an MBJ-neutrosophic ideal of X since

$$M_A(d) < \min\{M_A(d * c), M_A(c)\},$$

$$\tilde{B}_A(d) \prec \text{rmin}\{\tilde{B}_A(d * c), \tilde{B}_A(c)\},$$

and/or

$$J_A(d) > \max\{J_A(d * c), J_A(c)\}.$$

Table 5: Cayley table for the binary operation “ $*$ ”.

$*$	0	a	b	c	d	e
0	0	0	c	b	c	c
a	a	0	c	b	c	c
b	b	b	0	c	0	0
c	c	c	b	0	b	b
d	d	b	a	c	0	a
e	e	b	a	c	a	0

Table 6: MBJ-neutrosophic set $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$.

X	$M_A(x)$	$\tilde{B}_A(x)$	$J_A(x)$
0	0.7	[0.4, 0.9]	0.3
a	0.4	[0.04, 0.45]	0.6
b	0.7	[0.4, 0.9]	0.3
c	0.7	[0.4, 0.9]	0.3
d	0.4	[0.04, 0.45]	0.6
e	0.4	[0.04, 0.45]	0.6

Theorem 3.18. In a p -semisimple BCI-algebra X , the following are equivalent.

- (1) $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is a closed MBJ-neutrosophic ideal of X .
- (2) $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic subalgebra of X .

Proof. (1) \Rightarrow (2). See Theorem 3.14.

(2) \Rightarrow (1). Suppose that $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic subalgebra of X . For any $x \in X$, we get

$$M_A(0) = M_A(x * x) \geq \min\{M_A(x), M_A(x)\} = M_A(x),$$

$$\tilde{B}_A(0) = \tilde{B}_A(x * x) \succeq \text{rmin}\{\tilde{B}_A(x), \tilde{B}_A(x)\} = \tilde{B}_A(x),$$

and

$$J_A(0) = J_A(x * x) \leq \max\{J_A(x), J_A(x)\} = J_A(x).$$

Hence $M_A(0 * x) \geq \min\{M_A(0), M_A(x)\} = M_A(x)$, $\tilde{B}_A(0 * x) \succeq \text{rmin}\{\tilde{B}_A(0), \tilde{B}_A(x)\} = \tilde{B}_A(x)$ and $J_A(0 * x) \leq \max\{J_A(0), J_A(x)\} = J_A(x)$ for all $x \in X$. Let $x, y \in X$. Then

$$\begin{aligned} M_A(x) &= M_A(y * (y * x)) \geq \min\{M_A(y), M_A(y * x)\} \\ &= \min\{M_A(y), M_A(0 * (x * y))\} \\ &\geq \min\{M_A(x * y), M_A(y)\}, \end{aligned}$$

$$\begin{aligned} \tilde{B}_A(x) &= \tilde{B}_A(y * (y * x)) \succeq \text{rmin}\{\tilde{B}_A(y), \tilde{B}_A(y * x)\} \\ &= \text{rmin}\{\tilde{B}_A(y), \tilde{B}_A(0 * (x * y))\} \\ &\succeq \text{rmin}\{\tilde{B}_A(x * y), \tilde{B}_A(y)\} \end{aligned}$$

and

$$\begin{aligned} J_A(x) &= J_A(y * (y * x)) \leq \max\{J_A(y), J_A(y * x)\} \\ &= \max\{J_A(y), J_A(0 * (x * y))\} \\ &\leq \max\{J_A(x * y), J_A(y)\}. \end{aligned}$$

Therefore $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is a closed MBJ-neutrosophic ideal of X . □

Since every associative BCI-algebra is p -semisimple, we have the following corollary.

Corollary 3.19. In an associative BCI-algebra X , the following are equivalent.

- (1) $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is a closed MBJ-neutrosophic ideal of X .
- (2) $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic subalgebra of X .

Corollary 3.20. In a BCI-algebra X , consider the following conditions

- (1) every element x in X is minimal.
- (2) $X = \{0 * x \mid x \in X\}$.
- (3) $(\forall x, y \in X) (x * (0 * y) = y * (0 * x))$.
- (4) $(\forall x \in X) (0 * x = 0 \Rightarrow x = 0)$.
- (5) $(\forall a, x \in X) (a * (a * x) = x)$.
- (6) $(\forall a \in X) X = \{a * x \mid x \in X\}$.
- (7) $(\forall x, y, a, b \in X) ((x * y) * (a * b) = (x * a) * (y * b))$.
- (8) $(\forall x, y \in X) (0 * (y * x) = x * y)$.

If one of the conditions above is valid, then the following are equivalent.

- (1) $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is a closed MBJ-neutrosophic ideal of X .
- (2) $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic subalgebra of X .

Definition 3.21. Let X be an (S)-BCK-algebra. An MBJ-neutrosophic set $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ in X is called an MBJ-neutrosophic \circ -subalgebra of X if the following assertions are valid.

$$\begin{aligned} M_A(x \circ y) &\geq \min\{M_A(x), M_A(y)\}, \\ \tilde{B}_A(x \circ y) &\succeq \text{rmin}\{\tilde{B}_A(x), \tilde{B}_A(y)\}, \\ J_A(x \circ y) &\leq \max\{J_A(x), J_A(y)\} \end{aligned} \quad (3.10)$$

for all $x, y \in X$.

Lemma 3.22. Every MBJ-neutrosophic ideal of a BCK/BCI-algebra X satisfies the following assertion.

$$(\forall x, y \in X) (x \leq y \Rightarrow M_A(x) \geq M_A(y), \tilde{B}_A(x) \succeq \tilde{B}_A(y), J_A(x) \leq J_A(y)). \quad (3.11)$$

Proof. Assume that $x \leq y$ for all $x, y \in X$. Then $x * y = 0$, and so

$$M_A(x) \geq \min\{M_A(x * y), M_A(y)\} = \min\{M_A(0), M_A(y)\} = M_A(y),$$

$$\tilde{B}_A(x) \succeq \text{rmin}\{\tilde{B}_A(x * y), \tilde{B}_A(y)\} = \text{rmin}\{\tilde{B}_A(0), \tilde{B}_A(y)\} = \tilde{B}_A(y),$$

and

$$J_A(x) \leq \max\{J_A(x * y), J_A(y)\} = \max\{J_A(0), J_A(y)\} = J_A(y).$$

This completes the proof. \square

Theorem 3.23. In an (S)-BCK-algebra, every MBJ-neutrosophic ideal is an MBJ-neutrosophic \circ -subalgebra.

Proof. Let $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ be an MBJ-neutrosophic ideal of an (S)-BCK-algebra X . Note that $(x \circ y) * x \leq y$ for all $x, y \in X$. Using Lemma 3.22 and (3.2) implies that

$$M_A(x \circ y) \geq \min\{M_A((x \circ y) * x), M_A(x)\} \geq \min\{M_A(y), M_A(x)\},$$

$$\tilde{B}_A(x \circ y) \succeq \text{rmin}\{\tilde{B}_A((x \circ y) * x), \tilde{B}_A(x)\} \succeq \text{rmin}\{\tilde{B}_A(y), \tilde{B}_A(x)\},$$

and

$$J_A(x \circ y) \leq \max\{J_A((x \circ y) * x), J_A(x)\} \leq \max\{J_A(y), J_A(x)\}.$$

Therefore $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic \circ -subalgebra of X . \square

We provide a characterization of an MBJ-neutrosophic ideal in an (S)-BCK-algebra.

Theorem 3.24. Let $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ be an MBJ-neutrosophic set in an (S)-BCK-algebra X . Then $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic ideal of X if and only if the following assertions are valid.

$$M_A(x) \geq \min\{M_A(y), M_A(z)\}, \tilde{B}_A(x) \succeq \text{rmin}\{\tilde{B}_A(y), \tilde{B}_A(z)\}, J_A(x) \leq \max\{J_A(y), J_A(z)\} \quad (3.12)$$

for all $x, y, z \in X$ with $x \leq y \circ z$.

Proof. Assume that $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic ideal of X and let $x, y, z \in X$ be such that $x \leq y \circ z$. Using (3.1), (3.2) and Theorem 3.23, we have

$$\begin{aligned} M_A(x) &\geq \min\{M_A(x * (y \circ z)), M_A(y \circ z)\} \\ &= \min\{M_A(0), M_A(y \circ z)\} \\ &= M_A(y \circ z) \geq \min\{M_A(y), M_A(z)\}, \end{aligned}$$

$$\begin{aligned} \tilde{B}_A(x) &\succeq \text{rmin}\{\tilde{B}_A(x * (y \circ z)), \tilde{B}_A(y \circ z)\} \\ &= \text{rmin}\{\tilde{B}_A(0), \tilde{B}_A(y \circ z)\} \\ &= \tilde{B}_A(y \circ z) \succeq \text{rmin}\{\tilde{B}_A(y), \tilde{B}_A(z)\}, \end{aligned}$$

and

$$\begin{aligned} J_A(x) &\leq \max\{J_A(x * (y \circ z)), J_A(y \circ z)\} \\ &= \max\{J_A(0), J_A(y \circ z)\} \\ &= J_A(y \circ z) \leq \max\{J_A(y), J_A(z)\}. \end{aligned}$$

Conversely, let $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ be an MBJ-neutrosophic set in an (S)-BCK-algebra X satisfying the condition (3.12) for all $x, y, z \in X$ with $x \leq y \circ z$. Since $0 \leq x \circ x$ for all $x \in X$, it follows from (3.12) that

$$M_A(0) \geq \min\{M_A(x), M_A(x)\} = M_A(x),$$

$$\tilde{B}_A(0) \succeq \text{rmin}\{\tilde{B}_A(x), \tilde{B}_A(x)\} = \tilde{B}_A(x),$$

and

$$J_A(0) \leq \max\{J_A(x), J_A(x)\} = J_A(x).$$

Note that $x \leq (x * y) \circ y$ for all $x, y \in X$. Hence we have

$$M_A(x) \geq \min\{M_A(x * y), M_A(y)\}, \tilde{B}_A(x) \succeq \min\{\tilde{B}_A(x * y), \tilde{B}_A(y)\} \text{ and } J_A(x) \leq \max\{J_A(x * y), J_A(y)\}.$$

Therefore $\mathcal{A} = (M_A, \tilde{B}_A, J_A)$ is an MBJ-neutrosophic ideal of X . □

References

- [1] Zadeh L.A., Fuzzy sets, *Information and Control*, 1965, 8(3), 338–353.
- [2] Smarandache F., *Neutrosophy, Neutrosophic Probability, Set, and Logic*, ProQuest Information & Learning, Ann Arbor, Michigan, USA, 1998.
- [3] Smarandache F., *A unifying field in logics, Neutrosophy: Neutrosophic probability, set and logic*, Rehoboth: American Research Press, 1999.
- [4] Smarandache F., Neutrosophic set, a generalization of intuitionistic fuzzy sets, *International Journal of Pure and Applied Mathematics*, 2005, 24(5), 287–297.
- [5] Borzooei R.A., Mohseni Takallo M., Smarandache F., Jun Y.B., Positive implicative neutrosophic ideals in BCK-algebras, *Neutrosophic Sets and Systems*, 2018, 23, 126–141.

- [6] Borzooei R.A., Zhan X.H., Smarandache F., Jun Y.B., Commutative generalized neutrosophic ideals in BCK -algebras, *Symmetry*, 2018, 10, 350, doi:10.3390/sym10080350.
- [7] Jun Y.B., Neutrosophic subalgebras of several types in BCK/BCI -algebras, *Ann. Fuzzy Math. Inform.*, 2017, 14(1), 75–86.
- [8] Jun Y.B., Kim S.J., Smarandache F., Interval neutrosophic sets with applications in BCK/BCI -algebra, *Axioms*, 2018, 7, 23.
- [9] Jun Y.B., Smarandache F., Bordbar H., Neutrosophic \mathcal{N} -structures applied to BCK/BCI -algebras, *Information*, 2017, 8, 128.
- [10] Jun Y.B., Smarandache F., Song S.Z., Khan M., Neutrosophic positive implicative \mathcal{N} -ideals in BCK/BCI -algebras, *Axioms*, 2018, 7, 3.
- [11] Khan M., Anis S., Smarandache F., Jun Y.B., Neutrosophic \mathcal{N} -structures and their applications in semigroups, *Ann. Fuzzy Math. Inform.*, 2017, 14(6), 583–598.
- [12] Öztürk M.A., Jun Y.B., Neutrosophic ideals in BCK/BCI -algebras based on neutrosophic points, *J. Inter. Math. Virtual Inst.*, 2018, 8, 1–17.
- [13] Saeid A.B., Jun Y.B., Neutrosophic subalgebras of BCK/BCI -algebras based on neutrosophic points, *Ann. Fuzzy Math. Inform.*, 2017, 14(1), 87–97.
- [14] Song S.Z., Khan M., Smarandache F., Jun Y.B., A novel extension of neutrosophic sets and F_s application in BCK/BI -algebras, *New Trends in Neutrosophic Theory and Applications (Volume II)*, Pons Editions, Brussels, Belgium, EU 2018, 308–326.
- [15] Song S.Z., Smarandache F., Jun Y.B., Neutrosophic commutative \mathcal{N} -ideals in BCK -algebras, *Information*, 2017, 8, 130.
- [16] Mohseni Takallo M., Borzooei R.A., Jun Y.B., MJB-neutrosophic structures and its applications in BCK/BCI -algebras, *Neutrosophic Sets and Systems*, 2018, 23, 72–84.
- [17] Huang Y.S., *BCI-algebra*, Beijing: Science Press, 2006.
- [18] Meng J., Jun Y.B., *BCK-algebras*, Kyung Moon Sa Co., Seoul, 1994.