



Research Article

Muhammad Aamir Ali, Necmettin Alp, Hüseyin Budak, Yu-Ming Chu*, and Zhiyue Zhang

On some new quantum midpoint-type inequalities for twice quantum differentiable convex functions

<https://doi.org/10.1515/math-2021-0015>

received August 28, 2020; accepted January 22, 2021

Abstract: The present paper aims to find some new midpoint-type inequalities for twice quantum differentiable convex functions. The consequences derived in this paper are unification and generalization of the comparable consequences in the literature on midpoint inequalities.

Keywords: Hermite-Hadamard inequality, q -integral, quantum calculus, convex function

MSC 2020: 26D10, 26D15, 26A51

1 Introduction

Quantum calculus, sometimes called calculus without limits, is equivalent to the traditional infinitesimal calculus without the notion of limits. In the field of q -analysis, many studies have recently been carried out. Euler started this field because of the very high demand of mathematics that models quantum computing q -calculus appeared as a connection between physics and mathematics. It has applications in numerous areas of mathematics, such as combinatorics, number theory, basic hypergeometric functions, and orthogonal polynomials, and in fields of other sciences, such as mechanics, theory of relativity, and quantum theory [1–7]. Apparently, Euler was the founder of this branch of mathematics, by using the parameter q in Newton's work on infinite series. Later, the q -calculus was first given by Jackson [8]. In 1908–1909, Jackson defined the general q -integral and q -difference operator [3]. In 1969, Agarwal described the q -fractional derivative for the first time [9]. In 1966–1967, Al-Salam introduced a q -analog of the Riemann-Liouville fractional integral operator and q -fractional integral operator [10]. In 2004, Rajkovic gave a definition of the Riemann-type q -integral which generalized to Jackson q -integral. In 2013, Tariboon introduced ${}_aD_q$ -difference operator [11].

* Corresponding author: Yu-Ming Chu, Department of Mathematics, Huzhou University, Huzhou, 313000, China,
e-mail: chuyuming2005@126.com

Muhammad Aamir Ali: Jiangsu Key Laboratory for NSLSCS, School of Mathematical Sciences, Nanjing Normal University, Nanjing, 210023, China, e-mail: mahr.muhammad.aamir@gmail.com

Zhiyue Zhang: Jiangsu Key Laboratory for NSLSCS, School of Mathematical Sciences, Nanjing Normal University, Nanjing, 210023, China, e-mail: 05289@njnu.edu.cn

Necmettin Alp: Department of Mathematics, Faculty of Science and Arts, Düzce University, Düzce-Turkey, e-mail: placenn@gmail.com

Hüseyin Budak: Department of Mathematics, Faculty of Science and Arts, Düzce University, Düzce-Turkey, e-mail: hsyn.budak@gmail.com

Many integral inequalities well known in classical analysis, such as Hölder inequality, Simpson's inequality, Newton's inequality, Hermite-Hadamard inequality and Ostrowski inequality, Cauchy-Bunyakovsky-Schwarz, Gruss, Gruss-Cebysev, and other integral inequalities, have been proved and applied in q -calculus using classical convexity. Many mathematicians have carried out studies in q -calculus analysis, and the interested reader can check [11–23].

In this paper, motivated through q -calculus, we found bounds for q -midpoint integral inequalities. We first obtain an identity for twice q -differentiable functions. After that, through the derived identity, we attain some new outcomes for q -midpoint inequalities. With all this, we revealed the results we found in the classical analysis by using $q \rightarrow 1^-$.

2 Preliminaries and definitions of q -calculus

Throughout this paper, let $a < b$ and let $0 < q < 1$ be a constant. The definitions and theorems for q -derivative and q -integral of a function f on $[a, b]$ are as follows.

Definition 1. [4,11] For a function $f : [a, b] \rightarrow \mathbb{R}$, the q_a -derivative of f at $x \in [a, b]$ is characterized by the expression

$${}_aD_q f(x) = \frac{f(x) - f(qx + (1-q)a)}{(1-q)(x-a)}, \quad x \neq a. \quad (1)$$

If $x = a$, we define ${}_aD_q f(a) = \lim_{x \rightarrow a} {}_aD_q f(x)$ if it exists and it is finite.

In 2004, Rajkovic et al. [24] gave the following definition of the Riemann-type q -integral which was generalized to Jackson q -integral on $[a, b]$:

$$\int_a^x f(s) {}_a d_q s = (1-q)(x-a) \sum_{n=0}^{\infty} q^n f(q^n x + (1-q^n)a), \quad x \in [a, b]. \quad (2)$$

Definition 2. If $a = 0$ in (2), then $\int_0^x f(s) {}_0 d_q s = \int_0^x f(s) d_q s$, where $\int_0^x f(s) d_q s$ is the familiar q -definite integral on $[0, x]$ defined by the expression (see [4])

$$\int_0^x f(s) {}_0 d_q s = \int_0^x f(s) d_q s = (1-q)x \sum_{n=0}^{\infty} q^n f(q^n x). \quad (3)$$

If $c \in (a, x)$, then the q -definite integral on $[c, x]$ is expressed as

$$\int_c^x f(s) {}_a d_q s = \int_a^x f(s) {}_a d_q s - \int_a^c f(s) {}_a d_q s. \quad (4)$$

Alp et al. [11] proved the q -Hermite-Hadamard inequality; in [25], the authors proved the same inequality by removing the differentiability assumptions as follows:

Theorem 1. (q -Hermite-Hadamard inequality) Let $f : [a, b] \rightarrow \mathbb{R}$ be a convex function on $[a, b]$ and $0 < q < 1$. Then, we have

$$f\left(\frac{qa+b}{1+q}\right) \leq \frac{1}{b-a} \int_a^b f(x) {}_a d_q x \leq \frac{qf(a) + f(b)}{1+q}.$$

On the other hand, in [26], Bermudo et al. gave the following new definitions of quantum integral and derivative. In the same paper, the authors also proved a new variant of the quantum Hermite-Hadamard-type inequality linked to their newly defined quantum integral:

Definition 3. [26] Let $f : [a, b] \rightarrow \mathbb{R}$ be a function. Then, the q^b -definite integral on $[a, b]$ is given by

$$\int_a^b f(x) {}_q^b d_q x = (1-q)(b-a) \sum_{n=0}^{\infty} q^n f(q^n a + (1-q^n)b) = (b-a) \int_0^1 f(sa + (1-s)b) d_q s.$$

Definition 4. [26] Let $f : [a, b] \rightarrow \mathbb{R}$ be a function. Then, the q^b -derivative of f at $x \in [a, b]$ is given by

$${}_q^b D_q f(x) = \frac{f(qx + (1-q)b) - f(x)}{(1-q)(b-x)}, \quad x \neq b.$$

Theorem 2. [26] If $f : [a, b] \rightarrow \mathbb{R}$ is a convex function on $[a, b]$ and $0 < q < 1$, then, q -Hermite-Hadamard inequalities are given as follows:

$$f\left(\frac{a+qb}{1+q}\right) \leq \frac{1}{b-a} \int_a^b f(x) {}_q^b d_q x \leq \frac{f(a) + qf(b)}{1+q}. \quad (5)$$

We frequently used the following notations:

$$[n]_q = \begin{cases} \frac{q^n - 1}{q - 1} = \sum_{i=0}^{n-1} q^i, & n \in N, \\ \frac{q^n - 1}{q - 1}, & n \in C, \end{cases}$$

and

$$(1-s)_q^n = (s, q)_n = \prod_{i=0}^{n-1} (1 - q^i s). \quad (6)$$

Lemma 1. [11] For $\alpha \in \mathbb{R} \setminus \{-1\}$, the following formula holds:

$$\int_a^x (s-a)^\alpha {}_a d_q s = \frac{(x-a)^{\alpha+1}}{[\alpha+1]_q}. \quad (7)$$

Lemma 2. [27] The following equality is valid:

$$\int_{\frac{1}{[2]_q}}^1 (1 - qs)_q^n {}_q d_q s = \frac{\left(1 - \frac{1}{[2]_q}\right)_q^{n+1}}{[n+1]_q}.$$

Lemma 3. [27] The following equality is valid:

$$\int_{\frac{1}{[2]_q}}^1 s(1 - qs)_q^2 {}_q d_q s = \frac{2q + 4q^2 + q^3 - q^4 - q^5}{[2]_q^4 [3]_q [4]_q}. \quad (8)$$

Lemma 4. [27] The following equality is valid:

$$\int_{\frac{1}{[2]_q}}^1 (1 - s)(1 - qs)_q^2 {}_q d_q s = \frac{-q - q^2 + 2q^3 + 3q^4 + 2q^5 - q^6 - q^7}{[2]_q^4 [3]_q [4]_q}.$$

3 An identity for q^b -integrals

In this section, we will prove an equality which will help to obtain our main results.

Lemma 5. Let $f : [a, b] \rightarrow \mathbb{R}$ be a q -differentiable function on (a, b) and $q \in (0, 1)$. If ${}^bD_q^2f$ is continuous and integrable on $[a, b]$, then we attain the identity

$$\frac{(b-a)^2}{[2]_q} \left[\int_0^{\frac{1}{[2]_q}} q^3 s^2 {}^bD_q^2 f(sa + (1-s)b) d_q s + \int_{\frac{1}{[2]_q}}^1 (1-qs)^2 {}^bD_q^2 f(sa + (1-s)b) d_q s \right] = \frac{1}{b-a} \int_a^b f(s) {}^bD_q s - f\left(\frac{a+qb}{[2]_q}\right).$$

Proof. From Definition 4, we have the following equality:

$$\begin{aligned} & {}^bD_q^2 f(sa + (1-s)b) \\ &= {}^bD_q({}^bD_q(f(sa + (1-s)b))) \\ &= {}^bD_q\left(\frac{f(qsa + (1-qs)b) - f(sa + (1-s)b)}{(1-q)(b-a)s}\right) \\ &= \frac{1}{(1-q)(b-a)s} \left[\frac{f(q^2sa + (1-sq^2)b) - f(qsa + (1-qs)b)}{(1-q)q(b-a)s} - \frac{f(qsa + (1-qs)b) - f(sa + (1-s)b)}{(1-q)(b-a)s} \right] \\ &= \frac{f(q^2sa + (1-sq^2)b) - f(qsa + (1-qs)b)}{(1-q)^2q(b-a)^2s^2} - \frac{f(qsa + (1-qs)b) - f(sa + (1-s)b)}{(1-q)^2(b-a)^2s^2} \\ &= \frac{f(q^2sa + (1-sq^2)b) - (1+q)f(qsa + (1-qs)b) + qf(sa + (1-s)b)}{(1-q)^2q(b-a)^2s^2}. \end{aligned} \quad (9)$$

From (9) and the fundamental properties of q -integrals, we have

$$\begin{aligned} & \int_0^{\frac{1}{[2]_q}} q^3 s^2 {}^bD_q^2 f(sa + (1-s)b) d_q s + \int_{\frac{1}{[2]_q}}^1 (1-qs)^2 {}^bD_q^2 f(sa + (1-s)b) d_q s \\ &= \int_0^{\frac{1}{[2]_q}} q^3 s^2 {}^bD_q^2 f(sa + (1-s)b) d_q s + \int_0^1 (1-qs)^2 {}^bD_q^2 f(sa + (1-s)b) d_q s - \int_0^{\frac{1}{[2]_q}} (1-qs)^2 {}^bD_q^2 f(sa + (1-s)b) d_q s \\ &= \int_0^1 (1-qs)^2 {}^bD_q^2 f(sa + (1-s)b) d_q s + \int_0^{\frac{1}{[2]_q}} (q^3 s^2 - (1-qs)^2) {}^bD_q^2 f(sa + (1-s)b) d_q s \\ &= \frac{1}{(1-q)^2(b-a)^2} \int_0^1 \frac{(1-qs)_q^2}{s^2} \left(\frac{1}{q} f((q^2sa + (1-q^2s)b)) - \frac{1+q}{q} f(qsa + (1-qs)b) + f(sa + (1-s)b) \right) d_q s \\ &\quad + \frac{1}{(1-q)^2(b-a)^2} \int_0^{\frac{1}{[2]_q}} (q^3 s^2 - (1-qs)_q^2) \left(\frac{1}{q} f((q^2sa + (1-q^2s)b)) - \frac{1+q}{q} f(qsa + (1-qs)b) + f(sa + (1-s)b) \right) d_q s \\ &= \frac{1}{(1-q)^2(b-a)^2} [I_1 + I_2]. \end{aligned} \quad (10)$$

Now, let us calculate the values of integrals I_1 and I_2 as follows:

$$\begin{aligned} I_1 &= \int_0^1 \frac{(1-qs)_q^2}{s^2} \left(\frac{1}{q} f((q^2sa + (1-q^2s)b)) - \frac{1+q}{q} f(qsa + (1-qs)b) + f(sa + (1-s)b) \right) d_q s \\ &= (1-q) \sum_{n=0}^{\infty} q^n \frac{(1-qq^n)_q^2}{q^{2n}} \left\{ \frac{1}{q} f((q^2q^n a + (1-q^2q^n)b)) - \frac{1+q}{q} f(qq^n a + (1-qq^n)b) + f(q^n a + (1-q^n)b) \right\} \end{aligned}$$

$$\begin{aligned}
&= (1-q) \left\{ \frac{1}{q} \sum_{n=0}^{\infty} \frac{(1-qq^n)_q^2}{q^n} f((q^{n+2}a + (1-q^{n+2})b)) - \frac{1+q}{q} \sum_{n=0}^{\infty} \frac{(1-qq^n)_q^2}{q^n} f(q^{n+1}a + (1-q^{n+1})b) \right. \\
&\quad \left. + \sum_{n=0}^{\infty} \frac{(1-qq^n)_q^2}{q^n} f(q^n a + (1-q^n)b) \right\} \\
&= \frac{(1-q)}{q} \sum_{n=2}^{\infty} \frac{(1-qq^{n-2})_q^2}{q^{n-2}} f((q^n a + (1-q^n)b)) - \frac{(1-q)(1+q)}{q} \sum_{n=1}^{\infty} \frac{(1-qq^{n-1})_q^2}{q^{n-1}} f(q^n a + (1-q^n)b) \\
&\quad + (1-q) \sum_{n=0}^{\infty} \frac{(1-qq^n)_q^2}{q^n} f(q^n a + (1-q^n)b) \\
&= \frac{(1-q)}{q} \sum_{n=0}^{\infty} \frac{(1-qq^{n-2})_q^2}{q^{n-2}} f((q^n a + (1-q^n)b)) - \frac{(1-q)(1+q)}{q} \sum_{n=0}^{\infty} \frac{(1-qq^{n-1})_q^2}{q^{n-1}} f(q^n a + (1-q^n)b) \\
&\quad + (1-q) \sum_{n=0}^{\infty} \frac{(1-qq^n)_q^2}{q^n} f(q^n a + (1-q^n)b) + \frac{(1-q)(1+q)}{q} \frac{(1-qq^{-1})_q^2}{q^{-1}} f(a) \\
&\quad - \frac{(1-q)}{q} \frac{(1-qq^{-2})_q^2}{q^{-2}} f(a) - \frac{(1-q)}{q} \frac{(1-qq^{-1})_q^2}{q^{-1}} f((qa + (1-q)b)) \\
&= (1-q) \sum_{n=0}^{\infty} \left\{ \frac{(1-qq^{n-2})_q^2}{q^{n-1}} - \frac{(1+q)}{q} \frac{(1-qq^{n-1})_q^2}{q^{n-1}} + \frac{(1-qq^n)_q^2}{q^n} \right\} f(q^n a + (1-q^n)b) \\
&\quad + [(1-q^2)(1-qq^{-1})_q^2 - q(1-q)(1-qq^{-2})_q^2] f(a) - (1-q)(1-qq^{-1})_q^2 f(qa + (1-q)b) \\
&= (1-q) \sum_{n=0}^{\infty} \left\{ \frac{q(1-qq^{n-2})_q^2 - (1+q)(1-qq^{n-1})_q^2 + (1-qq^n)_q^2}{q^n} \right\} f(q^n a + (1-q^n)b) \\
&\quad + [(1-q^2)(1-qq^{-1})_q^2 - q(1-q)(1-qq^{-2})_q^2] f(a) - (1-q)(1-qq^{-1})_q^2 f(qa + (1-q)b) \\
&= (1+q)(1-q)^2(1-q) \sum_{n=0}^{\infty} q^n f(q^n a + (1-q^n)b) \\
&= \frac{(1+q)(1-q)^2}{b-a} (1-q)(b-a) \sum_{n=0}^{\infty} q^n f(q^n a + (1-q^n)b) \\
&= \frac{(1+q)(1-q)^2}{b-a} \int_a^b f(s) {}_b d_q s.
\end{aligned}$$

By similar operations, we can establish

$$\begin{aligned}
I_2 &= \int_0^{\frac{1}{[2]_q}} \frac{(q^3 s^2 - (1-qs)_q^2)}{s^2} \left(\frac{1}{q} f(q^2 sa + (1-q^2 s)b) - \frac{1+q}{q} f(qsa + (1-qs)b) + f(sa + (1-s)b) \right) d_q s \\
&= \frac{(1-q)}{[2]_q} \sum_{n=0}^{\infty} q^n \frac{\left(q^3 \left(\frac{q^n}{[2]_q} \right)^2 - \left(1 - q \frac{q^n}{[2]_q} \right)_q^2 \right)}{\left(\frac{q^n}{[2]_q} \right)^2} \\
&\quad \times \left(\frac{1}{q} f \left(\frac{q^{n+2}}{[2]_q} a + \left(1 - \frac{q^{n+2}}{[2]_q} \right) b \right) - \frac{1+q}{q} f \left(\frac{q^{n+1}}{[2]_q} a + \left(1 - \frac{q^{n+1}}{[2]_q} \right) b \right) + f \left(\frac{q^n}{[2]_q} a + \left(1 - \frac{q^n}{[2]_q} \right) b \right) \right) \\
&= \frac{(1-q)}{[2]_q} \sum_{n=0}^{\infty} \frac{\left(q^3 q^{2n} - [2]_q^2 \left(1 - q \frac{q^n}{[2]_q} \right)_q^2 \right)}{q^n} \\
&\quad \times \left(\frac{1}{q} f \left(\frac{q^{n+2}}{[2]_q} a + \left(1 - \frac{q^{n+2}}{[2]_q} \right) b \right) - \frac{1+q}{q} f \left(\frac{q^{n+1}}{[2]_q} a + \left(1 - \frac{q^{n+1}}{[2]_q} \right) b \right) + f \left(\frac{q^n}{[2]_q} a + \left(1 - \frac{q^n}{[2]_q} \right) b \right) \right)
\end{aligned}$$

$$\begin{aligned}
&= \frac{(1-q)}{[2]_q} \sum_{n=0}^{\infty} \frac{\left(q^3 q^{2n} - [2]_q^2 \left(1 - q \frac{q^n}{[2]_q} \right)_q^2 \right)}{q^{n+1}} f\left(\frac{q^{n+2}}{[2]_q} a + \left(1 - \frac{q^{n+2}}{[2]_q} \right) b \right) \\
&\quad - \frac{(1-q)(1+q)}{[2]_q} \sum_{n=0}^{\infty} \frac{\left(q^3 q^{2n} - [2]_q^2 \left(1 - q \frac{q^n}{[2]_q} \right)_q^2 \right)}{q^{n+1}} f\left(\frac{q^{n+1}}{[2]_q} a + \left(1 - \frac{q^{n+1}}{[2]_q} \right) b \right) \\
&\quad + \frac{(1-q)}{[2]_q} \sum_{n=0}^{\infty} \frac{\left(q^3 q^{2n} - [2]_q^2 \left(1 - q \frac{q^n}{[2]_q} \right)_q^2 \right)}{q^n} f\left(\frac{q^n}{[2]_q} a + \left(1 - \frac{q^n}{[2]_q} \right) b \right) \\
&= \frac{(1-q)}{[2]_q} \sum_{n=2}^{\infty} \frac{\left(q^3 q^{2n-4} - [2]_q^2 \left(1 - q \frac{q^{n-2}}{[2]_q} \right)_q^2 \right)}{q^{n-1}} f\left(\frac{q^n}{[2]_q} a + \left(1 - \frac{q^n}{[2]_q} \right) b \right) \\
&\quad - \frac{(1-q)(1+q)}{[2]_q} \sum_{n=1}^{\infty} \frac{\left(q^3 q^{2n-2} - [2]_q^2 \left(1 - q \frac{q^{n-1}}{[2]_q} \right)_q^2 \right)}{q^n} f\left(\frac{q^n}{[2]_q} a + \left(1 - \frac{q^n}{[2]_q} \right) b \right) \\
&\quad + \frac{(1-q)}{[2]_q} \sum_{n=0}^{\infty} \frac{\left(q^3 q^{2n} - [2]_q^2 \left(1 - q \frac{q^n}{[2]_q} \right)_q^2 \right)}{q^n} f\left(\frac{q^n}{[2]_q} a + \left(1 - \frac{q^n}{[2]_q} \right) b \right) \\
&= \frac{(1-q)}{[2]_q} \sum_{n=0}^{\infty} \frac{\left(q^3 q^{2n-4} - [2]_q^2 \left(1 - q \frac{q^{n-2}}{[2]_q} \right)_q^2 \right)}{q^{n-1}} f\left(\frac{q^n}{[2]_q} a + \left(1 - \frac{q^n}{[2]_q} \right) b \right) \\
&\quad - \frac{(1-q)(1+q)}{[2]_q} \sum_{n=0}^{\infty} \frac{\left(q^3 q^{2n-2} - [2]_q^2 \left(1 - q \frac{q^{n-1}}{[2]_q} \right)_q^2 \right)}{q^n} f\left(\frac{q^n}{[2]_q} a + \left(1 - \frac{q^n}{[2]_q} \right) b \right) \\
&\quad + \frac{(1-q)}{[2]_q} \sum_{n=0}^{\infty} \frac{\left(q^3 q^{2n} - [2]_q^2 \left(1 - q \frac{q^n}{[2]_q} \right)_q^2 \right)}{q^n} f\left(\frac{q^n}{[2]_q} a + \left(1 - \frac{q^n}{[2]_q} \right) b \right) \\
&\quad - \frac{(1-q)}{[2]_q} \frac{\left(q^3 q^{-4} - [2]_q^2 \left(1 - q \frac{q^{-2}}{[2]_q} \right)_q^2 \right)}{q^{-1}} f\left(\frac{a + qb}{[2]_q} \right) - \frac{(1-q)}{[2]_q} \left(q^3 q^{-2} - [2]_q^2 \left(1 - q \frac{q^{-1}}{[2]_q} \right)_q^2 \right) f\left(\frac{qa + b}{[2]_q} \right) \\
&\quad + \frac{(1-q)(1+q)}{[2]_q} \left(q^3 q^{-2} - [2]_q^2 \left(1 - q \frac{q^{-1}}{[2]_q} \right)_q^2 \right) f\left(\frac{a + qb}{[2]_q} \right) \\
&= \frac{(1-q)}{[2]_q} \sum_{n=0}^{\infty} \left\{ \frac{\left(q^3 q^{2n-4} - [2]_q^2 \left(1 - q \frac{q^{n-2}}{[2]_q} \right)_q^2 \right)}{q^{n-1}} - \frac{(1+q) \left(q^3 q^{2n-2} - [2]_q^2 \left(1 - q \frac{q^{n-1}}{[2]_q} \right)_q^2 \right)}{q^n} \right\} f\left(\frac{q^n}{[2]_q} a + \left(1 - \frac{q^n}{[2]_q} \right) b \right) \\
&\quad + \frac{\left(q^3 q^{2n} - [2]_q^2 \left(1 - q \frac{q^n}{[2]_q} \right)_q^2 \right)}{q^n} f\left(\frac{q^n}{[2]_q} a + \left(1 - \frac{q^n}{[2]_q} \right) b \right)
\end{aligned}$$

$$\begin{aligned}
& + \frac{(1-q)}{[2]_q} \left\{ (1+q) \left(q^3 q^{-2} - [2]_q^2 \left(1 - q \frac{q^{-1}}{[2]_q} \right)_q^2 \right) - q \left(q^3 q^{-4} - [2]_q^2 \left(1 - q \frac{q^{-2}}{[2]_q} \right)_q^2 \right) \right\} f \left(\frac{a+qb}{[2]_q} \right) \\
& - \frac{(1-q)}{[2]_q} \left(q^3 q^{-2} - [2]_q^2 \left(1 - q \frac{q^{-1}}{[2]_q} \right)_q^2 \right) f \left(\frac{qa+b}{[2]_q} \right) \\
& = \frac{(1-q)}{[2]_q} \sum_{n=0}^{\infty} \frac{1}{q^n} \left\{ \begin{aligned} & q \left(q^3 q^{2n-4} - [2]_q^2 \left(1 - q \frac{q^{n-2}}{[2]_q} \right)_q^2 \right) - (1+q) \left(q^3 q^{2n-2} - [2]_q^2 \left(1 - q \frac{q^{n-1}}{[2]_q} \right)_q^2 \right) \\ & + \left(q^3 q^{2n} - [2]_q^2 \left(1 - q \frac{q^n}{[2]_q} \right)_q^2 \right) \end{aligned} \right\} f \left(\frac{q^n}{[2]_q} a + \left(1 - \frac{q^n}{[2]_q} \right) b \right) \\
& + \frac{(1-q)}{[2]_q} \left\{ (1+q) \left(q^3 q^{-2} - [2]_q^2 \left(1 - q \frac{q^{-1}}{[2]_q} \right)_q^2 \right) - q \left(q^3 q^{-4} - [2]_q^2 \left(1 - q \frac{q^{-2}}{[2]_q} \right)_q^2 \right) \right\} f \left(\frac{a+qb}{[2]_q} \right) \\
& - \frac{(1-q)}{[2]_q} \left(q^3 q^{-2} - [2]_q^2 \left(1 - q \frac{q^{-1}}{[2]_q} \right)_q^2 \right) f \left(\frac{qa+b}{[2]_q} \right) \\
& = \frac{(1-q)}{[2]_q} \left\{ (1+q) \left(q - [2]_q^2 \left(1 - \frac{1}{[2]_q} \right)_q^2 \right) - q \left(\frac{1}{q} - [2]_q^2 \left(1 - \frac{q^{-1}}{[2]_q} \right)_q^2 \right) \right\} f \left(\frac{a+qb}{[2]_q} \right) \\
& = \frac{(1-q)}{[2]_q} \{[-1 + q^2 + q^3 - q]\} f \left(\frac{a+qb}{[2]_q} \right) \\
& = \frac{(1-q)}{1+q} (1+q) \{q^2 - 1\} f \left(\frac{a+qb}{[2]_q} \right) \\
& = -(1-q)^2 (1+q) f \left(\frac{a+qb}{[2]_q} \right).
\end{aligned}$$

By substituting the values of integrals I_1 and I_2 in (10), we attain

$$\begin{aligned}
& \int_0^{\frac{1}{[2]_q}} q^3 s^2 b D_q^2 f(sa + (1-s)b) d_q s + \int_{\frac{1}{[2]_q}}^1 (1-qs)_q^2 b D_q^2 f(sa + (1-s)b) d_q s \\
& = \frac{1}{(1-q)^2(b-a)^2} \left(\frac{(1+q)(1-q)^2}{b-a} \int_a^b f(s) b d_q s - (1-q)^2 (1+q) f \left(\frac{a+qb}{[2]_q} \right) \right) \\
& = \frac{[2]_q}{(b-a)^2} \left(\frac{1}{b-a} \int_a^b f(s) b d_q s - f \left(\frac{a+qb}{[2]_q} \right) \right),
\end{aligned}$$

which completes the proof. \square

Remark 1. If we take the limit $q \rightarrow 1^-$ in Lemma 5, then we have the equality

$$\frac{(b-a)^2}{2} \left[\int_0^{\frac{1}{2}} s^2 f''(sa + (1-s)b) ds + \int_{\frac{1}{2}}^1 (1-s)^2 f''(sa + (1-s)b) ds \right] = \frac{1}{b-a} \int_a^b f(s) ds - f \left(\frac{a+b}{2} \right)$$

which was given by Sarikaya et al. in [24].

4 New midpoint-type inequalities for quantum integrals

In this section, we obtain some new midpoint-type inequalities for newly defined quantum integrals utilizing twice q -differentiable convex functions.

Theorem 3. Suppose that the assumptions of Lemma 5 hold. If $|^bD_q^2 f|$ is convex on $[a, b]$, then we have the inequality

$$\begin{aligned} & \left| \frac{1}{b-a} \int_a^b f(s) {}^b d_q s - f\left(\frac{a+qb}{[2]_q}\right) \right| \\ & \leq \frac{(b-a)^2}{[2]_q^5 [3]_q [4]_q} [(2q + 4q^2 + 2q^3) |^b D_q^2 f(a)| + (-q - q^2 + 2q^3 + 4q^4 + 3q^5 + q^6) |^b D_q^2 f(b)|]. \end{aligned} \quad (11)$$

Proof. By taking the modulus in Lemma 5, we have

$$\begin{aligned} & \left| \frac{1}{b-a} \int_a^b f(s) {}^b d_q s - f\left(\frac{a+qb}{[2]_q}\right) \right| \\ & \leq \frac{(b-a)^2}{[2]_q} \left[\int_0^{\frac{1}{[2]_q}} q^3 s^2 |^b D_q^2 f(sa + (1-s)b)| d_q s + \int_{\frac{1}{[2]_q}}^1 (1-qs)_q^2 |^b D_q^2 f(sa + (1-s)b)| d_q s \right]. \end{aligned}$$

By using the convexity of $|^b D_q^2 f|$, we can write

$$\begin{aligned} & \left| \frac{1}{b-a} \int_a^b f(s) {}^b d_q s - f\left(\frac{a+qb}{[2]_q}\right) \right| \leq \frac{q^3(b-a)^2}{[2]_q} \int_0^{\frac{1}{[2]_q}} (s^3 |^b D_q^2 f(a)| + (s^2 - s^3) |^b D_q^2 f(b)|) d_q s \\ & \quad + \frac{(b-a)^2}{[2]_q} \int_{\frac{1}{[2]_q}}^1 (s(1-qs)_q^2 |^b D_q^2 f(a)| + (1-s)(1-qs)_q^2 |^b D_q^2 f(b)|) d_q s. \end{aligned}$$

We have the fact that

$$\begin{aligned} \int_0^{\frac{1}{[2]_q}} s^3 d_q s &= \frac{1}{[2]_q^4 [4]_q}, \\ \int_0^{\frac{1}{[2]_q}} (s^2 - s^3) d_q s &= \left(\frac{1}{[2]_q^3 [3]_q} - \frac{1}{[2]_q^4 [4]_q} \right), \\ \int_{\frac{1}{[2]_q}}^1 s(1-qs)_q^2 d_q s &= \frac{2q + 4q^2 + q^3 - q^4 - q^5}{[2]_q^4 [3]_q [4]_q}, \end{aligned}$$

and

$$\int_{\frac{1}{[2]_q}}^1 (1-s)(1-qs)_q^2 |^b D_q^2 f(b)| d_q s = \frac{-q - q^2 + 2q^3 + 3q^4 + 2q^5 - q^6 - q^7}{[2]_q^4 [3]_q [4]_q}.$$

By these equalities, we can write

$$\begin{aligned} & \left| \frac{1}{b-a} \int_a^b f(s) {}^b d_q s - f\left(\frac{a+qb}{[2]_q}\right) \right| \\ & \leq \frac{q^3(b-a)^2}{[2]_q} \left(\frac{(1+q+q^2)|{}^b D_q^2 f(a)|}{[2]_q^4 [3]_q [4]_q} + \frac{q+q^2+2q^3+q^4}{[2]_q^4 [3]_q [4]_q} |{}^b D_q^2 f(b)| \right) \\ & + \frac{(b-a)^2}{[2]_q} \left[\frac{2q+4q^2+q^3-q^4-q^5}{[2]_q^4 [3]_q [4]_q} |{}^b D_q^2 f(a)| + \frac{-q-q^2+2q^3+3q^4+2q^5-q^6-q^7}{[2]_q^4 [3]_q [4]_q} |{}^b D_q^2 f(b)| \right] \\ & = \frac{(b-a)^2}{[2]_q^5 [3]_q [4]_q} [(2q+4q^2+2q^3)|{}^b D_q^2 f(a)| + (-q-q^2+2q^3+4q^4+3q^5+q^6)|{}^b D_q^2 f(b)|]. \end{aligned}$$

This completes the proof. \square

Remark 2. If we take the limit $q \rightarrow 1^-$ in Theorem 3, then we obtain the inequality

$$\left| \frac{1}{b-a} \int_a^b f(s) ds - f\left(\frac{a+b}{2}\right) \right| \leq \frac{(b-a)^2}{48} (|f''(a)| + |f''(b)|)$$

which was proved by Sarikaya et al. in [28, Theorem 3].

Theorem 4. Suppose that the assumptions of Lemma 5 hold. If $|{}^b D_q^2 f|^{r_1}$, $r_1 > 1$, is convex function on $[a, b]$, then we have the inequality

$$\begin{aligned} & \left| \frac{1}{b-a} \int_a^b f(s) {}^b d_q s - f\left(\frac{a+qb}{[2]_q}\right) \right| \\ & \leq \frac{q^3(b-a)^2}{[2]_q^{\frac{3+r_1}{r_1} + \frac{2r_2+1}{r_2}} [2r_2+1]_q^{\frac{1}{r_2}}} (|{}^b D_q^2 f(a)|^{r_1} + (2q+q^2)|{}^b D_q^2 f(b)|^{r_1})^{\frac{1}{r_1}} \\ & + \frac{q^{\frac{1}{r_1}}(b-a)^2 \left(1 - \frac{1}{[2]_q}\right)^{\frac{2r_2+1}{r_2}}}{[2]_q^{\frac{3+r_1}{r_1}} [2r_2+1]_q^{\frac{1}{r_2}}} ((2+q)|{}^b D_q^2 f(a)|^{r_1} + (q+q^2-1)|{}^b D_q^2 f(b)|^{r_1})^{\frac{1}{r_1}}, \end{aligned}$$

where $\frac{1}{r_1} + \frac{1}{r_2} = 1$.

Proof. From Lemma 5 and using well-known q -Hölder's integral inequality, we get

$$\begin{aligned} & \left| \frac{1}{b-a} \int_a^b f(s) {}^b d_q s - f\left(\frac{a+qb}{[2]_q}\right) \right| \leq \frac{(b-a)^2}{[2]_q} q^3 \left(\int_0^{\frac{1}{[2]_q}} s^{2r_2} {}^b d_q s \right)^{\frac{1}{r_2}} \left(\int_0^{\frac{1}{[2]_q}} |{}^b D_q^2 f(sa + (1-s)b)|^{r_1} {}^b d_q s \right)^{\frac{1}{r_1}} \\ & + \frac{(b-a)^2}{[2]_q} \left(\int_{\frac{1}{[2]_q}}^1 (1-qs)^{2r_2} {}^b d_q s \right)^{\frac{1}{r_2}} \left(\int_{\frac{1}{[2]_q}}^1 |{}^b D_q^2 f(sa + (1-s)b)|^{r_1} {}^b d_q s \right)^{\frac{1}{r_1}}. \end{aligned}$$

Since $|{}^b D_q^2 f|^{r_1}$ is convex on $[a, b]$, we have

$$|{}^b D_q^2 f(sa + (1-s)b)|^{r_1} \leq s |{}^b D_q^2 f(a)|^{r_1} + (1-s) |{}^b D_q^2 f(b)|^{r_1}.$$

By Lemma 2, we get

$$\begin{aligned}
& \left| \frac{1}{b-a} \int_a^b f(s) {}^b d_q s - f\left(\frac{a+qb}{[2]_q}\right) \right| \\
& \leq \frac{(b-a)^2}{[2]_q} q^3 \left(\frac{1}{[2]_q^{\frac{2r_2+1}{r_2}} [2r_2+1]_q^{\frac{1}{r_2}}} \right)^{\frac{1}{r_2}} \left(\int_0^{\frac{1}{[2]_q}} (s |{}^b D_q^2 f(a)|^{r_1} + (1-s) |{}^b D_q^2 f(b)|^{r_1}) d_q s \right)^{\frac{1}{r_1}} \\
& \quad + \frac{(b-a)^2}{[2]_q} \left(\frac{(1-\frac{1}{[2]_q})_q^{2r_2+1}}{[2r_2+1]_q} \right)^{\frac{1}{r_2}} \left(\int_{\frac{1}{[2]_q}}^1 (s |{}^b D_q^2 f(a)|^{r_1} + (1-s) |{}^b D_q^2 f(b)|^{r_1}) d_q s \right)^{\frac{1}{r_1}} \\
& = \left(\frac{q^3(b-a)^2}{[2]_q [2]_q^{\frac{2r_2+1}{r_2}} [2r_2+1]_q^{\frac{1}{r_2}}} \right) \left(\frac{|{}^b D_q^2 f(a)|^{r_1}}{[2]_q^3} + \frac{(2q+q^2) |{}^b D_q^2 f(b)|^{r_1}}{[2]_q^3} \right)^{\frac{1}{r_1}} \\
& \quad + \frac{(b-a)^2}{[2]_q} \frac{\left(1-\frac{1}{[2]_q}\right)_q^{\frac{2r_2+1}{r_2}}}{[2r_2+1]_q^{\frac{1}{r_2}}} \left(\frac{(2q+q^2) |{}^b D_q^2 f(a)|^{r_1}}{[2]_q^3} + \frac{(q^2+q^3-q) |{}^b D_q^2 f(b)|^{r_1}}{[2]_q^3} \right)^{\frac{1}{r_1}} \\
& = \left(\frac{q^3(b-a)^2}{[2]_q^{\frac{3+r_1}{r_1}} [2]_q^{\frac{2r_2+1}{r_2}} [2r_2+1]_q^{\frac{1}{r_2}}} \right) (|{}^b D_q^2 f(a)|^{r_1} + (2q+q^2) |{}^b D_q^2 f(b)|^{r_1})^{\frac{1}{r_1}} \\
& \quad + \frac{(b-a)^2 \left(1-\frac{1}{[2]_q}\right)_q^{\frac{2r_2+1}{r_2}}}{[2]_q^{\frac{3+r_1}{r_1}} [2r_2+1]_q^{\frac{1}{r_2}}} ((2q+q^2) |{}^b D_q^2 f(a)|^{r_1} + (q^2+q^3-q) |{}^b D_q^2 f(b)|^{r_1})^{\frac{1}{r_1}} \\
& = \frac{q^3(b-a)^2}{[2]_q^{\frac{3+r_1}{r_1}+\frac{2r_2+1}{r_2}} [2r_2+1]_q^{\frac{1}{r_2}}} (|{}^b D_q^2 f(a)|^{r_1} + (2q+q^2) |{}^b D_q^2 f(b)|^{r_1})^{\frac{1}{r_1}} \\
& \quad + \frac{q^{\frac{1}{r_1}}(b-a)^2 \left(1-\frac{1}{[2]_q}\right)_q^{\frac{2r_2+1}{r_2}}}{[2]_q^{\frac{3+r_1}{r_1}} [2r_2+1]_q^{\frac{1}{r_2}}} ((2q+q^2) |{}^b D_q^2 f(a)|^{r_1} + (q+q^2-1) |{}^b D_q^2 f(b)|^{r_1})^{\frac{1}{r_1}}.
\end{aligned}$$

Thus, the proof is completed. \square

Remark 3. If we take the limit $q \rightarrow 1^-$ in Theorem 4, then we obtain the inequality

$$\left| \frac{1}{b-a} \int_a^b f(s) ds - f\left(\frac{a+b}{2}\right) \right| \leq \frac{(b-a)^2}{2^{4+\frac{2}{r_1}} (2r_2+1)^{\frac{1}{r_2}}} \left\{ (3|f''(a)|^{r_1} + |f''(b)|^{r_1})^{\frac{1}{r_1}} + (|f''(a)|^{r_1} + 3|f''(b)|^{r_1})^{\frac{1}{r_1}} \right\}$$

which was proved by Noor and Awan in [29, Theorem 3 (for $\alpha = s = 1$)].

Theorem 5. Suppose that the assumptions of Lemma 5 hold. If $|{}^b D_q^2 f|^{r_1}$, $r_1 \geq 1$, is convex on $[a, b]$, then we have the inequality

$$\begin{aligned}
& \left| \frac{1}{b-a} \int_a^b f(s) {}^b d_q s - f\left(\frac{a+qb}{[2]_q}\right) \right| \\
& \leq \frac{q^3(b-a)^2}{[2]_q^{4-\frac{3}{r_1}} [3]_q^{1-\frac{1}{r_1}}} \left(\frac{|{}^b D_q^2 f(a)|^{r_1}}{[2]_q^4 [4]_q} + \frac{(q+q^2+2q^3+q^4) |{}^b D_q^2 f(b)|^{r_1}}{[2]_q^4 [3]_q [4]_q} \right)^{\frac{1}{r_1}} + \frac{(q+q^2-q^3)^{\frac{1}{r_1}} (b-a)^2}{[2]_q^{4-\frac{3}{r_1}} [3]_q^{1-\frac{1}{r_1}}} \\
& \quad \times \left(\frac{(2q+4q^2+q^3-q^4-q^5) |{}^b D_q^2 f(a)|^{r_1}}{[2]_q^4 [3]_q [4]_q} + \frac{(-q-q^2+2q^3+3q^4+2q^5-q^6-q^7) |{}^b D_q^2 f(b)|^{r_1}}{[2]_q^4 [3]_q [4]_q} \right)^{\frac{1}{r_1}}.
\end{aligned}$$

Proof. Using Lemma 5 and power-mean integral inequality, we get

$$\begin{aligned} \left| \frac{1}{b-a} \int_a^b f(s) {}^b d_q s - f\left(\frac{a+qb}{[2]_q}\right) \right| &\leq \frac{(b-a)^2}{[2]_q} \left[q^3 \left(\int_0^{\frac{1}{[2]_q}} s^2 {}^b d_q s \right)^{1-\frac{1}{r_1}} \left(\int_0^{\frac{1}{[2]_q}} s^2 |{}^b D_q^2 f(sa + (1-s)b)|^{r_1} d_q s \right)^{\frac{1}{r_1}} \right] \\ &+ \frac{(b-a)^2}{[2]_q} \left(\int_{\frac{1}{[2]_q}}^1 (1-qs)_q^2 {}^b d_q s \right)^{1-\frac{1}{r_1}} \left(\int_{\frac{1}{[2]_q}}^1 (1-qs)_q^2 |{}^b D_q^2 f(sa + (1-s)b)|^{r_1} d_q s \right)^{\frac{1}{r_1}}. \end{aligned}$$

By using the convexity of $|{}^b D_q^2 f|^{r_1}$ and Lemmas 2–4, we get

$$\begin{aligned} &\left| \frac{1}{b-a} \int_a^b f(s) {}^b d_q s - f\left(\frac{a+qb}{[2]_q}\right) \right| \\ &\leq \frac{(b-a)^2}{[2]_q} \frac{q^3}{[2]_q^{3-\frac{3}{r_1}} [3]_q^{1-\frac{1}{r_1}}} \left(|{}^b D_q^2 f(a)|^{r_1} \int_0^{\frac{1}{[2]_q}} s^3 {}^b d_q s + |{}^b D_q^2 f(b)|^{r_1} \int_0^{\frac{1}{[2]_q}} (s^2 - s^3) {}^b d_q s \right)^{\frac{1}{r_1}} \\ &+ \frac{(b-a)^2}{[2]_q} \frac{\left[\left(1 - \frac{1}{[2]_q}\right)_q^3 \right]^{1-\frac{1}{r_1}}}{[3]_q^{1-\frac{1}{r_1}}} \left(|{}^b D_q^2 f(a)|^{r_1} \int_{\frac{1}{[2]_q}}^1 s(1-qs)_q^2 {}^b d_q s + |{}^b D_q^2 f(b)|^{r_1} \int_{\frac{1}{[2]_q}}^1 (1-s)(1-qs)_q^2 {}^b d_q s \right)^{\frac{1}{r_1}} \\ &= \frac{q^3(b-a)^2}{[2]_q^{4-\frac{3}{r_1}} [3]_q^{1-\frac{1}{r_1}}} \left(\frac{|{}^b D_q^2 f(a)|^{r_1}}{[2]_q^4 [4]_q} + \frac{(q+q^2+2q^3+q^4)|{}^b D_q^2 f(b)|^{r_1}}{[2]_q^4 [3]_q [4]_q} \right)^{\frac{1}{r_1}} + \frac{(q+q^2-q^3)^{\frac{1}{r_1}}(b-a)^2}{[2]_q^{4-\frac{3}{r_1}} [3]_q^{1-\frac{1}{r_1}}} \\ &\times \left(\frac{(2q+4q^2+q^3-q^4-q^5)|{}^b D_q^2 f(a)|^{r_1}}{[2]_q^4 [3]_q [4]_q} + \frac{(-q-q^2+2q^3+3q^4+2q^5-q^6-q^7)|{}^b D_q^2 f(b)|^{r_1}}{[2]_q^4 [3]_q [4]_q} \right)^{\frac{1}{r_1}} \end{aligned}$$

which completes the proof. \square

Remark 4. If we take the limit $q \rightarrow 1^-$ in Theorem 5, then we obtain the inequality

$$\left| \frac{1}{b-a} \int_a^b f(s) ds - f\left(\frac{a+b}{2}\right) \right| \leq \frac{(b-a)^2}{3(2^{4+\frac{3}{r_1}})} \left\{ (5|f''(a)|^{r_1} + 3|f''(b)|^{r_1})^{\frac{1}{r_1}} + (3|f''(a)|^{r_1} + 5|f''(b)|^{r_1})^{\frac{1}{r_1}} \right\}$$

which was proved by Noor and Awan in [29, Theorem 4 (for $\alpha = s = 1$)].

5 Concluding remarks

In this investigation, some new estimates of midpoint-type inequalities for quantum twice differentiable convex functions are attained. It is also proved that the outcomes of the present paper are the strong generalization of the existing comparable consequences in the literature. Upcoming researchers can find the Simpson-like inequalities, Newton-like inequalities, Ostrowski-like inequalities, and the same inequalities by using different kinds of convexities in their future work.

Acknowledgment: The authors would like to express their sincere thanks to the editor and the anonymous reviewers for their helpful comments and suggestions.

Funding information: The work was supported by the Natural Science Foundation of China (Grant Nos. 61673169, 11301127, 11701176, 11626101, 11601485, 11971241).

Authors contribution: All authors contributed equally to the writing of this paper. All authors read and approved the final manuscript.

Conflict of interest: Authors state no conflict of interest.

Data availability statement: Data sharing not applicable to this paper as no data sets were generated or analyzed during the current study.

References

- [1] T. Ernst, *The History of q-Calculus and New Method*, Department of Mathematics, Uppsala University, Sweden, 2000.
- [2] H. Gauchman, *Integral inequalities in q-calculus*, Comput. Math. Appl. **47** (2004), no. 2–3, 281–300.
- [3] F. H. Jackson, *On a q-definite integrals*, Quart. J. Pure Appl. Math. **41** (1910), 193–203.
- [4] V. Kac and P. Cheung, *Quantum Calculus*, Universitext, Springer-Verlag, New York, 2002.
- [5] H. Budak, *Some trapezoid and midpoint-type inequalities for newly defined quantum integrals*, Proyecciones **40** (2021), no. 1, 199–215.
- [6] T. Ernst, *A Comprehensive Treatment of q-Calculus*, Springer, Basel Heidelberg, New York Dordrecht London, 2012.
- [7] T. Ernst, *A method for q-calculus*, J. Nonlinear Math. Phys. **10** (2003), no. 4, 487–525, DOI: <https://doi.org/10.2991/jnmp.2003.10.4.5>.
- [8] F. H. Jackson, *q-difference equations*, Amer. J. Math. **32** (1910), no. 4, 305–314.
- [9] R. Agarwal, *A propos d'une note de M. Pierre Humbert*, Comptes rendus de l'Academie des Sciences **236** (1953), no. 21, 2031–2032.
- [10] W. Al-Salam, *Some fractional q-integrals and q-derivatives*, Proc. Edinb. Math. Soc. (2) **15** (1966), no. 2, 135–140.
- [11] N. Alp, M. Z. Sarikaya, M. Kunt, and I. Iscan, *q-Hermite-Hadamard inequalities and quantum estimates for midpoint-type inequalities via convex and quasi-convex functions*, J. King Saud Univ. Sci. **30** (2018), no. 2, 193–203, DOI: <https://doi.org/10.1016/j.jksus.2016.09.007>.
- [12] M. A. Ali, H. Budak, Z. Zhang, and H. Yıldırım, *Some new Simpsonas type inequalities for co-ordinated convex functions in quantum calculus*, Math. Methods Appl. Sci. **44** (2020), no. 6, 4515–4540, DOI: <https://doi.org/10.1002/mma.7048>.
- [13] M. A. Ali, H. Budak, M. Abbas, and Y.-M. Chu, *Quantum Hermite-Hadamard-type inequalities for functions with convex absolute values of second q^b -derivatives*, Adv. Differ. Equ. **2021** (2021), 7, DOI: <https://doi.org/10.1186/s13662-020-03163-1>.
- [14] M. A. Ali, M. Abbas, H. Budak, P. Agarwal, G. Murtaza, and Y.-M. Chu, *New quantum boundaries for quantum Simpson's and quantum Newton's type inequalities for preinvex functions*, Adv. Differ. Equ. **2021** (2021), 64, DOI: <https://doi.org/10.1186/s13662-021-03226-x>.
- [15] M. A. Ali, Y.-M. Chu, H. Budak, A. Akkurt, H. Yıldırım, and M. A. Zahid, *Quantum variant of Montgomery identity and Ostrowski-type inequalities for the mappings of two variables*, Adv. Differ. Equ. **2021** (2021), 25, DOI: <https://doi.org/10.1186/s13662-020-03195-7>.
- [16] H. Budak, M. A. Ali, and M. Tarhanaci, *Some new quantum Hermite-Hadamard-like inequalities for coordinated convex functions*, J. Optim. Theory Appl. **186** (2020), 899–910, DOI: <https://doi.org/10.1007/s10957-020-01726-6>.
- [17] H. Budak, S. Erden, and M. A. Ali, *Simpson and Newton type inequalities for convex functions via newly defined quantum integrals*, Math. Methods Appl. Sci. **44** (2020), no. 1, 378–390, DOI: <https://doi.org/10.1002/mma.6742>.
- [18] H. Budak, M. A. Ali, and T. Tunç, *Quantum Ostrowski-type integral inequalities for functions of two variables*, Math. Methods Appl. Sci. **44** (2021), no. 7, 5857–5872, DOI: <https://doi.org/10.1002/mma.7153>.
- [19] S. Erden and M. Z. Sarikaya, *New Hermite Hadamard type inequalities for twice differentiable convex mappings via Green function and applications*, Moroccan J. Pure and Appl. Anal. **2** (2016), no. 2, 107–117, DOI: <https://doi.org/10.7603/s40956-016-0009-x>.
- [20] M. A. Khan, M. Noor, E. R. Nwaeze, and Y.-M. Chu, *Quantum Hermite-Hadamard inequality by means of a Green function*, Adv. Difference Equ. **2020** (2020), 99, DOI: <https://doi.org/10.1186/s13662-020-02559-3>.
- [21] M. A. Noor, K. I. Noor, and M. U. Awan, *Some quantum estimates for Hermite-Hadamard inequalities*, Appl. Math. Comput. **251** (2015), 675–679, DOI: <https://doi.org/10.1016/j.amc.2014.11.090>.
- [22] M. A. Noor, K. I. Noor, and M. U. Awan, *Some quantum integral inequalities via preinvex functions*, Appl. Math. Comput. **269** (2015), 242–251, DOI: <https://doi.org/10.1016/j.amc.2015.07.078>.

- [23] M. A. Noor, K. I. Noor, and M. U. Awan, *Quantum Ostrowski inequalities for q -differentiable convex functions*, J. Math. Inequal. **10** (2016), no. 4, 1013–1018, DOI: <http://dx.doi.org/10.7153/jmi-10-81>.
- [24] P. M. Rajkovic, M. S. Stankovic, and S. D. Marinkovic, The zeros of polynomials orthogonal with respect to q -integral on several intervals in the complex plane, *Proceedings of The Fifth International Conference on Geometry, Integrability and Quantization*, Institute of Biophysics and Biomedical Engineering, Bulgarian Academy of Sciences, 2004.
- [25] Y. Zhang, T.-S. Du, H. Wang, and Y.-J. Shen, *Different types of quantum integral inequalities via (α, m) -convexity*, J. Inequal. Appl. **2018** (2018), 264, DOI: <https://doi.org/10.1186/s13660-018-1860-2>.
- [26] S. Bermudo, P. Kórus, and J. N. Valdés, *On q -Hermite-Hadamard inequalities for general convex functions*, Acta Math. Hungar. **162** (2020), 364–374, DOI: <https://doi.org/10.1007/s10474-020-01025-6>.
- [27] N. Alp, H. Budak, S. Erden, and M. Z. Sarikaya, *New bounds q -midpoint-type inequalities for twice q -differentiable convex functions on quantum calculus*, submitted, 2020.
- [28] M. Z. Sarikaya, A. Saglam, and H. Yıldırım, *New inequalities of Hermite-Hadamard type for functions whose second derivatives absolute values are convex and quasi-convex*, Int. J. Open Problems Comput. Math. **5** (2012), no. 3.
- [29] M. A. Noor and M. U. Awan, *Some integral inequalities for two kinds of convexities via fractional integrals*, TJMM **5** (2013), no. 2, 129–136.