


Gamal A. El-Hiti*, Keith Smith, Amany S. Hegazy, Saud A. Alanazi and Benson M. Kariuki

Crystal structure of 3-*tert*-butyl-3-hydroxy-1,3-dihydro-2*H*-pyrrolo[3,2-*c*]pyridin-2-one, $C_{11}H_{14}N_2O_2$



DOI 10.1515/ncrs-2015-0276

Received December 1, 2015; accepted March 1, 2016; available online April 2, 2016

Abstract

$C_{11}H_{14}N_2O_2$, orthorhombic, $P2_12_12_1$ (no. 19), $a = 7.5411(2)$ Å, $b = 11.5148(2)$ Å, $c = 12.5370(2)$ Å, $V = 1088.64(4)$ Å³, $Z = 4$, $R_{gt}(F) = 0.0301$, $wR_{ref}(F^2) = 0.0826$, $T = 296$ K.

CCDC no.: 1456766

The crystal structure is shown in the figure, Tables 1–3 contain details of the measurement method and a list of the atoms including atomic coordinates and displacement parameters.

Source of material

The title compound was prepared as previously reported *via* double lithiation of 4-pivaloylaminopyridine with *n*-BuLi (2.1 mole equivalents) in anhydrous THF at 0°C for 3 h followed by

Table 1: Data collection and handling.

Crystal:	Colourless, block, size 0.081 × 0.135 × 0.298 mm
Wavelength:	CuKα radiation (1.54184 Å)
μ:	7.17 cm ^{−1}
Diffractometer, scan mode:	SuperNova, Dual, Cu at zero, Atlas, ω scans
2θ _{max} :	148.2°
$N(hkl)_{measured}$, $N(hkl)_{unique}$:	3668, 2123
$N(param)_{refined}$:	141
Programs:	CrysAlis ^{PRO} [5], SHELX [6], WinGX [7], ChemDraw [8]

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
H(5)	4 <i>a</i>	0.1302	1.0553	0.0934	0.0480
H(6)	4 <i>a</i>	0.3050	1.0206	−0.0539	0.0520
H(7)	4 <i>a</i>	0.3790	0.6972	0.02510	0.0500
H(9A)	4 <i>a</i>	0.5447	0.7417	0.3790	0.1000
H(9B)	4 <i>a</i>	0.3600	0.8025	0.3922	0.1000
H(9C)	4 <i>a</i>	0.4705	0.8209	0.2877	0.1000
H(10A)	4 <i>a</i>	0.3884	0.5571	0.4235	0.0970
H(10B)	4 <i>a</i>	0.2357	0.5112	0.3498	0.0970
H(10C)	4 <i>a</i>	0.2024	0.6179	0.4249	0.0970
H(11A)	4 <i>a</i>	0.5359	0.6524	0.1737	0.1030
H(11B)	4 <i>a</i>	0.4211	0.5392	0.1812	0.1030
H(11C)	4 <i>a</i>	0.5708	0.5625	0.2653	0.1030
H(1)	4 <i>a</i>	0.0055	0.9476	0.2712	0.0480
H(2)	4 <i>a</i>	0.0242	0.6203	0.1404	0.0660

a reaction with carbon monoxide. The crude product obtained was purified by crystallization from ethyl acetate to give the title compound in 65% yield as colourless crystals [1].

Discussion

Synthesis of azaindoles is of great interest due to their growing biological applications [2]. The asymmetric unit consists of one molecule of $C_{11}H_{14}N_2O_2$. In the crystal structure, the hydroxyl group accepts one N–H···O hydrogen bond from a neighbouring molecule (N1–H1···O2 angle = 169.63°,

*Corresponding author: Gamal A. El-Hiti, Cornea Research Chair, Department of Optometry, College of Applied Medical Sciences, King Saud University, P. O. Box 10219, Riyadh 11433, Saudi Arabia, e-mail: gelhiti@ksu.edu.sa

Keith Smith, Amany S. Hegazy and Benson M. Kariuki: School of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, UK

Saud A. Alanazi: Department of Optometry, College of Applied Medical Sciences, King Saud University, P. O. Box 10219, Riyadh 11433, Saudi Arabia

Table 3: Fractional atomic coordinate and displacement parameters (Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> ₁₁	<i>U</i> ₂₂	<i>U</i> ₃₃	<i>U</i> ₁₂	<i>U</i> ₁₃	<i>U</i> ₂₃
C(1)	4a	0.0581(3)	0.7795(2)	0.2890(1)	0.0419(9)	0.0402(9)	0.0364(8)	0.0029(8)	0.0009(8)	0.0024(8)
C(2)	4a	0.1868(2)	0.7014(1)	0.2231(1)	0.0392(9)	0.0281(8)	0.0367(8)	−0.0005(7)	−0.0012(7)	0.0017(6)
C(3)	4a	0.2355(2)	0.7836(1)	0.1328(1)	0.0358(8)	0.0278(7)	0.0363(8)	0.0001(7)	−0.0026(7)	0.0000(7)
C(4)	4a	0.1578(2)	0.8916(1)	0.1517(1)	0.0356(8)	0.0300(8)	0.0354(8)	0.0007(7)	−0.0030(7)	−0.0021(6)
C(5)	4a	0.1812(3)	0.9829(2)	0.0818(2)	0.0428(9)	0.0275(8)	0.049(1)	0.0017(7)	−0.0024(9)	0.0034(7)
C(6)	4a	0.2847(3)	0.9599(2)	−0.0064(2)	0.046(1)	0.0376(9)	0.047(1)	−0.0022(8)	0.0019(9)	0.0118(8)
C(7)	4a	0.3312(3)	0.7698(2)	0.0404(2)	0.048(1)	0.0336(9)	0.0426(9)	0.0069(8)	0.0063(9)	0.0004(7)
C(8)	4a	0.3488(3)	0.6613(2)	0.2921(2)	0.0434(9)	0.0393(9)	0.044(1)	0.0023(8)	−0.0085(8)	0.0025(7)
C(9)	4a	0.4394(4)	0.7663(2)	0.3424(2)	0.062(1)	0.060(1)	0.078(2)	−0.004(1)	−0.027(1)	−0.004(1)
C(10)	4a	0.2882(4)	0.5793(2)	0.3808(2)	0.062(1)	0.071(2)	0.061(1)	0.000(1)	−0.017(1)	0.027(1)
C(11)	4a	0.4814(4)	0.5980(3)	0.2216(2)	0.067(2)	0.073(2)	0.067(2)	0.034(1)	−0.006(1)	0.004(1)
N(1)	4a	0.0594(2)	0.8886(1)	0.2445(1)	0.0487(8)	0.0339(7)	0.0385(8)	0.0088(7)	0.0043(7)	−0.0017(6)
N(2)	4a	0.3582(2)	0.8572(2)	−0.0288(1)	0.0509(9)	0.0437(9)	0.0443(9)	0.0044(7)	0.0099(8)	0.0072(7)
O(1)	4a	−0.0295(2)	0.7513(1)	0.3652(1)	0.071(1)	0.060(1)	0.0525(8)	0.0116(8)	0.0230(8)	0.0138(7)
O(2)	4a	0.0957(2)	0.6015(1)	0.1865(1)	0.0555(8)	0.0306(6)	0.0455(7)	−0.0087(6)	−0.0133(6)	0.0068(5)

N1⋯O2 distance = 2.851 Å) and donates one O—H⋯N bond to another molecule (O2—H2⋯N2 angle = 172.48°, O2⋯N2 distance = 2.710 Å) leading to a 3-D network. An extremely weak C—H⋯O contact is also observed (C5—H5⋯O1 angle = 173.13°, C⋯O distance = 3.362 Å) resulting in R²₂(9) graph set descriptor for this motif [3] between two molecules. The shortest C—H⋯O contact in the related 3-isopropyl-1-methyl-1,3-dihydro-2*H*-pyrrolo[3,2-*c*]pyridin-2-one [4] involves a tertiary C atom (C—H⋯O angle = 144.18° and C⋯O distance = 3.292 Å).

Experimental details

H atoms were placed in calculated positions and refined using a riding model, with *U*_{iso}(H) set to 1.2*U*_{eq}(C) and C—H and N—H distance of 0.93 and 0.86 Å respectively. The exceptions were the methyl (C—H = 0.96 Å) and hydroxyl (O—H = 0.86 Å) groups which were allowed to rotate around the C—C bond (HFIX 137 in SHELX [6]) and C—O bond (HFIX 147), with *U*_{iso}(H) set to 1.5*U*_{eq}(C/O).

Acknowledgements: The authors extend their appreciation to the Deanship of Scientific Research at King Saud University

for its funding for this research through the research group project RGP-VPP-239 and to Cardiff University for continued support.

References

- Smith, K.; El-Hiti, G. A.; Pritchard, G. J.; Hamilton, A.: Carbonylation of various organolithium reagents. A novel approach to heterocycles *via* intramolecular trapping of aromatic acyllithiums. *J. Chem. Soc., Perkin Trans. 1* (1999) 2299–2303.
- Popowycz, F.; Routier, S.; Joseph, B.; Mérou, J.-Y.: Synthesis and reactivity of 7-azaindole (1*H*-pyrrolo[2,3-*b*]pyridine). *Tetrahedron* **63** (2007) 1031–1064.
- Etter, M. C.; MacDonald, J. C.; Bernstein, J.: Graph-set analysis of hydrogen-bond patterns in organic crystals. *Acta Crystallogr. B* **46** (1990) 256–262.
- Harrison, W. T. A.; Male, L.; Hursthouse, M. B.: 3-isopropyl-1-methyl-1,3-dihydro-2*H*-pyrrolo[3,2-*c*]pyridin-2-one. University of Southampton, Crystal Structure Report Archive (2007) 337.
- Agilent: CrysAlis^{PRO}. Agilent Technologies, Yarnton, England, 2014.
- Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr. A* **64** (2008) 112–122.
- Farrugia, L. J.: WinGX and ORTEP for Windows: an update. *J. Appl. Crystallogr.* **45** (2012) 849–854.
- Cambridge Soft: ChemDraw Ultra. Cambridge Soft Corporation, Cambridge, Massachusetts, USA, 2001.