



Hans Reuter* and Tobias Gieschen

Crystal structure of (4-vinylpyridine- κN)triphenyl tin(IV) chloride, $C_{25}H_{22}ClNSn$



DOI 10.1515/ncrs-2016-0076

Received March 14, 2016; accepted June 23, 2016; available online August 10, 2016

Abstract

$C_{25}H_{22}ClNSn$, monoclinic, $P2_1/c$ (no. 14), $a = 17.8179(8)$ Å, $b = 15.6453(6)$ Å, $c = 17.4603(7)$ Å, $\beta = 116.588(2)^\circ$, $V = 4352.6(3)$ Å³, $Z = 8$, $R_{gt}(F) = 0.0297$, $wR_{ref}(F^2) = 0.0681$, $T = 100$ K.

CCDC no.: 1487302

The two crystallographically independent molecules forming the asymmetric unit of the crystal structure are shown in the figure. Tables 1 and 2 contain details of the measurement method and a list of the atoms including atomic coordinates and displacement parameters.

*Corresponding author: Hans Reuter, Institut für Chemie neuer Materialien, Universität Osnabrück, Barbarastr. 7, D-49069 Osnabrück, Germany, e-mail: hreuter@uos.de

Tobias Gieschen: Institut für Chemie neuer Materialien, Universität Osnabrück, Barbarastr. 7, D-49069 Osnabrück, Germany

Table 1: Data collection and handling.

Crystal:	Colourless, Blocks
	Size $0.21 \times 0.13 \times 0.12$ mm
Wavelength:	Mo $K\alpha$ radiation (0.71073 Å)
μ :	13.1 cm^{-1}
Diffractometer, scan mode:	Bruker APEX-II, φ and ω
$2\theta_{\text{max}}$, completeness:	56° , >99%
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$, R_{int} :	209691, 10503, 0.067
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 9081
$N(\text{param})_{\text{refined}}$:	512
Programs:	Bruker programs [12], SHELX [13, 14], DIAMOND [15]

Source of material

The title compound was obtained as a side product of the hydrostannylation of 4-vinylpyridine with triphenyltin(IV) hydride, Ph_3SnH , freshly prepared without purification from Ph_3SnCl and $\text{Li}[\text{AlH}_4]$.

Experimental details

All hydrogen atoms were identified in difference Fourier syntheses. All hydrogen atoms were included in the latest stages of the refinement by the use of a riding model. Common U_{iso}

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} [*] / <i>U</i> _{eq}
Sn1	0.54653(2)	0.65454(2)	0.24563(2)	0.01583(5)
Cl1	0.65677(4)	0.72810(4)	0.21916(4)	0.02173(12)
C111	0.56709(15)	0.73239(16)	0.35423(15)	0.0176(5)
C112	0.59856(16)	0.81499(16)	0.35912(17)	0.0223(5)
H112	0.6123	0.8350	0.3156	0.032(4)*
C113	0.60995(17)	0.86823(18)	0.42736(19)	0.0289(6)
H113	0.6311	0.9244	0.4299	0.032(4)*
C114	0.59065(17)	0.8398(2)	0.49150(18)	0.0303(6)
H114	0.5973	0.8768	0.5373	0.032(4)*
C115	0.56180(17)	0.7577(2)	0.48857(17)	0.0295(6)
H115	0.5497	0.7376	0.5332	0.032(4)*
C116	0.55017(16)	0.70392(18)	0.42060(16)	0.0232(5)
H116	0.5305	0.6473	0.4194	0.032(4)*
C121	0.44824(15)	0.68066(16)	0.12128(15)	0.0185(5)
C122	0.40999(17)	0.61406(18)	0.06424(17)	0.0250(6)
H122	0.4267	0.5568	0.0818	0.032(4)*
C123	0.34771(17)	0.6299(2)	−0.01805(18)	0.0286(6)
H123	0.3225	0.5837	−0.0564	0.032(4)*
C124	0.32246(17)	0.7129(2)	−0.04405(17)	0.0279(6)
H124	0.2796	0.7241	−0.1001	0.032(4)*
C125	0.36016(18)	0.7794(2)	0.01221(18)	0.0313(6)
H125	0.3431	0.8366	−0.0054	0.032(4)*
C126	0.42262(17)	0.76364(17)	0.09418(17)	0.0246(6)
H126	0.4481	0.8100	0.1321	0.032(4)*
C131	0.60506(15)	0.53290(16)	0.26762(16)	0.0185(5)
C132	0.65793(16)	0.50844(16)	0.35166(16)	0.0207(5)
H132	0.6674	0.5465	0.3975	0.029(2)*
C133	0.69679(16)	0.42900(17)	0.36884(17)	0.0244(5)
H133	0.7332	0.4133	0.4262	0.029(2)*
C134	0.68267(17)	0.37248(17)	0.30256(18)	0.0251(6)
H134	0.7091	0.3181	0.3144	0.029(2)*
C135	0.62967(17)	0.39587(17)	0.21883(18)	0.0243(5)
H135	0.6192	0.3571	0.1733	0.029(2)*
C136	0.59212(16)	0.47548(17)	0.20166(16)	0.0212(5)
H136	0.5569	0.4914	0.1440	0.029(2)*
N1	0.43663(13)	0.58054(13)	0.27428(13)	0.0174(4)
C141	0.37800(15)	0.62836(16)	0.28328(15)	0.0191(5)
H141	0.3714	0.6863	0.2651	0.029(2)*
C142	0.32724(15)	0.59712(16)	0.31765(16)	0.0200(5)
H142	0.2874	0.6334	0.3235	0.029(2)*
C143	0.33462(15)	0.51212(16)	0.34368(16)	0.0197(5)
C144	0.39198(17)	0.46202(17)	0.33018(18)	0.0244(5)
H143	0.3973	0.4030	0.3443	0.029(2)*
C145	0.44126(16)	0.49809(16)	0.29619(17)	0.0228(5)
H145	0.4803	0.4626	0.2880	0.029(2)*
C146	0.28546(17)	0.47389(19)	0.38371(17)	0.0261(6)
H146	0.2959	0.4154	0.3994	0.029(2)*
C147	0.22838(17)	0.5125(2)	0.40000(18)	0.0291(6)
H147	0.2156	0.5711	0.3855	0.029(2)*
H148	0.2000	0.4819	0.4262	0.029(2)*
Sn2	−0.04023(2)	0.31642(2)	0.23562(2)	0.01704(5)
Cl2	−0.14010(4)	0.24288(4)	0.28112(4)	0.02202(12)
C211	−0.05498(15)	0.21841(16)	0.14588(16)	0.0184(5)
C212	−0.10279(16)	0.23337(17)	0.05871(16)	0.0230(5)
H212	−0.1280	0.2878	0.0398	0.034(4)*
C213	−0.11414(17)	0.16992(18)	−0.00086(17)	0.0256(6)

Table 2 (continued)

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} [*] / <i>U</i> _{eq}
H213	−0.1467	0.1810	−0.0602	0.034(4)*
C214	−0.07798(16)	0.09020(17)	0.02617(17)	0.0236(5)
H214	−0.0867	0.0463	−0.0145	0.034(4)*
C215	−0.02935(16)	0.07484(17)	0.11221(17)	0.0240(5)
H215	−0.0039	0.0205	0.1306	0.034(4)*
C216	−0.01739(16)	0.13841(16)	0.17216(16)	0.0210(5)
H216	0.0165	0.1274	0.2313	0.034(4)*
C221	−0.11600(15)	0.42929(16)	0.19948(16)	0.0184(5)
C222	−0.12648(16)	0.48043(16)	0.12965(16)	0.0219(5)
H222	−0.1015	0.4636	0.0940	0.027(4)*
C223	−0.17287(17)	0.55533(17)	0.11179(17)	0.0260(6)
H223	−0.1790	0.5896	0.0644	0.027(4)*
C224	−0.21023(16)	0.58040(17)	0.16263(18)	0.0257(6)
H224	−0.2417	0.6319	0.1505	0.027(4)*
C225	−0.20162(16)	0.52997(17)	0.23141(18)	0.0247(6)
H225	−0.2273	0.5468	0.2664	0.027(4)*
C226	−0.15554(15)	0.45497(17)	0.24894(17)	0.0217(5)
H226	−0.1508	0.4203	0.2956	0.027(4)*
C231	0.06231(15)	0.31343(16)	0.36052(16)	0.0186(5)
C232	0.04892(16)	0.33591(17)	0.43085(17)	0.0228(5)
H232	−0.0059	0.3515	0.4221	0.041(4)*
C233	0.11439(17)	0.33589(18)	0.51340(17)	0.0261(6)
H233	0.1045	0.3529	0.5603	0.041(4)*
C234	0.19365(17)	0.31114(17)	0.52708(18)	0.0261(6)
H234	0.2382	0.3102	0.5836	0.041(4)*
C235	0.20822(16)	0.28768(17)	0.45865(18)	0.0258(6)
H235	0.2627	0.2699	0.4681	0.041(4)*
C236	0.14315(16)	0.29007(17)	0.37580(17)	0.0231(5)
H236	0.1541	0.2755	0.3289	0.041(4)*
N2	0.05415(13)	0.38467(13)	0.18790(14)	0.0196(4)
C241	0.07108(16)	0.46843(17)	0.20368(16)	0.0225(5)
H241	0.0471	0.4981	0.2349	0.033(3)*
C242	0.12123(16)	0.51308(17)	0.17688(16)	0.0225(5)
H242	0.1302	0.5725	0.1884	0.033(3)*
C243	0.15901(16)	0.47129(17)	0.13283(16)	0.0221(5)
C244	0.1437(2)	0.38428(19)	0.1193(2)	0.0347(7)
H244	0.1697	0.3524	0.0913	0.033(3)*
C245	0.0907(2)	0.34378(18)	0.1462(2)	0.0328(7)
H245	0.0799	0.2845	0.1347	0.033(3)*
C246	0.21274(17)	0.51564(19)	0.10180(17)	0.0273(6)
H246	0.2447	0.4813	0.0823	0.033(3)*
C247	0.22021(18)	0.5995(2)	0.09884(17)	0.0292(6)
H247	0.1893	0.6361	0.1178	0.033(3)*
H248	0.2565	0.6232	0.0778	0.033(3)*

parameters were refined for all H-atoms of any phenyl and vinyl group.

Discussion

In contrast to the structural chemistry of coordination compounds 4-vinylpyridine (4-ViPy) undergoes with many transition metals, *i.e.* ZnBr₂ · 2(4-ViPy) [1], Ni(NCS)₂ · 4(4-ViPy) [2], PtCl₂ · 2(4-ViPy) [3], Ag(ClO₃) · 2(4-ViPy) [4], and CuCl · 4(4-ViPy) [5], a research area strongly forced by the clathrate properties of TM(NCS)₂ · 4(4-ViPy)-complexes [6]

of transition metals TM, no structural information are available on its inorganic or organic tin(IV) complexes. Even the parent compound, pyridine (py), plays a minor role in the coordination chemistry of organotin(IV) compounds. Most structural information are available from octahedral 1:2 complexes of diorganotin(IV) dihalides of general formula $R_2SnHal_2 \cdot 2py$, i.e. $Et_2SnCl_2 \cdot 2(4-ViPy)$ [7], $Ph_2SnCl_2 \cdot 2(4-ViPy)$ [8], and $(CF_3-(CH_2)_2)_2SnCl_2 \cdot 2(4-ViPy)$ [9]. Moreover, only the structure of the 1:1 pyridine complex of the triorganotin(IV) monohalide, R_3SnHal with $R = \text{Methyl}$ and $Hal = Cl$ has been determined while from monoorganotin(IV) trihalides, R_3SnHal_3 , no complexes with this Lewis base are documented. As the structure determination of $Me_3SnCl \cdot py$ was determined from intensity data visually estimated from Weissenberg photographs the R -value isn't very low (18.6%). Consequently, structural parameters like bond lengths [i.e. $d(Sn-Cl) = 2.42(4) \text{ \AA}$] and angles are very imprecise [10]. The overall structure – a trigonal-bipyramidally coordinated tin atom with chlorine and pyridine in axial and the three methyl groups in equatorial positions, however, seems to be correct. This structural feature is also realised in the title structure, $Ph_3SnCl \cdot 4-ViPy$. The asymmetric unit consists of two crystallographic independent molecules with the tin atoms trigonal-bipyramidally coordinated from the three equatorially positioned phenyl groups and the chlorine atom and the 4-vinylpyridine molecule in axial positions. Within the equatorial plane, $Sn-C$ bond lengths differ from $2.121(2)$ – $2.143(2) \text{ \AA}$ in molecule 1 and from $2.123(2)$ – $2.139(2) \text{ \AA}$ in molecule 2 with a mean value over all six $Sn-C$ of $2.131(4) \text{ \AA}$. The range of bond angles between the phenyl groups is somewhat larger, ranging from $118.0(1)^\circ$ – $122.0(1)^\circ$ in molecule 1 and $117.4(1)^\circ$ – $121.7(1)^\circ$ in molecule 2. $C-C$ -bond lengths [mean value: $1.356(4) \text{ \AA}$] and angles within the phenyl groups are as expected, especially the *ipso*-effect [11] is well pronounced [$117.8(2)^\circ$ – $118.8(2)^\circ$]. Taking into account, that each of the two complexes exists as a pair of two enantiomers, the orientation of the three phenyl groups, defined by the angle between the normal vectors of the almost planar phenyl rings and the molecule axes, only slightly deviate from one molecule to another. Remarkable structural differences arise within the molecule axes, however. First of all, $Sn-Cl$ bond lengths differ significantly in both molecules as do $Sn-N$ bond lengths in opposite direction [$d(Sn-Cl) = 2.4918(6) \text{ \AA}/2.5272(6) \text{ \AA}$; $d(Sn-N) = 2.510(2) \text{ \AA}/2.432(2) \text{ \AA}$]. These values of $Sn-Cl$ bond lengths are much more longer than in case of $Me_3SnCl \cdot py$ (see above). Moreover, the orientation of the 4-vinylpyridine ligand in relation to the almost linear axis [$179.28(5)^\circ/178.28(5)^\circ$] as well as internal values for the rotation of the vinyl group around the $C-C$ single bond

to the pyridine moieties are strongly different. In molecule 1 the angle between the $Sn-N$ bond and the least-squares plane of the pyridine ligand has a value of $1.2(2)^\circ$, while in molecule 2 this angle has a value of $22.7(2)^\circ$. On the other side, the pyridine group and the vinyl groups show a dihedral angle of $2.3(2)^\circ$ in molecule 1, but $12.2(2)^\circ$ in molecule 2. All other bond lengths and angles within the pyridine moieties are as expected including the bond angle at nitrogen [$116.9(2)^\circ/117.3(2)^\circ$] and the $C-N$ bond lengths [$1.338(3) \text{ \AA}$ and $1.337(3) \text{ \AA}$ and $1.345(3) \text{ \AA}$]. The vinyl groups attached to the pyridine moieties are characterized by a $C-C$ single bond of $1.470(3)/1.458(4) \text{ \AA}$, a $C=C$ double bond of $1.319(4)/1.322(4) \text{ \AA}$, and a bond angle of $126.6(3)^\circ/125.3(3)^\circ$.

Acknowledgements: We acknowledge support by Deutsche Forschungsgemeinschaft (DFG) and Open Access Publishing Fund of Osnabrück University.

References

1. Hu, C.; Huster, J.; Englert, U.: Reversible crystal-to-crystal phase transition in dibromobis(4-vinylpyridine)zinc. *Z. Kristallogr.-Cryst. Mat.* **218** (2003) 761–765.
2. Lipkowski, J.; Suwińska, K.; Hatt, J.; Zielenkiewicz, A.; Zielenkiewicz, W.: On the clathratogenic properties of the $Ni(NCS)_2 \cdot (4\text{-Vinylpyridine})_4$ host. *J. Inclusion Phenom.* **2** (1984) 317–325.
3. Kubiak, M.; Kuduk-Jaworska, J.: Structure of cis-Dichlorobis(4-vinylpyridine)platinum(II). *Acta Crystallogr.* **C42** (1986) 1703–1705.
4. Georgiev, I. G.; Bučar, D.-K.; MacGillivray, L. R.: Stereospecific and quantitative photodimerisation of terminal olefins in the solid state. *Chem. Commun.* **46** (2010) 4956–4958.
5. Zhang, J.; Xiong, R.-G.; Chen, X.-T.; Xue, Z.; Peng, S.-M.; You, X.-Z.: Two-dimensional luminescent rectangular molecular grids containing copper(I)-olefin bonds vs. bridging spacers. *Organometallics* **21** (2002) 235–238.
6. Moore, M. H.; Nassimbeni, L. R.; Niven, M. L.; Taylor, M. W.: Studies in werner clathrates. Part 3. Structures of bis(isothiocyanato)tetra(4-vinylpyridine)nickel(II) and its clathrates with *ortho*-, *meta*-, and *para*-xylenes. *Inorg. Chim. Acta* **115** (1986) 211–217.
7. Casa, J. S.; García-Martínez, E.; Sánchez-González, A.; Sordo, J.; Villar, R.: Dichlorodiethylbis(pyridine-*N*)tin(IV). *Acta Crystallogr.* **C56** (2000) 299–300.
8. Yin, H.; Chen, S.; Wang, D.: Dichlorodiphenylbis(pyridine- κN)tin(IV). *Acta Crystallogr.* **E61** (2005) m2568–m2569.
9. Allouchi, H.; Cotrait, M.; Jousseau, B.; Rasclé, M.-C.; Toupance, T.: Dichlorobis(pyridine- κN)bis(3,3,3-trifluoropropyl- κC^1)tin(IV). *Acta Crystallogr.* **C58** (2002) m363–m364.

10. Hulme, R.: The crystal and molecular structure of chloro (trimethyl)-pyridinetin(IV). *J. Chem. Soc.* (1963) 1524–1527.
11. Jones, P. G.: The distortion of pentafluorophenyl groups in metal complexes: limitations of rigid-body refinement. *J. Organomet. Chem.* **345** (1988) 405–411.
12. Bruker: APEX2, SADABS, and SAINT. Bruker AXS Inc., Madison, USA 2009.
13. Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr.* **A64** (2008) 112–122.
14. Sheldrick, G. M.: Crystal Structure Refinement with SHELXL. *Acta Cryst.* **C71** (2015) 3–8.
15. Brandenburg, K.: DIAMOND. Visual Crystal Structure Information System. Crystal Impact, Bonn, Germany, 2006.