


Gamal A. El-Hiti*, Keith Smith, Amany S. Hegazy, Mohammad Hayal Alotaibi and Benson M. Kariuki

Crystal structure of 3-(2-bromophenyl)-1,1-dimethylthiourea, $C_9H_{11}BrN_2S$



DOI 10.1515/ncrs-2016-0116

Received April 13, 2016; accepted September 20, 2016; available online October 8, 2016

Abstract

$C_9H_{11}BrN_2S$, orthorhombic, $P2_12_12_1$ (no. 19), $a = 7.5187(3)$ Å, $b = 8.0634(3)$ Å, $c = 17.5320(6)$ Å, $V = 1062.90(7)$ Å³, $Z = 4$, $R_{gt}(F) = 0.0216$, $wR_{ref}(F^2) = 0.0536$, $T = 296(2)$ K.

CCDC no.: 1505381

The asymmetric unit of the title crystal structure is shown in the figure. Tables 1 and 2 contain details of the measurement method and a list of the atoms including atomic coordinates and displacement parameters.

Source of material

3-(2-Bromophenyl)-1,1-dimethylthiourea was synthesized from the dropwise addition of a solution of dimethylamine (1.1 equivalents) in ethanol to a stirred solution of 2-bromophenyl isothiocyanate in anhydrous dioxane over 5 min. The

Table 1: Data collection and handling.

Crystal:	Colourless needle
Size:	0.40 × 0.07 × 0.04 mm
Wavelength:	Cu $K\alpha$ radiation (1.54184 Å)
μ :	67.5 cm ⁻¹
Diffractometer, scan mode:	SuperNova, ω
$2\theta_{max}$, completeness:	147°, >99% up to 125.3°
$N(hkl)_{measured}$, $N(hkl)_{unique}$, R_{int} :	3493, 2052, 0.016
Criterion for I_{obs} , $N(hkl)_{gt}$:	$I_{obs} > 2 \sigma(I_{obs})$, 1969
$N(param)_{refined}$:	120
Programs:	CrysAlis ^{PRO} [12], SHELX [13], WinGX [14]

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	<i>x</i>	<i>y</i>	<i>z</i>	U_{iso}^*/U_{eq}
C1	−0.1300(4)	0.7897(4)	0.18698(16)	0.0385(6)
C2	−0.0864(4)	0.8683(4)	0.11919(15)	0.0397(6)
C3	−0.2084(6)	0.8842(5)	0.06025(18)	0.0516(9)
H3	−0.1772	0.9373	0.0151	0.062*
C4	−0.3767(6)	0.8197(5)	0.0699(2)	0.0580(10)
H4	−0.4603	0.8305	0.0312	0.070*
C5	−0.4225(5)	0.7389(6)	0.1369(3)	0.0606(10)
H5	−0.5360	0.6949	0.1427	0.073*
C6	−0.2998(5)	0.7239(5)	0.1949(2)	0.0513(8)
H6	−0.3310	0.6693	0.2397	0.062*
C7	−0.0060(4)	0.8368(4)	0.31579(16)	0.0383(6)
C8	0.3063(5)	0.7520(6)	0.3259(2)	0.0573(9)
H8A	0.2932	0.6364	0.3140	0.086*
H8B	0.4003	0.7659	0.3623	0.086*
H8C	0.3345	0.8123	0.2802	0.086*
C9	0.1465(7)	0.8677(6)	0.43742(19)	0.0663(11)
H9A	0.1900	0.9795	0.4402	0.100*
H9B	0.2242	0.7958	0.4656	0.100*
H9C	0.0291	0.8627	0.4588	0.100*
N1	0.0001(4)	0.7701(4)	0.24471(16)	0.0427(6)
H1	0.0914	0.7105	0.2337	0.051*
N2	0.1406(4)	0.8148(4)	0.35780(15)	0.0471(6)
S1	−0.18713(11)	0.94115(11)	0.34745(4)	0.0500(2)
Br1	0.14459(5)	0.95693(5)	0.10617(2)	0.05486(12)

*Corresponding author: Gamal A. El-Hiti, Cornea Research Chair, Department of Optometry, College of Applied Medical Sciences, King Saud University, P.O. Box 10219, Riyadh 11433, Saudi Arabia, e-mail: gelhiti@ksu.edu.sa

Keith Smith, Amany S. Hegazy and Benson M. Kariuki: School of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, UK

Mohammad Hayal Alotaibi: National Center for Petrochemicals Technology, King Abdulaziz City for Science and Technology, P.O. Box 6086, Riyadh 11442, Saudi Arabia

mixture was stirred for 1 h at room temperature. The solid obtained after work-up was purified by crystallization from a mixture of ethyl acetate and hexane (4:1 by volume) to give the title compound (92%) as light yellowish crystals, Mp 143–144 °C (lit. 142–143 °C) [1].

Experimental details

H atoms were positioned geometrically and refined using a riding model. $U_{\text{iso}}(\text{H})$ for aromatic and N–H hydrogens were set to 1.2 times U_{eq} of the parent atom. The values for the methyl groups were 1.5 times $U_{\text{eq}}(\text{C})$ with free rotation about the C–C bond. Aromatic C–H bonds were fixed at 0.93 Å, methyl C–H at 0.96 Å and N–H at 0.86 Å. The Flack parameter refined to a value of $-0.005(13)$ based on 790 quotients.

Discussion

Thiourea derivatives show various biological activities [2–5]. Therefore, the synthesis of such compounds is of general interest. The most common procedures for the synthesis of substituted thioureas involve reactions of amines with carbon disulfide in the presence of sodium or potassium hydroxide [6–8], of aliphatic amines with isocyanides in the presence of elemental sulfur [9] and of primary amines with isothiocyanates [10]. Thioureas can be used as precursors for the production of heterocycles, e.g. indigotin, *via* organolithium intermediates [1].

In the title structure the dimethylthiourea group is twisted from the plane of the bromophenyl moiety by 56.94(7)°. The amino groups are involved in intermolecular hydrogen bonds of the type N–H $\cdots$ S (with geometry: N $\cdots$ S = 3.410(3) Å, N–H $\cdots$ S = 141.5°) forming helical chains along [010]. The molecular conformation is similar to that found in the related 1-(2-bromo-4-chlorophenyl)-3,3-dimethylthiourea in which the intramolecular interplanar angle is 54.38(6)° and N–H $\cdots$ S hydrogen bonds also occur [11].

Acknowledgements: The authors extend their appreciation to the College of Applied Medical Sciences Research Centre and the Deanship of Scientific Research at King Saud University for their funding of this research and to Cardiff University for continued support.

References

- Smith, K.; Shukla, A. P.; Matthews, I.: A novel approach to indigotin and its substituted derivatives *via* carbonylation of doubly lithiated *N'*-aryl-*N,N*-dimethylthioureas. *Sulfur Lett.* **20** (1996) 121–137.
- Upadhyaya, R. S.; Kulkarni, G. M.; Vasireddy, N. R.; Vandavasi, J. K.; Dixit, S. S.; Sharma, V.; Chattapadhyaya, J.: Design, synthesis and biological evaluation of novel triazole, urea and thiourea derivatives of quinoline against *Mycobacterium tuberculosis*. *Bioorg. Med. Chem.* **17** (2009) 4681–4692.
- Khan, S. A.; Singh, N.; Saleem, K.: Synthesis, characterization and *in vitro* antibacterial activity of thiourea and urea derivatives of steroids. *Eur. J. Med. Chem.* **43** (2008) 2272–2277.
- Kocyigit-Kaymakcioglu, B.; Celen, A. O.; Tabanca, N.; Ali, A.; Khan, S. I.; Khan, I. A.; Wedge, D. E.: Synthesis and biological activity of substituted urea and thiourea derivatives containing 1,2,4-triazole moieties. *Molecules* **18** (2013) 3562–3576.
- Yao, J.; Chen, J.; Sun, W.; Xu, W.: Design, synthesis and biological activities of thiourea containing sorafenib analogs as antitumor agents. *Bioorg. Med. Chem.* **20** (2012) 2923–2929.
- Maddani, M. R.; Prabhu, K. R.: A concise synthesis of substituted thiourea derivatives in aqueous medium. *J. Org. Chem.* **75** (2010) 2327–2332.
- Abdel-Megeed, M. F.; Aly, Y. L.; Saleh, M. A.; Abdo, I. M.; El-Hiti, G. A.; Smith, K.: Novel one-pot procedure for the preparation of 3-substituted 2-thioxo-4(3*H*)quinazolones. *Sulfur Lett.* **19** (1995) 129–140.
- Boas, U.; Gertz, H.; Christensen, J. B.; Heegaard, P. M. H.: Facile synthesis of aliphatic isothiocyanates and thioureas on solid phase using peptide coupling reagents. *Tetrahedron Lett.* **45** (2004) 269–272.
- Nguyen, T. B.; Ermolenko, L.; Al-Mourabit, A.: Three-component reaction between isocyanides, aliphatic amines and elemental sulfur: Preparation of thioureas under mild conditions with complete atom economy. *Synthesis* **46** (2014) 3172–3179.
- Maki, T.; Tsuritani, T.; Yasukata, T.: A mild method for the synthesis of carbamate-protected guanidines using the Burgess reagent. *Org. Lett.* **16** (2014) 1868–1871.
- El-Hiti, G. A.; Smith, K.; Hegazy, A. S.; Alotaibic, M. H.; Kariuki, B. M.: 1-(2-Bromo-4-chlorophenyl)-3,3-dimethylthiourea. *Acta Crystallogr.* **E70** (2014) o704.
- Agilent. CrysAlis^{PRO}. Agilent Technologies, Yarnton, England (2014).
- Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr.* **A64** (2008) 112–122.
- Farrugia, L. J.: WinGX and ORTEP for Windows: an update. *J. Appl. Crystallogr.* **45** (2012) 849–854.