


Pan-Dong Mao, Ling-Ling Yan, Wei-Na Wu* and Hong-Xin Cai

N'-(5-ethoxycarbonyl-3,4-dimethyl-pyrrol-2-yl-methylidene)-4-hydroxybenzohydrazide monohydrate, C₁₇H₂₁N₃O₅



DOI 10.1515/ncrs-2017-0057

Received February 27, 2017; accepted August 15, 2017; available online September 13, 2017

Abstract

C₁₇H₂₁N₃O₅, monoclinic, *P*₂₁/*n* (no. 14), *a* = 9.2278(16) Å, *b* = 15.093(3) Å, *c* = 12.698(2) Å, β = 105.195(12)°, *V* = 1706.7(5) Å³, *Z* = 4, *R*_{gt}(*F*) = 0.0553, *wR*_{ref}(*F*²) = 0.1662, *T* = 296 K.

CCDC no.: 1569105

The asymmetric unit of the title crystal structure is shown in the figure. Tables 1 and 2 contain details of the measurement method and a list of the atoms including atomic coordinates and displacement parameters.

Source of materials

N'-(5-ethoxycarbonyl-3,4-dimethyl-pyrrol-2-yl-methylidene)-4-hydroxybenzohydrazide was prepared according to the

Table 1: Data collection and handling.

Crystal:	Yellow block
Size:	0.21 × 0.19 × 0.15 mm
Wavelength:	Mo Kα radiation (0.71073 Å)
μ:	1.0 cm ^{−1}
Diffractometer, scan mode:	Bruker APEX-II, φ and ω
2θ _{max} , completeness:	50°, >99%
<i>N</i> (<i>hkl</i>) _{measured} , <i>N</i> (<i>hkl</i>) _{unique} , <i>R</i> _{int} :	12944, 3000, 0.046
Criterion for <i>I</i> _{obs} , <i>N</i> (<i>hkl</i>) _{gt} :	<i>I</i> _{obs} > 2 σ(<i>I</i> _{obs}), 2016
<i>N</i> (<i>param</i>) _{refined} :	253
Programs:	SHELXL [1], Bruker programs [2]

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} [*] / <i>U</i> _{eq}
N2	0.6348(2)	0.02933(14)	0.93676(16)	0.0445(5)
N1	0.8876(2)	−0.06095(13)	1.05922(16)	0.0426(5)
H1A	0.8379	−0.0924	1.0052	0.051*
N3	0.5149(2)	0.07672(14)	0.87624(16)	0.0463(6)
H3A	0.5003	0.1306	0.8929	0.056*
O3	0.4397(2)	−0.04019(14)	0.76629(16)	0.0633(6)
C12	0.2949(3)	0.09113(17)	0.72426(19)	0.0418(6)
O4	−0.0472(2)	0.23575(14)	0.51712(15)	0.0651(6)
H4A	−0.0879	0.2690	0.5517	0.098*
C10	0.7236(3)	0.06866(17)	1.0172(2)	0.0426(6)
H10A	0.7049	0.1269	1.0337	0.051*
C14	0.1387(3)	0.22183(18)	0.6892(2)	0.0472(7)
H14A	0.1123	0.2767	0.7120	0.057*
C16	0.1013(3)	0.1067(2)	0.5556(2)	0.0547(7)
H16A	0.0500	0.0840	0.4881	0.066*
C15	0.0637(3)	0.18871(19)	0.5880(2)	0.0465(6)
O5	0.6797(2)	−0.15668(13)	0.87010(16)	0.0598(6)
C11	0.4201(3)	0.03682(18)	0.7897(2)	0.0443(6)
C13	0.2526(3)	0.17317(17)	0.7559(2)	0.0442(6)
H13A	0.3025	0.1958	0.8239	0.053*
C7	0.8525(3)	0.02272(16)	1.08231(19)	0.0402(6)
C4	1.0142(3)	−0.08758(18)	1.1352(2)	0.0442(6)
O2	1.1826(3)	−0.20519(16)	1.2033(2)	0.0844(7)
C6	0.9603(3)	0.05104(17)	1.1745(2)	0.0435(6)
C17	0.2150(3)	0.05843(19)	0.6233(2)	0.0519(7)
H17A	0.2388	0.0028	0.6010	0.062*
C5	1.0615(3)	−0.01860(18)	1.2083(2)	0.0439(6)
C9	0.9659(4)	0.1397(2)	1.2290(3)	0.0688(9)

*Corresponding author: Wei-Na Wu, College of Chemistry and Chemical Engineering, Henan Polytechnic University, Jiaozuo, 454000, P. R. China, e-mail: wwn08@hpu.edu.cn

Pan-Dong Mao: College of Chemistry and Chemical Engineering, Henan Polytechnic University, Jiaozuo, 454000, P. R. China

Ling-Ling Yan and Hong-Xin Cai: School of Physics and Electronic Information Engineering, Henan Polytechnic University, Jiaozuo, 454000, P. R. China

Table 2 (continued)

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} [*] / <i>U</i> _{eq}
H9A	1.0513	0.1419	1.2913	0.103*
H9B	0.8758	0.1483	1.2520	0.103*
H9C	0.9743	0.1855	1.1784	0.103*
O1	1.0057(3)	−0.21960(15)	1.0479(2)	0.0994(9)
C3	1.0783(4)	−0.1743(2)	1.1351(3)	0.0621(8)
C8	1.1961(3)	−0.0185(2)	1.3039(2)	0.0631(8)
H8A	1.2039	0.0377	1.3403	0.095*
H8B	1.2847	−0.0285	1.2793	0.095*
H8C	1.1865	−0.0647	1.3536	0.095*
C1 ^a	0.9520(14)	−0.3628(9)	1.0424(15)	0.110(5)
H1B ^a	0.9868	−0.3626	1.1206	0.165*
H1C ^a	0.9545	−0.4221	1.0158	0.165*
H1D ^a	0.8509	−0.3407	1.0207	0.165*
C1A ^b	0.9574(12)	−0.3590(6)	0.9697(12)	0.090(4)
H1A1 ^b	0.8638	−0.3672	0.9881	0.135*
H1A2 ^b	0.9984	−0.4157	0.9589	0.135*
H1A3 ^b	0.9408	−0.3248	0.9038	0.135*
C2A ^b	1.0609(13)	−0.3130(5)	1.0569(11)	0.079(4)
H2A1 ^b	1.1622	−0.3160	1.0483	0.095*
H2A2 ^b	1.0602	−0.3381	1.1271	0.095*
C2 ^a	1.050(2)	−0.3054(8)	0.9965(16)	0.118(7)
H2A ^a	1.0216	−0.3042	0.9173	0.141*
H2B ^a	1.1559	−0.3199	1.0236	0.141*
H5B	0.685(6)	−0.188(3)	0.815(3)	0.177*
H5A	0.626(5)	−0.111(2)	0.845(4)	0.177*

^aOccupancy: 0.464(18); ^bOccupancy: 0.536(18).

literature method [3]. The title crystals were obtained by slow evaporating of a THF/H₂O (1:1, v:v) solution at room temperature.

Experimental details

All H atoms were situated into idealized positions with the carrier atom-H distances = 0.93 Å for aryl, 0.96 Å for the methyl and 0.86 Å for the secondary amine H atoms. The *U*_{iso} values were constrained to be 1.5*U*_{eq} of the carrier atom for the methyl H atoms and 1.2*U*_{eq} for the remaining H atoms. Difference Fourier maps indicate methyl H atoms disorder, which has not been modelled.

Comment

Acylhydrazones are an important class of ligands in coordination chemistry and have found extensive application

in different fields [4]. Our previous work shows that acylhydrazone ligands bearing pyrrole units and their Cu(II) complexes exhibit considerable antibacterial and antitumor activity [3, 5, 6]. The structure of *N'*-(5-ethoxycarbonyl-3,4-dimethyl-pyrrol-2-yl-methylidene)-4-hydroxybenzohydrazide DMF adduct has been reported [3], while its monohydrate was characterized by X-ray diffraction in this work.

In the title crystal structure, the acylhydrazone molecule is in a ketone form and adopts an E configuration at the C = N double bond, in which all the bond lengths are comparable with those observed in the DMF adduct [6]. The dihedral angle between the pyrrole (N1/C4–C7, r.m.s. deviation 0.0034 Å) and the phenyl rings (C12–C17, r.m.s. deviation 0.0070 Å) is 10.7°. The torsion angles of N3–N2–C10–C7 and C11–N3–N2–C10 are 179.3(2)° and −178.95(19)°, respectively. In the solid state, the acylhydrazone molecules are linked into a two-dimensional supermolecular network by the crystal water molecules *via* intermolecular N–H⋯O and O–H⋯O hydrogen bonds.

References

- Sheldrick, G. M.: Crystal structure refinement with SHELXL. *Acta Crystallogr.* **C71** (2015) 3–8.
- Bruker. APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA, (2012).
- Ye, X.-P.; Zhu, T.-F.; Wu, W.-N.; Ma, T.-L.; Xu, J.; Zhang, Z.-P.; Wang, Y.; Jia, L.: Syntheses, characterizations and biological activities of two Cu(II) complexes with acylhydrazone ligand bearing pyrrole unit. *Inorg. Chem. Commun.* **47** (2014) 60–62.
- Dong, W.-K.; Li, X.-L.; Wang, L.; Zhang, Y.; Ding, Y.-J.: A new application of salamo-type bisoximes: As a relay-sensor for Zn²⁺/Cu²⁺ and its novel complexes for successive sensing of H⁺/OH[−]. *Sens. Actuators* **B229** (2016) 370–378.
- Han, X.-F.; Cai, H.-X.; Jia, L.; Wu, W.-N.; Zhang, X.; Xu, J.; Zhang, Z.-P.; Wang, Y.: Syntheses, characterizations and antitumor activities of two copper(II) complexes with an acylhydrazone ligand bearing Pyrrole Unit. *Chin. Inorg. Chem. Commun.* **31** (2015) 1453–1459.
- Ye, X.-P.; Wang, G.-J.; Pan, P.; Zhang, Z.-P.; Wu, W.-N.; Wang, Y.: Syntheses, crystal structures and biological activities of two Cu(II) complexes with an acylhydrazone ligand bearing pyrrole unit. *Chin. Inorg. Chem. Commun.* **30** (2014) 2789–2795.