

Amanda-Lee Manicum*, Orbett Alexander, Marietjie Schutte-Smith, Hendrik G. Visser and Andreas Roodt

Crystal structure of *fac*-(acetylacetonato- κ^2O,O')tricarbonyl(benzylidiphenylphosphine- κP)rhenium(I), $C_{27}H_{24}O_5PRe$




Table 1: Data collection and handling.

Crystal:	Yellow cuboid
Size:	0.36 × 0.19 × 0.11 mm
Wavelength:	Mo K α radiation (0.71073 Å)
μ :	50.3 cm ⁻¹
Diffractometer, scan mode:	Bruker APEX-II, φ and ω
$2\theta_{\max}$, completeness:	56.8°, 99%
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$, R_{int} :	6098, 6098, 0.052
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 5829
$N(\text{param})_{\text{refined}}$:	325
Programs:	Bruker programs [1], SHELX [2], SIR97 [3], DIAMOND [4], ORTEP [5], OLEX2 [6]

<https://doi.org/10.1515/ncrs-2017-0097>

Received April 6, 2017; accepted August 15, 2017; available online September 16, 2017

Abstract

$C_{27}H_{24}O_5PRe$, triclinic, $P\bar{1}$ (no. 2), $a = 9.0305(17)$ Å, $b = 12.084(2)$ Å, $c = 12.403(2)$ Å, $\alpha = 89.688(6)^\circ$, $\beta = 70.083(6)^\circ$, $\gamma = 76.315(6)^\circ$, $V = 1232.2(4)$ Å³, $Z = 2$, $R_{\text{gt}}(F) = 0.0180$, $wR_{\text{ref}}(F^2) = 0.0432$, $T = 100(2)$ K.

CCDC no.: 1569117

The asymmetric unit of the title crystal structure is shown in the figure. Tables 1 and 2 contain details of the measurement method and a list of the atoms including atomic coordinates and displacement parameters.

Source of materials

The starting complex *fac*-[Re(CO)₃(H₂O)(acac)] was prepared according to a published procedure [6]. *fac*-[Re(CO)₃(H₂O)(acac)] (39 mg; 0.1 mmol) was dissolved in MeOH (3 mL) and benzylidiphenylphosphine (PPh₂Bz) (0.028 g; 0.1 mmol) dissolved in methanol (2 mL) was added. The solution was stirred for 6 h at room temperature. The light yellow solution was left to crystallise and yellow crystals of the title complex were collected. Yield = 40 mg, 62%.

Experimental details

In the structure, all the H atoms were positioned geometrically and refined discernible using a riding model, with C–H_{methine} = 0.98 Å; C–H_{methylene} = 0.99 Å; C–H_{methyl} = 0.96 Å; C–H_{aromatic} = 0.93 Å. The U_{iso} parameters of the H atoms were fixed; $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for aromatic and methine and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$.

A limited number of inner reflections, which are cut by the primary beam stop have been omitted manually.

Comment

Some radionuclides of some metals are of significant interest for its rich coordination abilities and nuclear properties. The radiometal that has received significant attention is ^{99m}Tc, because it is easily accessible and possesses almost near perfect nuclear properties; ($t_{1/2} = 6.02$ h; $\gamma = 140$ keV, 100%).

*Corresponding author: Amanda-Lee Manicum, Department of Chemistry, University of the Free State, P.O. Box 339, Bloemfontein 9300, South Africa, e-mail: volminkae@ufs.ac.za
Orbett Alexander, Marietjie Schutte-Smith, Hendrik G. Visser and Andreas Roodt: Department of Chemistry, University of the Free State, P.O. Box 339, Bloemfontein 9300, South Africa

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	x	y	z	<i>U</i> _{iso} [*] / <i>U</i> _{eq}
Re1	0.19351(2)	0.29170(2)	0.12807(2)	0.01370(3)
P1	0.18998(7)	0.20541(5)	0.30979(5)	0.01253(11)
O1	−0.1769(2)	0.38212(16)	0.22543(17)	0.0255(4)
O2	0.1399(3)	0.07837(17)	0.02860(17)	0.0271(4)
O3	0.1887(3)	0.3977(2)	−0.0991(2)	0.0434(6)
O5	0.2380(2)	0.43764(15)	0.19616(16)	0.0196(4)
O4	0.4521(2)	0.22899(15)	0.06561(15)	0.0187(3)
C24	−0.1320(3)	−0.0925(2)	0.2441(2)	0.0246(5)
H24	−0.1588	−0.1519	0.2103	0.029*
C25	−0.0147(3)	−0.1183(2)	0.2949(2)	0.0232(5)
H25	0.0388	−0.1958	0.2966	0.028*
C26	0.0252(3)	−0.0316(2)	0.3435(2)	0.0186(5)
H26	0.1055	−0.0505	0.3786	0.022*
C23	−0.2101(3)	0.0208(2)	0.2428(2)	0.0253(6)
H23	−0.2918	0.0391	0.2089	0.030*
C16	0.1095(3)	0.3554(2)	0.5081(2)	0.0219(5)
H16	−0.0008	0.3555	0.5227	0.026*
C21	−0.0506(3)	0.0828(2)	0.3417(2)	0.0162(4)
C22	−0.1693(3)	0.1078(2)	0.2908(2)	0.0210(5)
H22	−0.2230	0.1851	0.2888	0.025*
C27	−0.0077(3)	0.1770(2)	0.3957(2)	0.0171(5)
H27A	−0.0050	0.1562	0.4725	0.021*
H27B	−0.0944	0.2481	0.4072	0.021*
C35	0.4843(3)	−0.1168(2)	0.2006(2)	0.0199(5)
H35	0.5311	−0.1651	0.1312	0.024*
C36	0.3999(3)	−0.0040(2)	0.2010(2)	0.0157(4)
H36	0.3902	0.0246	0.1316	0.019*
C14	0.3090(4)	0.4196(2)	0.5622(2)	0.0259(6)
H14	0.3360	0.4617	0.6142	0.031*
C34	0.4997(3)	−0.1583(2)	0.3018(2)	0.0201(5)
H34	0.5554	−0.2356	0.3018	0.024*
C31	0.3297(3)	0.06701(19)	0.3030(2)	0.0140(4)
C32	0.3492(3)	0.0252(2)	0.4040(2)	0.0169(5)
H32	0.3044	0.0738	0.4732	0.020*
C33	0.4339(3)	−0.0871(2)	0.4034(2)	0.0187(5)
H33	0.4468	−0.1153	0.4722	0.022*
C5	0.5586(3)	0.2839(2)	0.0614(2)	0.0185(5)
C11	0.2311(3)	0.2923(2)	0.4111(2)	0.0160(4)
C12	0.3919(3)	0.2954(2)	0.3892(2)	0.0196(5)
H12	0.4755	0.2538	0.3228	0.023*
C8	0.3628(4)	0.5744(2)	0.2351(3)	0.0324(7)
H8A	0.2486	0.6115	0.2786	0.049*
H8B	0.4086	0.6240	0.1769	0.049*
H8C	0.4241	0.5606	0.2876	0.049*
C7	0.3735(3)	0.4622(2)	0.1767(2)	0.0205(5)
C6	0.5261(3)	0.3947(2)	0.1098(2)	0.0220(5)
H6	0.6162	0.4274	0.0959	0.026*
C4	0.7328(3)	0.2191(3)	−0.0006(3)	0.0272(6)
H4A	0.7354	0.1430	−0.0294	0.041*
H4B	0.7930	0.2113	0.0526	0.041*
H4C	0.7831	0.2607	−0.0654	0.041*
C3	0.1949(4)	0.3613(2)	−0.0148(2)	0.0254(6)
C15	0.1486(4)	0.4185(2)	0.5836(2)	0.0271(6)
H15	0.0651	0.4606	0.6498	0.033*
C2	0.1598(3)	0.1578(2)	0.0673(2)	0.0180(5)
C1	−0.0364(3)	0.3486(2)	0.1884(2)	0.0183(5)
C13	0.4306(3)	0.3592(2)	0.4645(2)	0.0244(5)
H13	0.5404	0.3614	0.4489	0.029*

The 3rd row congener and 5d analogue of technetium, namely rhenium has been a significant focus, due to its β-emitting isotopes ¹⁸⁸Re and ¹⁸⁶Re that have a potential use in radiotherapy. Since its development by Alberto *et al.* [7–9], the *fac*-[M(H₂O)₃(CO)₃]⁺ (M = Tc, Re) has been extensively used because it opens innovative prospects for novel labelling strategies for the development of new diagnostic and therapeutic radiopharmaceutical drugs [7–13]. The primary reason for its popularity is because of the inert nature of the *fac*-[M(CO)₃]⁺ core in aqueous media and also the lability of the aqua ligands to be substituted by combinations of various mono- and bidentate ligands (*OO'*-, *NO*-, *NN'*-, *NS*-donating, etc.) *via* the '2 + 1' mixed ligand approach [14–21].

The geometry around the title complex, *fac*-[Re(Acac)(CO)₃(PPh₂Bz)] is typically octahedral and the central rhenium metal is bound to three *fac* carbonyl ligands, a chelate β-diketonato ligand, acetylacetonato (Acac) through its oxygen atoms and to a monodentate benzylidiphenylphosphine ligand (PPh₂Bz) through its phosphorus atom. In the coordination sphere, the Re1–P1 bond distance of 2.4718(7) Å is the longest and is consistent with those found in similar Re(I)-tricarbonyl complexes. All the bond distances and angles are considered to be within the normal range [11–24].

Acknowledgements: Financial assistance from the University of the Free State is gratefully acknowledged. We also express our gratitude towards NTeMBI, PETLabs Pharmaceuticals, SASOL and the South African National Research Foundation (SA-NRF/THRIP). This work is based on the research supported in part by the National Research Foundations of South-Africa and the Thuthuka programme (Grant specific unique reference numbers (UID) 84913 and 94142).

References

1. Bruker. SADABS and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA (2004).
2. Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr. A* **64** (2008) 112–122.
3. Altomare, A.; Burla, M. C.; Camalli, M.; Cascarano, G. L.; Giacovazzo, C.; Guagliardi, A.; Moliterni, A. G. G.; Polidori, G.; Spagna, R.: SIR97: a new tool for crystal structure determination and refinement. *J. Appl. Crystallogr.* **32** (1999) 115–119.
4. Brandenburg, K.; Putz, H.: DIAMOND. Visual crystal structure information system. Version 3.2i. Crystal Impact, Bonn, Germany (2012).
5. Farrugia, L. J.: WinGX and ORTEP for windows: an update. *J. Appl. Crystallogr.* **45** (2012) 849–854.
6. Dolomanov, O. V.; Bourhis, L. J.; Gildea, R. J.; Howard, J. A. K.; Puschmann, H.: OLEX2: a complete structure solution,

- refinement and analysis program. *J. Appl. Crystallogr.* **42** (2009) 339–341.
7. Schibli, R.; Alberto, R.; Abram, U.; Abram, S.; Schubiger, P. A.; Kaden, T. A.: Structural and ^{99}Tc NMR investigations of complexes with *fac*-[Tc(CO) $_3$] $^+$ moieties and macrocyclic thioethers of various ring sizes: synthesis and X-ray structure of the complexes *fac*-[Tc(9-ane-S $_3$)(CO) $_3$]Br, *fac*-[Tc $_2$ (tosylate) $_2$ (18-ane-S $_6$)(CO) $_6$], and *fac*-[Tc $_2$ (20-ane-S $_6$ OH)(CO) $_6$][tosylate] $_2$. *Inorg. Chem.* **37** (1998) 3509–3516.
 8. Alberto, R.; Schibli, R.; Egli, A.; Schubiger, A. P.; Abram, U.; Kaden, T. A.: A novel organometallic aqua complex of technetium for the labeling of biomolecules: synthesis of [$^{99m}Tc(OH_2)_3(CO)_3$] $^+$ from [$^{99m}TcO_4$] $^-$ in aqueous solution and its reaction with a bifunctional ligand. *J. Am. Chem. Soc.* **120** (1998) 7987–7988.
 9. Alberto, R.; Schibli, R.; Schubiger, A. P.; Abram, U.; Pietzsch, H. J.; Johannsen, B.: First application of *fac*-[$^{99m}Tc(OH_2)_3(CO)_3$] $^+$ in bioorganometallic chemistry: design, structure and *in vitro* affinity of 5-HT $_1A$ receptor ligand labeled with Tc-99m. *J. Am. Chem. Soc.* **121** (1999) 6076–6078.
 10. Botha, J. M.; Roodt, A.: Kinetic and high-pressure mechanistic investigation of the aqua substitution in the *Trans*-Aquaotetracyano Complexes of Re(V) and Tc(V): some implications for nuclear medicine. *Met.-Based Drugs.* **2008** (2008) 745989–745998.
 11. Visser, H. G.; Roodt, A.; Volmink, A.; Kemp, G.: *fac*-tricarbonyl-(pyridine- κN)(1,1,1-trifluoroacetylacetonato- $\kappa^2 O, O'$)rhenium(II). *Acta Crystallogr.* **E67** (2011) m1631–m1631.
 12. Manicum, A.; Schutte-Smith, M.; Kemp, G.; Visser, H. G.: Illustration of the electronic influence of coordinated β -diketone type ligands: a kinetic and structural study. *Polyhedron* **85** (2015) 190–195.
 13. Manicum, A.; Schutte-Smith, M.; Visser, H. G.; Pretorius, C.; Roodt, A.: Crystal structure of *fac*-tricarbonyl(hexafluoroacetylacetonato- $\kappa^2 O, O'$)-(nitrate- κO) rhenium(II), $C_{16}H_{21}O_8N_2F_6Re$. *Z. Kristallogr. -NCS* **231** (2016) 263–266.
 14. Sagnou, M.; Tsoukalas, C.; Triantis, C.; Raptopoulou, C. P.; Terzis, A.; Pirmettis, I.; Pelecanou, M.; Papadopoulos, M.: A new tricarbonyl *fac*-[M(acac)(isc)(CO) $_3$] complex (M = Re, ^{99m}Tc) with acetylacetonate (acac) and isocyanide (isc) in a 2 + 1 combination. *Inorg. Chim. Acta* **363** (2010) 1649–1653.
 15. Brink, A.; Visser, H. G.; Roodt, A.: Tetraethylammonium (acetylacetonato)-bromidotricarbonylrhenate(II). *Acta Crystallogr.* **E67** (2011) m34–m35.
 16. Schutte, M.; Roodt, A.; Visser, H. G.: Coordinated aqua vs methanol substitution kinetics in *fac*-Re(II) tricarbonyl tropolonato complexes. *Inorg. Chem.* **51** (2012) 11996–12006.
 17. Brink, A.; Visser, H. G.; Roodt, A.: Novel imino rhenium(II) tricarbonyl complexes of salicylidene-derived ligands: synthesis, X-ray crystallographic studies, spectroscopic characterization and DFT calculations. *Polyhedron* **52** (2013) 416–423.
 18. Schutte, M.; Kemp, G.; Visser, H. G.; Roodt, A.: Tuning the reactivity in classic low-spin d^6 rhenium(II) radiopharmaceutical synthon by selective bidentate ligand variation (L, L' -Bid; $L, L' = N, N', N, O$, and O, O' donor atom sets) in *fac*-[Re(CO) $_3$ (L, L' -Bid)(MeOH)] n complexes. *Inorg. Chem.* **50** (2011) 12486–12498.
 19. Schutte, M.; Roodt, A.; Visser, H. G.: Coordinated aqua vs methanol substitution kinetics in *fac*-Re(II) tricarbonyl tropolonato complexes. *Inorg. Chem.* **51** (2012) 11996–12006.
 20. Benny, P. D.; Fugate, G. A.; Barden, A. O.; Morley, J. E.; Silva-Lopez, E.; Twamley, B.: Metal-assisted in situ formation of a tridentate acetylacetonate ligand for complexation of *fac*-Re(CO) $_3$ $^+$ for radiopharmaceutical applications. *Inorg. Chem.* **47** (2008) 2240–2242.
 21. Sagnou, M.; Benaki, D.; Triantis, C.; Tsotakos, T.; Psycharis, V.; Raptopoulou, C. P.; Pirmettis, I.; Papadopoulos, M.; Pelecanou, M.: Curcumin as the OO bidentate ligand in '2 + 1' complexes with the [M(CO) $_3$] $^+$ (M = Re, ^{99m}Tc) tricarbonyl core for radiodiagnostic applications. *Inorg. Chem.* **50** (2011) 1295–1303.
 22. Manicum, A.; Visser, H. G.; Engelbrecht, I.; Roodt, A.: Crystal structure of *fac*-(acetylacetonato- $\kappa^2 O, O'$)tricarbonyl(cyclohexyldiphenylphosphine- κP) rhenium(II), $C_{26}H_{28}O_5PRE$. *Z. Kristallogr. -NCS* **230** (2015) 150–152.
 23. Triantis, C.; Tsotakos, T.; Tsoukalas, C.; Sagnou, M.; Raptopoulou, C.; Terzis, A.; Psycharis, V.; Pelecanou, M.; Pirmettis, I.; Papadopoulos, M.: Synthesis and characterization of *fac*-[M(CO) $_3$ (P)(OO)] and *cis-trans*-[M(CO) $_2$ (P) $_2$ (OO)] complexes (M = Re, ^{99m}Tc) with acetylacetonate and curcumin as OO donor bidentate ligands. *Inorg. Chem.* **52** (2013) 12995–13003.
 24. Hayes, T. R.; Kasten, B. B.; Barnes, C. L.; Benny, P. D.: Rhenium and technetium bi- and tricarbonyl complexes in a new strategy for biomolecule incorporation using click chemistry. *Dalton Trans.* **43** (2014) 6998–7001.