


Wang Li-Hua, Liang Lei, Li Xiang-Tao, Cao Shu-Hua and Tai Xi-Shi*

The crystal structure of bis(6-phenylpyridine-2-carboxylato- κ^2N,O)copper(II), $C_{24}H_{16}N_2O_4Cu$



<https://doi.org/10.1515/ncrs-2021-0293>

Received July 8, 2021; accepted September 9, 2021;
published online September 22, 2021

Abstract

$C_{24}H_{16}N_2O_4Cu$, triclinic, $P\bar{1}$ (no. 2), $a = 11.2454(6)$ Å, $b = 13.5960(7)$ Å, $c = 14.0076(8)$ Å, $V = 1958.9(2)$ Å³, $\alpha = 79.019(5)^\circ$, $\beta = 71.729(5)^\circ$, $\gamma = 76.126(5)^\circ$, $Z = 4$, $R_{gt}(F) = 0.0438$, $wR_{ref}(F^2) = 0.1030$, $T = 150$ K.

CCDC no.: 2108727

*Corresponding author: Tai Xi-Shi, College of Chemistry and Chemical Engineering, Weifang University, Weifang, Shandong 261061, P. R. China, E-mail: taixs@wfu.edu.cn. <https://orcid.org/0000-0002-0050-1900>

Wang Li-Hua, Li Xiang-Tao and Cao Shu-Hua, College of Chemistry and Chemical Engineering, Weifang University, Weifang, Shandong 261061, P. R. China

Liang Lei, School of Harbin Light Industry, Harbin, Heilongjiang 150040, P. R. China

Table 1: Data collection and handling.

Crystal:	Green block
Size:	0.12 × 0.11 × 0.10 mm
Wavelength:	Mo K α radiation (0.71073 Å)
μ :	1.15 mm ⁻¹
Diffractometer, scan mode:	SuperNova, ω
θ_{max} , completeness:	25.0°, 99%
$N(hkl)_{measured}$, $N(hkl)_{unique}$, R_{int} :	12,959, 6850, 0.037
Criterion for I_{obs} , $N(hkl)_{gt}$:	$I_{obs} > 2 \sigma(I_{obs})$, 5525
$N(param)_{refined}$:	559
Programs:	Bruker [1], Olex2 [2], SHELX [3], Diamond [4]

One of the two crystallographically independent Cu(II) complexes is shown in the figure. Table 1 contains crystallographic data and Table 2 contains the list of the atoms including atomic coordinates and displacement parameters.

Source of material

About 0.5 mmol (0.0996 g) of 6-phenylpyridine-2-carboxylic acid, 0.5 mmol (0.020 g) of NaOH and 0.25 mmol of (0.0499 g) copper(II) acetate monohydrate were added to 20 ml of water-ethanol (v:v = 1:1) at room temperature. After 10 min, precipitation appeared in the solution. The mixture was continuously stirred for 4.5 h at 85 °C. The solution was cooled to room temperature and filtered. After 16 days, the green crystals of the title compound were obtained from the filtrate.

Experimental details

The hydrogen atoms were positioned geometrically (C–H = 0.93 Å). Their U_{iso} values were set to $1.2U_{eq}$ of the parent atoms.

Comment

Some copper complexes with ligands containing a pyridine-carboxylate moiety exhibit novel coordination compounds [5]

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} */ <i>U</i> _{eq}
Cu1	0.47135 (4)	0.36703 (3)	−0.00940 (3)	0.01507 (12)
O1	0.8121 (2)	0.38318 (17)	−0.00452 (18)	0.0234 (6)
O2	0.6290 (2)	0.41365 (17)	−0.04643 (17)	0.0198 (5)
O3	0.4163 (2)	0.43253 (17)	−0.12634 (16)	0.0183 (5)
O4	0.2494 (2)	0.4538 (2)	−0.18717 (18)	0.0298 (6)
N1	0.5682 (2)	0.24241 (19)	0.05580 (19)	0.0140 (6)
N2	0.2892 (2)	0.36682 (19)	0.05673 (19)	0.0134 (6)
C1	0.5310 (3)	0.1537 (2)	0.1029 (2)	0.0176 (7)
C2	0.5941 (3)	0.0918 (3)	0.1708 (3)	0.0236 (8)
H2	0.565617	0.033002	0.207054	0.028*
C3	0.6980 (4)	0.1174 (3)	0.1840 (3)	0.0249 (8)
H3	0.738621	0.077277	0.230657	0.030*
C4	0.7423 (3)	0.2033 (3)	0.1278 (2)	0.0189 (8)
H4	0.816359	0.218867	0.132232	0.023*
C5	0.6743 (3)	0.2651 (2)	0.0650 (2)	0.0144 (7)
C6	0.7116 (3)	0.3618 (2)	0.0008 (2)	0.0162 (7)
C7	0.4264 (3)	0.1268 (2)	0.0767 (3)	0.0174 (7)
C8	0.3309 (3)	0.0833 (3)	0.1505 (3)	0.0247 (8)
H8	0.336002	0.066381	0.216851	0.030*
C9	0.2291 (4)	0.0650 (3)	0.1260 (3)	0.0307 (9)
H9	0.164233	0.038210	0.176016	0.037*
C10	0.2241 (4)	0.0870 (3)	0.0265 (3)	0.0355 (10)
H10	0.155572	0.074778	0.009917	0.043*
C11	0.3197 (4)	0.1267 (3)	−0.0482 (3)	0.0274 (9)
H11	0.316553	0.139810	−0.115079	0.033*
C12	0.4203 (3)	0.1470 (3)	−0.0233 (3)	0.0209 (8)
H12	0.484474	0.174284	−0.073585	0.025*
C13	0.2995 (3)	0.4290 (2)	−0.1177 (3)	0.0193 (8)
C14	0.2240 (3)	0.3922 (2)	−0.0132 (2)	0.0158 (7)
C15	0.0965 (3)	0.3869 (2)	0.0119 (3)	0.0197 (8)
H15	0.053908	0.400949	−0.037827	0.024*
C16	0.0334 (3)	0.3603 (3)	0.1119 (3)	0.0259 (9)
H16	−0.052350	0.355959	0.130661	0.031*
C17	0.0993 (3)	0.3405 (3)	0.1832 (3)	0.0222 (8)
H17	0.056855	0.325009	0.251158	0.027*
C18	0.2283 (3)	0.3432 (2)	0.1553 (2)	0.0166 (7)
C19	0.3021 (3)	0.3258 (2)	0.2304 (2)	0.0163 (7)
C20	0.2830 (3)	0.2501 (3)	0.3139 (2)	0.0216 (8)
H20	0.221951	0.210798	0.323719	0.026*
C21	0.3546 (3)	0.2334 (3)	0.3819 (3)	0.0272 (9)
H21	0.343072	0.181918	0.436272	0.033*
C22	0.4434 (3)	0.2934 (3)	0.3690 (3)	0.0283 (9)
H22	0.491755	0.282182	0.414472	0.034*
C23	0.4601 (3)	0.3703 (3)	0.2880 (3)	0.0221 (8)
H23	0.519048	0.411157	0.279836	0.026*
C24	0.3898 (3)	0.3866 (3)	0.2196 (2)	0.0176 (7)
H24	0.401320	0.438634	0.165839	0.021*
Cu2	1.00839 (4)	0.13203 (3)	−0.47829 (3)	0.01778 (12)
O5	0.9162 (2)	0.11547 (18)	−0.72097 (17)	0.0250 (6)
O6	1.0172 (2)	0.08432 (18)	−0.60066 (17)	0.0220 (6)
O7	1.1771 (2)	0.07280 (17)	−0.46650 (16)	0.0192 (5)
O8	1.2948 (2)	0.0688 (2)	−0.36298 (19)	0.0321 (6)
N3	0.9029 (2)	0.2582 (2)	−0.53186 (19)	0.0151 (6)
N4	0.9638 (3)	0.1379 (2)	−0.3318 (2)	0.0168 (6)
C25	0.8637 (3)	0.2375 (3)	−0.6061 (2)	0.0186 (8)

Table 2: (continued)

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} */ <i>U</i> _{eq}
C26	0.9351 (3)	0.1376 (3)	−0.6471 (2)	0.0199 (8)
C27	0.7712 (3)	0.3043 (3)	−0.6439 (3)	0.0233 (8)
H27	0.745555	0.288584	−0.695283	0.028*
C28	0.7174 (4)	0.3956 (3)	−0.6033 (3)	0.0283 (9)
H28	0.652838	0.441333	−0.625788	0.034*
C29	0.7603 (3)	0.4182 (3)	−0.5294 (3)	0.0242 (8)
H29	0.724771	0.479233	−0.501804	0.029*
C30	0.8567 (3)	0.3495 (3)	−0.4962 (2)	0.0187 (8)
C31	0.9184 (3)	0.3733 (2)	−0.4269 (2)	0.0184 (8)
C32	1.0510 (3)	0.3503 (3)	−0.4491 (3)	0.0226 (8)
H32	1.099580	0.322956	−0.508520	0.027*
C33	1.1108 (4)	0.3680 (3)	−0.3831 (3)	0.0287 (9)
H33	1.199345	0.352157	−0.398076	0.034*
C34	1.0390 (4)	0.4094 (3)	−0.2945 (3)	0.0343 (10)
H34	1.079088	0.420635	−0.249784	0.041*
C35	0.9079 (4)	0.4339 (3)	−0.2732 (3)	0.0302 (9)
H35	0.859501	0.461271	−0.213801	0.036*
C36	0.8481 (3)	0.4178 (2)	−0.3397 (3)	0.0238 (8)
H36	0.759896	0.437046	−0.325909	0.029*
C37	1.1925 (3)	0.0842 (3)	−0.3819 (3)	0.0208 (8)
C38	1.0703 (3)	0.1178 (2)	−0.3015 (3)	0.0183 (7)
C39	1.0671 (4)	0.1298 (3)	−0.2052 (3)	0.0247 (8)
H39	1.141907	0.117809	−0.186041	0.030*
C40	0.9499 (4)	0.1601 (3)	−0.1378 (3)	0.0320 (10)
H40	0.944929	0.169499	−0.072553	0.038*
C41	0.8409 (4)	0.1761 (3)	−0.1674 (3)	0.0264 (9)
H41	0.761823	0.194825	−0.121485	0.032*
C42	0.8480 (3)	0.1647 (2)	−0.2657 (2)	0.0190 (8)
C43	0.7331 (3)	0.1776 (2)	−0.3004 (2)	0.0186 (7)
C44	0.6271 (3)	0.2538 (3)	−0.2698 (3)	0.0243 (8)
H44	0.628896	0.298754	−0.228092	0.029*
C45	0.5194 (3)	0.2635 (3)	−0.3006 (3)	0.0287 (9)
H45	0.449866	0.315749	−0.281008	0.034*
C46	0.5146 (4)	0.1952 (3)	−0.3609 (3)	0.0307 (9)
H46	0.441200	0.200751	−0.380364	0.037*
C47	0.6193 (3)	0.1189 (3)	−0.3921 (3)	0.0274 (9)
H47	0.616644	0.074029	−0.433531	0.033*
C48	0.7275 (3)	0.1094 (3)	−0.3619 (2)	0.0222 (8)
H48	0.797163	0.057487	−0.382312	0.027*

and excellent properties in many ways such as non-linear optical (NLO) property, antitumor activity, electrical conductance property, magnetic property [6–9]. Therefore, studies on structure and property of copper complexes have been one of the hot topics. In our previous work, some metal complexes with ligands containing pyridine-carboxylate moieties have been synthesized and structural characterized [10–12]. In order to further enrich the structure of the copper complexes with ligands such, a new Cu(II) complex was synthesized using 6-phenylpyridine-2-carboxylic acid, NaOH and cupric acetate monohydrate as raw materials.

The asymmetric unit of Cu(II) complex contains two Cu(II) ion and four bidentate 6-phenylpyridine-2-carboxylate

ligands. Each Cu(II) ion is four-coordinated with two O atoms and two N atoms from two different 6-phenylpyridine-2-carboxylate ligands. The sum of bond angles around Cu1, for example is 372.47° , showing that the O2, N1, N2, O3 is almost coplanar. The complex molecules form a 1D chained structure by the π - π stacking of phenyl and pyridine rings.

Author contributions: All the authors have accepted responsibility for the entire content of this submitted manuscript and approved submission.

Research funding: This project was supported by the National Natural Science Foundation of China (No. 21171132, <http://dx.doi.org/10.13039/501100001809>), the Natural Science Foundation of Shandong (ZR2014BL003, <http://dx.doi.org/10.13039/501100007129>), the Project of Shandong Province Higher Educational Science and Technology Program (J14LC01, <http://dx.doi.org/10.13039/501100015642>) and Science Foundation of Weifang.

Conflict of interest statement: The authors declare no conflicts of interest regarding this article.

References

1. Bruker. SAINT and SADABS; Bruker AXS Inc.: Madison, Wisconsin, USA, 2000.
2. Dolomanov O. V., Bourhis L. J., Gildea R. J., Howard J. A. K., Puschmann H. OLEX2: a complete structure solution, refinement and analysis program. *J. Appl. Crystallogr.* 2009, 42, 339–341.
3. Sheldrick G. M. Crystal structure refinement with SHELXL. *Acta Crystallogr.* 2015, C71, 3–8.
4. Brandenburg K. DIAMOND. Visual Crystal Structure Information System (Ver. 3.2); Crystal Impact: Bonn, Germany, 2012.
5. Lv B.-L. Crystal structure of *catena*-poly[aqua-(4-iodopyridine-2,6-dicarboxylato- κ^2N,O,O')-(4-amino-4H-1,2,4-triazole- $\kappa^2N:N'$)copper(II)], $C_9H_8N_5O_5CuI$. *Z. Kristallogr. N. Cryst. Struct.* 2021, 236, 873–875.
6. Vural H., Orbay M. Synthesis, crystal structure, spectroscopic investigations and DFT calculations of the copper(II) complex of 4-(trifluoromethyl)pyridine-2-carboxylic acid. *J. Mol. Struct.* 2017, 1146, 669–676.
7. Li C.-P., Huang T.-F., Fu Y., Liu Y.-X., Zhou S.-F., Qi Z.-Y., Li C.-Z. Interaction of di-2-pyridylketone 2-pyridine carboxylic acid hydrazone and its copper complex with BSA: effect on antitumor activity as revealed by spectroscopic studies. *Molecules* 2016, 21, 563.
8. Pathak S., Chakraborty K., Ghosh S., Roy K., Jana B., Konar S. Synthesis and crystal structure of new copper(II) metal complex: noncovalent interactions and electrical conductance properties. *J. Mol. Struct.* 2018, 1152, 96–100.
9. Zhang X.-Y., Wen H.-F., Yang Q.-F., Zhou R.-S., Song J.-F. Eight new coordination polymers containing rigid 4-(4-carboxyphenyl)-pyridine-2-carboxylic acid: synthesis, structural diversity, fluorescence and magnetic properties. *Inorg. Chim. Acta.* 2020, 507, 119600.
10. Tai X.-S., Jiang G.-Y., Yu Y.-T., Yao H., Wang A.-J. Bis(ethanol- κO)-bis(6-aminopicolinato- κ^2N,O) magnesium(II), $C_{16}H_{22}N_4O_6Mg$. *Z. Kristallogr. N. Cryst. Struct.* 2019, 234, 289–290.
11. Tai X.-S., Zhou X.-J., Liu L.-L., Cao S.-H., Wang L.-H. The crystal structure of diaqua-bis(6-aminopicolinato- κ^2N,O)magnesium(II), $C_{12}H_{14}O_6N_4Mg$. *Z. Kristallogr. N. Cryst. Struct.* 2020, 235, 605–606.
12. Tai X.-S., Zhang L.-L., Liu L.-L., Cao S.-H., Wang L.-H. The crystal structure of bis(ethanol- κO)-bis(6-aminopicolinato- κ^2N,O) manganese(II), $C_{16}H_{22}O_6N_4Mn$. *Z. Kristallogr. N. Cryst. Struct.* 2021, 236, 551–552.