

Research Article

Arash Pashrashid, Cesar A. Gómez S., Seyed M. Mirhosseini-Alizamini,
Seyed Navid Motevalian, M. Daher Albalwi, Hijaz Ahmad, and Shao-Wen Yao*

On traveling wave solutions to Manakov model with variable coefficients

<https://doi.org/10.1515/phys-2022-0235>

received November 20, 2021; accepted February 20, 2023

Abstract: We use a general transformation, to find exact solutions for the Manakov system with variable coefficients (depending on the time ε) using an improved tanh-coth method. The solutions obtained in this work are more general compared to those in other works because they involve variable coefficients. The implemented computational method is applied in a direct way on the reduced system, avoid in this way the reduction to only one equation, as occurs in the works respect to exact solutions, made by other authors. Clearly, from the solutions obtained here, new solutions are derived for the standard model (constant coefficients), complementing in this way the results obtained by other authors mentioned here. Finally, we give some discussion on the results and give the respective conclusions.

Keywords: traveling wave solutions, Manakov equation, improved tanh-coth method

1 Introduction

The soliton theory is actually one of the most important branch of the applied mathematics. In the physical sciences, especially in nonlinear optic and from some year ago, in the communications theory, the study of optical solitons is the great relevance. In recent years, several models have been used to modelling optical solitons, and the nonlinear Schrödinger equation (NLSE) is one of the most important; however, other models such as the Chen-Lee-Liu equation, the Gerdjikov-Ivanov equation, the Ginzburg-Landau equation, the Manakov system, and several other models have been used to generate optical solitons to be applied in process of communication. Our interest in this work is the study, from of point of view of it exact solutions, of the following Manakov system with variable coefficients, given in the following form:

$$\begin{cases} iq_\varepsilon + \Gamma_1(\varepsilon)q_{xx} + \Theta_1(\varepsilon)(|q|^2 + |r|^2)q = 0 \\ ir_\varepsilon + \Gamma_2(\varepsilon)r_{xx} + \Theta_2(\varepsilon)(|q|^2 + |r|^2)r = 0, \end{cases} \quad (1.1)$$

where $q = q(x, \varepsilon)$, $r = r(x, \varepsilon)$ represent a complex-value function and are depending on the spacial variable x and the temporal variable ε . The coefficients Γ_1 , Γ_2 , Θ_1 , Θ_2 are functions depending on ε . In the case of constant coefficients, the model (1.1) is known as a system of coupled NLSEs, which are considered as important models in nonlinear optic [1–3]. In this last case, it has been found that the Manakov system is integrable *via* the inverse scattering method [1]; furthermore, this is very important in the design of new technology that require the use of optical pulses [3]. The model presented here, clearly includes the constant coefficients; therefore, model (1.1) is a generalization of the classical Manakov equation presented in references [1–3] and in many others works on the classical Manakov system [4–6]. On the other hand, differential equations with variable coefficients (depending on variable ε) are generalized models that include the standard models (constant coefficients). From the mathematical point of view, the study of that models has a great relevance because it includes a variety of considerations that

* **Corresponding author: Shao-Wen Yao**, School of Mathematics and Information Science, Henan Polytechnic University, Jiaozuo 454000, China, e-mail: yaoshaowen@hpu.edu.cn

Arash Pashrashid: Department of Computer Engineering, Sharif University of Technology, Tehran, Iran

Cesar A. Gómez S.: Department of Mathematics, Universidad Nacional de Colombia, Bogotá, Colombia

Seyed M. Mirhosseini-Alizamini: Department of Mathematics, Payame Noor University (PNU), Tehran, Iran

Seyed Navid Motevalian: Faculty of Civil and Environmental Engineering, Shahid Beheshti University, Tehran, Iran

M. Daher Albalwi: Yanbu Industrial College, The Royal Commission for Jubail and Yanbu, 30436, Saudi Arabia

Hijaz Ahmad: Operational Research Center in Healthcare, Near East University, Near East Boulevard, PC: 99138 Nicosia/Mersin 10, Turkey; Section of Mathematics, International Telematic University Uninettuno, Corso Vittorio Emanuele II, 39, 00186 Roma, Italy, e-mail: hijaz555@gmail.com

can be important in the phenomena described by the model. Furthermore, from the solutions of the generalized model, we can derive solutions for the standard, with the advantage that we can consider new types of solitons that can be used to understand the phenomena described by the respective model and can be used to implement new technology. In recent years, models with variable coefficients have been presented by several authors as equations with practical applications, for instance, the following equation:

$$u_\varepsilon + k_1 \varepsilon^n u^2 + k_2 \varepsilon^m u_{xxx} = 0,$$

studied in [7], and other equations considered for instance in [8,9] are types of this class of models.

When new equations are appearing to model the pulses transmission, a great variety of computational method have been implemented to solve it. We can found several models and computational methods for solving nonlinear partial differential equation in references [10–37]. Recently, nonlinear partial differential equations with fractional exponents have used for modeling several phenomena of the nature, so that, as in the classical models mentioned in the previous references, new techniques to solve them, specially from the point of view of exact and numerical solutions, have implemented [38–48]. However, one of the methods used in a satisfactory way and that did not appear in the aforementioned references is the improved tanh-coth method (ITCM) presented in refs [49,50], which we will use to solve (1.1). The advantage of this method, compared with others, is that it can be implemented in a mathematical software as Mathematica and Maples, can be applied into systems directly, do not require a special computer, and can be implemented easily, and finally, it is a generalization of the tanh-coth method [51] and the Kudryashov method [52] and is used widely in the literature. The work is organized as follows: In Section 2, we present a brief description of the ITCM to solve a system. In Section 3, we solve (1.1), and we obtain conditions on the coefficients with the aim to derive non-trivial solutions and we present the solutions obtained. Finally, we present a discussion on the results obtained, and we present some final conclusions.

2 Solving Eq. (1.1)

With the aim to solve the Manakov system given by Eq. (1.1), we need, at first, to use the following transformation:

$$\begin{cases} q(x, \varepsilon) = q(\eta) e^{i(x + \int \rho(\varepsilon) d\varepsilon + \eta_0)}, \\ r(x, \varepsilon) = r(\eta) e^{i(x + \int \rho(\varepsilon) d\varepsilon + \eta_0)}, \\ \eta = x + \int \lambda(\varepsilon) d\varepsilon + \eta_1, \end{cases} \quad (2.1)$$

where, $\rho(\varepsilon)$ and $\lambda(\varepsilon)$ are functions of ε to be determined later, and η_0, η_1 arbitrary constants. With the use of (2.1), we have the following system of ordinary differential equations in the unknowns $q(\eta), r(\eta)$ (where by simplicity, we have used the same variables q, r):

$$\begin{cases} -\rho(\varepsilon)q(\eta) - \Gamma_1(\varepsilon)q(\eta) + \Gamma_1(\varepsilon)q''(\eta) \\ \quad + \Theta_1(\varepsilon)(q^2(\eta) + r^2(\eta))u(\eta) \\ \quad + i(\lambda(\varepsilon) + 2\Gamma_1(\varepsilon))q'(\eta) = 0, \\ -\rho(\varepsilon)r(\eta) - \Gamma_2(\varepsilon)r(\eta) + \Gamma_2(\varepsilon)r''(\eta) \\ \quad + \Theta_2(\varepsilon)(q^2(\eta) + r^2(\eta))r(\eta) + i(\lambda(\varepsilon) + 2\Gamma_2(\varepsilon))r'(\eta) \\ \quad = 0. \end{cases} \quad (2.2)$$

Here, $q'(\varepsilon) = \frac{dq}{d\varepsilon}$, $r'(\varepsilon) = \frac{dr}{d\varepsilon}$. As we mentioned previously, one of the functions that we need to obtain is $\lambda(\varepsilon)$, so that, in the first equation of (2.2), we consider:

$$\lambda(\varepsilon) = -2\Gamma_1(\varepsilon). \quad (2.3)$$

With this selection, the imaginary part of the first equation of (2.2) is eliminated. Now, replacing (2.3), in the second equation of (2.2) and considering the following condition on system (1.1)

$$\Gamma_1(\varepsilon) = \Gamma_2(\varepsilon), \quad (2.4)$$

we can eliminate the imaginary part of the second equation of (2.2), so that, finally, (2.2) converts to

$$\begin{cases} -\rho(\varepsilon)q(\eta) - \Gamma_1(\varepsilon)q(\eta) + \Gamma_1(\varepsilon)q''(\eta) \\ \quad + \Theta_1(\varepsilon)(q^2(\eta) + r^2(\eta))q(\eta) = 0, \\ -\rho(\varepsilon)r(\eta) - \Gamma_1(\varepsilon)r(\eta) + \Gamma_1(\varepsilon)r''(\eta) \\ \quad + \Theta_2(\varepsilon)(q^2(\eta) + r^2(\eta))r(\eta) = 0. \end{cases} \quad (2.5)$$

2.1 The ITCM method

A full description of the ITCM can be found in reference [49]. However, similar to all computational methods, the first step is transformation of the nonlinear partial differential equation (or system of nonlinear partial differential equations) to an ordinary differential equation (or system of ordinary differential equations), by means of adequate transformation. In our case, we have used (2.1) to reduce (1.1) into (2.2). In the following step of the ITCM method,

we illustrate them by applying directly to our system. We consider solutions of (2.5) in the following form:

$$\begin{cases} q(\eta) = \sum_{i=0}^M a_i(\varepsilon)\phi(\eta)^i + \sum_{i=M+1}^{2M} a_i(\varepsilon)\phi(\eta)^{M-i}, \\ r(\eta) = \sum_{i=0}^N b_i(\varepsilon)\phi(\eta)^i + \sum_{i=N+1}^{2N} b_i(\varepsilon)\phi(\eta)^{N-i}, \end{cases} \quad (2.6)$$

where $\phi(\eta)$ is solution of the Riccati equation [50]:

$$\phi'(\eta) = \sigma(\varepsilon)\phi^2(\eta) + \varrho(\varepsilon)\phi(\eta) + \chi(\varepsilon). \quad (2.7)$$

By substituting (2.6) into (2.5) and applying the balancing technique, we have $M + 2 = 3M$ and $N + 2 = 3N$, respectively, thus obtaining $M = 1$, $N = 1$. With this values, (2.6) reduces to

$$\begin{cases} q(\eta) = a_0(\varepsilon) + a_1(\varepsilon)\phi(\eta) + a_2(\varepsilon)\phi(\eta)^{-1}, \\ r(\eta) = b_0(\varepsilon) + b_1(\varepsilon)\phi(\eta) + b_2(\varepsilon)\phi(\eta)^{-1}. \end{cases} \quad (2.8)$$

By substituting (2.8) into (2.5) and considering (2.7), we have an algebraic system in the unknowns $\chi(\varepsilon)$, $\varrho(\varepsilon)$, $\sigma(\varepsilon)$, $\rho(\varepsilon)$, $a_0(\varepsilon)$, $a_1(\varepsilon)$, $a_2(\varepsilon)$, $b_0(\varepsilon)$, $b_1(\varepsilon)$, $b_2(\varepsilon)$:

$$\begin{cases} 3a_1\Gamma_1(\varepsilon)\varrho(\varepsilon)\sigma(\varepsilon) + 2a_1b_0b_1\Theta_1(\varepsilon) + a_0b_1^2\Theta_1(\varepsilon) + 3a_0a_1^2\Theta_1(\varepsilon) = 0, \\ 2a_1\Gamma_1(\varepsilon)\sigma^2(\varepsilon) + a_1b_1^2\Theta_1(\varepsilon) + a_1^3\Theta_1(\varepsilon) = 0, \\ a_1^2b_0\Theta_2(\varepsilon) + 2a_0a_1b_1\Theta_2(\varepsilon) + 3\Gamma_1(\varepsilon)\varrho(\varepsilon)b_1\sigma(\varepsilon) + 3b_0b_1^2\Theta_2(\varepsilon) = 0, \\ a_1^2b_1\Theta_2(\varepsilon) + 2\Gamma_1(\varepsilon)b_1\sigma^2(\varepsilon) + b_1^3\Theta_2(\varepsilon) = 0, \\ 3\chi(\varepsilon)a_2\Gamma_1(\varepsilon)\varrho(\varepsilon) + 2a_2b_0b_2\Theta_1(\varepsilon) + a_0b_2^2\Theta_1(\varepsilon) + 3a_0a_2^2\Theta_1(\varepsilon) = 0, \\ 2\chi^2(\varepsilon)a_2\Gamma_1(\varepsilon) + a_2b_2^2\Theta_1(\varepsilon) + a_2^3\Theta_1(\varepsilon) = 0, \\ a_2^2b_0\Theta_2(\varepsilon) + 2a_0a_2b_2\Theta_2(\varepsilon) + 3\chi(\varepsilon)\Gamma_1(\varepsilon)\varrho(\varepsilon)b_2 + 3b_0b_2^2\Theta_2(\varepsilon) = 0, \\ a_2^2b_2\Theta_2(\varepsilon) + 2\chi(\varepsilon)^2\Gamma_1(\varepsilon)b_2 + b_2^3\Theta_2(\varepsilon) = 0, \\ \chi(\varepsilon)a_1\Gamma_1(\varepsilon)\varrho(\varepsilon) + a_2\Gamma_1(\varepsilon)\varrho(\varepsilon)\sigma(\varepsilon) - a_0\Gamma_1(\varepsilon) + a_0b_0^2\Theta_1(\varepsilon) + 2a_0b_1b_2\Theta_1(\varepsilon) \\ + 2a_2b_0b_1\Theta_1(\varepsilon) + 2a_1b_0b_2\Theta_1(\varepsilon) + a_0^3\Theta_1(\varepsilon) + 6a_1a_2a_0\Theta_1(\varepsilon) - a_0\rho(\varepsilon) = 0, \\ 2\chi(\varepsilon)a_1\Gamma_1(\varepsilon)\sigma(\varepsilon) + a_1\Gamma_1(\varepsilon)\varrho^2(\varepsilon) - a_1\Gamma_1(\varepsilon) + a_1b_0^2\Theta_1(\varepsilon) + a_2b_1^2\Theta_1(\varepsilon) \\ + 2a_0b_0b_1\Theta_1(\varepsilon) + 2a_1b_1b_2\Theta_1(\varepsilon) + 3a_0^2a_1\Theta_1(\varepsilon) + 3a_1^2a_2\Theta_1(\varepsilon) - a_1\rho(\varepsilon) = 0, \\ 2\chi(\varepsilon)a_2\Gamma_1(\varepsilon)\sigma(\varepsilon) + a_2\Gamma_1(\varepsilon)\varrho^2(\varepsilon) - a_2\Gamma_1(\varepsilon) + a_2b_0^2\Theta_1(\varepsilon) + a_1b_2^2\Theta_1(\varepsilon) \\ + 2a_0b_0b_2\Theta_1(\varepsilon) + 2a_2b_1b_2\Theta_1(\varepsilon) + 3a_1a_2^2\Theta_1(\varepsilon) + 3a_2^2a_2\Theta_1(\varepsilon) - a_2\rho(\varepsilon) = 0, \\ a_0^2b_0\Theta_2(\varepsilon) + 2a_1a_2b_0\Theta_2(\varepsilon) + 2a_0a_2b_1\Theta_2(\varepsilon) + 2a_0a_1b_2\Theta_2(\varepsilon) \\ + \chi(\varepsilon)\Gamma_1(\varepsilon)\varrho(\varepsilon)b_1 + \Gamma_1(\varepsilon)\varrho(\varepsilon)b_2\sigma(\varepsilon) - \Gamma_1(\varepsilon)b_0 + b_0^3\Theta_2(\varepsilon) + 6b_1b_2b_0\Theta_2(\varepsilon) - b_0\rho(\varepsilon) = 0, \\ 2a_0a_1b_0\Theta_2(\varepsilon) + a_0^2b_1\Theta_2(\varepsilon) + 2a_1a_2b_1\Theta_2(\varepsilon) + a_1^2b_2\Theta_2(\varepsilon) \\ + 2\chi(\varepsilon)\Gamma_1(\varepsilon)b_1\sigma(\varepsilon) + \Gamma_1(\varepsilon)b_1\varrho^2(\varepsilon) - \Gamma_1(\varepsilon)b_1 + 3b_0^2b_1\Theta_2(\varepsilon) + 3b_1^2b_2\Theta_2(\varepsilon) - b_1\rho(\varepsilon) = 0, \\ 2a_0a_2b_0\Theta_2(\varepsilon) + a_2^2b_1\Theta_2(\varepsilon) + a_0^2b_2\Theta_2(\varepsilon) + 2a_1a_2b_2\Theta_2(\varepsilon) \\ + 2\chi(\varepsilon)\Gamma_1(\varepsilon)b_2\sigma(\varepsilon) + \Gamma_1(\varepsilon)b_2\varrho^2(\varepsilon) - \Gamma_1(\varepsilon)b_2 + 3b_1b_2^2\Theta_2(\varepsilon) + 3b_2^2b_2\Theta_2(\varepsilon) - b_2\rho(\varepsilon) = 0. \end{cases} \quad (2.9)$$

$\Gamma_1(\varepsilon)$, $\Theta_1(\varepsilon)$, and $\Theta_2(\varepsilon)$ are the coefficients of Eq. (1.1). The rest of parameters in the previous system depend on variable ε ($a_1 = a_1(\varepsilon), \dots$). By using a mathematical software such as Maple or Mathematica, we obtain the following solution of the system (2.9), from which, we can obtain nontrivial solutions for (1.1):

$$\begin{cases} a_0(\varepsilon) = b_0(\varepsilon) = \chi(\varepsilon) = \sigma(\varepsilon) = 0, \\ a_1(\varepsilon) = \pm ib_1(\varepsilon), \\ a_2(\varepsilon) = \pm ib_2(\varepsilon), \\ \rho(\varepsilon) = \Gamma(\varepsilon)(-1 + \varrho^2(\varepsilon)). \end{cases} \quad (2.10)$$

With these values, the solution of (2.7) is expressed as follows:

$$\phi(\eta) = e^{\varrho(\varepsilon)\eta}. \quad (2.11)$$

Here $\varrho(\varepsilon)$ is the arbitrary function. According to (2.8), and (2.10), we have


$$\begin{cases} q(\eta) = \pm ib_1(\varepsilon)\phi(\eta) \pm ib_2(\varepsilon)\phi(\eta)^{-1}, \\ r(\eta) = b_1(\varepsilon)\phi(\eta) + b_2(\varepsilon)\phi(\eta)^{-1}, \end{cases} \quad (2.12)$$

where $\phi(\eta)$ is earlier given in (2.11) and $b_1(\varepsilon)$, $b_2(\varepsilon)$ arbitrary functions. Finally, taking into account (2.1), (2.3),

and (2.12), under condition (2.4), we have the following solution of system (1.1):

$$\begin{cases} q(x, \varepsilon) = \pm i r(\eta) e^{i \left(x + \int \Gamma_1(\varepsilon) (-1 + \varrho^2(\varepsilon)) dt + \eta_0 \right)} \\ r(x, \varepsilon) = r(\eta) e^{i \left(x + \int \Gamma_1(\varepsilon) (-1 + \varrho^2(\varepsilon)) dt + \eta_0 \right)} \\ \eta = x + \int -2\Gamma_1(\varepsilon) dt + \eta_1, \end{cases} \quad (2.13)$$


where $\varrho(\varepsilon)$ is arbitrary function, η_0 and η_1 arbitrary constants, and $r(\eta)$ is given in (2.12).

(a) r_c

$$\Theta_1(\varepsilon) = \Theta_2(\varepsilon). \quad (2.14)$$

With (2.14), the system (2.5) reduces to

$$\begin{cases} -\rho(\varepsilon)q(\eta) - \Gamma_1(\varepsilon)q(\eta) + \Gamma_1(\varepsilon)q''(\eta) \\ \quad + \Theta_1(\varepsilon)(q^2(\eta) + r^2(\eta))q(\eta) = 0, \\ -\rho(\varepsilon)r(\eta) - \Gamma_1(\varepsilon)r(\eta) + \Gamma_1(\varepsilon)r''(\eta) \\ \quad + \Theta_1(\varepsilon)(q^2(\eta) + r^2(\eta))r(\eta) = 0. \end{cases} \quad (2.15)$$

(b) r_v

r_c corresponds to graph of $r(\eta)$ given in (2.12), for the following constant values: $b_1(\varepsilon) = 2$, $b_2(\varepsilon) = 3$, $\varrho(\varepsilon) = 4$, and $\Gamma_1(\varepsilon) = 2$, for $[x, t] \in [0.6, 1] \times [0.1, 0.3]$ and $\eta_1 = 0$.

r_v corresponds to graph of $r(\eta)$ given in (2.12), for the following variable values: $b_1(\varepsilon) = 2t$, $b_2(\varepsilon) = 3t^2$, $\varrho(\varepsilon) = 4t^3$, and $\Gamma_1(\varepsilon) = 2t^2$, for $[x, t] \in [0.6, 5] \times [0.1, 1]$ and $\eta_1 = 0$.

2.2 A second case

In Section 2.1, we have obtained solutions for system (1.1), for the case given by (2.4). In this section, we obtain new solutions, taking into account the following additionally condition:

Under these new conditions, the following are new solutions of the system (2.9):

$$\begin{cases} \rho(\varepsilon) = \pm \frac{2\Theta_1(\varepsilon)a_2^2(\varepsilon)b_1(\varepsilon) - \Gamma_1(\varepsilon)b_2(\varepsilon) + 2\Theta_1(\varepsilon)b_1(\varepsilon)b_2^2(\varepsilon)}{b_2(\varepsilon)}, \\ \chi(\varepsilon) = \pm \frac{i\sqrt{\Theta_1(\varepsilon)}\sqrt{a_2^2(\varepsilon) + b_2^2(\varepsilon)}}{\sqrt{2\Gamma_1(\varepsilon)}}, \\ \sigma(\varepsilon) = \pm \frac{i\sqrt{\Theta_1(\varepsilon)}b_1(\varepsilon)\sqrt{a_2^2(\varepsilon) + b_2^2(\varepsilon)}}{\sqrt{2\Gamma_1(\varepsilon)}b_2(\varepsilon)}, \\ a_1(\varepsilon) = \frac{a_2(\varepsilon)b_1(\varepsilon)}{b_2(\varepsilon)}, \\ \varrho(\varepsilon) = a_0(\varepsilon) = b_0(\varepsilon) = 0, \end{cases} \quad (2.16)$$

$a_2(\varepsilon)$, $b_1(\varepsilon)$, and $b_2(\varepsilon)$ arbitrary functions (depending on variable ε).

$$\begin{cases}
\rho(\varepsilon) = \frac{4\Theta_1(\varepsilon)a_2^2(\varepsilon)b_1(\varepsilon) - \Gamma_1(\varepsilon)b_2(\varepsilon) + 4\Theta_1(\varepsilon)b_1(\varepsilon)b_2^2(\varepsilon)}{b_2(\varepsilon)}, & b_1(\varepsilon) \text{ and } b_2(\varepsilon) \text{ arbitrary functions (depending on variable } \varepsilon\text{).} \\
\chi(\varepsilon) = \pm \frac{i\sqrt{\Theta_1}\sqrt{a_2^2(\varepsilon) + b_2^2(\varepsilon)}}{\sqrt{2\Gamma_1(\varepsilon)}}, \\
\sigma(\varepsilon) = \mp \frac{i\sqrt{\Theta_1(\varepsilon)}b_1(\varepsilon)\sqrt{a_2^2(\varepsilon) + b_2^2(\varepsilon)}}{\sqrt{2\Gamma_1(\varepsilon)}b_2(\varepsilon)}, \\
a_1(\varepsilon) = \frac{a_2(\varepsilon)b_1(\varepsilon)}{b_2(\varepsilon)}, \\
\rho(\varepsilon) = a_0(\varepsilon) = b_0(\varepsilon) = 0,
\end{cases} \quad (2.17)$$

$a_2(\varepsilon)$, $b_1(\varepsilon)$, and $b_2(\varepsilon)$ arbitrary functions (depending on variable ε).

$$\begin{cases}
\rho(\varepsilon) = \Gamma(\varepsilon)(-1 + \varrho^2(\varepsilon)), \\
a_1(\varepsilon) = \pm ib_1(\varepsilon), \\
a_2(\varepsilon) = \pm ib_2(\varepsilon), \chi(\varepsilon) = \sigma(\varepsilon) = a_0(\varepsilon) = b_0(\varepsilon) \\
= 0,
\end{cases} \quad (2.18)$$

$$\begin{cases}
\rho(\varepsilon) = -\Gamma(\varepsilon) + \Gamma(\varepsilon)\varrho^2(\varepsilon) + 4\Theta_1(\varepsilon)b_1(\varepsilon)b_2(\varepsilon), \\
a_1(\varepsilon) = ib_1(\varepsilon), \\
a_2(\varepsilon) = -ib_2(\varepsilon), \chi(\varepsilon) = \sigma(\varepsilon) = a_0(\varepsilon) = b_0(\varepsilon) \\
= 0,
\end{cases} \quad (2.19)$$

$b_1(\varepsilon)$ and $b_2(\varepsilon)$ arbitrary functions (depending on variable ε).

We can obtain several expressions, which include periodic and hyperbolic functions, for solutions of (2.15), using the following classifications of solutions of Eq. (2.7) [50] and the previous set of solutions of (2.9), given from (2.16) until (2.19):

(1) If $\chi(\varepsilon) \neq 0$, $\sigma(\varepsilon) \neq 0$, and $\varrho(\varepsilon) \neq 0$:

$$\begin{cases}
\frac{1}{\sigma(\varepsilon)} \left(-\frac{1}{\eta} - \frac{\varrho(\varepsilon)}{2} \right), & \varrho^2(\varepsilon) = 4\sigma(\varepsilon)\chi(\varepsilon) \\
\frac{1}{\sigma(\varepsilon)} \left(\frac{\sqrt{4\chi(\varepsilon)\sigma(\varepsilon) - \varrho^2(\varepsilon)}}{2} \tan \left[\frac{\sqrt{4\chi(\varepsilon)\sigma(\varepsilon) - \varrho^2(\varepsilon)}}{2} \eta \right] - \frac{\varrho(\varepsilon)}{2} \right), & 4\chi(\varepsilon)\sigma(\varepsilon) - \varrho(\varepsilon)^2 > 0 \\
\frac{1}{\sigma(\varepsilon)} \left(-\frac{\sqrt{4\chi(\varepsilon)\sigma(\varepsilon) - \varrho^2(\varepsilon)}}{2} \cot \left[\frac{\sqrt{4\chi(\varepsilon)\sigma(\varepsilon) - \varrho^2(\varepsilon)}}{2} \eta \right] - \frac{\varrho(\varepsilon)}{2} \right), & 4\chi(\varepsilon)\sigma(\varepsilon) - \varrho^2(\varepsilon) > 0 \\
\frac{1}{\sigma(\varepsilon)} \left(-\frac{\sqrt{\varrho^2(\varepsilon) - 4\chi(\varepsilon)\sigma(\varepsilon)}}{2} \tanh \left[\frac{\sqrt{\varrho^2(\varepsilon) - 4\chi(\varepsilon)\sigma(\varepsilon)}}{2} \eta \right] - \frac{\varrho(\varepsilon)}{2} \right), & 4\chi(\varepsilon)\sigma(\varepsilon) - \varrho^2(\varepsilon) < 0 \\
\frac{1}{\sigma(\varepsilon)} \left(-\frac{\sqrt{\varrho^2(\varepsilon) - 4\chi(\varepsilon)\sigma(\varepsilon)}}{2} \coth \left[\frac{\sqrt{\varrho^2(\varepsilon) - 4\chi(\varepsilon)\sigma(\varepsilon)}}{2} \eta \right] - \frac{\varrho(\varepsilon)}{2} \right), & 4\chi(\varepsilon)\sigma(\varepsilon) - \varrho^2(\varepsilon) < 0
\end{cases} \quad (2.20)$$

(2) If $\chi(\varepsilon) = 0$, $\sigma(\varepsilon) \neq 0$:

$$\begin{cases}
\frac{\varrho(\varepsilon)}{-\sigma(\varepsilon) + \varrho(\varepsilon)e^{-\varrho(\varepsilon)\eta}}, & \varrho(\varepsilon) \neq 0 \\
-\frac{1}{\sigma(\varepsilon)\eta}, & \varrho(\varepsilon) = 0 \\
\frac{1}{\sigma(\varepsilon)} \left(\frac{\sqrt{-\varrho^2(\varepsilon)}}{2} \tan \left[\frac{\sqrt{-\varrho^2(\varepsilon)}}{2} \eta \right] - \frac{\varrho(\varepsilon)}{2} \right), & \varrho^2(\varepsilon) < 0 \\
\frac{1}{\sigma(\varepsilon)} \left(-\frac{\sqrt{-\varrho^2(\varepsilon)}}{2} \cot \left[\frac{\sqrt{-\varrho^2(\varepsilon)}}{2} \eta \right] - \frac{\varrho(\varepsilon)}{2} \right), & \varrho^2(\varepsilon) < 0 \\
\frac{1}{\sigma(\varepsilon)} \left(-\frac{\varrho(\varepsilon)}{2} \tanh \left[\frac{\varrho(\varepsilon)}{2} \eta \right] - \frac{\varrho(\varepsilon)}{2} \right), & \varrho^2(\varepsilon) > 0 \\
\frac{1}{\sigma(\varepsilon)} \left(-\frac{\varrho(\varepsilon)}{2} \coth \left[\frac{\varrho(\varepsilon)}{2} \eta \right] - \frac{\varrho(\varepsilon)}{2} \right), & \varrho^2(\varepsilon) > 0
\end{cases} \quad (2.21)$$

(3) If $\varrho(\varepsilon) = 0$, $\sigma(\varepsilon) \neq 0$

$$\phi(\eta) = \begin{cases} -\frac{1}{\sigma(\varepsilon)\eta}, & \chi(\varepsilon) = 0 \\ \frac{1}{\sigma(\varepsilon)}(\sqrt{\chi(\varepsilon)\sigma(\varepsilon)} \tan[\sqrt{\chi(\varepsilon)\sigma(\varepsilon)}\eta]), & \chi(\varepsilon)\sigma(\varepsilon) > 0 \\ \frac{1}{\sigma(\varepsilon)}(-\sqrt{\chi(\varepsilon)\sigma(\varepsilon)} \cot[\sqrt{\chi(\varepsilon)\sigma(\varepsilon)}\eta]), & \chi(\varepsilon)\sigma(\varepsilon) > 0 \\ \frac{1}{\sigma(\varepsilon)}(-\sqrt{-\chi(\varepsilon)\sigma(\varepsilon)} \tanh[\sqrt{-\chi(\varepsilon)\sigma(\varepsilon)}\eta]), & \chi(\varepsilon)\sigma(\varepsilon) < 0 \\ \frac{1}{\sigma(\varepsilon)}(-\sqrt{-\chi(\varepsilon)\sigma(\varepsilon)} \coth[\sqrt{-\chi(\varepsilon)\sigma(\varepsilon)}\eta]), & \chi(\varepsilon)\sigma(\varepsilon) < 0. \end{cases} \quad (2.22)$$

(4) If $\sigma(\varepsilon) = 0$, $\varrho(\varepsilon) \neq 0$

$$\phi(\eta) = \frac{-\chi(\varepsilon) + \varrho(\varepsilon)e^{\varrho(\varepsilon)\eta}}{\varrho(\varepsilon)}. \quad (2.23)$$

We illustrate solutions of (2.7), using the values given by (2.16) and (2.22) (taking the first sign in each case): First, we note that $(\chi(\varepsilon))(\sigma(\varepsilon)) = -\frac{b_1(\varepsilon)\Theta_1(\varepsilon)(a_2(\varepsilon)^2 + b_2(\varepsilon)^2)}{2\Gamma_1(\varepsilon)b_2(\varepsilon)}$, so that, after simplifications, we have:

where $\rho(\varepsilon) = \frac{2\Theta_1(\varepsilon)a_2^2(\varepsilon)b_1(\varepsilon) - \Gamma_1(\varepsilon)b_2(\varepsilon) + 2\Theta_1(\varepsilon)b_1(\varepsilon)b_2^2(\varepsilon)}{b_2(\varepsilon)}$, and $\eta = x + \int -2\Gamma_1(\varepsilon)d\varepsilon + \eta_1$, and $a_2(\varepsilon)$, $b_1(\varepsilon)$, $b_2(\varepsilon)$ are arbitrary functions depending on the variable ε , and $\phi(\eta)$ is given by (2.24). In the previous expressions,



$$\begin{cases} q(\eta) = \frac{a_2(\varepsilon)b_1(\varepsilon)}{b_2(\varepsilon)}\phi(\eta) + a_2(\varepsilon)\phi(\eta)^{-1} \\ r(\eta) = b_1(\varepsilon)\phi(\eta) + b_2(\varepsilon)\phi(\eta)^{-1} \end{cases} \quad (2.26)$$

$$\phi(\eta) = \begin{cases} -\frac{\sqrt{2\Gamma_1(\varepsilon)b_2(\varepsilon)}}{(\sqrt{-\Theta_1(\varepsilon)b_1(\varepsilon)\sqrt{a_2^2(\varepsilon) + b_2^2(\varepsilon)}})\eta}, & \text{if } \chi(\varepsilon) = 0, \\ \sqrt{\frac{b_2(\varepsilon)}{b_1(\varepsilon)}} \tan\left[\sqrt{\frac{b_1(\varepsilon)\Theta_1(\varepsilon)(a_2(\varepsilon)^2 + b_2(\varepsilon)^2)}{2\Gamma_1(\varepsilon)b_2(\varepsilon)}}\eta\right] & \text{if } \chi(\varepsilon)\sigma(\varepsilon) > 0 \\ -\sqrt{\frac{b_2(\varepsilon)}{b_1(\varepsilon)}} \cot\left[\sqrt{\frac{b_1(\varepsilon)\Theta_1(\varepsilon)(a_2(\varepsilon)^2 + b_2(\varepsilon)^2)}{2\Gamma_1(\varepsilon)b_2(\varepsilon)}}\eta\right] & \text{if } \chi(\varepsilon)\sigma(\varepsilon) > 0 \\ -\sqrt{\frac{b_2(\varepsilon)}{b_1(\varepsilon)}} \tanh\left[\sqrt{\frac{b_1(\varepsilon)\Theta_1(\varepsilon)(a_2(\varepsilon)^2 + b_2(\varepsilon)^2)}{2\Gamma_1(\varepsilon)b_2(\varepsilon)}}\eta\right] & \text{if } \chi(\varepsilon)\sigma(\varepsilon) < 0 \\ -\sqrt{\frac{b_2(\varepsilon)}{b_1(\varepsilon)}} \coth\left[\sqrt{\frac{b_1(\varepsilon)\Theta_1(\varepsilon)(a_2(\varepsilon)^2 + b_2(\varepsilon)^2)}{2\Gamma_1(\varepsilon)b_2(\varepsilon)}}\eta\right] & \text{if } \chi(\varepsilon)\sigma(\varepsilon) < 0 \end{cases} \quad (2.24)$$



Therefore, according to (2.1), (2.8), and (2.16), we have the following solution for (1.1) (under two conditions $\Gamma_1(\varepsilon) = \Gamma_2(\varepsilon)$ and $\Theta_1(\varepsilon) = \Theta_2(\varepsilon)$):

$$\begin{cases} q(x, \varepsilon) = \left(\frac{a_2(\varepsilon)b_1(\varepsilon)}{b_2(\varepsilon)}\phi(\eta) + a_2(\varepsilon)\phi(\eta)^{-1}\right)e^{l(x + \int \rho(\varepsilon)d\varepsilon + \eta_0)}, \\ r(x, \varepsilon) = (b_1(\varepsilon)\phi(\eta) + b_2(\varepsilon)\phi(\eta)^{-1})e^{l(x + \int \rho(\varepsilon)d\varepsilon + \eta_0)}, \end{cases} \quad (2.25)$$



are solutions of (2.8). By using the expression \tanh in (2.24), the following are graphics corresponding to (2.26), for adequate values of the respective variables:

(c) q_1 (d) r_1

q_1 and r_1 correspond to values: $\Gamma(\varepsilon) = 2$, $\Theta(\varepsilon) = -2$, $a_2(\varepsilon) = 1$, $b_1(\varepsilon) = -1$, $b_2(\varepsilon) = 1$, and $(x, \varepsilon) \in [0, 5] \times [0, 2]$.

(e) q_2 (f) r_2

q_2 and r_2 correspond to values of the parameters: $\Gamma(\varepsilon) = 2\varepsilon$, $\Theta(\varepsilon) = -2\varepsilon^2$, $a_2(\varepsilon) = 1$, $b_1(\varepsilon) = -1$, $b_2(\varepsilon) = 1$, and $(x, \varepsilon) \in [0, 5] \times [0, 2]$. Here, we have taken variable coefficients for Γ and Θ .

(g) q_3 (h) r_3

Finally, q_3 and r_3 are the graphs of (2.26) corresponding to values $\Gamma(\varepsilon) = 2\varepsilon$, $\Theta(\varepsilon) = -2\varepsilon^2$, $a_2(\varepsilon) = 1$, $b_1(\varepsilon) = -\varepsilon$, $b_2(\varepsilon) = \varepsilon^2$, and $(x, \varepsilon) \in [0, 5] \times [0, 2]$. We have taken variable coefficients for all free parameters.

For all previous graphics, we have used $\eta_1 = 0$.

3 Results and discussion

In this work, we have used the ITCM to obtain solutions for a system with variable coefficients, applying directly the method to the respective system (after reduction to ordinary equations), thus avoiding the reduction to only one equation, as presented in the previous study. The first observation deals with the effectiveness of the method. A second observation, all solutions obtained here, is in terms of variable coefficients, which is new in the literature for the model studied in this work. The model considered here is known as the Manakov system with variable coefficients (1.1). We have obtained solutions for it in two particular cases: In the first case, we have considered (2.4) to obtain the solution given by (2.13). As we have mentioned early, the solutions are new in the literature due to existence of variable coefficients; however, if we take the coefficients as constants, as a particular case, the solutions given by (2.13) are again new. The graphs showed by r_c and r_v illustrate the two cases, taking constant coefficients (in the case of r_c) where we have obtained a solitary wave with smooth perfil, and by

taking variable coefficients (in the case of r_v), we can obtain a new structure of the solution, where we can note the evolution of the previous wave to a wave with a truncate perfil. We think that this fact can be taken into account in a real application of the model. In a second case, we have assumed additionally the condition (2.14), and we have obtained solutions for (1.1). In this last case, the solutions obtained are new in the literature due to variable coefficients. However, if we take constant coefficients, the solutions are new, compared with those obtained in references [3–5]. The solutions obtained and expressed by (q_1, r_1) have the structure of a dark soliton, but in the case were we have used some constant coefficients combined with variable coefficient, the wave have a relevant change, and the new wave have a part smooth as r_c , r_v , with special evolutions. In the third case, where we have taken all variable coefficients, q_3 , v_3 , the waves have the structure similar to dark solitons; however in this case, it is clear that q_3 is different to r_3 . The method used here gives us solutions for (1.1), with a more general structure compared, for instance, with those obtained in ref. [3], which can be seen in the solutions obtained for (2.15), which include a sum of tanh and coth functions, using the solutions mentioned in Eq. (2.7). The results obtained in this study are complementary to those found by other researchers. These results contribute to the development of the solitons theory, especially since the use of variable coefficients has become more important recently.

4 Conclusion

We have obtained exact solutions for the Manakov system with variable coefficients (1.1), which, due to its variable coefficients, are new in the literature. The ITCM has been applied directly on the ordinary differential system (2.2), (2.15), for obtaining exact solutions of (1.1) in a satisfactory way. The case in which the system has constant coefficients can be derived as a particular case. New solutions for the classical Manakov model (constant coefficients) are derived from the solutions obtained here. From the graphs r_c and r_v , we can see that with the use of variable coefficients, we can have different structures of the solutions, compared with the solutions in the case of constants coefficients. We have found new solutions for the standard model by obtaining optical solutions for the model considered here under two conditions: firstly, when $\Gamma_1(\varepsilon) = \Gamma_2(\varepsilon)$, and secondly, when $\Theta_1(\varepsilon) = \Theta_2(\varepsilon)$. Comparing that solutions with those obtained by others authors [3–5], we can conclude that our results are complementary to those obtained in the mentioned previously mentioned references. By using (2.20), (2.21), (2.22), and (2.23), new type of solutions can be derived to complement our work, and to keep it short, we have omitted here.

Funding information: The Fundamental Research Funds for the Universities of Henan Province (No. NSFRF210314), and Innovative Research Team of Henan Polytechnic University (No. T2022-7).

Author contributions: All authors have accepted responsibility for the entire content of this manuscript and approved its submission.

Conflict of interest: The authors declare no conflict of interest.

References

- [1] Manakov SV. On the theory of two-dimensional stationary self-focusing of electromagnetic waves. *Zh Eksp Fiz.* 1973;65:505–16.
- [2] Takayuki T. N-soliton collision in the Manakov model. *Progr Theo Phys.* 2004;111(2):151–82.
- [3] Yildirim Y. Optical soliton molecules of Manakov model by modified simple equation technique. *Optik.* 2019;185:1182–8.
- [4] Priya NV, Senthivelan M, Lakshmanan M. Akhmediev breathers, Ma solitons, and general breathers from rogue waves: a case study in the Manakov system. *Phys Rev E.* 2013;88(2):022918.
- [5] Chen S, Mihalache D. Vector rogue waves in the Manakov system: diversity and compossibility. *J Phys A Math Theor.* 2015;48(21):215202.
- [6] Steiglitz K. Multistable collision cycles of Manakov spatial solitons. *Phys Rev E.* 2001;63(4):046607.
- [7] Nirmala N, Vedan MJ, Baby BV. Auto-Bäcklund transformation, Lax pairs, and Painlevé property of a variable coefficient Korteweg-de Vries equation. I. *J Math Phys.* 1986;27(11):2640–3.
- [8] Miura RM. Korteweg-de Vries equation and generalizations. I. A remarkable explicit nonlinear transformation. *J Math Phys.* 1968;9(8):1202–4.
- [9] Yang LC. The applications of bifurcation method to a higher-order KdV equation. *Math Anal Appl.* 2012;275:1–12.
- [10] Ghanbari B. Abundant exact solutions to a generalized nonlinear Schrödinger equation with local fractional derivative. *Math Meth Appl Sci.* 2021;44(11):8759–74.
- [11] Ghanbari B, Nisar KS, Aldhaifallah M. Abundant solitary wave solutions to an extended nonlinear Schrödinger equation with conformable derivative using an efficient integration method. *Adv Differ Equ.* 2020;2020(1):1–25.
- [12] Islam MT, Akbar MA, Ahmad H, Ilhan OA, Gepreel KA. Diverse and novel soliton structures of coupled nonlinear Schrödinger type equations through two competent techniques. *Modern Phys Lett B.* 2022 Apr 20;36(11):2250004.
- [13] Zahran EH, Bekir A, Ahmad H. New unexpected perceptions for the optical solitary wave solution to the cubic-order nonlinear Schrödinger equation. *Optical Quantum Electr.* 2022 Jun;54(6):1–6.
- [14] Zahran EH, Ahmad H, Askar S, Botmart T, Shehata MS. Dark-soliton behaviors arising from a coupled nonlinear Schrödinger system. *Results Phys.* 2022 May 1;36:105459. doi: 10.1016/j.rinp.2022.105459.
- [15] Rezazadeh H, Kumar D, Neirameh A, Eslami M, Mirzazadeh M. Applications of three methods for obtaining optical soliton solutions for the Lakshmanan-Porsezian-Daniel model with Kerr law nonlinearity. *Pramana.* 2020;94(1):1–11.
- [16] Abbagari S, Houwe A, Rezazadeh H, Bekir A, Bouetou TB, Crepin KT. Optical soliton to multi-core (coupling with all the neighbors) directional couplers and modulation instability. *European Phys J Plus.* 2021;136(3):1–19.
- [17] Islam M, Akbar M, Ahmad H. Diverse optical soliton solutions of the fractional coupled (2+1)-dimensional nonlinear Schrödinger equations. *Optical Quantum Electr.* 2022 Feb;54(2):1–23.
- [18] Vahidi J, Zabihi A, Rezazadeh H, Ansari R. New extended direct algebraic method for the resonant nonlinear Schrödinger equation with Kerr law nonlinearity. *Optik.* 2021;227:165936.
- [19] Rezazadeh H, Odabasi M, Tariq KU, Abazari R, Baskonus HM. On the conformable nonlinear Schrödinger equation with second order spatiotemporal and group velocity dispersion coefficients. *Chinese J Phys.* 2021;72:403–14.
- [20] Liu X, Ahsan M, Ahmad M, Nisar M, Liu X, Ahmad I, et al. Applications of Haar wavelet-finite difference hybrid method and its convergence for hyperbolic nonlinear Schrödinger equation with energy and mass conservation. *Energies.* 2021 Jan;14(23):7831.
- [21] Tariq H, Sadaf M, Akram G, Rezazadeh H, Baili J, Lv YP, et al. Computational study for the conformable nonlinear

- Schrodinger equation with cubic-quintic-septic nonlinearities. *Results Phys.* 2021 Nov 1;30:104839. doi: 10.1016/j.rinp.2021.104839.
- [22] Jhangeer A, Rezazadeh H, Seadawy A. A study of travelling, periodic, quasiperiodic and chaotic structures of perturbed Fokas-Lenells model. *Pramana.* 2021;95(1):1–11.
- [23] Akinyemi L, Rezazadeh H, Shi QH, Inc M, Khater MM, Ahmad H, et al. New optical solitons of perturbed nonlinear Schrodinger-Hirota equation with spatio-temporal dispersion. *Results Phys.* 2021 Aug 23;29:104656. doi: 10.1016/j.rinp.2021.104656.
- [24] Baskonus HM, Gao W, Rezazadeh H, Mirhosseini-Alizamini SM, Baili J, Ahmad H, et al. New Classifications of nonlinear Schrodinger Model with group velocity dispersion via new extended method. *Results Phys.* 2021 Oct 14;31:104910. doi: 10.1016/j.rinp.2021.104910.
- [25] Darvishi MT, Najafi M, Wazwaz AM. Some optical soliton solutions of space-time conformable fractional Schrödinger-type models. *Phys Scr.* 2021;96(6):065213.
- [26] Ahmad H, Khan TA, Stanimirovic PS, Shatanawi W, Botmart T. New approach on conventional solutions to nonlinear partial differential equations describing physical phenomena. *Results Phys.* 2022;41:105936.
- [27] Sulaiman TA, Yusuf A, Abdel-Khalek S, Bayram M, Ahmad H. Nonautonomous complex wave solutions to the (2+1)-dimensional variable-coefficients nonlinear Chiral Schrodinger equation. *Results Phys.* 2020;19:103604. doi: 10.1016/j.rinp.2020.103604.
- [28] Rezazadeh H, Ullah N, Akinyemi L, Shah A, Mirhosseini SM, Chu YM, et al. Optical soliton solutions of the generalized non-autonomous nonlinear Schrodinger equations by the new Kudryashovas method. *Results Phys.* 2021;2021:104179.
- [29] Akinyemi L, Senol M, Rezazadeh H, Ahmad H, Wang H. Abundant optical soliton solutions for an integrable (2+1)-dimensional nonlinear conformable Schrodinger system. *Results Phys.* 2021;21:104177.
- [30] Rahman Z, Ali MZ, Ullah MS, Wen XY. Dynamical structures of interaction wave solutions for the two extended higher-order KdV equations. *Pramana.* 2021;95(3):1–14.
- [31] Roshid HO, Ali MZ. Asymptotic solutions of fifth order overdamped-oscillatory nonlinear systems. *Contemporary Math.* 2020;1(4):243–52.
- [32] Ullah MS, Ali MZ, Roshid HO, Hoque M. Collision phenomena among lump, periodic and stripe soliton solutions to a (2+1)-dimensional Benjamin-Bona-Mahony-Burgers Model. *European Phys J Plus.* 2021;136(4):1–9.
- [33] Arshed S, Raza N. Optical solitons perturbation of Fokas-Lenells equation with full nonlinearity and dual dispersion. *Chinese J Phys.* 2020;63:314–24.
- [34] Raza N, Osman MS, Abdel-Aty AH, Abdel-Khalek S, Besbes HR. Optical solitons of space-time fractional Fokas-Lenells equation with two versatile integration architectures. *Adv Differ Equ.* 2020;2020(1):1–15.
- [35] Afzal U, Raza N, Murtaza IG. On soliton solutions of time fractional form of Sawada-Kotera equation. *Nonlinear Dynamics.* 2019;95(1):391–405.
- [36] Yokus A, Durur H, Nofal TA, Abu-Zinadah H, Tuz M, Ahmad H. Study on the applications of two analytical methods for the construction of traveling wave solutions of the modified equal width equation. *Open Phys.* 2020;18(1):1003–10.
- [37] Yokus A. On the exact and numerical solutions to the FitzHugh-Nagumo equation. *Int J Modern Phys B.* 2020;34(17):2050149.
- [38] Yokus A, Durur H, Kaya D, Ahmad H, Nofal TA. Numerical comparison of Caputo and Conformable derivatives of time fractional Burgers-Fisher equation. *Results Phys.* 2021;25:104247.
- [39] Hammouch Z, Yavuz M, Osdemir N. Numerical results and synchronization of variable-order fractional chaotic system. *Modell Numer Simul Appl.* 2021;1(1):11–23.
- [40] Yussuf A, Abdulkadir T, Abdebanad T, Yavuz M, Sulaiman TA. The Schrödinger-KdV equation of fractional order with Mittag-Leffler nonsingular kernel. *Alexand Eng J.* 2021;60(2):2715–24.
- [41] Alvarado A, Cortes ER, Hernandez A, Martinez FJ. Bi-dimensional crime model based on anomalous diffusion with law enforcement effect. *Math Modell Numer Simul Appl.* 2022;2(1):26–40.
- [42] Veerasha P, Yavuz M, Baishya Ch. A computational approach for shallow water forced Korteweg de-Vries equation on critical flow over a hole with three fractional operators. *Int J Opt Contr Theor Appl (IJOCTA).* 2021;11(3):52–67.
- [43] Yokus A. Construction of differnts types of traveling wave solutions of the relativistic wave equation associated with the Schrödinger equation. *Math Modell Siml Appl.* 2021;1(1):24–31.
- [44] Yavuz M. European option pricing models described by fractional operators with classical and generalized Mittag-Leffler kernels. *Numer Meth Part Differ Equ.* 2022;38(3):434–56. doi: 10.1002/num.22645.
- [45] Yokus A, Yavuz M. Novel comparison of numerical and analytical methods for fractional Burger-Fisher equation. *Discrete Contin Dyn Sys-S.* 2021;14(7):2591–606.
- [46] Veerasha P. A numerical approach to the coupled atmospheric ocean model using a fractional operator. *Math Modell Numer Simul Appl.* 2021;1(1):1–10.
- [47] Yavuz M, Sene N. Approximate solutions of the model describing fluid flow using generalized p-Laplace transform method and heat balance integral method. *Axioms.* 2020;9(4):123.
- [48] Mohan JR, Chakraverty S, Yavuz M. Two-hybrid techniques coupled with an integral transform for Caputo time-fractional Navier-Stokes Equations. *Prog Fract Differentiation Appl.* 2020;6(3):201–13.
- [49] Gomez S CA, Salas AH. The Cole-Hopf transformation and improved tanh-coth method applied to new integrable system (KdV6). *Appl Math Comput.* 2008;204(2):957–62.
- [50] Salas A. Special symmetries to standard Riccati equations and applications. *Appl Math Comput.* 2010;216(10):3089–96.
- [51] Wazwaz AM. The tanh-coth method for solitons and kink solutions for nonlinear parabolic equations. *Appl Math Comput.* 2007;188(2):1467–75.
- [52] Kudryashov NA. One method for finding exact solutions of nonlinear differential equations. *Commun Nonlinear Sci Numer Simul.* 2012;17(6):2248–53.