

Review

Jan Franko*

Therapeutic efficacy of systemic therapy for colorectal peritoneal carcinomatosis: surgeon's perspective

<https://doi.org/10.1515/pap-2018-0102>

Received February 10, 2018; accepted February 11, 2018;
previously published online March 16, 2018

Abstract: Treatment choices for colorectal peritoneal carcinomatosis/metastases include systemic therapy and increasingly cytoreductive surgery with intraperitoneal chemotherapy delivery. These options are best considered as complementary and not exclusive alternatives. Two prospective randomized trials support use of peritonectomy procedures and intraperitoneal chemotherapy for colorectal peritoneal carcinomatosis. This overview examines efficacy, limitations and landscape of systemic therapy focusing on colorectal peritoneal carcinomatosis. Observations from literature support notions that (1) systemic therapy provides survival benefit for all prototypical patients with mCRC irrespective of metastatic disease site; (2) the magnitude of this benefit is considerably reduced among patients with peritoneal metastases who consequently experience significantly shorter overall survival; (3) efficacy of systemic therapy improved over time but at a slower pace for those with carcinomatosis; (4) this therapeutic difference has not diminished with introduction of targeted therapy, but perhaps widened; (5) further research of cytoreductive surgery and/or intraperitoneal regional therapies is thus a multidisciplinary responsibility of the entire oncology community; (6) peritonectomy procedures with intraperitoneal regional therapy are not scientifically supported in absence of systemic therapies.

Keywords: cytoreduction, hyperthermia, intraperitoneal therapy, intraperitoneal chemotherapy, peritoneal surface, peritonectomy

Introduction

Contemporary landscape of treatment options in metastatic colorectal cancer (mCRC) includes systemic agents (cytotoxic, targeted and immunotherapy) and regional therapies (hepatic resection, ablation, regional chemo/radiotherapy delivery and similar). Comprehensive management of metastatic colorectal cancer patients is based on individual patient performance status, metastatic disease extent and treatment-related risk–benefit ratio. Thus, in context of systemic tumor dissemination (or risk thereof), prescription of systemic therapy until disease progression, unacceptable toxicity or disease complications is experienced by most patients. Outlook of mCRC patients has vastly improved in past decades, yet cure (long-term disease-free survival) remains restricted to a limited subgroup of those with resectable liver or lung metastases [1–6].

Peritoneal metastases in colorectal cancer (pmCRC) develop after coelomic metastatic spread as opposed to hematogenous route in hepatic or pulmonary metastases. Peritoneal metastases (or carcinomatosis) are associated with markedly worsened prognosis [7–15], perhaps due to progressive gastrointestinal dysfunction culminating in cachexia from carcinomatosis syndrome [16, 17]. Peritoneal surface lesions are difficult to radiologically detect, often do not meet RECIST criteria to qualify for “measurable disease” and thus are often excluded from response evaluation. Carcinomatosis deposits have a dense extracellular matrix causing elevated interstitial fluid pressure. Such environment forms an unfavorable therapeutic gradient resulting in diminished drug bioavailability and amplified drug clearance from peritoneal nodules after intravenous delivery [16, 18]. Nevertheless, a minority of authors would consider patients with limited peritoneal carcinomatosis as potentially curable if approached by combination of systemic and regional therapy [19–21]. Peritonectomy, abdominal cytoreduction and regional chemotherapy have been cautiously yet increasingly accepted by major guidelines both in Europe [22] and the United States [3] after some 30 years

*Corresponding author: Jan Franko, Chair of Surgical Oncology, Mercy Medical Center, 411 Laurel Street, Suite 2100, Des Moines, IA 50314, USA, E-mail: jan.franko@gmail.com

Table 1: Overall survival of colorectal peritoneal carcinomatosis patients treated with systemic chemotherapy alone or in combination with cytoreductive surgery and intraperitoneal chemotherapy. Listed are all four available comparative studies with internal control.

Study type Origin [citation]	Overall median survival (months)		Hazard ratio
	Systemic therapy-only	Systemic therapy and CS + IP therapy	
Prospective Amsterdam [21]	12.6	22.3	0.55
Retrospective France [19]	23.9	62.7	0.38 (estimate)
Retrospective Pittsburgh [24]	16.8	34.7	0.42
Prospective Sweden [20] ^a	18.0	25.0	0.51

^aScandinavian trial delivered intraperitoneal and systemic therapy through intraperitoneal port and did not use hyperthermia, which was used in other three studies. CS + IP, cytoreductive surgery plus intraperitoneal therapy.

of liberal clinical experience, nevertheless controversy on this treatment modality is far from over [23].

Comparative studies of systemic therapy and its combination with surgical cytoreduction demonstrate reliable improvement in clinical outcomes (Table 1) [19–21, 24, 25], despite differences in histological type and disease volume burden [26–28]. Another important, yet often overlooked feature of colorectal peritoneal metastases/carcinomatosis, is its consistently worse survival as compared to unselected mCRC patients or those with absence of peritoneal involvement. Meaningful (by about 30–40 %) and statistically significant overall survival reduction was

observed in population reports, retrospective institutional series as well as pooled studies of prospective randomized trials (Table 2, Figures 1, 2, 3) [7–11, 29–32].

While systemic therapy approaches are largely agnostic of mCRC disease site, regional approaches, such as hepatectomy, lung resection, and peritoneal cytoreduction with intraperitoneal chemotherapy capitalize on subtle phenotypic regions-specific characteristics. Optimal treatment of pmCRC must therefore require a combination of systemic and regional therapies in harmony to ensure synergy and long-term disease control [2, 19–21, 33]. To this end it is important to study effectiveness of systemic

Table 2: Overall survival of patients with colorectal peritoneal metastases (pmCRC) treated in prospective studies by systemic therapy only (no regional therapies).

Population sample/study	n pmCRC only	Systemic therapy	Hazard ratio pmCRC vs. other mCRC	Median OS, months	Weighted average (median OS)
[12]	45	5-FU	NR	6.0	7.2 months
Royal Marsden Hospital [7]	91	5-FU ± interferon	1.46	NR	HR = 1.27
EVOCAPE-1 [15]	118	5-FU	NR	5.2	
[21, 41] (some appendix cancers)	51	5-FU (±IRI)	NR	12.6	
[8]	326	5-FU	1.38	6.9–7.8	
[8]	71	5-FU + IRI	1.19	17.9	13.2 months
[9]	364	5-FU/OX/IRI	1.32	12.7	HR = 1.27
CAIRO [11]	34	Cap ±IRI → Cap + Ox	1.66 ^E	10.4	
CAIRO2 [11]	47	Cap + Ox + Bev ± cetuximab	1.32 ^E	15.2	
Peritoneum-only ARCAD [10]	193	5-FU/Ox/IRI/bio	1.28 vs. non-pmCRC with 1 site	16.3	
Peritoneum + another site(s) [10]	1181	5-FU/Ox/IRI/bio	1.24 vs. non-pmCRC with ≥ 1 site	12.6	

5-FU, 5-fluorouracil; IRI, irinotecan; OX, oxaliplatin; Cap, capecitabine; Bev, bevacizumab; bio, biologic/targeted agent; ^Eestimated hazard ratio; OS, overall survival; NR, not reported.




Figure 1: Overall survival in patients with metastatic colorectal cancer with metastases in a single organ. Adapted with permission from [10].




Figure 2: Overall survival of mCRC treated by first-line systemic therapy in 14 prospective randomized trials. Individual groups are selected by peritoneal involvement (solid lines – mCRC with peritoneal involvement; dashed lines – mCRC without peritoneal involvement) and number of metastatic disease sites (1 or ≥2). Adapted with permission from [10].




Figure 3: Median overall survival in selected studies of systemic chemotherapy published after adoption of oxaliplatin and irinotecan. All presented studies are secondary/retrospective analysis of individual patient data collected prospectively on randomized trials of systemic therapy for mCRC. Larger size of circle corresponds to larger dataset. Blue circles = mCRC patients without peritoneal involvement, red circles = patients with colorectal peritoneal carcinomatosis (pmCRC+).

therapy specifically for peritoneal metastases, if one desires to meticulously examine the role of peritoneal-directed regional approaches, such as cytoreduction ± hyperthermic intraperitoneal chemotherapy (HIPEC) [19, 21, 24], extended postoperative intraperitoneal chemotherapy (EPIC) [20] or pressurized intraperitoneal chemotherapy (PIPAC) [33]. This overview examines efficacy, characteristics and limitations of systemic therapy for colorectal peritoneal carcinomatosis.

Era of 5-fluorouracil

5-fluorouracil has been a backbone of cytotoxic chemotherapy for colorectal cancer for decades and the first widely used drug for metastatic colorectal cancer. Folprecht et al. provided excellent evidence of clinically meaningful outcome differences between cases with and without peritoneal metastases in 2007 [8]. Analyzing 2,568 patients treated by first-line 5-FU chemotherapy in randomized prospective trials an overall survival benefit was observed for infusional 5-FU as compared to patients treated by 5-FU

bolus (14.6 versus 10.8 months, $p < 0.0001$). However, no survival difference between infusional and bolus 5-FU application was observed among patients with peritoneal metastases (7.8 versus 6.9 months, $p = 0.44$). Additionally, pmCRC patients demonstrated markedly shorter median survival when compared to mCRC without carcinomatosis, although it was not commented upon by authors. In agreement with survival data, there was a significant difference in response rates based on peritoneal status and mode of 5-FU delivery. Patients without peritoneal involvement enjoyed substantially better objective response rates (36.2% and 19.9% for infusional and bolus 5-FU, respectively; $p < 0.001$) as compared to pmCRC cases (19% and 12.6% for infusional and bolus 5-FU, respectively; $p = 0.14$).

Many authors observed that peritoneal disease sites were radiologically less responsive as compared to hepatic metastases in mCRC [7, 27, 34, 35]. Assersohn et al. [7] pooled data obtained from trials of 5-FU based therapy conducted before approval of modern cytostatics. This analysis of Royal Marsden Hospital prospective data demonstrated substantially lower objective response of metastases in peritoneum as compared to other metastatic

Site-specific response w/ 5-FU

Royal Marsden Hospital pooled prospective trials of 5-FU therapy; n=497




Figure 4: Response rates depending on site of metastases in mCRC patients treated with historical 5-FU combinations. Data from Royal Marsden Hospital studies [7].

sites. Any response was observed among 10% of peritoneal metastases, while liver metastases had 40% response rate. Additionally, progressive disease was noted among 20% of pmCRC patients over the study period as compared to 10% of colorectal liver metastases (Figure 4).

Era of modern cytostatics: oxaliplatin and irinotecan

Widespread approval of oxaliplatin and irinotecan around 2004 led to a meaningful improvement in survival of patient population with mCRC as a whole (Figure 3). Importantly, there is virtually no clinical difference observed when administering all cytotoxic drugs concurrently in a very intense regimen as compared to sequential treatment started with less toxic combination first [36–38]. Survival is improved as long as all available cytotoxic agents are used during disease course [39].

However, the Eindhoven cancer registry study suggested that while survival for patients with colorectal liver metastases was longer in 2005–2008 period as compared to earlier periods, there was no significant improvement in overall survival among registered patients with isolated peritoneal disease, even when analysis was limited to chemotherapy recipients [29]. Because only 10 out of 904 patients with peritoneal-

only metastases received peritoneal cytoreduction and HIPEC, this regional therapy has not confounded study results. A nearly identical conclusion came from a two-decade retrospective German institutional review with 2,406 patients; while mCRC patients all together experienced substantial survival gain over time, this benefit was not realized among those with carcinomatosis [32].

Survival advantage of irinotecan combined with 5-FU was suggested by separating survival curves, but was not statistically significant in a limited sample of patients with peritoneal carcinomatosis (n=71; 17.9 versus 9.8 months for 5-FU + irinotecan versus 5-FU only, $p=0.17$). Response rates were again higher among non-peritoneal mCRC and irinotecan combination as compared to pmCRC or 5-FU only [8]. Reported objective response rate among mCRC with no peritoneal metastases were 56% for 5-FU+irinotecan versus 32% for 5-FU ($p<0.001$). Lower response rates were reported among those with colorectal peritoneal metastases (39% for 5-FU + irinotecan versus 14% 5-FU alone; $p=0.03$).

A larger and detailed analysis of clinical outcomes of modern cytotoxic chemotherapy among patients with peritoneal involvement was provided by our group in an analysis of the North Central Cancer Treatment Group trials (N9741 trial, first-line therapy and N9841 trial, second-line therapy after first progression) [9]. There were 364 patients with peritoneal metastases among 2,101 patients extracted from

databases of these prospective randomized studies. Hazard ratio for earlier death among pmCRC patients was consistently some 30 % higher as compared to cases without carcinomatosis, even after multiple adjustments. Of note, peritoneal metastases were more frequent among more advanced disease patients (22.6 % among patients in second-line versus 15.9 % in first-line treatment trial, $p < 0.001$). Even greater survival difference between patients with and without peritoneal metastases was observed subsequently by re-analysis of CAIRO (estimated HR 1.6 for pmCRC patients as compared to those without peritoneal involvement) and CAIRO2 studies (estimated HR 1.4). This survival difference was judged to be related to intrinsic features of peritoneal involvement and not undertreatment, because median number of chemotherapy cycles was not statistically different between patients with and without peritoneal carcinomatosis [11].

Histological subtype has been a recognized survival predictor among CS-HIPEC patients [24, 26]. Regrettably, there is a remarkable paucity of information on histological features of mCRC among patients enrolled to randomized trials. Nonetheless, adjusted retrospective analyses suggest that non-mucinous tumors are 3.4 times more likely to respond as compared to mucinous and that non-peritoneal metastases are 2.7 times more likely to respond as compared to peritoneal metastases [27]. Based on nearly 6000 autopsies mucinous

and signet-ring cell tumors metastasize to peritoneum more frequently as compared to more favorable adenocarcinoma with no other specification [28]. Possible link between proximal colon cancer, mucinous or signet-ring-cell histology and peritoneal dissemination represents an opportunity for further research [10, 28, 40].

Era of targeted therapies

A comprehensive evaluation of cytotoxic chemotherapy alone and in combination with targeted (biologic) therapy was facilitated by the availability of the ARCAD Project (Aide et Recherche en Cancérologie Digestive; www.fondationarcad.org).

One ARCAD endeavor adopted included prospective randomized studies of first-line systemic therapy, which either solicited request for peritoneal involvement in their protocol or performed peritoneum-specific review of original computed tomography scans to ascertain whether peritoneum was or was not involved [10]. Trial inclusion criteria scrutiny was so tall that we did not include N9741 trial forming basis of our prior report of colorectal peritoneal carcinomatosis [9]. The final study involved individual patient-level data from 14 prospective randomized studies with known peritoneal status and included 10,533 patients [10]. There were 1181 patient with peritoneal




Figure 5: Hazard ratios for overall survival by site of metastases in cytotoxic-only therapy (left panel) and combined therapy including at least one targeted agent (right panel).

*Denotes statistical significance at least $p < 0.05$. Data from [10].

metastases in addition to other metastatic disease sites and 194 patients with metastases isolated to peritoneum. We reconfirmed in the largest dataset that peritoneal involvement among mCRC patients recruited to randomized trials is associated with shortened overall survival and progression free survival. Additionally, survival difference between patients with peritoneal involvement and disease-free peritoneum widened (Figures 1, 2, 5), data which was previously suggested by secondary analysis of CAIRO2 trial [11]. Median overall survival of patients with isolated peritoneal-only metastases was 16.8 months, whether cytotoxic chemotherapy was used alone or in combination with targeted agents (of note they were only 25 patients with peritoneal-only metastases and targeted therapy). On the contrary, there was trend toward longer survival among colorectal liver metastases patients treated by targeted agent as compared to those treated by cytotoxic chemotherapy only (20.4 months versus 18 months). Interestingly, remarkably similar survival of cetuximab-treated patients without peritoneal metastases (20.7 months) was observed in CAIRO2 study [11].

Discussion

There are multiple prospective data based [7–12, 15, 21, 41] and purely retrospective studies [27, 29, 32, 42] examining therapeutic efficacy of systemic therapy for colorectal peritoneal carcinomatosis. The largest prospective data-based studies confirmed both therapeutic efficacy of systemic chemotherapy but also consistently demonstrated inferior survival of patients affected by colorectal carcinomatosis in early 5-FU period [8], oxaliplatin/irinotecan era [9, 11] and current era of targeted biologic therapy [10, 11]. Nevertheless, the literature recorded survival improvement of all forms of mCRC, including for those with peritoneal surface metastases (Table 2, Figure 3).

Meaningful survival improvements, and sometimes even cure, have been observed among those with resected or ablated liver metastases, typically in combination with systemic therapy [1, 2, 4–6]. On the other hand, failure to demonstrate clear survival benefit of well-designed liver-directed approaches, like Y90-based selective internal radiation or hepatic arterial infusion, reveals the complexity of metastatic colorectal cancer [43, 44].

Systemic therapy approaches are agnostic of metastatic site. On the contrary, regional approaches involving surgery and regional chemotherapy have been designed for specific metastatic site since their conception. Clinicians faced with historically poor prognosis developed peritoneal

surface surgery combined with hyperthermic intraperitoneal chemotherapy [17, 21, 25, 45]. After years of research both the European and the United States National Comprehensive Cancer Network guidelines carefully recognized a restricted role of peritoneal cytoreduction and HIPEC in colorectal peritoneal carcinomatosis in addition to established role of systemic therapy [3, 22].

There are notable limitations of this review. Data from most studies examining peritoneal status come from older studies, and thus we lack information on efficacy of modern cytostatics (e.g. TAS-102). On the contrary, fairly good agent-specific data are available for 5-FU, oxaliplatin and irinotecan [8, 9, 37–39]. Much less granularity is available for six approved targeted therapies by the end of 2017: bevacizumab, cetuximab, panitumumab, ziv-aflibercept, regorafenib, ramucirumab. The ARCAD agreements make it impossible to study drug versus drug, and therefore only class of drugs may be compared within ARCAD projects, i.e. antiangiogenic class as opposed to individual effect of bevacizumab or ziv-aflibercept. No peritoneum-specific facts are available on immunotherapy for advanced mCRC, such as approved pembrolizumab or not-yet-approved ipilimumab. Notably, contemporaneous immunotherapy is applicable exclusively to patients with mismatch repair deficient genome.

There are no uniformly accepted and clinically useful prognostic tumor biomarkers for mCRC. *Consensus Molecular Subtypes* classification has identified clinical differences in overall survival, relapse-free survival and survival after relapse among four defined subtypes, but has not been applied to peritoneal or other site-specific metastases [46]. While *BRAF* mutations have been associated with worsened survival among mCRC patients, it became clear that survival shortening is specifically associated with mutations in codon 600 (^{V600E}*BRAF*-mutant mCRC), while ^{non-V600}*BRAF* mutations feature clinical course superior to that seen among wild-type *BRAF* mCRC patients [47]. Recent data, however, support even simpler clinical characteristics as powerful biomarkers – both peritoneal involvement and sidedness of primary colon tumor, with right-sided tumors featuring shorter survival and inefficacy of epithelial growth factor blocking therapies. Increased proportion of right-sided tumors and peritoneal carcinomatosis was observed in the ARCAD study in the whole population and wild-type *BRAF* mCRC patients alike [10].

In the author's interpretation, systemic treatment forms a backbone of modern therapy for metastatic colorectal cancer, including among patients with peritoneal metastases. Two randomized trials of surgical cytoreduction and

intraperitoneal chemotherapy were conducted on background of systemic chemotherapy [20, 21, 41], as were both retrospective studies with internal control supporting the therapeutic role of CS-HIPEC in colorectal peritoneal carcinomatosis [19, 24]. Despite proven efficacy of systemic therapy in all forms of mCRC, clinical outcome among those with peritoneal metastases is consistently inferior to unselected mCRC population [7–11]. Therefore, it is imperative to explore adjuncts in peritoneal carcinomatosis treatment such as surgical cytoreduction and various intraperitoneal therapies (HIPEC, EPIC, PIPAC). Yet those adjunct approaches must evolve in addition to and not instead of systemic therapy. Moreover, surgeons must maintain knowledge of therapeutic effectiveness of colorectal systemic therapy, and related risks and benefits for patients in surgical consideration.

Author contribution: The author has accepted responsibility for the entire content of this submitted manuscript and approved submission.

Research funding: None declared.

Employment or leadership: None declared.

Honorarium: None declared.

Competing interests: None declared.

References

- Nordlinger B, Sorbye H, Glimelius B, Poston GJ, Schlag PM, Rougier P, et al. Group EG-ITC, Cancer Research UK, Arbeitsgruppe Lebermetastasen und-tumoren in der Chirurgischen Arbeitsgemeinschaft O, Australasian Gastro-Intestinal Trials G, Federation Francophone de Cancerologie D. Perioperative FOLFOX4 chemotherapy and surgery versus surgery alone for resectable liver metastases from colorectal cancer (EORTC 40983): long-term results of a randomised, controlled, phase 3 trial. *Lancet Oncol* 2013;14:1208–15.
- Ruers T, Van Coevorden F, Punt CJ, Pierie JE, Borel-Rinkes I, Ledermann JA, et al. European Organisation for R, Treatment of C, Gastro-Intestinal Tract Cancer G, Arbeitsgruppe Lebermetastasen und tumoren in der Chirurgischen Arbeitsgemeinschaft O, National Cancer Research Institute Colorectal Clinical Study G. Local Treatment of Unresectable Colorectal Liver Metastases: results of a Randomized Phase II Trial. *J Natl Cancer Inst* 2017;109.
- NCCN Guidelines: Colon Cancer. 2018. Accessed February 4, 2018, 2018. http://www.nccn.org/professionals/physician_gls/pdf/colon.pdf.
- Abdalla EK, Vauthey JN, Ellis LM, Ellis V, Pollock R, Broglio KR, et al. Recurrence and outcomes following hepatic resection, radiofrequency ablation, and combined resection/ablation for colorectal liver metastases. *Ann Surg* 2004;239:818–25. discussion 25–7.
- Adam R, Pascal G, Castaing D, Azoulay D, Delvart V, Paule B, et al. Tumor progression while on chemotherapy: a contraindication to liver resection for multiple colorectal metastases? *Ann Surg* 2004;240:1052–61. discussion 61–4.
- Fong Y, Fortner J, Sun RL, Brennan MF, Blumgart LH. Clinical score for predicting recurrence after hepatic resection for metastatic colorectal cancer: analysis of 1001 consecutive cases. *Ann Surg* 1999;230:309–18. discussion 18–21.
- Assersohn L, Norman A, Cunningham D, Benepal T, Ross PJ, Oates J. Influence of metastatic site as an additional predictor for response and outcome in advanced colorectal carcinoma. *Br J Cancer* 1999;79:1800–5.
- Folprecht G, Köhne C-H LM. Systemic chemotherapy in patients with peritoneal carcinomatosis from colorectal cancer. In: Ceelen WP, editor. *Peritoneal carcinomatosis: a multidisciplinary approach*. New York: Springer, 2007:425–40.
- Franko J, Shi Q, Goldman CD, Pockaj BA, Nelson GD, Goldberg RM, et al. Treatment of colorectal peritoneal carcinomatosis with systemic chemotherapy: a pooled analysis of north central cancer treatment group phase III trials N9741 and N9841. *J Clin Oncol* 2012;30:263–7.
- Franko J, Shi Q, Meyers JP, Maughan TS, Adams RA, Seymour MT, et al. Analysis, research in cancers of the digestive system G. prognosis of patients with peritoneal metastatic colorectal cancer given systemic therapy: an analysis of individual patient data from prospective randomised trials from the analysis and research in cancers of the digestive system (ARCAD) database. *Lancet Oncol* 2016;17:1709–19.
- Klaver YL, Simkens LH, Lemmens VE, Koopman M, Teerenstra S, Bleichrodt RP, et al. Outcomes of colorectal cancer patients with peritoneal carcinomatosis treated with chemotherapy with and without targeted therapy. *Eur J Surg Oncol* 2012;38: 617–23.
- Chu DZ, Lang NP, Thompson C, Osteen PK, Westbrook KC. Peritoneal carcinomatosis in nongynecologic malignancy. A prospective study of prognostic factors. *Cancer* 1989;63: 364–7.
- Jayne DG, Fook S, Loi C, Seow-Choen F. Peritoneal carcinomatosis from colorectal cancer. *Br J Surg* 2002;89:1545–50.
- Koppe MJ, Boerman OC, Oyen WJ, Bleichrodt RP. Peritoneal carcinomatosis of colorectal origin: incidence and current treatment strategies. *Ann Surg* 2006;243:212–22.
- Sadeghi B, Arvieux C, Glehen O, Beaujard AC, Rivoire M, Baulieux J, et al. Peritoneal carcinomatosis from non-gynecologic malignancies: results of the EVOCAPE 1 multicentric prospective study. *Cancer* 2000;88:358–63.
- Ceelen WP, Bracke ME. Peritoneal minimal residual disease in colorectal cancer: mechanisms, prevention, and treatment. *Lancet Oncol* 2009;10:72–9.
- Sugarbaker PH, Jablonski KA. Prognostic features of 51 colorectal and 130 appendiceal cancer patients with peritoneal carcinomatosis treated by cytoreductive surgery and intraperitoneal chemotherapy. *Ann Surg* 1995;221:124–32.
- Francescutti V, Rivera L, Seshadri M, Kim M, Haslinger M, Camoriano M, et al. The benefit of intraperitoneal chemotherapy for the treatment of colorectal carcinomatosis. *Oncol Rep* 2013;30:35–42.
- Elias D, Lefevre JH, Chevalier J, Brouquet A, Marchal F, Classe JM, et al. Complete cytoreductive surgery plus intraperitoneal chemohyperthermia with oxaliplatin for peritoneal carcinomatosis of colorectal origin. *J Clin Oncol* 2009;27: 681–5.
- Cashin PH, Mahteme H, Spang N, Syk I, Frodin JE, Torkzad M, et al. Cytoreductive surgery and intraperitoneal chemotherapy

- versus systemic chemotherapy for colorectal peritoneal metastases: A randomised trial. *Eur J Cancer* 2016;53:155–62.
21. Verwaal VJ, Van Ruth S, De Bree E, Van Sloothen GW, Van Tinteren H, Boot H, et al. Randomized trial of cytoreduction and hyperthermic intraperitoneal chemotherapy versus systemic chemotherapy and palliative surgery in patients with peritoneal carcinomatosis of colorectal cancer. *J Clin Oncol* 2003;21:3737–43.
 22. Van Cutsem E, Cervantes A, Adam R, Sobrero A, Van Krieken JH, Aderka D, et al. ESMO consensus guidelines for the management of patients with metastatic colorectal cancer. *Ann Oncol* 2016;27:1386–422.
 23. Ryan D. Cytoreductive surgery and hyperthermic intraperitoneal chemotherapy: history repeating itself or a new standard? In: ASCO 2011 educational book. American Society of Clinical Oncology 2011:127–8.
 24. Franko J, Ibrahim Z, Gusani NJ, Holtzman MP, Bartlett DL, HJ Z. 3rd. Cytoreductive surgery and hyperthermic intraperitoneal chemoperfusion versus systemic chemotherapy alone for colorectal peritoneal carcinomatosis. *Cancer* 2010;116:3756–62.
 25. Elias D, Delpero JR, Sideris L, Benhamou E, Pocard M, Baton O, et al. Treatment of peritoneal carcinomatosis from colorectal cancer: impact of complete cytoreductive surgery and difficulties in conducting randomized trials. *Ann Surg Oncol* 2004;11:518–21.
 26. Verwaal VJ, Van Tinteren H, Van Ruth S, Zoetmulder FA. Predicting the survival of patients with peritoneal carcinomatosis of colorectal origin treated by aggressive cytoreduction and hyperthermic intraperitoneal chemotherapy. *Br J Surg* 2004;91:739–46.
 27. Catalano V, Loupakis F, Graziano F, Torresi U, Bisonni R, Mari D, et al. Mucinous histology predicts for poor response rate and overall survival of patients with colorectal cancer and treated with first-line oxaliplatin- and/or irinotecan-based chemotherapy. *Br J Cancer* 2009;100:881–7.
 28. Hugen N, Van De Velde CJ, De Wilt JH, Nagtegaal ID. Metastatic pattern in colorectal cancer is strongly influenced by histological subtype. *Ann Oncol* 2014;25:651–7.
 29. Klaver YL, Lemmens VE, Creemers GJ, Rutten HJ, Nienhuijs SW, De Hingh IH. Population-based survival of patients with peritoneal carcinomatosis from colorectal origin in the era of increasing use of palliative chemotherapy. *Ann Oncol* 2011;22:2250–6.
 30. Lemmens VE, Klaver YL, Verwaal VJ, Rutten HJ, Coebergh JW, De Hingh IH. Predictors and survival of synchronous peritoneal carcinomatosis of colorectal origin: a population-based study. *Int J Cancer* 2011;128:2717–25.
 31. Segelman J, Granath F, Holm T, Machado M, Mahteme H, Martling A. Incidence, prevalence and risk factors for peritoneal carcinomatosis from colorectal cancer. *Br J Surg* 2012;99:699–705.
 32. Kersch AG, Chua TC, Gasser M, Maeder U, Kunzmann V, Isbert C, et al. Impact of peritoneal carcinomatosis in the disease history of colorectal cancer management: a longitudinal experience of 2406 patients over two decades. *Br J Cancer* 2013;108:1432–9.
 33. Demtröder C, Solass W, Zieren J, Strumberg D, Giger-Pabst U, Raymond MA. Pressurized intraperitoneal aerosol chemotherapy with oxaliplatin in colorectal peritoneal metastasis. *Colorectal Dis* 2016;18:364–71.
 34. Hill M, Norman A, Cunningham D, Findlay M, Watson M, Nicolson V, et al. Impact of protracted venous infusion fluorouracil with or without interferon alfa-2b on tumor response, survival, and quality of life in advanced colorectal cancer. *J Clin Oncol* 1995;13:2317–23.
 35. Kouri M, Pyrhonen S, Kuusela P. Elevated CA19-9 as the most significant prognostic factor in advanced colorectal carcinoma. *J Surg Oncol* 1992;49:78–85.
 36. Tournigand C, Andre T, Achille E, Lledo G, Flesh M, Mery-Mignard D, et al. FOLFIRI followed by FOLFOX6 or the reverse sequence in advanced colorectal cancer: a randomized GERCOR study. *J Clin Oncol* 2004;22:229–37.
 37. Koopman M, Antonini NF, Douma J, Wals J, Honkoop AH, Erdkamp FL, et al. Sequential versus combination chemotherapy with capecitabine, irinotecan, and oxaliplatin in advanced colorectal cancer (CAIRO): a phase III randomised controlled trial. *Lancet* 2007;370:135–42.
 38. Seymour MT, Maughan TS, Ledermann JA, Topham C, James R, Gwyther SJ, et al. National cancer research institute colorectal clinical studies G. different strategies of sequential and combination chemotherapy for patients with poor prognosis advanced colorectal cancer (MRC FOCUS): a randomised controlled trial. *Lancet* 2007;370:143–52.
 39. Grothey A, Sargent D. Overall survival of patients with advanced colorectal cancer correlates with availability of fluorouracil, irinotecan, and oxaliplatin regardless of whether doublet or single-agent therapy is used first line. *J Clin Oncol* 2005;23:9441–2.
 40. Hugen N, Verhoeven RH, Lemmens VE, Van Aart CJ, Elferink MA, Radema SA, et al. Colorectal signet-ring cell carcinoma: benefit from adjuvant chemotherapy but a poor prognostic factor. *Int J Cancer* 2015;136:333–9.
 41. Verwaal VJ, Bruin S, Boot H, Van Slooten G, Van Tinteren H. 8-Year follow-up of randomized trial: cytoreduction and hyperthermic intraperitoneal chemotherapy versus systemic chemotherapy in patients with peritoneal carcinomatosis of colorectal cancer. *Ann Surg Oncol* 2008;15:2426–32.
 42. Kennecke H, Yu J, Gill S, Cheung WY, Blanke CD, Speers C, et al. Effect of M1a and M1b category in metastatic colorectal cancer. *Oncologist* 2014;19:720–6.
 43. Wasan HS, Gibbs P, Sharma NK, Taieb J, Heinemann V, Ricke J, et al. First-line selective internal radiotherapy plus chemotherapy versus chemotherapy alone in patients with liver metastases from colorectal cancer (FOXFIRE, SIRFLOX, and FOXFIRE-Global): a combined analysis of three multicentre, randomised, phase 3 trials. *Lancet Oncol* 2017;18:1159–71.
 44. Kemeny NE, Chou JF, Boucher TM, Capanu M, DeMatteo RP, Jarnagin WR, et al. Updated long-term survival for patients with metastatic colorectal cancer treated with liver resection followed by hepatic arterial infusion and systemic chemotherapy. *J Surg Oncol* 2016;113:477–84.
 45. Shen P, Hawksworth J, Lovato J, Loggie BW, Geisinger KR, Fleming RA, et al. Cytoreductive surgery and intraperitoneal hyperthermic chemotherapy with mitomycin C for peritoneal carcinomatosis from nonappendiceal colorectal carcinoma. *Ann Surg Oncol* 2004;11:178–86.
 46. Guinney J, Dienstmann R, Wang X, De Reynies A, Schlicker A, Soneson C, et al. The consensus molecular subtypes of colorectal cancer. *Nat Med* 2015;21:1350–6.
 47. Jones JC, Renfro LA, Al-Shamsi HO, Schrock AB, Rankin A, Zhang BY, et al. (Non-V600) BRAF mutations define a clinically distinct molecular subtype of metastatic colorectal cancer. *J Clin Oncol* 2017;35:2624–30.