Supplementary material

S1 Preliminary searches

The search terms used in the preliminary searches are reported in Table S1. The reviews that were examined are detailed on Table S2 (published reviews) and Table S3 (ongoing reviews).

Table S1: Preliminary searches terms

Database	Date	Terms	N hits	N review on EIH
				and reliability
COCHRANE	2021-09-12	hypoalgesia	72	0
	2021-09-12	exercise-induced analgesia	32	0
	2021-09-12	exercise-induced analgesia	28	0
	2021-09-19	exercise-induced analgesia OR exercise-induced hypoalgesia	77	0
EPISTEMONIKOS	2021-09-12	exercise-induced hypoalgesia	8	0
	2021-09-12	hypoalgesia	25	0
	2021-09-12	exercise-induced analgesia	2	0
	2021-09-14	(title:(exercise-induced hypoalgesia) OR abstract:(exercise-	9	0
		induced hypoalgesia)) OR (title:(exercise-induced analgesia) OR		
		abstract:(exercise-induced analgesia))		
	2021-09-19	exercise induced hypoalgesia OR exercise induced analgesia	3	0
PROSPERO	2021-09-12	exercise-induced hypoalgesia	11	0
	2021-09-12	hypoalgesia	20	0
	2021-09-12	exercise-induced analgesia	2	0
	2021-09-13	exercise-induced hypoalgesia OR hypoalgesia OR exercise-	21	0
		induced analgesia		

EIH, Exercise-induced hypoalgesia.

Table S2: Published reviews found during preliminary searches

	Reliability EIH Author (Yea	ar) SR	R N stu	idies D	Author (Year) SR N studies Design studies	Type of exercise	Population	Years of	Language	Databases
								included	limit	
								searches		
No Yes	Wewege	Ye	Yes 13	2	Randomized controlled studies	Single bout of	healthy human,	1996 to 2019	English	Scopus, EMBASE,
	et al. (2020)			<u> </u>	(cross-over or paralell)	standardized	human with			PsycINFO,CINAHL, SPORTDiscus,
						exercise	msk pain			and MEDLINE
No Yes	Munneke	Yes	es 9	U	Cross-sectional	Single bout of	Adults: healthy and	2002 to 2018	Not found	MEDLINE Ovid, EMBASE, Web of
	et al. (2019)					exercise	with msk pain			Science, PsycINFO Ovid and
										CINAHL EBSCOhost
No Yes	Naugle	No	0 25	⋖	All published	Aerobic, Isometric,	Adults, healthy or	1900 to 2012	Not reported	PubMED, Medline, PsychINFO,
	et al. (2012)					resistance exercise	chronic pain			and Academic Search
No Yes	Pacheco-	Yes	es 36	~	RCT including: included parallel-	exercise	Healthy subjects	1995 to 2019	English	MEDLINE, EMBASE, Web of
	Barrios			б	group, crossover designs and	intervention				Science, Lilacs, and Scopus
	et al. (2020)			d	pilot studies and quasi-					
				Φ	experimental studies					
No Unc	Unclear Baehr	ž	No 196	~	RCT, quasi-experimental,	Not found	Healthy adults and	Not found	Not found	Medline (Pubmed), CINAHL,
	et al. (2021)			0	observational		adults with chronic			Web of Science, Embase
							medical conditions			
No	Polaski	Ye	Yes 75	~	RCT	75.0	See Green et al 2017	See Green et	English	See Green et al. 2017
	et al. (2019)							al 2017		
No Yes	Bonello	Ye	Yes 13	~	RCT, non-randomized pre-post	Isometric exercise	Persons with	2002 to 2020	English	MEDLINE, EMBASE, CINAHL &
	et al. (2021)						localised msk pain			SportDiscus

Table S3: Ongoing reviews found during preliminary searches

Psychometric EIH	Ħ	Author	S.	N studies	N studies Design studies	Type of	Population	Years of	Langnage	Databases
		(Year)				exercise		included	limit	
								searches		
No	Unclear	Unclear RB Rodriguez Yes	Yes		RCT	Aerobic	Patients with		Not reported	Not reported PubMed, PEDro, EBSCO, WOS
		et al. (2021)				exercise	musculoskeletal pain			
No	Yes	Lemley	Yes	_	Randomized controlled	Retrowalking	Adults with lower extrmity		English	Academic Search Premier,
		et al. (2018)		7	trials, cohort, case control,		osteoarthitis			MEDLINE Complete, CINAHL,
				5	case series, case reports,					ScienceDirect, ClinicalKey,
				J	quasi-experimental designs,					ProQuest, PubMed
				J	clinical trials					
No	Unclear	Unclear Kuithan	Yes	J	Observational studies	Physical activity	Physical activity Adults with spinal pain		English	MEDLINE (OVID interface),
		et al. (2019)								EMBASE (OVID interface),
										CINAHL (EBSCO interface),
										Web of Science, PubMed and
										Google Scholar, Grey
										literature, encompassing
										dissertation abstracts and
										Ethos, OpenGrey, ZETOC and
										British National bibliography
No	Yes	Rio	Yes	•	All	Isometric	Healthy human		English	MEDLINE via Ovid, Web of
		et al. (2018)				exercise				Science, EMBASE, Scopus and
										CINAHL
No	Yes	Senarath	Yes	_	RCT, observational studies	Exercise for	Adults with neck pain		Enlgish	PEDro, MEDLINE, EMBASE,
		et al. (2021)				neck pain				CINAHL, SPORTDiscuss, AMED
										through Ovid
Yes	No	Alqarni	Yes	J	Cross-sectional, cohort	No exercise	Adults with		English	CINAHL, ScienceDirect,
		et al. (2016)		01	studies		musculoskeletal pain			MEDLINE, PubMed, Scopus,
										PsycINFO, Web of Knowledge

Table S3: Continued

Psychometric EIH	Author (Year)	SR N stuc	SR N studies Design studies	Type of exercise	Population	Years of included searches	Language limit	Databases
No Yes	Cayrol	Yes	Observational studies	Single bout of	Adults (18 or older) with	From 1992	Not limited	MEDLINE/PubMed, EMBASE
	et al. (2021)			exercise	chronic pain-related			(including the "MEDLINE
					myofascial			search"), Scopus, CINAHL,
					temporomandibular disorders			PsycINFO
No Yes	Deere	Yes	RCT, non-randomized	Exercise	Adults(>40y) with loer	From 2000	English	PubMed, Scopus, Web of
	et al. (2020)		pre-post	training	extremity OA)			Science
No	Delavy	Yes	Experimental design, full-	ll- Exercise	Adults(>18y)		Not limited	MEDLINE, CINAHL,
	et al. (2019)		text report. Parallel arm	training alone				SPORTDiscus, EMBASE,
			(individual- or cluster-					psycINFO, CENTRAL
			designed) randomised					
			controlled trials					
No Yes	Trinder	Yes	All but observationnal	Isometric	People with local msk pain		English	MEDLINE, Scopus, CINAHL and
	et al. (2017)		studies & pre_post without	out exercise				AMED via the EBSCOhost
			comparison					interface

S2 Completed PRISMA 2020 item checklist

Section and Topic	Item #	Checklist item	Location where item is reported
TITLE			
Title	1	Identify the report as a systematic review	P1
ABSTRACT	•	tuentily the report as a systematic review	
Abstract	2	See the PRISMA 2020 for Abstracts checklist	P2
INTRODUCTION	_	occ die Fidomi 2020 for Abottucto electrist	12
Rationale	3	Describe the rationale for the review in the context of existing knowledge	P3-4
Objectives		Provide an explicit statement of the objective(s) or question(s) the review addresses	
METHODS	•	110viae an explicit statement of the objective(s) of question(s) the review addresses	
Eligibility criteria	5	Specify the inclusion and exclusion criteria for the review and how studies were	P5-6
211.6121111) 0111.01111	J	grouped for the syntheses	
Information sources	6	Specify all databases, registers, websites, organisations, reference lists and other	P6-7
mornaton sources	Ü	sources searched or consulted to identify studies. Specify the date when each source	107
		was last searched or consulted	
Search strategy	7	Present the full search strategies for all databases, registers and websites, including	Annendix 3
ocuren strategy	,	any filters and limits used	ripperium o
Selection process	8	Specify the methods used to decide whether a study met the inclusion criteria of the	P7
october process	· ·	review, including how many reviewers screened each record and each report	
		retrieved, whether they worked independently, and if applicable, details of auto-	
		mation tools used in the process	
Data collection process	9	Specify the methods used to collect data from reports, including how many	P7
Data concensis process	3	reviewers collected data from each report, whether they worked independently,	1,
		any processes for obtaining or confirming data from study investigators, and if	
		applicable, details of automation tools used in the process	
Data items	10a	List and define all outcomes for which data were sought. Specify whether all results	P7-8 Annendix 4
Duta Iteliis	100	that were compatible with each outcome domain in each study were sought (e.g. for	170 Appellaix 1
		all measures, time points, analyses), and if not, the methods used to decide which	
		results to collect	
	10h	List and define all other variables for which data were sought (e.g. participant and	P7-8 Annendix 4
	102	intervention characteristics, funding sources). Describe any assumptions made	17 o ripperium 1
		about any missing or unclear information	
Study risk of bias	11	Specify the methods used to assess risk of bias in the included studies, including	P8-9 Appendix 6
assessment		details of the tool(s) used, how many reviewers assessed each study and whether	10 5 ripperium 0
assessment		they worked independently, and if applicable, details of automation tools used in	
		the process	
Effect measures	12	Specify for each outcome the effect measure(s) (e.g. risk ratio, mean difference) used	NΔ
Effect fileasures	12	in the synthesis or presentation of results	IVII
Synthesis methods	132	Describe the processes used to decide which studies were eligible for each synthesis	NΔ
Synthesis methods	130	(e.g. tabulating the study intervention characteristics and comparing against the	IVII
		planned groups for each synthesis (item #5))	
	12h	Describe any methods required to prepare the data for presentation or synthesis,	P9-10
	131)	such as handling of missing summary statistics, or data conversions	1.0-10
	120		DQ.10
	130	Describe any methods used to tabulate or visually display results of individual	P9-10
	124	studies and syntheses	DQ.10
	130	Describe any methods used to synthesize results and provide a rationale for the	P9-10
		choice(s). If meta-analysis was performed, describe the model(s), method(s) to	

		identify the presence and extent of statistical heterogeneity, and software package	
	10-	(s) used	NA
	136	Describe any methods used to explore possible causes of heterogeneity among study	NA
	10f	results (e.g. subgroup analysis, meta-regression)	NIA
	131	Describe any sensitivity analyses conducted to assess robustness of the synthesized results	NA
Reporting bias assessment	14	Describe any methods used to assess risk of bias due to missing results in a synthesis	NA
		(arising from reporting biases)	
Certainty assessment	15	Describe any methods used to assess certainty (or confidence) in the body of evi-	P9-10 Appendix 8
		dence for an outcome	
RESULTS			
Study selection	16a	Describe the results of the search and selection process, from the number of records	P11
		identified in the search to the number of studies included in the review, ideally	
		using a flow diagram	
	16b	Cite studies that might appear to meet the inclusion criteria, but which were	P11 Appendix 9
		excluded, and explain why they were excluded	
Study characteristics		Cite each included study and present its characteristics	P11-17
Risk of bias in studies		Present assessments of risk of bias for each included study	P18-19
Results of individual	19	For all outcomes, present, for each study: (a) summary statistics for each group	P20-23
studies		(where appropriate) and (b) an effect estimate and its precision (e.g. confidence/	
Results of syntheses	200	credible interval), ideally using structured tables or plots For each synthesis, briefly summarise the characteristics and risk of bias among	NA
Results of Syntheses	20a	contributing studies	NA
	20b	Present results of all statistical syntheses conducted. If meta-analysis was done,	NA
		present for each the summary estimate and its precision (e.g. confidence/credible	
		interval) and measures of statistical heterogeneity. If comparing groups, describe	
		the direction of the effect	
	20c	Present results of all investigations of possible causes of heterogeneity among study	NA
		results	
	20d	Present results of all sensitivity analyses conducted to assess the robustness of the	NA
		synthesized results	
Reporting biases	21	Present assessments of risk of bias due to missing results (arising from reporting	NA
		biases) for each synthesis assessed	
Certainty of evidence	22	Present assessments of certainty (or confidence) in the body of evidence for each	P 20, 23 Appendix 8
		outcome assessed	
DISCUSSION			
Discussion		Provide a general interpretation of the results in the context of other evidence	P24-25
		Discuss any limitations of the evidence included in the review	P25-26
		Discuss any limitations of the review processes used	P25-26 P27
OTHER INFORMATION	23u	Discuss implications of the results for practice, policy, and future research	P21
Registration and protocol	2/12	Provide registration information for the review, including register name and	P2&P5
Registration and protocor	240	registration number, or state that the review was not registered	12015
	24h	Indicate where the review protocol can be accessed, or state that a protocol was not	P5
	-12	prepared	-
	24c	Describe and explain any amendments to information provided at registration or in	P28
		the protocol	
Support	25	Describe sources of financial or non-financial support for the review, and the role of	P28
-		the funders or sponsors in the review	
Competing interests	26	Declare any competing interests of review authors	P28

Availability of data, code and other materials

27 Report which of the following are publicly available and where they can be found: P6,7,11 Appendix 3, template data collection forms; data extracted from included studies; data used for 4, 5, 9 all analyses; analytic code; any other materials used in the review

S3 Search strategy

1) PUBMED

• Date of search: 16/02/2024

• Database searched from 1978 to February 2024

• Total results: 1106

I ODINIE	D 16/02/2024									
Search	Domain	Query	Hits							
#1	Construct	"Exercise"[MeSH Terms] OR "Circuit-Based Exercise"[MeSH Terms] OR "Exercise Test"[MeSH Terms]	814,485							
	Instruments	OR "Running"[MeSH Terms] OR "Isometric contraction"[MeSH Terms] OR "Walking"[MeSH Terms] OR								
		"Exercise therapy"[MeSH Terms] OR "cycling"[Title/Abstract] OR "Exercise test*"[Title/Abstract] OR								
		"Running"[Title/Abstract] OR "isometric contraction"[Title/Abstract] OR "exercise induced								
		hypoalgesia"[Title/Abstract] OR "EIH"[Title/Abstract] OR "exercise induced analgesia"[Title/Abstract]								
		OR "EIA"[Title/Abstract] OR "walk*"[Title/Abstract] OR "jogging"[Title/Abstract] OR "bicycle"[Title/								
		Abstract] OR "bicycling"[Title/Abstract] OR "rowing"[Title/Abstract] OR "swim*"[Title/Abstract] OR								
		"circuit training"[Title/Abstract] OR "bicycle exercise test*"[Title/Abstract] OR "kinesiotherapy"[Title/								
		Abstract] OR "muscle isometric contraction"[Title/Abstract] OR "aerobic exercise"[Title/Abstract] OR								
		"aerobic exercises"[Title/Abstract] OR "Exercise"[Title/Abstract] OR "Exercises"[Title/Abstract] OR								
	"aerobic exercises"[Title/Abstract] OR "Exercise"[Title/Abstract] OR "isometric exercises"[Title/Abstract] OR "isometric exercises"[Title/Abstract] OR "isotonic exercises"[Title/Abstract] OR "isotonic exercises"[Title/Abstract] OR "isotonic exercises"[Title/Abstract] OR "isokinetic exercises"[Title/Abstract] OR "isodynamic exercises"[Title/Abstract] OR "anaerobic exercises"[Title/Abstract] OR "anaerobic exercises"[Title/Abstract] OR "high intensity exercises"[Title/Abstract] OR "high intensity exercises"[Title/Abstract] OR "low intensity exercises"[Title/Abstract] OR "moderate intensity exercises"[Title/Abstract] OR "static exercise"[Title/Abstract] OR "stretching exercises"[Title/Abstract] OR "stretching exercises"[Title/Abstract] OR "stretching exercises"[Title/Abstract] OR "therapeutic exercises"[Title/Abstract] OR "therapeutic exercises"[Title/Abstract] OR "pressure pain thresholds"[Title/Abstract] OR "pressure threshold"[Title/Abstract] OR "pain pressure threshold"[Title/Abstract] OR "pain pressure thresholds"[Title/Abstract] OR "pain pressure threshold"[Title/Abstract] OR "pain pressure thresholds"[Title/Abstract] OR "pain pressure threshold"[Title/Abstract] OR "pain pressure threshold"[Title/Abstract] OR "pain pressure thresholds"[Title/Abstract] OR "pain pressure threshold"[Title/Abstract] OR "pain pressure threshold"[Title/Ab									
"aerobic exercises" [Title/Abstract] OR "Exe "isometric exercise" [Title/Abstract] OR "iso exercise" [Title/Abstract] OR "isotonic exercise" [Title/Abstract] OR "isokinetic exercises" [Title/Abstract] OR "Circuit-Based Exercises" [Title/Abstract] OR "Circuit-Based Exercises" [Title/Abstract] OR "anaerobic exercises" [Title/Abstract] OR "dynamic exercise" [Title/Abstract] OR "high exercise" [Title/Abstract] OR "low intensity exercise" [Title/Abstract] OR "moderate into Abstract] OR "static exercises" [Title/Abstract] OR "therapeutic exercises" [Title/Abstract] OR "therapeutic exercises" [Title/Abstract] OR "therapeutic exercises" [Title/Abstract] OR "pain puthresholds" [Title/Abstract] [Title/Abstract] OR "pain puth	exercise"[Title/Abstract] OR "isotonic exercises"[Title/Abstract] OR "isokinetic exercise"[Title/Abstract]									
		OR "isokinetic exercises"[Title/Abstract] OR "isodynamic exercise"[Title/Abstract] OR "isodynamic								
		exercises"[Title/Abstract] OR "Circuit-Based Exercise"[Title/Abstract] OR "Circuit-Based								
		Exercises"[Title/Abstract] OR "anaerobic exercise"[Title/Abstract] OR "anaerobic exercises"[Title/								
Exercises"[Title/Abstract] OR "anaerobic exercise"[Title/Abstract] OR "anaerobic exercises"[Title/Abstract] OR "dynamic exercises"[Title/Abstract] OR "dynamic exercises"[Title/Abstract] OR "high intensity exercises"[Title/Abstract] OR "high intensity exercises"[Title/Abstract] OR "low intensity exercises"[Title/Abstract] OR "moderate intensity exercises"[Title/Abstract] OR "moderate intensity exercises"[Title/Abstract] OR "static exercise"[Title/Abstract] OR "static exercise"[Title/Abstract] OR "stretching exercises"[Title/Abstract] OR "stretching exercises"[Title/Abstract] OR "stretching exercises"[Title/Abstract] OR "therapeutic exercises"[Title/Abstract] OR "therapeutic exercises"[Title/Abstract] OR "pressure pain threshold"[Title/Abstract] OR "pressure pain threshold"[Tit										
						exercise"[Title/Abstract] OR "moderate intensity exercises"[Title/Abstract] OR "static exercise"[Title/Abstract] OR "stretching exercise"[Title/Abstract] OR "stretching exercises"[Title/Abstract] OR "stretching exercises"[Title/Abstract] OR "therapeutic exercise"[Title/Abstract] OR "therapeutic exercises"[Title/Abstract] OR "therapeutic exercises"[Title/Abstract] OR "pressure pain threshold"[Title/Abstract] OR "pressure pa				
								exercises"[Title/Abstract] OR "therapeutic exercise"[Title/Abstract] OR "therapeutic exercises"[Title/		
								Abstract]	tle/Abstract] OR "stretching nerapeutic exercises"[Title/ itle/Abstract] OR "pressure 99,760	
exercise"[Title/Abstract] OR "moderate intensity exercises"[Title/Abstract] OR "static exercise"[Title Abstract] OR "static exercises"[Title Abstract] OR "stretching exercises"[Title Abstract] OR "stretching exercises"[Title Abstract] OR "therapeutic exercises"[Title Abstract] OR "therapeutic exercises"[Title Abstract] OR "therapeutic exercises"[Title Abstract] OR "pressure pain threshold"[Title Abstract] OR "pressure pain threshold"[Title Abstract] OR "pain pressure threshold"[Title Abstract] OR "pain pressure threshold"[Title Abstract] OR "pain threshold"[Title										
#2	Instruments	pain thresholds"[Title/Abstract] OR "pain pressure threshold"[Title/Abstract] OR "pain pressure								
		thresholds"[Title/Abstract] OR "pain threshold"[Title/Abstract] OR "heat temporal summation"[Title/								
		Abstract] OR "cold tolerance"[Title/Abstract] OR ("hydraul*"[Title/Abstract] AND ("algomet*"[Title/								
		Abstract] OR "pain ratings"[Title/Abstract])) OR "nociceptive flexion reflex"[Title/Abstract] OR "elec-								
		trical pain ratings"[Title/Abstract] OR "heat pain ratings"[Title/Abstract] OR "pressure pain								
		ratings"[Title/Abstract] OR "cold pain ratings"[Title/Abstract] OR "heat discrimination"[Title/Abstract]								
		OR "pain tolerance"[Title/Abstract] OR "pressure tolerance thresholds"[Title/Abstract] OR "pressure								
		tolerance threshold"[Title/Abstract] OR "experimental pain"[Title/Abstract] OR "pain perception"[Title/								
		Abstract] OR "electrical pain threshold"[Title/Abstract] OR "electrical pain thresholds"[Title/Abstract]								
		OR "perception threshold"[Title/Abstract] OR "perception thresholds"[Title/Abstract] OR "pain								
		thresholds"[Title/Abstract] OR "heat pain thresholds"[Title/Abstract] OR "pain threshold								

VIII — M. Vladimir Aron et al. DE GRUYTER

temperature"[Title/Abstract] OR "pain thresholds temperature"[Title/Abstract] OR "heat pain threshold"[Title/Abstract] OR "perceptive threshold"[Title/Abstract] OR "perceptive thresholds"[Title/Abstract] Abstract] OR "sensory threshold" [Title/Abstract] OR "sensory thresholds" [Title/Abstract] OR "sense thresholds"[Title/Abstract] OR "nociception threshold"[Title/Abstract] OR "nociception thresholds"[Title/Abstract] OR "nociceptive threshold"[Title/Abstract] OR "nociceptive thresholds"[Title/Abstract] OR "nociceptive pressure threshold"[Title/Abstract] OR "nociceptive pressure thresholds"[Title/Abstract] OR "pressure nociceptive threshold"[Title/Abstract] OR "pressure nociceptive thresholds"[Title/Abstract] OR "endogenous pain modulation"[Title/Abstract] OR "temporal summation"[Title/Abstract] OR "pain modulation"[Title/Abstract] OR "nociceptive withdrawal reflex"[Title/Abstract] OR "pain tolerance threshold"[Title/Abstract] OR "pain tolerance thresholds"[Title/Abstract] OR "pain inhibition"[Title/Abstract] OR "pain inhibitory process"[Title/ Abstract] OR "pain inhibitor" [Title/Abstract] OR "hypoalgesia" [Title/Abstract] OR "reduced pain sensitivity" [Title/Abstract] OR "hyposensitivity to pain" [Title/Abstract] OR "nociception" [Title/ Abstract] OR "pain sensation" [Title/Abstract] OR "pain sensitivity" [Title/Abstract] OR "quantitative sensory testing"[Title/Abstract] OR "quantitative sensation testing"[Title/Abstract] OR "quantitative sensory test"[Title/Abstract] OR "Sensory Thresholds"[MeSH Terms] OR "Pain Threshold"[MeSH Terms] OR "Nociception" [MeSH Terms] OR "Pain Perception" [MeSH Terms]

Population

#3

Terms] OR "Nociception" [MeSH Terms] OR "Pain Perception" [MeSH Terms]

"Rat" [Title] OR "Rats" [Title] OR "Mouse" [Title] OR "Mice" [Title] OR "Horse" [Title] OR "Horses" [Title] OR 2,801,217

"Gorilla" [Title] OR "Animal" [Title] OR "Animals" [Title] OR "manipulation" [Title] OR

"anaphylaxis" [Title] OR "acupuncture" [Title] OR "pigeons" [Title] OR "asthma" [Title] OR "plant" [Title]

OR "Deer" [Title] OR "children" [Title] OR "child" [Title] OR "heart disease" [Title] OR "coronary disease" [Title] OR "myocardial ischemia" [Title] OR "asthmatic" [Title] OR "rodent" [Title] OR "rodents" [Title] OR "Angina" [Title] OR "coronary artery disease" [Title] OR "paraplegics" [Title] OR "spinal cord lesion" [Title] OR "asthmatics" [Title] OR "chronic heart failure" [Title] OR "hypoxemia" [Title]

#4 Reliab

"psychometrics"[MeSH] OR psychometr*[tiab] OR clinimetr*[tw] OR clinometr*[tw] OR "observer" 4,173,736 variation"[MeSH] OR "observer variation"[tiab] OR "reproducibility of results"[MeSH] OR reproducib* [tiab] OR"agreement"[Title/Abstract] OR "precision"[Title/Abstract] OR "imprecision"[Title/Abstract] OR "precise values" [Title/Abstract] OR "test-retest" [Title/Abstract] OR ("test" [Title/Abstract] AND "retest"[Title/Abstract]) OR ("reliab*"[Title/Abstract] AND ("test"[Title/Abstract] OR "retest"[Title/ Abstract])) OR "stability"[Title/Abstract] OR "interrater"[Title/Abstract] OR "inter-rater"[Title/Abstract] OR "intrarater"[Title/Abstract] OR "intra-rater"[Title/Abstract] OR "intertester"[Title/Abstract] OR "inter-tester"[Title/Abstract] OR "intratester"[Title/Abstract] OR "intra-tester"[Title/Abstract] OR "interobserver"[Title/Abstract] OR "inter-observer"[Title/Abstract] OR "intraobserver"[Title/Abstract] OR "intra-observer"[Title/Abstract] OR "intertechnician"[Title/Abstract] OR "inter-technician"[Title/Abstract] Abstract] OR "intratechnician" [Title/Abstract] OR "intra-technician" [Title/Abstract] OR "interexaminer"[Title/Abstract] OR "inter-examiner"[Title/Abstract] OR "intraexaminer"[Title/ Abstract] OR "intra-examiner" [Title/Abstract] OR "interassay" [Title/Abstract] OR "inter-assay" [Title/Abst Abstract] OR "intraassay" [Title/Abstract] OR "intra-assay" [Title/Abstract] OR "interindividual" [Title/Abstract] OR "intraassay" [Abstract] OR "inter-individual"[Title/Abstract] OR "intraindividual"[Title/Abstract] OR "intraindividual"[Title/Abstract] OR "interparticipant"[Title/Abstract] OR "inter-participant"[Title/Abstract] OR "intraparticipant"[Title/Abstract] OR "intra-participant"[Title/Abstract] OR "kappa"[Title/Abstract] OR "kappas"[Title/Abstract] OR "kappas"[Title/Abstract] OR "repeatab*"[Title/Abstract] OR (("replicab*"[Title/Abstract] OR "repeated"[Title/Abstract]) AND ("measure"[Title/Abstract] OR "measures"[Title/Abstract] OR "findings"[Title/Abstract] OR "result"[Title/Abstract] OR "results"[Title/ Abstract] OR "test"[Title/Abstract] OR "tests"[Title/Abstract])) OR "generaliza*"[Title/Abstract] OR "generalisa*"[Title/Abstract] OR "concordance"[Title/Abstract] OR ("intraclass"[Title/Abstract] AND "correlation*"[Title/Abstract]) OR "discriminative"[Title/Abstract] OR "known group"[Title/Abstract] OR "factor analysis" [Title/Abstract] OR "factor analyses" [Title/Abstract] OR "dimensionality" [Title/Abstract] Abstract] OR "subscale*"[Title/Abstract] OR "multitrait scaling analysis"[Title/Abstract] OR "multitrait

scaling analyses"[Title/Abstract] OR "item discriminant"[Title/Abstract] OR "interscale correlation"[Title/Abstract] OR "interscale correlations"[Title/Abstract] OR (("error"[Title/Abstract] OR "errors"[Title/Abstract]) AND ("measure*"[Title/Abstract] OR "correlat*"[Title/Abstract] OR "evaluat*"[Title/Abstract] OR "accuracy"[Title/Abstract] OR "accurate"[Title/Abstract] OR "precision"[Title/Abstract] OR "mean"[Title/Abstract])) OR "individual variability"[Title/Abstract] OR "variability analysis" [Title/Abstract] OR ("uncertainty" [Title/Abstract] AND ("measurement" [Title/ Abstract] OR "measuring"[Title/Abstract])) OR "standard error of measurement"[Title/Abstract] OR "sensitiv*"[Title/Abstract] OR "responsive*"[Title/Abstract] OR ("small*"[Title/Abstract] AND ("real"[Title/Abstract] OR "detectable"[Title/Abstract]) AND ("change"[Title/Abstract] OR "difference"[Title/Abstract])) OR "meaningful change"[Title/Abstract] OR "minimal important change"[Title/Abstract] OR "minimal important difference"[Title/Abstract] OR "minimally important change"[Title/Abstract] OR "minimally important difference"[Title/Abstract] OR "minimal detectable change"[Title/Abstract] OR "minimal detectable difference"[Title/Abstract] OR "minimally detectable change"[Title/Abstract] OR "minimally detectable difference"[Title/Abstract] OR "minimal real difference"[Title/Abstract] OR "ceiling effect"[Title/Abstract] OR "floor effect"[Title/Abstract] OR "g theory"[Title/Abstract] OR "generalizability theory"[Title/Abstract] OR "generalisability theory"[Title/ Abstract] OR "equivalence"[Title/Abstract] #1 AND #2 AND #4 NOT #3 1.106

2) EMBASE

#5

• Date of search: 15/02/2024

Database searched from 1974 to February 2024

• Total results: 1,339

Search	Domain	Query	Hits
#1	Construct	1. exp leg exercise/ or exp aquatic exercise/ or exp isotonic exercise/ or exp stretching exercise/ or exp	647,379
	Instruments	anaerobic exercise/ or exp low intensity exercise/ or exp aerobic exercise/ or exp isometric exercise/ or	
		exp treadmill exercise/ or exp dynamic exercise/ or exp exercise/ or exp "squatting (exercise)"/ or exp	
		static exercise/ or exp open kinetic chain exercise/ or exp arm exercise/ or exp muscle exercise/ or exp	
		isokinetic exercise/ or exercise.mp. or exp moderate intensity exercise/	
		2. (cycling or exercise test* or running or isometric contraction or exercise induced hypoalgesia or eih	866,216
		or exercise induced analgesia or eia or walk* or jogging or bicycle or bicycling or rowing or swim* or	
		circuit training or bicycle exercise test* or kinesiotherapy or muscle isometric contraction or aerobic	
		exercise or aerobic exercises or exercise or exercises or isometric exercise or isometric exercises or	
		isotonic exercise or isotonic exercises or isokinetic exercise or isokinetic exercises or isodynamic	
		exercise or isodynamic exercises or circuit-cased exercise or circuit-based exercises or anaerobic	
		exercise or anaerobic exercises or dynamic exercise or dynamic exercises or high intensity exercise or	
		high intensity exercises or low intensity exercise or low intensity exercises or moderate intensity	
		exercise or moderate intensity exercises or static exercise or static exercises or stretching exercise or	
		stretching exercises or therapeutic exercise or therapeutic exercises).ab,ti.	
		1 or 2	1,041,442
#2	Construct	3 pain threshold.mp. or exp pain threshold/	26,358
	Instruments	4 exp heat pain threshold/	960
		5 exp nociception/	89,741
		6 exp nociceptive stimulation/	7,311

62,006

7 (nociceptive stimulation or pressure pain threshold or pressure pain thresholds or pain pressure threshold or pain pressure thresholds or pain threshold or heat temporal summation or cold tolerance or (hydraul* and (algomet* or pain ratings)) or nociceptive flexion reflex or electrical pain ratings or heat pain ratings or pressure pain ratings or cold pain ratings or heat discrimination or pain tolerance or pressure tolerance thresholds or pressure tolerance threshold or experimental pain or pain perception or electrical pain threshold or electrical pain thresholds or perception threshold or perception thresholds or pain thresholds or pain thresholds or pain threshold temperature or pain thresholds temperature or heat pain threshold or perceptive threshold or perceptive thresholds or sensory threshold or sensory thresholds or sense thresholds or nociception threshold or nociception thresholds or nociceptive threshold or nociceptive thresholds or nociceptive pressure threshold or nociceptive pressure thresholds or pressure nociceptive threshold or pressure nociceptive thresholds or endogenous pain modulation or temporal summation or pain modulation or nociceptive withdrawal reflex or pain tolerance threshold or pain tolerance thresholds or pain inhibition or pain inhibitory process or pain inhibitor or hypoalgesia or reduced pain sensitivity or hyposensitivity to pain or nociception or pain sensation or pain sensitivity or quantitative sensory testing or quantitative sensation testing or quantitative sensory test).ab,ti.

19 3 or 4 or 5 or 6 or 7

113,083 3,132,888

#3 Population

(rat or rats or mouse or mice or horse or horses or gorilla or animal or animals or manipulation or anaphylaxis or acupuncture or pigeons or asthma or plant or deer or children or child or heart disease or coronary disease or myocardial ischemia or asthmatic or rodent or rodents or angina or coronary artery disease or paraplegics or spinal cord lesion or asthmatics or chronic heart failure or hypoxemia).ti.

#4 Reliab

10 exp observer variation/ 21,102
11 exp reproducibility/ 265,438
12 exp test retest reliability/ 24545
13 exp intrarater reliability/ or exp interrater reliability/ or exp reliability/ 243105
14 exp minimum detectable change/ 1,135
15 exp correlation coefficient/ 117,454
16 exp minimal clinically important difference/ 4,848

17 (equivalence or psychometrics or psychometr* or clinimetr* or clinometr* or observer variation or 4,842,515 reproducib* or agreement or precision or imprecision or precise values or test-retest or stability or interrater or inter-rater or intrarater or intra-rater or intertester or inter-tester or intratester or intratester or interobserver or inter-observer or intra-observer or intra-observer or intertechnician or intertechnician or intratechnician or intra-technician or interexaminer or inter-examiner or intraexaminer or intra-examiner or interassay or inter-assay or intra-assay or intra-assay or interindividual or interindividual or intraindividual or intra-individual or interparticipant or inter-participant or intraparticipant or intra-participant or kappa or kappas or repeatab* or generaliza* or generalisa* or concordance or discriminative or known group or factor analysis or factor analyses or dimensionality or subscale* or multitrait scaling analysis or multitrait scaling analyses or item discriminant or interscale correlation or interscale correlations or individual variability or variability analysis or standard error of measurement or sensitiv* or responsive* or meaningful change or minimal important change or minimal important difference or minimally important change or minimally important difference or minimal detectable change or minimal detectable difference or minimally detectable change or minimally detectable difference or minimal real difference or ceiling effect or floor effect or g theory or generalizability theory or generalisability theory or (reliab* and (test or retest)) or ((replicab* or repeated) and (measure or measures or findings or result or results or test or tests)) or (intraclass and correlation*) or ((error or errors) and (measure* or correlat* or evaluat* or accuracy or accurate or precision or mean)) or (uncertainty and (measurement or measuring)) or (small* and (real or detectable) and (change or difference))).ab,ti.

 $20\ 9$ or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17

5,136,999

#1 AND #2 AND #4 NOT #3

1,339

3) SCOPUS

• Date of search: 16/02/2024

• Database searched from 1966 to February 2024

• Total results: 1509

SCOPUS	: 16/02/2024		
Search	Domain	Query	Hits
#1	Construct Instruments	TITLE-ABS-KEY("cycling" OR "exercise test*" OR "running" OR {isometric contraction} OR "exercise induced hypoalgesia" OR {EIH} OR "exercise induced analgesia" OR {EIA} OR "walk*" OR "jogging" OR "bicycle" OR "bicycling" OR "rowing" OR "swim*" OR {circuit training} OR "bicycle exercise test*" OR {kinesiotherapy} OR {muscle isometric contraction} OR {aerobic exercise} OR {aerobic exercises} OR {exercise} OR {isometric exercise} OR {isotonic exercise} OR {isotonic exercise} OR {isotonic exercise} OR {isotonic exercise} OR {isodynamic exercise} OR {isodynamic exercise} OR {isodynamic exercise} OR {isodynamic exercise} OR {anaerobic exercise} OR {dynamic exercise} OR {dynamic exercise} OR {high intensity exercise} OR {high intensity exercise} OR {low intensity exercise} OR {moderate intensity exercise} OR {static exercise} OR {therapeutic exercise} OR {therapeuti	
#2	Construct Instruments	TITLE-ABS-KEY({nociceptive stimulation} OR {pressure pain threshold} OR {pressure pain thresholds} OR {pain pressure threshold} OR {pain pressure threshold} OR {pain ratings})) OR {nociceptive summation} OR {cold tolerance} OR ("hydraul*" AND ("algomet*" OR {pain ratings})) OR {nociceptive flexion reflex} OR {electrical pain ratings} OR {heat pain ratings} OR {pressure pain ratings} OR {cold pain ratings} OR {heat discrimination} OR {pain tolerance} OR {pressure tolerance threshold} OR {pressure tolerance threshold} OR {experimental pain} OR {pain perception} OR {electrical pain threshold} OR {perception threshold} OR {pain threshold} OR {perceptive threshold} OR {perceptive threshold} OR {perceptive threshold} OR {nociceptive threshold} OR {nociceptive threshold} OR {nociceptive threshold} OR {nociceptive threshold} OR {pressure nociceptive threshold} OR {pain modulation} OR {pain modulation} OR {pain tolerance threshold} OR {pain inhibition} OR {pain inhibitor} OR {pain inhibitor} OR {pain sensitivity} OR {pain sensitivity} OR {pain sensitivity} OR {pain sensitivity to pain} OR {quantitative sensory testing} OR {quantitative sensory testing	
#3	Population	TITLE("rat" OR "rats" OR "mouse" OR "mice" OR "horse" OR "horses" OR "gorilla" OR "animal" OR "animals" OR "manipulation" OR "anaphylaxis" OR "acupuncture" OR "pigeons" OR "asthma" OR "plant" OR "deer" OR "children" OR "child" OR {heart disease} OR {coronary disease} OR {myocardial ischemia} OR "asthmatic" OR "rodent" OR "rodents" OR "angina" OR {coronary artery disease} OR "paraplegics" OR {spinal cord lesion} OR "asthmatics" OR {chronic heart failure} OR "hypoxemia")	4,023,412
#4	Reliab	TITLE-ABS-KEY("equivalence" OR "psychometrics" OR "psychometr*" OR "clinimetr*" OR "clinometr*" OR {observer variation} OR "reproducib*" OR "agreement" OR "precision" OR "imprecision" OR "precise values" OR {test-retest} OR "stability" OR "interrater" OR {inter-rater} OR "intrarater" OR {intra-rater} OR "intertester" OR {intra-tester} OR "interobserver" OR {inter-observer} OR "interobserver" OR {inter-observer} OR "intertechnician" OR {inter-observer} OR "inter-observer] OR "intertechnician" OR {inter-observer} OR "intertec	12,399,519

XII — M. Vladimir Aron et al. DE GRUYTER

technician} OR "intratechnician" OR {intra-technician} OR "inter-examiner" OR "inter-examiner" OR "intraexaminer" OR "intra-examiner" OR "interassay" OR "inter-assay" OR "intraassay" OR [intraassay} OR "interindividual" OR {inter-individual} OR "intraindividual" OR {intra-individual} OR "interparticipant" OR {inter-participant} OR "intraparticipant" OR {intra-participant} OR "kappa" OR "kappas" OR {kappas} OR "repeatab*" OR "generaliza*" OR "generalisa*" OR "concordance" OR "discriminative" OR {known group} OR {factor analysis} OR {factor analyses} OR "dimensionality" OR "subscale*" OR {multitrait scaling analysis} OR {multitrait scaling analyses} OR {item discriminant} OR (interscale correlation) OR (interscale correlations) OR (individual variability) OR (variability analysis} OR {standard error of measurement} OR "sensitiv*" OR "responsive*" OR {meaningful change) OR (minimal important change) OR (minimal important difference) OR (minimally important change} OR {minimally important difference} OR {minimal detectable change} OR {minimal detectable difference} OR {minimally detectable change} OR {minimally detectable difference} OR {minimal real difference} OR {ceiling effect} OR {floor effect} OR {g theory} OR {generalizability theory} OR {generalisability theory} OR ("reliab*" AND ("test" OR "retest")) OR (("replicab*" OR "repeated") AND ("measure" OR "measures" OR "findings" OR "result" OR "results" OR "test" OR "tests")) OR ("intraclass" AND "correlation*") OR (("error" OR "errors") AND ("measure*" OR "correlat*" OR "evaluat*" OR "accuracy" OR "accurate" OR "precision" OR "mean")) OR ("uncertainty" AND ("measurement" OR "measuring")) OR ("small*" AND ("real" OR "detectable") AND ("change" OR "difference")))

#5 #1 AND #2 AND #4 AND NOT #3 1,509

4) CINAHL Complete

• Date of search: 16/02/2024

• Database searched from 1983 to February 2024

Total results: 572

CINAHL	: 16/02/2024		
Search	Domain	Query	Hits
#1	Construct	((MH "Exercise+") OR (MH "Therapeutic Exercise+") OR (MH "Exercise Test+") OR (MH "Aerobic	298,624
	Instruments	Exercises+") OR (MH "Running+") OR (MH "Walking+") OR (MM "Isokinetic Exercises") OR (MM	
		"Isometric Exercises") OR (MM "Isotonic Exercises") OR (MM "Isometric Contraction")) OR TI ("cycling"	
		OR "Exercise test*" OR "Running" OR "isometric contraction" OR "exercise induced hypoalgesia" OR	
		"EIH" OR "exercise induced analgesia" OR "EIA" OR "walk*" OR "jogging" OR "bicycle" OR "bicycling"	
		OR "rowing" OR "swim*" OR "circuit training" OR "bicycle exercise test*" OR "kinesiotherapy" OR	
		"muscle isometric contraction" OR "aerobic exercise" OR "aerobic exercises" OR "Exercise" OR	
		"Exercises" OR "isometric exercise" OR "isometric exercises" OR "isotonic exercise" OR "isotonic	
		exercises" OR "isokinetic exercise" OR "isokinetic exercises" OR "isodynamic exercise" OR "isodynamic	
		exercises" OR "Circuit-Based Exercise" OR "Circuit-Based Exercises" OR "anaerobic exercise" OR	
		"anaerobic exercises" OR "dynamic exercise" OR "dynamic exercises" OR "high intensity exercise" OR	
		"high intensity exercises" OR "low intensity exercise" OR "low intensity exercises" OR "moderate	
		intensity exercise" OR "moderate intensity exercises" OR "static exercise" OR "static exercises" OR	
		"stretching exercise" OR "stretching exercises" OR "therapeutic exercise" OR "therapeutic exercises")	
		OR AB ("cycling" OR "Exercise test*" OR "Running" OR "isometric contraction" OR "exercise induced	
		hypoalgesia" OR "EIH" OR "exercise induced analgesia" OR "EIA" OR "walk*" OR "jogging" OR "bicycle"	
		OR "bicycling" OR "rowing" OR "swim*" OR "circuit training" OR "bicycle exercise test*" OR "kine-	
		siotherapy" OR "muscle isometric contraction" OR "aerobic exercise" OR "aerobic exercises" OR	
		"Exercise" OR "Exercises" OR "isometric exercise" OR "isometric exercises" OR "isotonic exercise" OR	

"isotonic exercises" OR "isokinetic exercise" OR "isokinetic exercises" OR "isodynamic exercise" OR "isodynamic exercises" OR "Circuit-Based Exercise" OR "Circuit-Based Exercises" OR "anaerobic exercise" OR "anaerobic exercises" OR "dynamic exercise" OR "dynamic exercises" OR "high intensity exercise" OR "high intensity exercises" OR "low intensity exercise" OR "low intensity exercises" OR "moderate intensity exercise" OR "moderate intensity exercises" OR "static exercise" OR "static exercises" OR "stretching exercise" OR "stretching exercises" OR "therapeutic exercise" OR "therapeutic exercises")

#2 Construct Instruments (MM "Pain Threshold") OR TI ("nociceptive stimulation" OR "pressure pain threshold" OR "pressure 13.621 pain thresholds" OR "pain pressure threshold" OR "pain pressure thresholds" OR "pain threshold" OR "heat temporal summation" OR "cold tolerance" OR ("hydraul*" AND ("algomet*" OR "pain ratings")) OR "nociceptive flexion reflex" OR "electrical pain ratings" OR "heat pain ratings" OR "pressure pain ratings" OR "cold pain ratings" OR "heat discrimination" OR "pain tolerance" OR "pressure tolerance thresholds" OR "pressure tolerance threshold" OR "experimental pain" OR "pain perception" OR "electrical pain threshold" OR "electrical pain thresholds" OR "perception threshold" OR "perception thresholds" OR "pain thresholds" OR "heat pain thresholds" OR "pain threshold temperature" OR "pain thresholds temperature" OR "heat pain threshold" OR "perceptive threshold" OR "perceptive thresholds" OR "sensory threshold" OR "sensory thresholds" OR "sense thresholds" OR "nociception threshold" OR "nociception thresholds" OR "nociceptive threshold" OR "nociceptive thresholds" OR "nociceptive pressure threshold" OR "nociceptive pressure thresholds" OR "pressure nociceptive threshold" OR "pressure nociceptive thresholds" OR "endogenous pain modulation" OR "temporal summation" OR "pain modulation" OR "nociceptive withdrawal reflex" OR "pain tolerance threshold" OR "pain tolerance thresholds" OR "pain inhibition" OR "pain inhibitory process" OR "pain inhibitor" OR "hypoalgesia" OR "reduced pain sensitivity" OR "hyposensitivity to pain" OR "nociception" OR "pain sensation" OR "pain sensitivity" OR "quantitative sensory testing" OR "quantitative sensation testing" OR "quantitative sensory test") OR AB ("nociceptive stimulation" OR "pressure pain threshold" OR "pressure pain thresholds" OR "pain pressure threshold" OR "pain pressure thresholds" OR "pain threshold" OR "heat temporal summation" OR "cold tolerance" OR ("hydraul*" AND ("algomet*" OR "pain ratings")) OR "nociceptive flexion reflex" OR "electrical pain ratings" OR "heat pain ratings" OR "pressure pain ratings" OR "cold pain ratings" OR "heat discrimination" OR "pain tolerance" OR "pressure tolerance thresholds" OR "pressure tolerance threshold" OR "experimental pain" OR "pain" perception" OR "electrical pain threshold" OR "electrical pain thresholds" OR "perception threshold" OR "perception thresholds" OR "pain thresholds" OR "heat pain thresholds" OR "pain threshold temperature" OR "pain thresholds temperature" OR "heat pain threshold" OR "perceptive threshold" OR "perceptive thresholds" OR "sensory threshold" OR "sensory thresholds" OR "sense thresholds" OR "nociception threshold" OR "nociception thresholds" OR "nociceptive threshold" OR "nociceptive thresholds" OR "nociceptive pressure threshold" OR "nociceptive pressure thresholds" OR "pressure nociceptive threshold" OR "pressure nociceptive thresholds" OR "endogenous pain modulation" OR "temporal summation" OR "pain modulation" OR "nociceptive withdrawal reflex" OR "pain tolerance threshold" OR "pain tolerance thresholds" OR "pain inhibition" OR "pain inhibitory process" OR "pain inhibitor" OR "hypoalgesia" OR "reduced pain sensitivity" OR "hyposensitivity to pain" OR "nociception" OR "pain sensation" OR "pain sensitivity" OR "quantitative sensory testing" OR "quantitative sensation testing" OR "quantitative sensory test")

#3 **Population** TI ("rat" OR "rats" OR "mouse" OR "mice" OR "horse" OR "horses" OR "gorilla" OR "animal" OR "animals" 460,209 OR "manipulation" OR "anaphylaxis" OR "acupuncture" OR "pigeons" OR "asthma" OR "plant" OR "deer" OR "children" OR "child" OR "heart disease" OR "coronary disease" OR "myocardial ischemia" OR "asthmatic" OR "rodent" OR "rodents" OR "angina" OR "coronary artery disease" OR "paraplegics" OR "spinal cord lesion" OR "asthmatics" OR "chronic heart failure" OR "hypoxemia") OR TI ("rat" OR "rats" OR "mouse" OR "mice" OR "horse" OR "horses" OR "gorilla" OR "animal" OR "animals" OR "manipulation" OR "anaphylaxis" OR "acupuncture" OR "pigeons" OR "asthma" OR "plant" OR "deer" OR "children" OR "child" OR "heart disease" OR "coronary disease" OR "myocardial ischemia" OR

Reliab

#4

#5

"spinal cord lesion" OR "asthmatics" OR "chronic heart failure" OR "hypoxemia") ((MM "Psychometrics") OR (MH "Reliability+") OR (MH "Stability+") OR (MM "Test-Retest Reliability") 1,439,006 OR (MH "Equivalence+") OR (MM "Interrater Reliability") OR (MM "Intrarater Reliability") OR (MM "Precision") OR (MH "Measurement Error+") OR (MH "Reproducibility of Results")) OR TI ("equivalence" OR "psychometrics" OR "psychometr*" OR "clinimetr*" OR "clinometr*" OR "observer variation" OR "reproducib*" OR "agreement" OR "precision" OR "imprecision" OR "precise values" OR "test-retest" OR "stability" OR "interrater" OR "inter-rater" OR "intra-rater" OR "intra-rater" OR "inter-rater" OR "inter-tester" OR "intratester" OR "intra-tester" OR "interobserver" OR "inter-observer" OR "intraobserver" OR "intra-observer" OR "intertechnician" OR "inter-technician" OR "intratechnician" OR "intratechnician" OR "interexaminer" OR "inter-examiner" OR "intraexaminer" OR "intra-examiner" OR "interassay" OR "inter-assay" OR "intraassay" OR "intra-assay" OR "interindividual" OR "inter-individual" OR "intraindividual" OR "intra-individual" OR "interparticipant" OR "inter-participant" OR "intraparticipant" OR "intra-participant" OR "kappa" OR "kappas" OR "repeatab*" OR "generaliza*" OR generalisa*" OR "concordance" OR "discriminative" OR "known group" OR "factor analysis" OR "factor" analyses" OR "dimensionality" OR "subscale*" OR "multitrait scaling analysis" OR "multitrait scaling analyses" OR "item discriminant" OR "interscale correlation" OR "interscale correlations" OR "individual variability" OR "variability analysis" OR "standard error of measurement" OR "sensitiv*" OR "responsive*" OR "meaningful change" OR "minimal important change" OR "minimal important difference" OR "minimally important change" OR "minimally important difference" OR "minimal detectable change" OR "minimal detectable difference" OR "minimally detectable change" OR "minimally detectable difference" OR "minimal real difference" OR "ceiling effect" OR "floor effect" OR "g theory" OR "generalizability theory" OR "generalisability theory" OR ("reliab*" AND ("test" OR "retest")) OR (("replicab*" OR "repeated") AND ("measure" OR "measures" OR "findings" OR "result" OR "results" OR "test" OR "tests")) OR ("intraclass" AND "correlation*") OR (("error" OR "errors") AND ("measure*" OR "correlat*" OR "evaluat*" OR "accuracy" OR "accurate" OR "precision" OR "mean")) OR ("uncertainty" AND ("measurement" OR "measuring")) OR ("small*" AND ("real" OR "detectable") AND ("change" OR "difference")) OR AB ("equivalence" OR "psychometrics" OR "psychometr*" OR "clinimetr*" OR "clinometr*" OR "observer variation" OR "reproducib*" OR "agreement" OR "precision" OR "imprecision" OR "precise values" OR "test-retest" OR "stability" OR "interrater" OR "inter-rater" OR "intrarater" OR "intra-rater" OR "intertester" OR "inter-tester" OR "intratester" OR "intra-tester" OR "interobserver" OR "inter-observer" OR "intraobserver" OR "intra-observer" OR "intertechnician" OR "inter-technician" OR "intratechnician" OR "intra-technician" OR "interexaminer" OR "inter-examiner" OR "intraexaminer" OR "intra-examiner" OR "interassay" OR "inter-assay" OR "intraassay" OR "intraassay" OR "interindividual" OR "inter-individual" OR "intraindividual" OR "intra-individual" OR "interparticipant" OR "inter-participant" OR "intraparticipant" OR "intra-participant" OR "kappa" OR kappas" OR "repeatab*" OR "generaliza*" OR "generalisa*" OR "concordance" OR "discriminative" OR" "known group" OR "factor analysis" OR "factor analyses" OR "dimensionality" OR "subscale*" OR "multitrait scaling analysis" OR "multitrait scaling analyses" OR "item discriminant" OR "interscale correlation" OR "interscale correlations" OR "individual variability" OR "variability analysis" OR "standard error of measurement" OR "sensitiv*" OR "responsive*" OR "meaningful change" OR "minimal important change" OR "minimal important difference" OR "minimally important change" OR "minimally important difference" OR "minimal detectable change" OR "minimal detectable difference" OR "minimally detectable change" OR "minimally detectable difference" OR "minimal real difference" OR "ceiling effect" OR "floor effect" OR "g theory" OR "generalizability theory" OR "generalisability theory" OR ("reliab*" AND ("test" OR "retest")) OR (("replicab*" OR "repeated") AND ("measure" OR "measures" OR "findings" OR "result" OR "results" OR "test" OR "tests")) OR ("intraclass" AND "correlation*") OR (("error" OR "errors") AND ("measure*" OR "correlat*" OR "evaluat*" OR "accuracy" OR "accurate" OR "precision" OR "mean")) OR ("uncertainty" AND ("measurement" OR "measuring")) OR ("small*" AND ("real" OR "detectable") AND ("change" OR "difference"))

"asthmatic" OR "rodent" OR "rodents" OR "angina" OR "coronary artery disease" OR "paraplegics" OR

#1 AND #2 AND #4 NOT #3 572

5) PSYCINFO

• Date of search: 16/02/2024

· Database searched from 1932 to February 2024

Total results: 236

PSYCHINFO:	16/02/2024
------------	------------

Search Domain Query Hits

#1 Construct Instruments (MJMAINSUBJECT.EXACT("Aerobic Exercise") OR MJMAINSUBJECT.EXACT("Exercise") OR MJMAINSUB- 133,216 JECT.EXACT("Walking")) OR ab("cycling" OR "Exercise test*" OR "Running" OR "isometric contraction" OR "exercise induced hypoalgesia" OR "EIH" OR "exercise induced analgesia" OR "EIA" OR "walk*" OR "jogging" OR "bicycle" OR "bicycling" OR "rowing" OR "swim*" OR "circuit training" OR "bicycle exercise test*" OR "kinesiotherapy" OR "muscle isometric contraction" OR "aerobic exercise" OR "aerobic exercises" OR "Exercise" OR "Exercises" OR "isometric exercise" OR "isometric exercises" OR "isometr exercise" OR "isotonic exercises" OR "isokinetic exercise" OR "isokinetic exercises" OR "isokynamic exercise" OR "isodynamic exercises" OR "Circuit-Based Exercise" OR "Circuit-Based Exercises" OR "anaerobic exercise" OR "anaerobic exercises" OR "dynamic exercise" OR "dynamic exercises" OR "high intensity exercise" OR "high intensity exercises" OR "low intensity exercise" OR "low intensity exercises" OR "moderate intensity exercise" OR "moderate intensity exercises" OR "static exercise" OR "static exercises" OR "stretching exercise" OR "stretching exercises" OR "therapeutic exercise" OR "therapeutic exercises") OR ti("cycling" OR "Exercise test*" OR "Running" OR "isometric contraction" OR "exercise induced hypoalgesia" OR "EIH" OR "exercise induced analgesia" OR "EIA" OR "walk*" OR "jogging" OR "bicycle" OR "bicycling" OR "rowing" OR "swim*" OR "circuit training" OR "bicycle exercise test*" OR "kinesiotherapy" OR "muscle isometric contraction" OR "aerobic exercise" OR "aerobic exercises" OR "Exercise" OR "Exercises" OR "isometric exercise" OR "isometric exercises" OR "isotonic exercise" OR "isotonic exercises" OR "isokinetic exercise" OR "isokinetic exercises" OR "isodynamic exercise" OR "isodynamic exercises" OR "Circuit-Based Exercise" OR "Circuit-Based Exercises" OR "anaerobic exercise" OR "anaerobic exercises" OR "dynamic exercise" OR "dynamic exercises" OR "high intensity exercise" OR "high intensity exercises" OR "low intensity exercise" OR "low intensity exercises" OR "moderate intensity exercise" OR "moderate intensity exercises" OR "static exercise" OR "static exercises" OR "stretching exercise" OR "stretching exercises" OR "therapeutic exercise" OR "therapeutic exercises")

#2 Construct Instruments (MJMAINSUBJECT.EXACT("Pain Thresholds") OR MJMAINSUBJECT.EXACT("Pain Perception")) OR ab ("nociceptive stimulation" OR "pressure pain threshold" OR "pressure pain thresholds" OR "pain pressure threshold" OR "pain pressure thresholds" OR "pain threshold" OR "heat temporal summation" OR "cold tolerance" OR ("hydraul*" AND ("algomet*" OR "pain ratings")) OR "nociceptive flexion reflex" OR "electrical pain ratings" OR "heat pain ratings" OR "pressure pain ratings" OR "cold pain ratings" OR "heat discrimination" OR "pain tolerance" OR "pressure tolerance thresholds" OR "pressure tolerance threshold" OR "experimental pain" OR "pain perception" OR "electrical pain threshold" OR "electrical pain thresholds" OR "perception threshold" OR "perception thresholds" OR "pain thresholds" OR "heat pain thresholds" OR "pain threshold temperature" OR "pain thresholds temperature" OR "heat pain threshold" OR "perceptive threshold" OR "perceptive thresholds" OR "sensory threshold" OR "sensory thresholds" OR "sense thresholds" OR "nociception threshold" OR "nociception thresholds" OR "nociceptive threshold" OR "nociceptive thresholds" OR "nociceptive pressure threshold" OR "nociceptive pressure thresholds" OR "pressure nociceptive threshold" OR "pressure nociceptive thresholds" OR "endogenous pain modulation" OR "temporal summation" OR "pain modulation" OR "nociceptive withdrawal reflex" OR "pain tolerance threshold" OR "pain tolerance thresholds" OR "pain inhibition" OR "pain inhibitory process" OR "pain inhibitor" OR "hypoalgesia" OR "reduced pain sensitivity" OR

20,293

"hyposensitivity to pain" OR "nociception" OR "pain sensation" OR "pain sensitivity" OR "quantitative sensory testing" OR "quantitative sensation testing" OR "quantitative sensory test") OR ti("nociceptive stimulation" OR "pressure pain threshold" OR "pressure pain thresholds" OR "pain pressure threshold" OR "pain pressure thresholds" OR "pain threshold" OR "heat temporal summation" OR "cold tolerance" OR ("hydraul*" AND ("algomet*" OR "pain ratings")) OR "nociceptive flexion reflex" OR "electrical pain ratings" OR "heat pain ratings" OR "pressure pain ratings" OR "cold pain ratings" OR "heat discrimination" OR "pain tolerance" OR "pressure tolerance thresholds" OR "pressure tolerance threshold" OR "experimental pain" OR "pain perception" OR "electrical pain threshold" OR "electrical pain thresholds" OR "perception threshold" OR "perception thresholds" OR "pain thresholds" OR "heat pain thresholds" OR "pain threshold temperature" OR "pain thresholds temperature" OR "heat pain threshold" OR "perceptive threshold" OR "perceptive thresholds" OR "sensory threshold" OR "sensory thresholds" OR "sense thresholds" OR "nociception threshold" OR "nociception thresholds" OR "nociceptive threshold" OR "nociceptive thresholds" OR "nociceptive pressure threshold" OR "nociceptive pressure thresholds" OR "pressure nociceptive threshold" OR "pressure nociceptive thresholds" OR "endogenous pain modulation" OR "temporal summation" OR "pain modulation" OR "nociceptive withdrawal reflex" OR "pain tolerance threshold" OR "pain tolerance thresholds" OR "pain inhibition" OR "pain inhibitory process" OR "pain inhibitor" OR "hypoalgesia" OR "reduced pain sensitivity" OR "hyposensitivity to pain" OR "nociception" OR "pain sensation" OR "pain sensitivity" OR "quantitative sensory testing" OR "quantitative sensation testing" OR "quantitative sensory test")

#3 **Population** ti("rat" OR "rats" OR "mouse" OR "mice" OR "horse" OR "horses" OR "gorilla" OR "animal" OR "animals" 496.430 OR "manipulation" OR "anaphylaxis" OR "acupuncture" OR "pigeons" OR "asthma" OR "plant" OR "deer" OR "children" OR "child" OR "heart disease" OR "coronary disease" OR "myocardial ischemia" OR "asthmatic" OR "rodent" OR "rodents" OR "angina" OR "coronary artery disease" OR "paraplegics" OR spinal cord lesion" OR "asthmatics" OR "chronic heart failure" OR "hypoxemia") OR ti("rat" OR "rats" OR" "mouse" OR "mice" OR "horse" OR "horses" OR "gorilla" OR "animal" OR "animals" OR "manipulation" OR "anaphylaxis" OR "acupuncture" OR "pigeons" OR "asthma" OR "plant" OR "deer" OR "children" OR "child" OR "heart disease" OR "coronary disease" OR "myocardial ischemia" OR "asthmatic" OR "rodent" OR "rodents" OR "angina" OR "coronary artery disease" OR "paraplegics" OR "spinal cord lesion" OR "asthmatics" OR "chronic heart failure" OR "hypoxemia")

#4 Reliab (MAINSUBJECT.EXACT.EXPLODE("Test-Retest Reliability") OR MJMAINSUBJECT.EXACT ("Psychometrics") OR MAINSUBJECT.EXACT.EXPLODE("Test Responsiveness")) OR ab("equivalence" OR "psychometrics" OR "psychometr*" OR "clinimetr*" OR "clinometr*" OR "observer variation" OR "reproducib*" OR "agreement" OR "precision" OR "imprecision" OR "precise values" OR "test-retest" OR "stability" OR "interrater" OR "inter-rater" OR "intrarater" OR "intra-rater" OR "intertester" OR "intertester" OR "intratester" OR "intra-tester" OR "inter-observer" OR "inter-observer" OR "intra-observer" OR "intra-observer" OR "intertechnician" OR "inter-technician" OR "intratechnician" OR "intra-technician" OR "interexaminer" OR "inter-examiner" OR "intraexaminer" OR "intra-examiner" OR "interassay" OR "inter-assay" OR "intraassay" OR "intra-assay" OR "interindividual" OR "inter-individual" OR "intraindividual" OR "intra-individual" OR "interparticipant" OR "inter-participant" OR "intraparticipant" OR "intra-participant" OR "kappa" OR "kappas" OR "repeatab*" OR "generaliza*" OR "generalisa*" OR "concordance" OR "discriminative" OR "known group" OR "factor analysis" OR "factor analyses" OR "dimensionality" OR "subscale*" OR "multitrait scaling analysis" OR "multitrait scaling analyses" OR "item discriminant" OR "interscale correlation" OR "interscale correlations" OR "individual variability" OR "variability analysis" OR "standard error of measurement" OR "sensitiv*" OR "responsive*" OR "meaningful change" OR "minimal important change" OR "minimal important difference" OR "minimally important change" OR "minimally important difference" OR "minimal detectable change" OR "minimal detectable difference" OR "minimally detectable change" OR "minimally detectable difference" OR "minimal real difference" OR "ceiling effect" OR "floor effect" OR "g theory" OR "generalizability theory" OR "generalisability theory" OR ("reliab*" AND ("test" OR "retest")) OR (("replicab*" OR "repeated") AND ("measure" OR "measures" OR "findings" OR "result" OR "results" OR "test" OR "tests")) OR ("intraclass" AND "correlation*") OR (("error" OR "errors") AND ("measure*" OR "correlat*" OR "evaluat*" OR "accuracy" OR "accurate" OR "precision" OR "mean")) OR ("uncertainty" AND ("measurement" OR "measuring")) OR ("small*"

687.955

AND ("real" OR "detectable") AND ("change" OR "difference"))) OR ti("equivalence" OR "psychometrics" OR "psychometr*" OR "clinimetr*" OR "clinometr*" OR "observer variation" OR "reproducib*" OR "agreement" OR "precision" OR "imprecision" OR "precise values" OR "test-retest" OR "stability" OR "interrater" OR "interrater" OR "intrarater" OR "intra-rater" OR "intertester" OR "inter-tester" OR "intra-tester" OR "intra-tester" OR "inter-observer" OR "inter-observer" OR "intra-observer" OR "intra-observer" OR "intertechnician" OR "intertechnician" OR "intratechnician" OR "intra-technician" OR "interexaminer" OR "inter-examiner" OR "intraexaminer" OR "intra-examiner" OR "interassay" OR "inter-assay" OR "intraassay" OR "intra-assay" OR "interindividual" OR "inter-individual" OR "intraindividual" OR "intra-individual" OR "interparticipant" OR "inter-participant" OR "intraparticipant" OR "intra-participant" OR "kappa" OR "kappas" OR "repeatab*" OR "generaliza*" OR "generalisa*" OR "concordance" OR "discriminative" OR "known group" OR "factor analysis" OR "factor analyses" OR "dimensionality" OR "subscale*" OR "multitrait scaling analysis" OR "multitrait scaling analyses" OR "item discriminant" OR "interscale correlation" OR "interscale correlations" OR "individual variability" OR "variability analysis" OR "standard error of measurement" OR "sensitiv*" OR "responsive*" OR "meaningful change" OR "minimal important change" OR "minimal important difference" OR "minimally important change" OR "minimally important difference" OR "minimal detectable change" OR "minimal detectable difference" OR "minimally detectable change" OR "minimally detectable difference" OR "minimal real difference" OR "ceiling effect" OR "floor effect" OR "g theory" OR "generalizability theory" OR "generalisability theory" OR ("reliab*" AND ("test" OR "retest")) OR (("replicab*" OR "repeated") AND ("measure" OR "measures" OR "findings" OR "result" OR "results" OR "test" OR "tests")) OR ("intraclass" AND "correlation*") OR (("error" OR "errors") AND ("measure*" OR "correlat*" OR "evaluat*" OR "accuracy" OR "accurate" OR "precision" OR "mean")) OR ("uncertainty" AND ("measurement" OR "measuring")) OR ("small*" AND ("real" OR "detectable") AND ("change" OR "difference")))

#5 #1 AND #2 AND #4 NOT #3 236

6) Manual searches

Google scholar

- Date of search: 27th of March 2022, 04th of September 2023, and 20th of February 2024
- Terms: ("exercise-induced hypoalgesia" OR "exercise-induced analgesia") AND ("reliability" OR "measurement error")
- Pages: 1–10
- Total results: 0

References of included studies

Date: 18th of March 2022

Author	Study title	Source	Date	Number of	Number of
(Year)				references	new reports
Gomolka	Assessing Endogenous Pain Inhibition: Test–Retest Reliability of Exercise-	Scopus	2022-	52	0
(2019)	Induced Hypoalgesia in Local and Remote Body Parts After Aerobic Cycling		03-18		
Vaegter	Hypoalgesia after bicycling at lactate threshold is reliable between sessions	Scopus	2022-	34	0
(2019)			03-18		
Vaegter	Exercise-Induced Hypoalgesia After Isometric Wall Squat Exercise: A Test-Retest	Scopus	2022-	34	0
(2019)	Reliabilty Study		03-18		
Vaegter	Test-retest reliability of exercise-induced hypoalgesia after aerobic exercise	Scopus	2022-	40	0
(2018)			03-18		
Hviid	Walking increases pain tolerance in humans: an experimental cross-over study	Scopus	2022-	39	0
(2019)			03-18		

S4 Data extraction form

Table S4 presents the code book of the data extraction form. It describes all the variables that were collected.

Table S4: Code book of the data extraction form

Extracted variables	Description	Expected information	In case of missing	
			information	
Article information				
email_first_author	Write the mail of the first author.	Per article	Not reported	
study_title	Write the study title in English.	Per article	Not reported	
ournal_publication	Write the full name of the journal	Per article	Not reported	
	where the article is published.			
ear_publication	Write year of publication. Should	Per article	Not reported	
	follow: "published in/ the:"			
ountry_study	Select the country where the study	Denmark, Brazil, Belgium,	Other	
	was done.	Canada, China, Germany, India,		
		Iran, Italy, Japan, Korea,		
		Netherlands, Sweden,		
		Switzerland, UK, USA, Other		
ountry_study_other	If the country name was not in the	Per article	Not reported	
	drop down list, write the name of			
	the country where the study was			
	conducted. The first letter should be			
	in capital case. (e.g., Belgium). If the			
	country was already selected in the			
	previous point then leave blank.			
anguage	Select the language of the study.	English,German, Portuguese,	Other	
		Spanish, French, Other		
anguage_other	If the language was not found in the	Per article	Not reported	
	drop down menu, write the name of			
	the language used to write the			
	article. The first letter should be in			
	capital case. (e.g., English). If the			
	language was already selected in			
	the previous point then leave blank.			
unding	Select the source of funding of the	Academic Foundation,	Other	
	study.	Government, Industry, No		
		Funding, Other, Unclear		
unding_other	Write the funding source with first	Per article	Not reported	
	letter in capital case if it was not in			
	the drop down list.			
rial_registration			Not reported	

Table S4: Continued

Extracted variables	Description	Expected information	In case of missing information
	Select among options . If not a trial,	Yes, No, Not Applicable, Unclear,	
	select "Not Applicable".	Not reported	
rial_registration_location	Select among options for where the	ClinicalTrial.gov, Other, Not	Other
	trial was registered. If not a trial,	Applicable, Unclear	
	select "Not Applicable".		
rial_registration_location_other	Write the trial registration location	Per article	Not reported
	with first letter in capital case if it		
	was not in the drop down list.		
ethical_approval	Select among options regarding the	Yes, No, Not Reported, Unclear	Not reported
	state of the ethical approval. The		·
	authors should whether ethical		
	approval was obtained for their		
	study.		
Study information	•		
study_design	Select among options.	Cross-over study, Parallel study,	Other
, <u> </u>	5 ,	Cross-sectional study,	
		Retrospective study, Case-control	
		study, Randomized controlled	
		trial, Clinical trial, Other	
study_setting	Select among options.	At Patients Home, Hospital,	Other
,- J	5 ,	Private Clinic, Rehab Centre,	
		Multicentre Governmental	
		Institute, Other, Not applicable,	
		Unclear	
study_setting_other	Describe study setting starting with	Per article	Not reported
	a capital letter.		•
sample_n	Insert the total size sample. It	Per article	Not reported
	should be a number. E.g., 40.		·
sample_age	Insert the sample age. It should be a	Per article	Not reported
5	number. E.g., 40.		·
sample_age_sd	Insert the sample age sd. It should	Per article	Not reported
, - 3 -	be a number. E.g., 40		·
sample_sex	Select among options.	Males, Females, Mixed	Not reported
ample_sex_n_female_male	If sample is gender mixed then state	Per article	Not reported
,	the ratio of females/males. It should		•
	be expressed as, for example, 3/10.		
sample_BMI	Insert the sample BMI. It should be	Per article	Not reported
· r -=-···-	a number. E.g., 40.		
sample_BMI_sd	Insert the sample age sd. It should	Per article	Not reported
	be a number. E.g., 40		

Table S4: Continued

Extracted variables	Description	Expected information	In case of missing information
sample_condition	Select among options.	Healthy, Low back pain, Neck pain, Shoulder pain, Knee pain, Knee oestoarthitis, Hip pain, Hip osteoarthitis, Other	Other
sample_condition_additional	Write sample condition starting with a capital letter if not found in the drop-down. If it was found in the previous drop-down, write "Not applicable".	Per article	Not reported OR Not applicable
diagnosis_tool_diagnosis	Describe diagnosis tool for condition described above. Start with a capital letter. If healthy, state "Not applicable".	Per article	Not reported
target_population	Descirbe target population. E.g., adults with low back pain	Per article	Not reported
outcomes_n	State the number of outcomes. Only "general outcomes" should be accounted for. For example if pressure pain threshold is used at two different sites, one should only count pressure pain threshold. It should be a number e.g., 3	Per article	Not reported
outcomes_list	List the outcomes (see previous point). Start each name with a capital letter and separate them with a coma.	Per article	Not reported
outcomes_main	List the main outcomes. Main outcome are identified according to the following decision rule: 1) According to the authors (e.g., "our primary outcomes are") 2) Used to estimate sample size 3) Based on the order of the results section. For example if the authors sate that A is their primary outcome but report it only last in the results, we should consider A as primary outcome; conversely, the first outcome—B—reported in the results will be	Per article	Not reported
	selected if the authors do not		

Table S4: Continued

Extracted variables	Description	Expected information	In case of missing information
	explicitly separate outcomes and		
	don't report sample size calculation.		
	Start each name with a capital letter		
	and separate them with a coma.		
	E.g., Pressure pain threshold, Pain		
	intensity.	Dan autiala	Networked
outcomes_main_assessor	identify the profile main assessor of	Per article	Not reported
	the outcomes. E.g., therapist,		
	therapis with X years of clinical experience.		
sampling_method	Select among options.	Random sample, Convenience	Other
		sample, Cluster sample,	
		Purposive sample, Unclear, Other	
sampling_method_other	If the sampling method was not	Per article	Not reported OR Not
	found in the drop down list then		applicable
	write it with first letter in caps. If it		
	was found in the previous drop-		
	down, write "Not applicable".		
sample_size_calculation	Select among options wether the	Yes, No, Unclear, Not reported	Not reported
	authors perfomed an apriori sample		
	size calculation.		
sample_size_adequate	Does the size of the recruited	Yes, No, Unclear, No prior sample	No prior sample size
	sample matches the a priori	size calculation	calculation
	calculation ? Select among options.		
recruitment_period	State the recruitment period as	Per article	Not reported
	described in the article.		
eligibility_in	Write the inclusion criteria for study	Per article	Not reported
	participation. Use the following		
	format: 1) First letter is capital , 2)		
	First letter is capital. E.g., 1) Adults 2)		
	Healthy		
eligibility_out	Write the exclusion criteria for study	Per article	Not reported
	participation. Use the following		
	format: 1) First letter is capital , 2)		
	First letter is capital. E.g., 1) Adults 2)		
	Healthy		
Exercise information			
exercise_mode	Select among options. See here	Aerobic, Anaerobic-Submax,	Not reported
	for:exercise defintions	Anaerobic-EE, Anaerobic-HIE,	
		Anaerobic-EIE, Resistance-CRT,	

Table S4: Continued

Extracted variables	Description	Expected information	In case of missing
			information
		Resistance-Concentric,	
		Resistance-DCER, Resistance-	
		Eccentric, Resistance-Isometric,	
		Resistance-Isokinetic, Other, Not	
		reported	
exercise_type	Describe exercise type starting with	Per article	Not reported
	a capital letter. Examples: Cycling,		
	Walking, Running. In cas of		
	isodynamic or isometric exercise,		
	enter the name of the movement		
	(e.g., Shoulder abduction).		
exercise_duration	Enter the duration of the exercise as	Per article	Not reported
	a number. E.g., 30		
exercise_duration_units	Select the units of the number	sec, min, hours, Not reported	Not reported
	stated above.		
exercise_intensity	Describe the exercise intensity.	Per article	Not reported
	Numeric value are preferred. Hence,		
	if an article states 30% VO2max, 30		
	should be entered. Otherwise, non-		
	numeric values can be entered, e.g.,		
	Maximal Intensity. If exercise does		
	not require that intensity is		
	prespecified, state: No intensity		
	control.		
exercise_intensity_determination	Select among options how exercise	Maximal exercise test, Sub-	Other
	intensity was targeted.	maximal exercise test, Age-based	
		formula, EMG Normalization	
		(MVC), Other, No intensity control	
exercise_intensity_determination_other	If exercise intensity determination	Per article	Not reported OR Not
	method was not found in the drop-		applicable
	down list then describe it starting		
	with a capital letter. If it was found		
	in the previous drop-down, write		
	"Not applicable".		
exercise_location	Select among options.	Laboratory, Outside, Not reported, Other	Not reported
exercise_supervisor	Insert information regarding who	Per article	Not reported
	supervised the exercise. If reported		
	the following should be extracted:		

Table S4: Continued

Extracted variables	Description	Expected information	In case of missing information
	profession, number of years of		
	experience, training. E.g., Physcial		
	therapist with 5 years of clinical		
	experience trained to apply		
exercise_support	Select among options to define	Yes, No, Not reported, Unclear	Not reported
	whether verbal support was given		
	while the participants exercised.		
	Thus, were the participants		
	motivated or encouraged by the		
	assessor/supervisor during the		
	exercise ?		
warm_up	Select among options.	Yes, No, Unclear, Not reported	Not reported
warm_up_mode	Select among options. See here	Aerobic, Anaerobic-Submax,	Not reported or Not
	for:exercise defintions	Anaerobic-EE, Anaerobic-HIE,	applicable if no
		Anaerobic-EIE, Resistance-CRT,	warm-up
		Resistance-Concentric,	
		Resistance-DCER, Resistance-	
		Eccentric, Resistance-Isometric,	
		Resistance-Isokinetic, Other, Not	
		reported	
warm_up_type	Describe warm-up exercise type	Per article	Not reported or Not
- 7 - 7.	starting with a capital letter.		applicable if no
	Examples: Cycling, Walking,		warm-up
	Running. In cas of isodynamic or		
	isometric exercise, enter the name		
	of the movement (e.g., Shoulder		
	abduction).		
warm_up_intensity	Describe the exercise intensity.	Per article	Not reported or Not
,	Numeric value are preferred. Hence,		applicable if no
	if an article states 30% VO2max, 30		warm-up
	should be entered. Otherwise, non-		,
	numeric values can be entered, e.g.,		
	Maximal Intensity.		
warm_up_duration	Enter the duration of the exercise as	Per article	Not reported or Not
ap_aaraaon	a number. E.g., 30	. c. dicici	applicable if no
	a namber. E.g., 50		warm-up
warm_up_duration_units	Select the units of the number	sec, min, hours, Not reported,	Not reported or Not
ap_aaradon_anno	stated above.	Not applicable	applicable if no
	Stated above.	ιτοι αργιιταμίτ	
control_mode	Select among options.		warm-up

Table S4: Continued

Extracted variables	Description	Expected information	In case of missing information
		Aerobic, Isodynamic, Isometric, Isokinetic, Not an exercise, Not reported, Not applicable	Not reported or Not applicable if no control
control_type	Descirbe control type starting with a capital letter. Examples: Rest in sitting position, Cycling	Per article	Not reported or Not applicable if no control
control_duration	Enter the duration of the control intrevntion as a number. E.g., 30	Per article	Not reported or Not applicable if no control
control_duration_units	Select the units of the number stated above.	sec, min, hours, Not applicable	Not reported or Not applicable if no control
control_intensity	Describe the control intensity. Only applicable if control is an exercise intrevention. Otherwise, state "Not Applicable".	Per article	Not reported or Not applicable if no control
control_location	Select among options.	Laboratory, Outside, Not reported, Other	Not reported
control_interval	State the time interval between the control condition and the exercise condition. It should be a numeric value. Only applicable if a cross-over design was used. Otherwise, state "Not Applicable".	Per article	Not reported
equipment_exercise	Describe the equipment used in the exercise protocol. State it in the following fashion: Equipment type_1 (brand_1); Equipment type_2 (brand_2); For example, Cycle ergometer(Monark,v20,USA); Cardiofrequencemeter(Polar,USA)	Per article	Not reported
exercise_prep	Describe prepatory actions that were conducted before the exercise. For example, a test session to determine the intensity for the upcoming exercise sessions or calibration of the material. Use the following form: Prep1;Prep2 E.g., Submaximal exercise test (lactate	Per article	Not reported

Table S4: Continued

Extracted variables	Description	Expected information	In case of missing information
	threshold determination); Cycle		
	ergometer calibration.		
Test modalities information			
stimu_type	Select among options.	Mechanical, Ischaemic, Thermal, Chemical, Electrical	Not reported
test_modality	Enter the abbreviated form of test modalities used. Use only upper cases. E.g., PPT, HPT	Per article	Not reported
test_modality_location	Select among options to describe the location of application of the test modality.	Quadriceps, Hand, Lower back, Forearm, Elbow, Deltoid, Upper trapezius, Calf, Other	Not reported
test_modality_location_other	Describe shortly the test modality location if it was not found in the drop-down menu above; the first letter should be in uppercase.	Per article	Not reported
test_modality_full_name	Enter, the full name of the test test modalities used. The first letter should be in uppercase. E.g., Pain pressue threshold.	Per article	Not reported
eih_session1	Select among options to determine whether exercise-indued hypoalgesia was manifested. This should be based on the results and discussion sections of the article.	Yes,No, Not reported, Unclear	Not reported
eih_session_1_computation	Select whether eih was computed from absolute or relative change . Relative change is computed using a proportion e.g., 10% increase of PPT; absolute change is computed using "raw" values e.g., 100 kpA increase. In the case of, for instance, EIH from relative change was reported, specify whether the authors also tried to compute absolute change (and perjhaps did not find EIH) or only assessed EIH from relative change. The same applies for	Relative change, Absolute change, Absolute change (relative change not computed), Relative change (absolute change not computed), Relative and absolute change, Not reported, Unclear	Not reported
	absolute change.		
eih_session2		Yes, No, Not reported, Unclear	Not reported

Table S4: Continued

Extracted variables	Description	Expected information	In case of missing information
	Select among options to determine		
	whether exercise-indued		
	hypoalgesia was manifested. This		
	should be based on the results and		
	discussion sections of the article.		
eih_session_2_computation	Select whether eih was computed	Relative change, Absolute	Not reported
	from absolute or relative change .	change, Absolute change (relative	
	Relative change is computed using a	change not computed),Relative	
	proportion e.g., 10% increase of PPT;	change (absolute change not	
	absolute change is computed using	computed), Relative and absolute	
	"raw" values e.g., 100 kpA increase.	change, Not reported, Unclear	
	In the case of, for instance, EIH from		
	relative change was reported,		
	specify whether the authors also		
	tried to compute absolute change		
	(and perjhaps did not find EIH) or		
	only assessed EIH from relative		
	change. The same applies for		
	absolute change.		
ntsruments_brand	State the brand of the instrument(s)	Per article	Not reported
	used for the test modality		·
instruments_description	Describe the instruments used for	Per article	Not reported
	the test modality.		·
test_modality_n_measurement	Select among options. A numeric	1,2,3,4,5, Not Applicable, Not	Not reported
•	value should be selected to report	reported	·
	the number of test for each test	·	
	modality. E.g., PPT was measured 3		
	times at the quadriceps.		
test_modality_rate	Enter a numeric value to define the	Per article	Not reported
- ,-	rate of application of the test		·
	modality. If the test modality		
	doesn't require a rate of application		
	then state: "Not applicable".		
test_modality_break	If several measurement of each test	Per article	Not reported or Not
- >-	modality were conducted, then		applicable if only one
	insert the inter-measurement		measurement per
	interval of the same test modality as		test modality
	a numeric value.		,
test_modality_between_interval		Per article	

Table S4: Continued

Extracted variables	Description	Expected information	In case of missing information
	If several test modalities have been		Not reported or Not
	applied, then insert the inter-		applicable if only one
	measurement interval of the		test modality.
	different test modalites as a		
	numeric value.		
test_modality_subject_position	Select among options to define the	Sitting, Prone, Supine, Standing,	Not reported
	subject position while test modality	Not reported, Unclear	
	was assessed.		
test_modaity_description	Insert here a summary of the test	Per article	Not reported
	modality procedure. Elements from		
	previous cells can be repeated.		
	Specify, when possible, the identity		
	of the assessor. In parenthesis, state		
	(position; rate; n measurement/site;		
	interval; sites order)		
rater_description	Insert information regarding rater.	Per article	Not reported
	If reported the following should be		
	extracted: profession, number of		
	years of experience, training. E.g.,		
	Physcial therapist with 5 years of		
	clinical experience trained to apply		
instructions_prep	Insert informations regarding the	Per article	Not reported
	instructions provided to the before		
	the experiment subjects. E.g.,		
	subjects were asked to refrain from		
	physcial activity on the day of		
	experiment.		
adherence_instructions	State whether the participants	Yes, No, Unclear, Not reported	Not reported
	adhered to the instructions and how		
	this was verified. Select among		
	options.		
adherence_instructions_verification	If "yes" was selected above, select	Per article	Not applicable or Not
	how this was assessed. E.g., verbal		reported
	verificatin on the day of		
	participation.		
familiarization	Select among options wheter a	Yes, No, Not reported	Not reported
	familiarization session was	·	•
	conducted. Familiarization relates to		
	physically experimenting the		
	procedures.		

Table S4: Continued

Extracted variables	Description	Expected information	In case of missing information
familiarization_details	If a familiarization session was	Per article	Not applicable or Not
	conducted, describe the part of the		reported
	procedures to which it applied. For		
	example if the subjects were		
	familiarized with the PPT, state "Test		
	modality" (starting with a capital		
	letter). If the subjcets were		
	familiarized with different parts of		
	the exercise, list them with a "," as		
	separator. For example, if the		
	subjects realized a exercise test on a		
	cycle ergometer and were		
	familiarized with the test modalities		
	prior to the sessions used to assess		
	reliability, state "Cycle ergometer,		
	Test modality". If the subjects were		
	familiarized with the entire		
	procedure,state "EIH procedure". If		
	the previous point was "no", then		
	state "Not applicable".		
explanation_explicit	Select among options whether an	Yes, No, Not reported	Not reported
	explanation regarding the exercise		
	and/or test modality was given to		
	the subjects.		
test_mod_prep	Describe prepatory actions for	Per article	Not reported
, '	experimental pain assessment that		•
	are not covered by previous items.		
	Use the following form:		
	Prep1;Prep2 E.g., Calibration of		
	algomter or rater training. If not		
	reported, state not reported.		
assessor_blinding_outcome description	If applicable, describe how blinding	Per article	Not reported or Not
abbesser_aag_aateae aebep.t.a	of the assessor was ensured.	, c. article	applicable if no
			blinding.
control_other	Describe any other from of control	Per article	Not reported
	reported in the study. If several		eported
	were included, then state them		
	while using "," as separator.		
Information regarding exercise and test m			
injormation regulating exercise and test in	ouunties		

Table S4: Continued

Extracted variables	Description	Expected information	In case of missing
			information
	These cells are only present in		
	database_2. They will auto-		
	complete. Ignore while extracting.		
Test modalities information			
equipment_test_mod	These cells are only present in	1	/
	database_2. They will auto-		
	complete. Ignore while extracting.		
adherence_instructions	These cells are only present in	1	/
	database_2. They will auto-		
	complete. Ignore while extracting.		
IH determination methods			
eih_data_processing_storage	This item relates to "All actions	Per article	Not reported
	undertaken on the raw data to store		
	it in a usable (electronic) form for		
	later data manipulation (such as		
	score assignment or statistical		
	analysis)". Describe thus how the		
	acquired data were stored and by		
	who. E.g., PPT values stored in a		
	computer under the guidance of		
	rater 1.		
eih_score	Describe here how eih was scored.	Per article	Not reported
	For example, Relative (insert		
	formula used) and absolute change		
	(insert fomrmula used)		
data_analysis_used	Describe the statistical approach to	Per article	Not reported
	assessing exercise-induced		
	hypoalgesia. NOT RELIABILITY at		
	this stage.		
significance_level	Select the significance level that was	0.05, 0.01, Not reported	Not reported
	set (alpha).	·	·
paseline_data_analyze_each_group	Was all the baseline data analyzed in	Yes, No, Not applicable, Unclear,	Not reported
,,	each group ? If cross-over design,	Not reported	·
	select "Not applicable".	•	
nformation regarding reliability or meas			
ater_roles_description	Summarize the role(s) of the rater(s)	Per article	Not reported
	according to the article. The		
	information extracted above can be		
	repated.		
	reputeu.		

Table S4: Continued

Extracted variables	Description	Expected information	In case of missing information
	Describe the component of the		
	procedure(s) that will be repeated.		
	The whole procedure can be		
	repeated (i.e., equipment		
	preparation, data collection, data		
	storage and score assignement) on		
	all occasions or for all raters or only		
	a part of it (e.g., score assignement).		
	A priroi, in studies examining		
	exercise-induced hypoalgesia, the		
	whole procedure should be		
	repeated. State also the focus of		
	interest (e.g., scoring EIH).		
variation_components_time	Select among options wheter the	Yes, No, Unclear, Not reported	Not reported
	occasion is varied across		
	measurement. Typically in a test-		
	retest or intrarater reliability study,		
	this should be the case, select		
	then "yes".		
variation_components_rater	Select among options whether the	Yes, No, Unclear, Not reported	Not reported
- ' -	rater component is varied across		·
	measurement. In a interrater study,		
	it should be the case. Select		
	then "yes".		
variation_components_instruments	Select among options whether the	Yes, No, Unclear, Not reported	Not reported
- ' -	instrument(s) are varied between	, , , , ,	·
	measurements. For example, if a		
	intrarater study uses algometer A		
	on occasion 1 and algometer B on		
	occasion 2 then select "yes".		
variation_components	Summarize the components being	Per article	Not reported
anation_components	varied using the information	Tel didec	rec reported
	extracted above. E.g., Occasion ($n =$		
	2, intrarater reliability); Rater (<i>n</i> = 3,		
	interrater reliability).		
Stat_continuous_scores_reliab	Select among options thetest	ICC, Pearson correlation	Not reported
otat_continuous_scores_reliab			Not reported
	statistic used to assess reliabillity of	coefficient, Other Not reported	
	continuous outcomes. If no	coefficient, Other, Not reported,	
	continuous outcome was assessed then select "Not applicable"	Not applicable	

Table S4: Continued

Extracted variables	Description	Expected information	In case of missing information
Stat_ordinal_scores_reliab	Select among options thetest	ICC, Weighted kappa,	Not reported
	statistic used to assess reliabillity of	Unweighted kappa, Other, Not	
	ordinal outcomes. If no ordinal	applicable, Not reported	
	outcome was assessed then select		
	"Not applicable"		
tat_dichotomous_scores_reliab	Select among options thetest	ICC, Weighted kappa,	Not reported
	statistic used to assess reliabillity of	Unweighted kappa, Other, Not	
	dichotomous or nominal outcomes.	applicable, Not reported	
	If no dichotomous or nominal		
	outcome was assessed then select		
	"Not applicable"		
Risk of bias assessment			
stability_patient	R & M (1) Were the patients stable in	very good, adequate, doubtful,	Leave blank
	the time betweenthe repeated	inadequate, NA	
	measurementson the construct to		
	be measured? Select among options		
	according to: COSMIN RoB tool		
tability_patient_justif	Justify previous item.	Per article	Leave blank
ime_interval	R &M (2) Was the time between the	very good, doubtful,	Leave blank
	measurements appropriate? Select	inadequate, NA	
	among options according to:		
	COSMIN RoB tool		
ime_interval_justif	Justify previous item based one the	Per article	Leave blank
	article. Ideally, provide page		
	number or article quote.		
imilar_measurements_conditions	R &M (3) Were the measurement	Very good, adequate, doubtful,	Leave blank
	conditions similar for the	inadequate, NA	
	measurements - except for the		
	condition being evaluated as a		
	source of variation? Select among		
	options according to: COSMIN		
	RoB tool		
imilar_measurements_conditions_justif	Justify previous item based one the	Per article	Leave blank
	article. Ideally, provide page		
	number or article quote.		
neasurements_administration	R &M (4) Did the professional(s)	Very good, adequate, doubtful,	Leave blank
	administer the measurement	inadequate	
	without knowledge of scores or		
	values of other repeated		
	measurement(s) in the same		

Table S4: Continued

Extracted variables	Description	Expected information	In case of missing information
	patients? Select among options		
	according to: COSMIN RoB tool		
measurements_administration_justif	Justify previous item based one the	Per article	Leave blank
	article. Ideally, provide page		
	number or article quote.		
score_assignements	R &M (5) Did the professional(s)	Very good, adequate, doubtful,	Leave blank
	assign scores or determine values	inadequate	
	without knowledge of the scores or		
	values of other repeated		
	measurement(s) in the same		
	patients? Select among options		
	according to: COSMIN RoB tool		
score_assignements_justif	Justify previous item based one the	Per article	Leave blank
	article. Ideally, provide page		
	number or article quote.		
other_flaws	R &M (6) Were there any other	Very good, doubtful, inadequate	Leave blank
	important flaws in the design or		
	statistical methods of the study?		
	Select among options according to:		
	COSMIN RoB tool		
other_flaws_justif	Justify previous item based one the	Per article	Leave blank
	article. Ideally, provide page		
	number or article quote.		
stat_continuous_scores_reliab	R (7) For continuous scores: was an	Very good, adequate,	Leave blank
	intraclass correlation coefficient	doubtful,NA	
	(ICC) calculated? Select among		
	options according to: COSMIN		
	RoB tool		
stat_continuous_scores_reliab_justif	Justify previous item based one the	Per article	Leave blank
	article. Ideally, provide page		
	number or article quote.		
stat_ordinal_scores_reliab	R (8) For ordinal scores: was a	Very good, adequate,NA	Leave blank
	(weighted) kappa calculated? Select	·	
	among options according to:		
	COSMIN RoB tool		
stat_ordinal_scores_reliab_justif		Per article	Leave blank

Table S4: Continued

Extracted variables	Description	Expected information	In case of missing information
	Justify previous item based one the		
	article. Ideally, provide page		
	number or article quote.		
stat_dichotomous_scores_reliab	R (9) For dichotomous/nominal	Very good	Leave blank
	scores: was Kappa calculated for		
	each category against the other		
	categories combined? Select among		
	options according to: COSMIN		
	RoB tool		
tat_dichotomous_scores_reliab_justif	Justify previous item based one the	Per article	Leave blank
	article. Ideally, provide page		
	number or article quote.		
stat_continuous_scores_reliab	M (7) For continuous scores: was the	Very good, adequate, doubtful,	Leave blank
	Standard Error of Measurement	inadequate,NA	
	(SEM), Smallest Detectable Change		
	(SDC), Limits of Agreement (LoA) or		
	Coefficient of Variation (CV)		
	calculated? Select among options		
	according to: COSMIN RoB tool		
tat_continuous_scores_reliab_justif	Justify previous item based one the	Per article	Leave blank
	article. Ideally, provide page		
	number or article quote.		
tat_dichotomous_ordinal_scores_reliab	M (8) For dichotomous/nominal/	Very good, adequate, NA	Leave blank
	ordinal scores: Was the percentage		
	specific (e.g. positive and negative)		
	agreement calculated? Select		
	among options according to:		
	COSMIN RoB tool		
stat_dichotomous_ordinal_scores_reliab_justif	Justify previous item based one the	Per article	Leave blank
	article. Ideally, provide page		
	number or article quote.		
nal_rating	At the end of the R & M risk of bias	Very good, adequate, doubtful,	Leave blank
	assessment, provide an overall	inadequate, NA	
	score using the worst-score counts	·	
	method (i.e., the overall score is		
	determined by the lowest value		
	upon all preceeding score). For		
	example, if a study is rated "very		

Table S4: Continued

Extracted variables	Description	Expected information	In case of missing
			information
	good" everywhere but the R_3		
	component is rated "doubtful", then		
	the final rating will be "doubtful".		
	For more information, see the		
	COSMIN long button (in the		
	reliability part).		
Quantitative data			
mean_pré_session1	Insert the mean value of the test	Per article	Not reported OR Not
	modality pre-exercise in session 1.		aplicable
sd_pré_session1	Insert the standard deviation value	Per article	Not reported OR Not
	of the test modality pre-exercise in		aplicable
	session 1.		
units_pré_session1	Specify the units of the above	Per article	Not reported OR Not
	mentionned values.		aplicable
mean_absolute_change_session1	Insert the mean absolute change	Per article	Not reported OR Not
	(i.e., expressed in the units of the		aplicable
	test modality) of the test modality		
	after exercise in session 1.		
sd_absolute_change_session1	Insert the standard deviation of	Per article	Not reported OR Not
	absolute change of the test modality		aplicable
	after exercise in session 1.		
mean_relative_change_session1	Insert the mean relative change (i.e.,	Per article	Not reported OR Not
•	expressed as decimal or percent) of		aplicable
	the test modality after exercise in		
	session 1. Express it in percent.		
	E.g, 14.5		
sd_relative_change_session1	Insert the standard deviation of	Per article	Not reported OR Not
•	relative change of the test modality		aplicable
	after exercise in session 1. Express it		·
	in percent. E.g, 14.5		
eih_session1	Was there eih in session 1 ? This cell	Don't insert anything	Don't insert anything
	value will be copied from the	, , , , , , , , , , , , , , , , , , ,	, , , , , , , , , , , , , , , , , , ,
	eih_procedures sheet.		
mean_pré_session2	Insert the mean value of the test	Per article	Not reported OR Not
	modality pre-exercise in session 2.		aplicable
sd_pré_session2	Insert the standard deviation value	Per article	Not reported OR Not
<u>-</u>	of the test modality pre-exercise in		aplicable
	session 2.		ирпсиоле

Table S4: Continued

Extracted variables	Description	Expected information	In case of missing information
units_pré_session2	Specify the units of the above mentionned values.	Per article	Not reported OR Not
mean_absolute_change_session2	Insert the mean absolute change of the test modality after exercise in session 2.	Per article	Not reported OR Not aplicable
sd_absolute_change_session2	Insert the standard deviation of absolute change (i.e., expressed in the units of the test modality) of the test modality after exercise in session 2.	Per article	Not reported OR Not aplicable
mean_relative_change_session2	Insert the mean relative change(i.e., expressed as decimal or percent) of the test modality after exercise in session 2. Express it in percent. E.g., 14.5	Per article	Not reported OR Not aplicable
sd_relative_change_session2	Insert the standard deviation of relative change of the test modality after exercise in session 2. Express it in percent. E.g., 14.5	Per article	Not reported OR Not aplicable
eih_session2	Was there eih in session 2 ? This cell value will be copied from the eih_procedures sheet.	Don't insert anything	Don't insert anything
ICC_absolute_change	Insert the value of the ICC computed for absolute change. It should be between 0 and 1.	Per article	Not reported OR Not aplicable
ICC_absolute_change_CI_low	Insert the value of the lower bound of the confidence interval of ICC computed for absolute change.	Per article	Not reported OR Not aplicable
ICC_absolute_change_CI_high	Insert the value of the upper bound of the confidence interval of ICC computed for absolute change.	Per article	Not reported OR Not aplicable
ICC_relative_change	Insert the value of the ICC computed for relative change. It should be between 0 and 1.	Per article	Not reported OR Not aplicable
ICC_relative_change_CI_low	Insert the value of the lower bound of the ICC computed for relative change.	Per article	Not reported OR Not aplicable
ICC_relative_change_CI_high		Per article	

Table S4: Continued

Extracted variables	Description	Expected information	In case of missing information
	Insert the value of the upper bound		Not reported OR Not
	of the ICC computed for relative		aplicable
	change.		
SEM_absolute_change	Insert the value of the SEM	Per article	Not reported OR Not
	(Standard Error of Measurement)		aplicable
	computed for absolute change.		
SEM_relative_change	Insert the value of the SEM	Per article	Not reported OR Not
	(Standard Error of Measurement)		aplicable
	computed for relative change.		
r_absolute_change	Insert the value of the Pearson	Per article	Not reported OR Not
-	correlation coefficient computed for		aplicable
	the absolute change.		•
r_absolute_change_CI_low	Insert the value of the lower bound	Per article	Not reported OR Not
	of the confidence interval of the		aplicable
	Pearson correlation coefficient for		.,
	absolute change.		
r_absolute_change_CI_high	Insert the value of the upper bound	Per article	Not reported OR Not
	of the confidence interval of the	i ci diticio	aplicable
	Pearson correlation coefficient for		ирпсиыс
	absolute change.		
r_relative_change	Insert the value of the Pearson	Per article	Not reported OR Not
I_relative_trialige	correlation coefficient computed for	rei aiticie	aplicable
	the relative change.		арпсавте
r_relative_change_CI_low	Insert the value of the lower bound	Per article	Not reported OP Not
T_Telative_Change_Ci_low	of the confidence interval of the	rei aiticie	Not reported OR Not
	Pearson correlation coefficient for		aplicable
a valativa abanana CI birah	relative change.	Den autiala	Net was asted OD Net
r_relative_change_CI_high	Insert the value of the upper bound	Per article	Not reported OR Not
	of the confidence interval of the		aplicable
	Pearson correlation coefficient for		
	relativechange.		
kappa_EIH_resp_between_sessions	Report the value of the kappa	Per article	Not reported OR Not
	coefficient calculated from the		aplicable
	agreement bteween EIH responders		
	in session 1 vs session 2.		
kappa_EIH_resp_between_sessions_CI_low	Insert the value of the lower bound	Per article	Not reported OR Not
	of the confidence interval of the		aplicable

Table S4: Continued

Extracted variables	Description	Expected information	In case of missing information
	kappa value determined in previous		
	point.		
kappa_EIH_resp_between_sessions_CI_high	Insert the value of the lower bound	Per article	Not reported OR Not
	of the confidence interval of the		aplicable
	kappa value determined in previous		
	point.		
EIH_resp_session1_2	If applicable (i.e., EIH was	Per article	Not applicable or Not
	dichotomized in two categories) ,		reported
	state the number of EIH responder		
	(as defined in the study) in session 1		
	AND in session 2. Enter a numeric		
	value.		
EIH_non_resp_session1_2	If applicable(i.e., EIH was	Per article	Not applicable or Not
	dichotomized in two categories) ,		reported
	state the number of EIH non		
	responder(as defined in the study)		
	in session 1 AND in session 2. Enter a		
	numeric value.		
EIH_resp_session1	If applicable(i.e., EIH was	Per article	Not applicable or Not
	dichotomized in two categories) ,		reported
	state the number of EIH responder		
	(as defined in the study) in session 1		
	only. Enter a numeric value.		
EIH_resp_session2	If applicable(i.e., EIH was	Per article	Not applicable or Not
	dichotomized in two categories) ,		reported
	state the number of EIH responder		
	(as defined in the study) in session 2		
	only. Enter a numeric value.		

S5 Correspondence with the authors

The questions asked to the authors and the status of their answers are provided in Table S5. The authors were all contacted on the 4th of May 2022, at the end of the data collection process.

Table S5: Correspondence with the authors

lviid (2019), Walking increases pain tolerance in humans: an exper	michal cross over stady
Question	Answer
Where did the study take place?	Laboratory at the Pain Center at Odense University Hospital, Odens
	Denmark
Who supervised the exercise?	Experienced Physio
Who assessed the pressure pain threshold (PPT), cuff pressure pain	Experienced Physio. Manual PPT and cuff algometry assessments we
hreshold (CPPT), and cuff pressure tolerance threshold (CPTT), and	trained and supervised by experienced assessor several times over a
what was his/her background and training status?	weeks period prior to the study. Instructions given to subjects were standardized
What was the order and the time interval between each test different	The order was first PPT assessment at the quad, first PPT assessment
nodality (e.g., PPT and CPPT)?	the trap, second PPT assessment at the quad, second PPT assessme
	at the trap. Then cPPT (threshold) and cPTT (tolerance) was determin
	(same test).
What was the time interval between the PPT (i.e., between each	For PPTs at least 20 sec between repeated assessments at the
epeated PPT assessment)?	same site.
as the adherence to the instructions (e.g., to refrain from physical	YES
ctivity on the dayss of participation) checked?	
low and by who were the data (PPT, CPTT, CPPT) processed and	PPT, cPPT and cPTT data was initially documented on a physical
tored?	datasheet (one sheet per session). Cuff data were also stored on cpa
	nocitech system PC. After sessions data were entered into an Excel
	spreadsheet and stored on secure server at Odense University Hospi
he ICC 3.1 that was used implies that the reliability results that were	We chose the ICC 3.1 with Two-way mixed, single measures,
ound cannot be generalized to other raters, could you elaborate on	consistency because: Two-way mixed: a fixed number of raters were
ne reasons of this choice?	used and each subject was measured by the fixed raters. Single
	measure: even though more than one measure at each site was tak
	we used the average of a single rater Consistency: The repeated
	measurements of two sessions were performed by the same rater,
	we decided that systematic errors of the rater were canceled and or
	the random residual error was kept.
hat was the formula of the ICC: consistency or agreement?	Consistency
ou classified subjects in EIH responder and non-responders. EIH was	This was done to enable us to expore the agreement in the
nus dichotomized. Could you elaborate on the reasons of this choice?	classification of EIH responses within a subject across the two session
	We used Intraclass correlation coefficients to investigate the ability

Table S5: Continued

Hviid (2019), Walking increases pain tolerance in humans: an expe	inicital cross over stady
Question	Answer
	EIH responses to differentiate values between subjects across the two
	sessions.
You determined the within session standard error of measurement	The reason for calculation of the PPT, cPPT and cPTT SEM was to be able
(SEM) of the PPT, CPTT, and CPPT. Why did not you also compute the	to classify subjects into EIH-responder and non-responders to
SEM of the EIH itself?	determine absolute reliability. We did not consider calculating SEM for
	EIH in this study.
Vaegter (2019), Exercise-Induced Hypoalgesia After Isometric Wall	Squat Exercise: A Test-Retest Reliabilty Study
Question	Answer
Were other medications than pain killers considered in the exclusion criteria?	NO
Where did the study take place?	Department of Physiotherapy, University College North Denmark,
	Aalborg, Denmark
What were the inclusion criteria?	Pain-free women and men with adequate Danish language skills
What was the recruitment period?	February 1st 2017 - May 1st 2017
Who supervised the exercise?	Physio student in final year of education
Were the participants motivated during the exercise?	Subjects were encouraged to complete the entire 3 min of the wall
	squat every 30 secs with standardized phrases; "Keep up the good work" or "You are doing well"
Who assessed the pressure pain threshold and what was his/her	Physio student in final year of education. PPT assessments were trained
background and training status?	and supervised by experienced assessor several times over a 3 weeks
	period prior to the study.
Were the subjects blinded to the study hypothesis?	In the subject information handed out prior to consent, subjects were
	given the following information: Physical activity and exercise are
	frequently used as treatment, but more knowledge about the effect of
	physical activity is needed to optimize the effect. Based on the
	hypothesis that physical activity and exercise lead to a change in the
	perception of pain, we have carried out a number of research projects
	exploring the effect of physical activity on the pain sensitivity. Whether
	a potential change in pain sensitivity is similar over repeated days of
	physical activity is unknown.
Was the examinator blinded to the score of the subjects?	NO
Was the adherence to the instructions (e.g., to refrain from physical	YES
activity on the dayss of participation) checked?	
How long lasted each testing session?	Approximately 30 min
How and by who was the data (PPT) processed and stored?	PPT data was initially documented on a physical datasheet (one sheet
	per session). After sessions data were entered into an Excel

spreadsheet and stored on secure server at Odense University Hospital

Table S5: Continued

Hviid (2019).	Walking increases	s pain tolerance	e in humans:	: an experimental	cross-over study

Question	Answer
Could you confirm that only one rater (PPT) was involved?	The same rater assessed PPTs in both sessions.
What was the the order of sites in the PPT assessments?	The order was first PPT assessment at the quad, first PPT assessment a
	the trap, second PPT assessment at the quad, second PPT assessmen
	at the trap.
The ICC 3.1 that was used implies that the reliability results that were	We chose the ICC 3.1 with Two-way mixed, single measures,
found cannot be generalized to other raters, could you elaborate on	consistency because: Two-way mixed: a fixed number of raters were
the reasons of this choice?	used and each subject was measured by the fixed raters. Single
	measure: even though more than one measure at each site was taker
	we used the average of a single rater Consistency: The repeated
	measurements of two sessions were performed by the same rater, so
	we decided that systematic errors of the rater were canceled and only
	the random residual error was kept.
The standard error of measurement (SEM) was reported as the square	The method for calculating SEM was based on the paper Quantifying
root of the mean square error term (MSE) of the ANOVA-RM. There are	Test-Retest Reliability Using The Intraclas Correlation Coefficient and
two possibilities of SEM, consistency and agreement, depending on	The SEM by Weir JP published in J Strength and Conditioning Research
whether variance due to bias was included in the MSE or not. Could	2005, 19(1):231-240. We used the formula: SEM = square root of the
you state the formula that was used?	mean square error (type III sum of squares in error for WithinSubjec
	effects divided by df for error in WithinSubject effect) from the ANOVA
	table
You classified subjects in EIH responder and non-responders. EIH was	This was done to enable us to expore the agreement in the
thus dichotomized. Could you elaborate on the reasons of this choice?	classification of EIH responses within a subject across the two sessions
	We used Intraclass correlation coefficients to investigate the ability o
	EIH responses to differentiate values between subjects across the two
	sessions
Vaegter (2019), Hypoalgesia after bicycling at lactate threshold is r	eliable between sessions.
Question	Answer
Were other medications than pain killers considered in the exclusion criteria?	NO

Question	Answer
Were other medications than pain killers considered in the exclusion	NO
criteria?	
What were the inclusion criteria?	Pain-free women and men with adequate Danish language skills
What was the recruitment period?	Sept 25 2017 - 15. nov 2017
Who supervised the exercise in sessions 2 and 3?	Physio student in final year of education
Were the participants motivated during the exercise in sessions 2	Subjects were encouraged to complete the entire 15 min of the
and 3?	bicycling exercise session 2 and 3 every 2 mins with standardized
	phrases; "Keep up the good work" or "You are doing well"
Where did the study take place?	Laboratory at the Pain Center at Odense University Hospital, Odense,
	Denmark
Were the people who did not finish the study similar in terms of	3 included subjects did not finish the study due to scheduling conflicts.
characteristics to the people staying in the study?	We did not explore whether they were different from the 34 subjects
	that completed all 3 sessions.

Table S5: Continued

What was the recruitment period?

Hviid (2019).	Walking ir	ncreases pair	i tolerance	n humans:	an experimental	cross-over study

Question	Answer
Who assessed the pressure pain threshold and what was his/her	Physio student in final year of education. PPT assessments were trained
background and training status?	and supervised by experienced assessor several times over a 3 weeks
	period prior to the study.
Was the adherence to the instructions (e.g., to refrain from physical	YES
activity on the dayss of participation) checked?	
Was the examinator blinded to the score of the subjects in some ways?	NO
Could you confirm that only one rater (PPT) was involved?	The same rater assessed PPTs in the same subject in sessions 2 and 3.
How and by who were the data (PPT) processed and stored?	PPT data was initially documented on a physical datasheet (one sheet
	per session). After sessions data were entered into an Excel
	spreadsheet and stored on secure server at Odense University Hospital
The ICC 3.1 that was used implies that the reliability results that were	We chose the ICC 3.1 with Two-way mixed, single measures,
found cannot be generalized to other raters, could you elaborate on	consistency because: Two-way mixed: a fixed number of raters were
the reasons of this choice?	used and each subject was measured by the fixed raters. Single
	measure: even though more than one measure at each site was taken
	we used the average of a single rater Consistency: The repeated
	measurements of two sessions were performed by the same rater, so
	we decided that systematic errors of the rater were canceled and only
	the random residual error was kept.
You classified subjects in EIH responder and non-responders. EIH was	This was done to enable us to expore the agreement in the
thus dichotomized. Could you elaborate on the reasons of this choice?	classification of EIH responses within a subject across the two sessions.
	We used Intraclass correlation coefficients to investigate the ability of
	EIH responses to differentiate values between subjects across the two
	sessions.
You determined the within session standard error of measurement	The reason for calculation of the PPT SEM was to be able to classify
(SEM) of the pressure pain threshold. Why did not you also compute	subjects into EIH-responder and non-responders to determine
the SEM of the EIH itself?	absolute reliability. We did not consider calculating SEM for EIH in this
	study.
How long lasted each session	Session 2 and session 3 lasted aprox. 45 min
Did you also compute ICC for EIH determined as relative change (and	No, we did not.
not only absolute change)? If so, what were the results?	
What was the standard deviation of the relative change (EIH) for each	I just calculated standard deviation for relative change in PPT (EIH):
site (quadriceps and upper trapezius) on each session?	Session 2 Quadriceps: 18.1 Trapezius: 13.9 Session 3 Quadriceps: 18.5
	Trapezius: 10.4
Vaegter (2018), Test-Retest Reliabilty of Exercise-Induced Hypoalge	sia After Aerobic Exercise
Question	Answer
Were other medications than pain killers considered in the exclusion	NO
criteria?	
What were the inclusion criteria?	Pain-free women and men with adequate Danish language skills

February 1st 2017 - April 5 2017

Table S5: Continued

	humans: an experimental	

Question	Answer
Where did the study take place?	Laboratory at the Pain Center at Odense University Hospital, Odense, Denmark
What was the time interval between the rest and exercise condition?	15 min
Were the participants motivated during the exercise?	Subjects were encouraged to complete the entire 15 min of the
	bicycling exercise session 1 and 2 every 2 mins with standardized
	phrases; "Keep up the good work" or "You are doing well"
Was the adherence to the instructions (e.g., to refrain from physical	YES
activity on the dayss of participation) checked?	
Who assessed the pressure pain threshold (PPT) and what was his/her	Physio student in final year of education. PPT assessments were trained
background and training status?	and supervised by experienced assessor several times over a 3 weeks
	period prior to the study.
Could you confirm that only one rater (PPT) was involved?	The same rater assessed PPTs in the same subject in sessions 1 and 2.
What was the time interval between the PPT (i.e., between each repeated PPT assessment)?	at least 20 sec between repeated assessments at the same site.
What was the the order of sites in the PPT assessments?	The order was first PPT assessment at the quad, first PPT assessment at
	the trap, second PPT assessment at the quad, second PPT assessment
	at the trap.
How and by who were the data (PPT) processed and stored?	PPT data was initially documented on a physical datasheet (one sheet
	per session). After sessions data were entered into an Excel
	spreadsheet and stored on secure server at Odense University Hospital
The ICC 3.1 that was used implies that the reliability results that were	We chose the ICC 3.1 with Two-way mixed, single measures,
found cannot be generalized to other raters, could you elaborate on	consistency because: Two-way mixed: a fixed number of raters were
the reasons of this choice?	used and each subject was measured by the fixed raters. Single
	measure: even though more than one measure at each site was taken
	we used the average of a single rater Consistency: The repeated
	measurements of two sessions were performed by the same rater, so
	we decided that systematic errors of the rater were canceled and only
	the random residual error was kept.
You classified subjects in EIH responder and non-responders. EIH was	This was done to enable us to expore the agreement in the
thus dichotomized. Could you elaborate on the reasons of this choice?	classification of EIH responses within a subject across the two sessions.
	We used Intraclass correlation coefficients to investigate the ability of
	EIH responses to differentiate values between subjects across the two sessions.
The standard error of measurement (SEM) was reported as the square	The method for calculating SEM was based on the paper Quantifying
root of the mean square error term (MSE) of the ANOVA-RM. There are	Test-Retest Reliability Using The Intraclas Correlation Coefficient and
two possibilities of SEM, consistency and agreement, depending on	The SEM by Weir JP published in J Strength and Conditioning Research
whether variance due to bias was included in the MSE or not. Could	2005, 19(1):231-240. We used the formula: SEM = square root of the
you state the formula that was used?	mean square error (type III sum of squares in error for WithinSubject
	effects divided by df for error in WithinSubject effect) from the ANOVA
	table

DE GRUYTER

Table S5: Continued

Question	Answer
How long lasted each session?	Approximately 60 min
Gomolka (2019), Assessing Endogenous Pain Inhibition: Test–Retest	Reliability of Exercise-Induced Hypoalgesia in Local and Remote Body
Parts After Aerobic Cycling	
Question	Answer
Were other medications than analgesics and corticosteroids	No.
considered in the exclusion criteria?	
What was the recruitment period?	November 2016 - March 2017
Where did the study take place?	Ruhr University of Bochum
How and by who was the data (PPT) processed and stored (i.e., after	During PPT assessment, PPT values were manually recorded by SG,
that SG assessed the pressure pain threshold, who strored this data)?	who transferred them into an SPSS file after the experiment. Data were
	processed by SG and CT, and securely stored within the Department of
	Medical Psychology and Medical Sociology of the Ruhr University
	Bochum
The ICC 3.1 that was used implies that the reliability results that were	With having investigated a rather homogeneous sample in terms of
found cannot be generalized to other raters, could you elaborate on	age and health status, we were interested in how close the absolute
the reasons of this choice?	and percent change scores were. When choosing the ICC, we followed
	Koo & Li (2016): "In addition, absolute agreement definition should
	always be chosen for both test-retest and intrarater reliability studies
	because measurements would be meaningless if there is no agreement
	between repeated measurements." Koo, T. K., & Li, M. Y. (2016). A
	guideline of selecting and reporting intraclass correlation coefficients
	for reliability research. Journal of chiropractic medicine, 15(2), 155–163
The standard error of measurement (SEM) was reported as the square	Variance due to bias was not included in the MSE
root of the mean square error term (MSE) of the ANOVA-RM. There are	
two possibilities of SEM, consistency and agreement, depending on	
whether variance due to bias was included in the MSE or not. Could	
you state the formula that was used?	

S6 COSMIN Risk of Bias tool

The following figures, from Mokkink et al. [1], display the standards and rating system for reliability and measurement error of the COSMIN RoB tool

Standards for studies on reliability

	Design requirements	very good	adequate	doubtful	inadequate	NA
1	Were patients stable in the time between the repeated measurements on the construct to be measured?	Yes (evidence provided)	Reasons to assume standard was met	Unclear	No (evidence provided)	Na
2	Was the time interval between the repeated measurements appropriate?	Yes		Doubtful, OR time interval not stated	No	Na
3	Were the measurement conditions similar for the repeated measurements – except for the condition being evaluated as a source of variation?	Yes (evidence provided)	Reasons to assume standard was met, OR change was unavoidable	Unclear	No (evidence provided)	Na
4	Did the professional(s) administer the measurement without knowledge of scores or values of other repeated measurement(s) in the same patients?	Yes (evidence provided)	Reasons to assume standard was met	Unclear	No (evidence provided)	
5	Did the professional(s) assign scores or determine values without knowledge of the scores or values of other repeated measurement(s) in the same patients?	Yes (evidence provided)	Reasons to assume standard was met	Unclear	No (evidence provided)	
6	Were there any other important flaws in the design or statistical methods of the study?	No		Minor methodological flaws	Yes	

St	atistical methods	very good	adequate	doubtful	inadequate
7	For continuous scores: was an intraclass correlation coefficient (ICC) calculated?	ICC calculated; the model or formula was described, and matches study design and the data	ICC calculated but model or formula was not described or does not optimally match the study design OR Pearson or Spearman correlation coefficient calculated WITH evidence provided that no systematic difference between measurements has occurred	Pearson or Spearman correlation coefficient calculated WITHOUT evidence provided that no systematic difference between measurements has occurred OR WITH evidence provided that systematic difference between measurements has occurred	
8	For ordinal scores: was a (weighted) kappa calculated?	Kappa calculated; the weighting scheme was described, and matches the study design and the data	Kappa calculated, but weighting scheme not described or does not optimally match the study design		
9	For dichotomous/nominal scores: was Kappa calculated for each category against the other categories combined?	Kappa calculated for each category against the other categories combined			

Standards for studies on measurement error

	Design requirements	very good	adequate	doubtful	inadequate	NA
1	Were patients stable in the time between the repeated measurements on the construct to be measured?	Yes (evidence provided)	Reasons to assume standard was met	Unclear	No (evidence provided)	Na
2	Was the time interval between the repeated measurements appropriate?	Yes		Doubtful, OR time interval not stated	No	Na
3	Were the measurement conditions similar for the repeated measurements – except for the condition being evaluated as a source of variation?	Yes (evidence provided)	Reasons to assume standard was met, OR change was unavoidable	Unclear	No (evidence provided)	Na
4	Did the professional(s) administer the measurement without knowledge of scores or values of other repeated measurement(s) in the same patients?	Yes (evidence provided)	Reasons to assume standard was met	Unclear	No (evidence provided)	
5	Did the professional(s) assign scores or determine values without knowledge of the scores or values of other repeated measurement(s) in the same patients?	Yes (evidence provided)	Reasons to assume standard was met	Unclear	No (evidence provided)	
6	Were there any other important flaws in the design or statistical methods of the study?	No		Minor methodological flaws	Yes	

St	atistical methods	very good	adequate	doubtful	inadequate
7	For continuous scores: was the Standard Error of Measurement (SEM), Smallest Detectable Change (SDC), Limits of Agreement (LoA) or Coefficient of Variation (CV) calculated?	SEM, SDC, LoA or CV calculated; the model or formula for the SEM/SDC is described; it matches the reviewer constructed research question and the data	SEM, SDC, LoA or CV calculated, but the model or formula is not described or does not optimally match the reviewer constructed research question* and evidence provided that no systematic difference has occurred	SEM _{consistency} SDC _{consistency} or LoA or CV calculated, without knowledge about systematic difference or with evidence provided that systematic difference has occurred	SEM calculated based on Cronbach's alpha, or using SD from another population
8	For dichotomous/nominal/ordinal scores: Was the percentage specific (e.g., positive and negative) agreement calculated?	% specific agreement calculated	% agreement calculated		

S7 COMSIN criteria for good reliability

The COMSIN initiative proposes to rate each individual result based on the following criteria [1]:

- "+": Sufficient
- "?": Indeterminate
- "-": Insufficient

M(C)IC stand for Minimal (Clinical) Important Change; It is defined by the COSMIN initiative as: "the smallest change in score in the construct to be measured which patients perceive as important" [2].

	+	ICC or (weighted) Kappa ≥ 0.70		
Reliability	?	ICC or (weighted) Kappa not reported		
	1	ICC or (weighted) Kappa < 0.70		
	+	SDC or LoA or CV* $\sqrt{2}$ *1.96 < M(C)IC¹; % specific agreement > $80\%^2$		
Measurement error	?	MIC not defined		
	-	SDC or LoA or CV* $\sqrt{2}$ *1.96 > M(C)IC¹; % specific agreement < $80\%^2$		

¹ the M(C)IC value may come from another study. ² Sometimes a higher percentage is more appropriate; when substantiated, this could be appropriate, too.

S8 COSMIN modified GRADE approach

The following elements, cited from Prinsen et al. [3] complete the information regarding the modified GRADE approach that was presented in the methods section. Table S6 summarizes the results of the modified GRADE approach for each study.

The modified GRADE approach defines 4 quality levels, which are summarized in the following figure (from Prinsen et al. [3]).

Quality level	Definition
High	We are very confident that the true measurement property lies
	close to that of the estimate* of the measurement property
Moderate	We are moderately confident in the measurement property estimate: the true measurement property is likely to be close to the estimate of the measurement property, but there is a possibility that it is substantially different
Low	Our confidence in the measurement property estimate is limited: the true measurement property may be substantially different from the estimate of the measurement property
Very low	We have very little confidence in the measurement property estimate: the true measurement property is likely to be substantially different from the estimate of the measurement property

st Estimate of the measurement property refers to the pooled or summarized result of the measurement property of a PROM.

These definitions were adapted from the GRADE approach (20)

These quality levels are defined according to four constructs: Risk of Bias (RoB), Inconsistency, Imprecision, and Indirectness. They are detailed below.

1. Risk of Bias (RoB)

The COSMIN group suggests downgrading the quality of evidence by one to three levels proportionally to the RoB – one level, "serious RoB"; two levels, "very Serious RoB"; three levels, "extremely serious RoB". The table below summarizes the criteria used to determine the level of RoB. In the case of our systematic review, all included studies were considered to have a doubtful RoB. Thus, a "serious RoB" was assigned, downgrading the quality by one level (Table S6).

Risk of bias	Downgrading for Risk of Bias
No	There are multiple studies of at least adequate quality, or
	there is one study of very good quality available
Serious	There are multiple studies of doubtful quality available,
	or there is only one study of adequate quality
Very serious	There are multiple studies of inadequate quality, or there
	is only one study of doubtful quality available
Extremely	There is only one study of inadequate quality available
serious	

2. Inconsistency

Inconsistency relates to the presence of both sufficient and insufficient results in the quantitative or qualitative data summary (i.e., results deemed sufficient or insufficient account for less than 75% of all ratings). The COSMIN group suggests to first explore potential causes of inconsistency by determining whether it arises from an inadequate grouping of studies (i.e., the variability in the results is explained by differences in the study population, methodology, or RoB). In the case where no explanation can be found, they suggest downgrading the quality of evidence with one or two levels based on whether the review team considers it as serious (1 level) or very serious (2 levels), according to the specific context of the included studies. In the case of our systematic review, we could not summarize data from the included studies and considered inconsistency to be non-applicable (Table S6).

3. Imprecision

Imprecision is linked with the sample size of the included studies. The COSMIN group recommends downgrading the quality of evidence by one or two levels when the total sample size of the data summary is respectively below 100 or 50. In our review, we could not summarize data from the included studies and none of them had a sample size above 50. We thus downgraded the quality of evidence by two levels for each study (Table S6).

4. Indirectness

Indirectness relates to the similarity in the population or construct of interest between the summarized studies and the review. The COSMIN group suggests that this assessment is context dependent and leaves to the research team the decision to consider it as serious or very serious, downgrading the evidence by respectively one or two levels. In our review, each of the included studies were deemed to appropriately match with the targeted construct of interest—reliability or measurement error—and population pain-free adults or adults with MSK pain. We thus did not downgrade the quality of evidence for indirectness (Table S6).

The following figure [3] summarizes how these constructs are used to GRADE the quality of evidence.

Quality of evidence	Lower if
High	Risk of bias
Moderate	-1 Serious
Low	-2 Very serious
Very low	-3 Extremely serious
	Inconsistency -1 Serious -2 Very serious
	Imprecision
	-1 total n=50-100
	-2 total n<50
	Indirectness -1 Serious -2 Very serious

n=sample size

Table S6: Results of the modified GRADE approach for each study

Author (Year)	GRADE factors						
	Imprecision (levels)	Inconsistency (levels)	Indirectness (levels)	Reliability (levels)	Measurement error		
Gomolka (2019)	Yes(2)	NA	No(0)	doubtful(1)	doubtful(1)		
Hviid (2019)	Yes(2)	NA	No(0)	doubtful(1)	NA		
Vaegter (2019a)	Yes(2)	NA	No(0)	doubtful(1)	NA		
Vaegter (2018)	Yes(2)	NA	No(0)	doubtful(1)	doubtful(1)		
Vaegter (2019b)	Yes(2)	NA	No(0)	doubtful(1)	doubtful(1)		

NA, Not applicable. Rating: Rating of the reliability value according to the COMSIN criteria, Grade: Grading of the quality of the evidence according to the modified GRADE approach from COSMIN, GRADE factors: Factors used to grade the evidence, Levels:number of downgraded levels

S9 Reasons for exclusion of reports

We excluded 53 reports after reading their texts in full; only the conference report was found for two reports. Table S7 presents the reasons for exclusion of each of them.

Table S7: Excluded reports after full text reading

Author (Year)	Tite	Journal (Volume, Issue, Pages, DOI)	Reason for	
			exclusion	
Adrian (2015)	Physical activity, pain responses to heat stimuli,	Menopause (22, 8, 816–825, 10.1097/	not EIH nor	
	and conditioned pain modulation in	GME.000000000000398)	reliab	
	postmenopausal women			
Alappattu (2011)	Stability of behavioral estimates of activity- dependent modulation of pain	Journal of Pain Research (4, 151–157, 10.2147/JPR.S18105)	not EIH	
Cabido (2014)	Acute effect of constant torque and angle	Journal of Strength and Conditioning Research (28, 4,	not EIH nor	
	stretching on range of motion, muscle passive properties, and stretch discomfort percept	1050–1057, 10.1519/JSC.0000000000000241)	reliab	
Choi (2013)	The effects of stability exercises on shoulder	Journal of Physical Therapy Science (25, 2, 155–158,	not EIH nor	
	pain and function of middle-aged women	10.1589/jpts.25.155)	reliab	
Cowley (2018)	Influence of remote pain on movement control	Experimental Brain Research (236, 8, 2309–2319,	not EIH nor	
	and muscle endurance during repetitive movements	10.1007/s00221-018-5303-6)	reliab	
Dannecker (2015)	Inducing a Blind Spot: Blinding Data Collectors	Pain Medicine (United States) (16, 6, 1145–1154, 10.1111/	not EIH nor	
	in an Investigation of Experimental Pain	pme.12706)	reliab	
Dawson (2012)	Assessment of proprioceptive allodynia after	J Orofac Pain (26, 1, 39–48)	not reliab	
	tooth-clenching exercises			
Oroste (1991)	Physical exercise, endogenous opiates and pain regulation	Der Schmerz (5, 3, 138–147, 10.1007/BF02528099)	not reliab	
Duncan (2013)	Acute caffeine ingestion enhances strength	European journal of sport science (13, 4, 392–399,	not EIH nor	
	performance and reduces perceived exertion and muscle pain perception during resistance exercise	10.1080/17461391.2011.635811)	reliab	
Eberhard (2018)	Quantitative sensory response of the SCM	Arch Oral Biol (86, 87–93, 10.1016/	not reliab	
,	muscle on sustained low level activation	j.archoralbio.2017.09.030)		
	simulating co-contractions during bruxing	,,		
isher (2010)	Carotid baroreflex control of arterial blood	American Journal of Physiology - Regulatory Integrative	not EIH nor	
, ,	pressure at rest and during dynamic exercise in	and Comparative Physiology (299, 5, R1241–R1247,	reliab	
	aging humans	10.1152/ajpregu.00462.2010)		
Friedman (1993)	The effect of epidural anaesthesia with 1%	The Journal of Physiology (470, 1, 681–691, 10.1113/	not reliab	
(/	lidocaine on the pressor response to dynamic	jphysiol.1993.sp019882)		
	exercise in man	2 2		
Fuller (1993)	A test of exercise analgesia using signal	Perceptual and motor skills (76, 3, Pt 2, 1,299–1,310,	not reliab	
,	detection theory and a within-subjects design	http://dx.doi.org/10.2466/pms.1993.76.3c.1299)		

Table S7: Continued

Author (Year)	Tite	Journal (Volume, Issue, Pages, DOI)	Reason for exclusion
Gagnon (2012)	Does sex have an independent effect on thermoeffector responses during exercise in the heat?	Journal of Physiology (590, 23, 5,963–5,973, 10.1113/ jphysiol.2012.240739)	review
Geisler (2019)	An fMRI study on runner's high and exercise- induced hypoalgesia after a 2-h-run in trained non-elite male athletes	Sport Sciences for Health (NA, 10.1007/s11332-019-00592-8)	not reliab
Geisler (2020)	An fMRI study on runner's high and exercise- induced hypoalgesia after a 2-h-run in trained non-elite male athletes	Sport Sciences for Health (16, 1, 159–167, 10.1007/s11332-019-00592-8)	not reliab
Graven- Nielsen (2013)	The interaction of musculoskeletal pain and physical activity: Human studies	Routledge Handbook of Physical Activity and Mental Health (NA, 359–371, 10.4324/9780203132678-39)	review
Hansen (2022)	Hypoalgesia after exercises with painful vs. non-painful muscles in healthy subjects - a randomized cross-over study	Scand J Pain (22, 3, 614–621, 10.1515/sjpain-2021-0161)	not reliab
Harden (2012)	Home-Based Aerobic Conditioning for Management of Symptoms of Fibromyalgia: A Pilot Study	Pain Medicine (United States) (13, 6, 835–842, 10.1111/j.1526-4637.2012.01384.x)	not EIH nor reliab
Hoffman (2004)	Intensity and duration threshold for aerobic exercise-induced analgesia to pressure pain	Archives of Physical Medicine and Rehabilitation (85, 7, 1,183–1,187, 10.1016/j.apmr.2003.09.010)	not reliab
Hollmann (2000)	Brain, psyche and physical activity	Orthopade (29, 11, 948–956, 10.1007/s001320050547)	review
Hudson (2008)	Effects of caffeine and aspirin on light resistance training performance, perceived exertion, and pain perception	Journal of Strength and Conditioning Research (22, 6, 1,950–1,957, 10.1519/JSC.0b013e31818219cb)	not EIH nor reliab
Hughes (2020)	The effect of blood flow restriction exercise on exercise-induced hypoalgesia and endogenous opioid and endocannabinoid mechanisms of pain modulation	Journal of Applied Physiology (128, 4, 914–924, 10.1152/ JAPPLPHYSIOL.00768.2019)	not reliab
anal (1996)	Pain sensitivity, exercise and stoicism	Journal of the Royal Society of Medicine (89, 7, 376–381, 10.1177/014107689608900706)	review
anal (1985)	Pain sensitivity, mood, and plasma endocrine levels in man following long-distance running: Effects of naloxone	(NA, 1374,)	not reliab
ang (2020)	The effects of acute resistance exercise intensity on pain perception and muscle mechanical properties of taekwondo athletes	Exercise Science (29, 3, 281–290, 10.15857/ ksep.2020.29.3.281)	not reliab
Kaluga (2006)	A comparative analysis of changes in tactile sensitivity in men and women practicing selected sports	Human Movement (7, 2, 153–161)	not EIH nor reliab
Kashima (2009)	•	Oral Science International (6, 1, 55–61, 10.11277/osi.6.55)	not reliab

Table S7: Continued

Author (Year)	Tite	Journal (Volume, Issue, Pages, DOI)	Reason for exclusion
			exclusion
	Mouth-opening Exercises Produce a Decrease		
	in Pain Perception in Patients with Disk		
	Displacement with Reduction		
Katz-	Aerobic Upper-Limb Exercise-Induced	Applied Sciences (Switzerland) (12, 22, 10.3390/	not reliab
Betzalel (2022)	Hypoalgesia: Does It Work?	app122211391)	
(etola (2012)	Constant, cycling, hot and cold thermal	Journal of Evolutionary Biology (25, 6, 1,209–1,215,	not humans
	environments: Strong effects on mean viability	10.1111/j.1420-9101.2012.02513.x)	
	but not on genetic estimates		
Kiyatkin (1990)	Endogenous opiate mechanisms in regulation	Izevestiya Akademii Nauk SSSR - Seriya Biologicheskaya	not humans
	of pain sensitivity and behavioral reactivity at	(NA, 5, 778–781)	
	aversive states of different nature		
(uithan (2020)	Exercise induced hypoalgesia: stability of	Physiotherapy (United Kingdom) (107, e49–e50, 10.1016/	conference
	measures with functional lumbar spine	j.physio.2020.03.068)	paper
	resistance training		
arbig (1994)	EEG correlates of pain control	EEG-EMG Zeitschrift fur Elektroenzephalographie	not EIH nor
		Elektromyographie und Verwandte Gebiete (25, 2,	reliab
		151–160,)	
_au (2015)	Changes in electrical pain threshold of fascia	European journal of applied physiology (115, 5,	not EIH nor
	and muscle after initial and secondary bouts of	959–968, 10.1007/s00421-014-3077-5)	reliab
	elbow flexor eccentric exercise		
Lee (2022)	Association of conditioned pain modulation	Osteoarthritis and Cartilage (30, Supplement 1, S383,	conference
	with exercise-induced hypoalgesia in knee	https://dx.doi.org/10.1016/j.joca.2022.02.515)	paper
	osteoarthritis		
Lewis (2018)	The Effect of Group Size and Synchrony on Pain	Small Group Research (49, 6, 723-738, 10.1177/	not reliab
	Threshold Changes	1046496418765678)	
Mahalakshmi	Possible neuroprotective mechanisms of	International Journal of Molecular Sciences (21, 16, 1–17,	review
2020)	physical exercise in neurodegeneration	10.3390/ijms21165895)	
McLoughlin (2011)	The relationship between physical activity and	Journal of Pain (12, 6, 640–651, 10.1016/	not EIH nor
	brain responses to pain in fibromyalgia	j.jpain.2010.12.004)	reliab
Meeus (2010)	Reduced pressure pain thresholds in response	Journal of Rehabilitation Medicine (Stiftelsen	not reliab
	to exercise in chronic fatigue syndrome but not	Rehabiliteringsinformation) (42, 9, 884–890, 10.2340/	
	in chronic low back pain: an experimental study	16501977-0595)	
Motl (2002)	Low intensity pain reported during elicitation of	Brain Research (951, 1, 53–58, 10.1016/S0006-8993(02)	not reliab
	the H-reflex: No effects of trait anxiety and high	03134-7)	
	intensity cycling exercise		
Muller (2011)	Pain and thermal sensation in the cold: the	Eur J Appl Physiol (111, 6, 979-87, 10.1007/s00421-010-	not EIH
	effect of interval versus continuous exercise	1726-x)	
Newcomb (2009)	The influence of a preferred versus a prescribed	Journal of Pain (10, 4, S16, 10.1016/j.jpain.2009.01.070)	not reliab
	exercise session on pain in women with		
	fibromyalgia		

Table S7: Continued

Author (Year)	Tite	Journal (Volume, Issue, Pages, DOI)	Reason for exclusion
Niwa (2022)	Effects of Exercise-Induced Hypoalgesia at Different Aerobic Exercise Intensities in Healthy Young Adults	Journal of Pain Research (15, 3615–3624, https://dx.doi.org/10.2147/JPR.S384306)	not reliab
O'Leary (2017)	High but not moderate-intensity endurance training increases pain tolerance: a randomised trial	European journal of applied physiology (117, 11, 2201-2210, 10.1007/s00421-017-3708-8)	not reliab
O'Neill (2019)	Acute sensory and motor response to 45-s heavy isometric holds for the plantar flexors in patients with Achilles tendinopathy	Knee Surgery, Sports Traumatology, Arthroscopy (27, 9, 2765–2773, 10.1007/s00167-018-5050-z)	not reliab
Rainville (2004)	The influence of intense exercise-based physical therapy program on back pain anticipated before and induced by physical activities	Spine Journal (4, 2, 176–183, 10.1016/ j.spinee.2003.08.031)	not EIH nor reliab
Sajedi (2016)	The evaluation of the aerobic exercise effects on pain tolerance	Sport Science (9, 7–11,)	not EIH nor reliab
Smith (1966)	An experimental pain method sensitive to morphine in man: the submaximum effort tourniquet technique	Journal of Pharmacology and Experimental Therapeutics (154, 2, 324–332,)	not EIH nor reliab
Tcheang (2005)	Systematic distortions of perceptual stability investigated using immersive virtual reality	Vision Research (45, 16, 2177–2189, 10.1016/ j.visres.2005.02.006)	not EIH nor reliab
Titze (2016)	Physical activity and musculoskeletal pain: A focus review within the MiSpEx research group	Schmerz (30, 5, 421–428, 10.1007/s00482-016-0152-3)	review
Vaegter (2017)	Exercise increases pressure pain tolerance but not pressure and heat pain thresholds in healthy young men	European Journal of Pain (21, 1, 73–81, 10.1002/ejp.901)	not reliab
Vallance (2021)	Contralateral mechanical hyperalgesia and altered pain modulation in men who have unilateral insertional Achilles tendinopathy: A cross-sectional study	Musculoskeletal Science and Practice (52, 10.1016/ j.msksp.2021.102353)	not EIH nor reliab
Wang (2020)	Correlation between aerobic exercise and improvement of physical sub-health based on multiple linear regression	Proceedings – 2020 12th International Conference on Measuring Technology and Mechatronics Automation, ICMTMA 2020 (NA, 1039–1043, 10.1109/ ICMTMA50254.2020.00223)	not reliab
Weiner (2006)	The relationship between pain, neuropsychological performance, and physical function in community-dwelling older adults with chronic low back pain	Pain Medicine (7, 1, 60–70, 10.1111/j.1526-4637.2006.00091.x)	not EIH nor reliab
Wu (2022)	Effects of Exercise-induced Hypoalgesia and Its Neural Mechanisms	Medicine and Science in Sports and Exercise (54, 2, 220–231, 10.1249/MSS.000000000002781)	not reliab

References

- [1] Mokkink LB, Boers M, van der Vleuten CPM, Bouter LM, Alonso J, Patrick DL, et al. COSMIN Risk of Bias tool to assess the quality of studies on reliability or measurement error of outcome measurement instruments: a Delphi study. BMC Med Res Methodol. 2020;20(1):293.
- [2] Vet HCW de. Measurement in medicine, Cambridge: Cambridge University Press; 2011. p. 350.
- [3] Prinsen CAC, Mokkink LB, Bouter LM, Alonso J, Patrick DL, de Vet HCW, et al. COSMIN guideline for systematic reviews of patientreported outcome measures. Qual Life Res. 2018;27(5):1147-57.