

Integration of them over y leads to

$$\begin{cases} \delta(z) = \beta e^{b\alpha z/a}, \\ \omega(z) = b a^2 z + \gamma, \end{cases} \quad (14)$$

where β and γ are constants. We thus obtain the result

$$u(x, y, z, t) = F - a \cdot \tanh \left[\frac{\beta e^{\alpha(t+bz/a)} + ax + by + a^2 bz + \gamma}{2} \right]. \quad (15)$$

Sample 2: Solitary waves are a special case of the family, since we are able to assume that

$$\phi(y, z) + \lambda(z, t) = by + cz + pt + h, \quad (16)$$

where b, c, p and h are constants. Constraint (10) then leads to

$$u(x, y, z, t) = F - a \cdot \tanh \left[\frac{ax + by + cz + a \left(\frac{c}{b} - a^2 \right) t + h}{2} \right]. \quad (17)$$

Acknowledgements

We thank Prof. Q. S. Lu and the referee for their valuable comments. YTG and BT were supported by the Fund for Excellent Young University Teachers and the Research Grants for Scholars Returning from Abroad, State Education Commission of China; and supported by the National Natural Science Foundation of China. WPH was supported by Catholic University of Taegu-Hyosung.

- [1] P. Clarkson and E. Mansfield, *Nonlinearity*, **7**, 975 (1994).
 [2] M. Jimbo and T. Miwa, *Publ. Res. Inst. Math. Sci., Kyoto Univ.* **19**, 943 (1983).
 [3] B. Dorizzi, B. Gammaticos, A. Ramani, and P. Winternitz, *J. Math. Phys.* **27**, 2848 (1986).
 [4] B. Tian and Y. T. Gao, *Z. Naturforsch.* **51a**, 171 (1996).

- [5] Y. T. Gao and B. Tian, *Computers Math. Applic.* **30**, 97 (1995).
 [6] B. Tian and Y. T. Gao, *J. Phys. A* **29**, 2895 (1996).
 [7] W.-H. Steeb, *Continuous Symmetries, Lie Algebras, Differential Equations and Computer Algebra*, World Scientific, Singapore 1996.

New Exact Solutions for a Generalized Breaking Soliton Equation

Yi-Tian Gao^{a,b}, Bo Tian^a, and Woopyo Hong^b

^a Depts. of Applied Physics and Mathematics, Beijing University of Aeronautics and Astronautics, Beijing 100083, China*

^b Dept. of Physics, Catholic University of Taegu-Hyosung, Hayang, Kyongsan, Kyungbuk 712-702, South Korea

Z. Naturforsch. **53a**, 807–808 (1998);
 received May 8, 1998

The breaking soliton equations are a class of nonlinear evolution equations of broad interest in physical and mathematical sciences. In this paper, the application of the generalized tanh method with symbolic computation leads to new exact solutions for a generalized breaking soliton equation, of which the previously-obtained solutions are the special cases.

Within a decade, a class of nonlinear evolution equations, called the breaking soliton equations, has

become a widely interesting subject in physical and mathematical sciences, as seen, e.g., in [1–5]. In addition to the rich mathematical properties, those equations are found to include the self-dual Yang-Mills equation, and to be of value in describing the (2+1)-dimensional interaction of the Riemann waves and long waves. A generalized breaking soliton equation [1, 2] reads as

$$(u_{xt} - 4u_x u_{xy} - 2u_y u_{xx} + u_{xxx})_x = -\alpha^2 u_{yyy}, \quad (1)$$

where α^2 is real. Two classes of exact solutions of (1) have been found, one of which is the solitary waves plus arbitrary functions of t [4], the other is linear with respect to y but independent of α^2 [5].

In this paper, we apply the computerized symbolic computation and generalized tanh method [6] to (1), assuming that the exact solutions are of the form

$$u(x, y, t) = \sum_{m=0}^N A_m(y, t) \cdot \tanh^m [\Psi(y, t)x + \Theta(y, t)], \quad (2)$$

Reprint request to Prof. Bo Tian;
 * Mailing address.

0932-0784 / 98 / 0900-0807 \$ 06.00 © – Verlag der Zeitschrift für Naturforschung, Tübingen · www.znaturforsch.com



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.

where the $A_m(y, t)$'s, $\Psi(y, t) \neq 0$ and $\Theta(y, t)$ are differentiable functions of y and t only, with $A_N(y, t) \neq 0$, while $N = 1$, determined by the leading-order analysis.

Inserting Ansatz (2) into (1) leads to a huge expression. Then we separate out the different terms. Equating to zero the coefficients of like powers of x , we obtain from the terms with x and x^3 the first constraint,

$$\Psi(y, t) = \psi = \text{constant}. \quad (3)$$

Equating to zero the coefficients of like powers of $\tanh(\Psi x + \Theta)$, we obtain more constraints on $\psi, A_1(y, t), A_0(y, t)$ and $\Theta(y, t)$, one of which,

$$\tanh^6(\Psi x + \Theta): [A_1(y, t) + 2\psi] \Theta_y(y, t) = 0, \quad (4)$$

leads to the following families of exact solutions of (1):

Family I:

With the condition that $A_1 = -2\psi$, we get the expressions for $\Theta(y, t)$ and $A_0(y, t)$ from the terms with $\tanh^3(\Psi x + \Theta)$ and $\tanh^4(\Psi x + \Theta)$, separately. The first family is thus found as

$$\begin{aligned} u^I(x, y, t) = & -2\psi \cdot \tanh[\psi x + \mu(t)y + \varpi(t)] \\ & + \frac{y^2 \mu_t(t)}{4\psi} + \frac{y \varpi_t(t)}{2\psi} \\ & + 2\psi y \mu(t) + \frac{\alpha^2 y \mu^3(t)}{2\psi^3} + \gamma(t), \end{aligned} \quad (5)$$

where the constant $\psi \neq 0$ as well as the differentiable functions $\mu(t)$, $\omega(t)$ and $\gamma(t)$ are all arbitrary.

The solutions obtained in [4, 5] are special cases of Family I.

Family II:

Starting from the assumption that $\Theta = \Theta(t)$ only, we obtain the expression for $A_0(y, t)$ from the terms with both $\tanh^2(\Psi x + \Theta)$ and $\tanh^4(\Psi x + \Theta)$, and get $A_1(y, t) = \tau = \text{constant}$ from the terms with $\tanh(\Psi x + \Theta)$, $\tanh^3(\Psi x + \Theta)$ and $\tanh^5(\Psi x + \Theta)$. The second family is calculated as

$$u^{II}(x, y, t) = \tau \cdot \tanh[\psi x + \Theta(t)] + \frac{y \Theta_t(t)}{2\psi} + \phi(t), \quad (6)$$

where the constants $\psi \neq 0$ and $\tau \neq 0$ as well as the differentiable functions $\Theta(t)$ and $\phi(t)$ are all arbitrary. This family is independent of α .

The solutions obtained in [5] are a special case of Family II.

Acknowledgements

We thank the referee for his/her valuable comments. YTG and BT were supported by the Fund for Excellent Young University Teachers and the Research Grants for Scholars Returning from Abroad, State Education Commission of China; and supported by the National Natural Science Foundation of China. WPH was supported by Catholic University of Taegu-Hyosung.

- [1] O. Bogoyavlenskii, *Izv. Akad. Nauk SSSR Ser. Mat.* **53**, 907 (1989).
- [2] O. Bogoyavlenskii, *Usp. Mat. Nauk* **45**, 17 (1990).
- [3] Y. Li, T. M. Φ. (a Russian journal) **99**, 441 (1994).
- [4] Y. T. Gao and B. Tian, *Computers Math. Applic.* **33**, 35 (1996).

- [5] Y. T. Gao and B. Tian, *Chaos, Solitons and Fractals* **8**, 879 (1997).
- [6] B. Tian and Y. T. Gao, *Mod. Phys. Lett. A* **10**, 2937 (1995); *Computer Phys. Comm.* **95**, 139 (1996).