

Structural Studies of N-(2,4,6-trisubstitutedphenyl)-Chloroacetamides, 2,4,6-X₃C₆H₂-NHCO-CH₂-Cl_y (X = CH₃ or Cl and y = 1 - 3).

B. Thimme Gowda*, I. Svoboda, and Hartmut Fuess

Darmstadt University of Technology, Institute of Materials Science,
Petersenstr.23, D-64287 Darmstadt

Permanent Address: Mangalore University Department of Chemistry,
Mangalagangothri-574 199, India

Reprint requests to Prof. B. T. G.; Fax: +91 824 742 367; E-mail: gowdabt@yahooO-Com
or Prof. H. F.; Fax: +49 6151 166023; E-mail: hfuess@tu-darmstadt.de

Z. Naturforsch. **55 a**, 779–790 (2000); received July 6, 2000

The effect of side chain and ring substitutions on the crystal structures of N-(2,4,6-substituted-phenyl)-chloroacetamides of the type, 2,4,6-X₃C₆H₂-NHCO-CH₂-Cl_y (X = CH₃ or Cl and 1 ≤ y ≤ 3), has been studied by determining the crystal structures of N-(2,4,6-trimethylphenyl)-2-chloroacetamide, 2,4,6-(CH₃)₃C₆H₂-NHCO-CH₂Cl (**TMPMCA**); N-(2,4,6-trichlorophenyl)-2-chloroacetamide, 2,4,6-Cl₃C₆H₂-NHCO-CH₂Cl (**TCPMCA**); N-(2,4,6-trichlorophenyl)-2,2-dichloroacetamide, 2,4,6-Cl₃C₆H₂-NHCO-CHCl₂ (**TCPDCA**) and N-(2,4,6-trichlorophenyl)-2,2,2-trichloroacetamide, 2,4,6-Cl₃C₆H₂-NHCO-CCl₃ (**TCPTCA**). The crystal type, space group, formula units and lattice constants in Å are: (**TMPMCA**): monoclinic, P2₁/n, Z = 8, a = 9.029(5), b = 15.688(2), c = 16.150(3), β = 96.77(2)°; (**TCPMCA**): orthorhombic, Pna2₁, Z = 8, a = 30.708(4), b = 4.685(1), c = 14.506(2); (**TCPDCA**) orthorhombic, P2₁2₁2, Z = 4, a = 16.459(3), b = 15.240(3), c = 4.640(1) and (**TCPTCA**): triclinic, P $\bar{1}$, Z = 4, a = 9.828(2), b = 11.953(2), c = 12.278(2), α = 71.05(1)°, β = 75.26(2)°, γ = 83.29(1)°. The results have been analysed along with the structures of the ring unsubstituted N-phenylacetamide (**PA**), N-phenyl-2,2,2-trichloroacetamide, side chain unsubstituted N-(2,4,6-trichlorophenyl)-acetamide, 2,4,6-Cl₃C₆H₂-NHCO-CH₃ (**TCPA**) and the corresponding di-substituted phenyl acetamides and chloroacetamides. The compounds **TMPMCA**, **TCPMCA** and **TCPTCA** have 2 molecules each in their asymmetric units. This is in agreement with the multiple lines observed in the ³⁵Cl NQR spectra of **TCPTCA**. Further, this compound shows disorder as indicated by the multiplicity of the C(side chain) - Cl bonds. The compound **TCPMCA** does not show ³⁵Cl NQR spectra, while the compound **TMPMCA** shows a single frequency which fades out above 200 K, indicating that it may undergo a phase transition well below room temperature. Conversion of 2,4,6-trichloroaniline into 2,4,6-trichlorophenyl acetamide and gradual replacement of H-atoms in the side chain of the latter decreases slightly the mean ring distances of the compounds, while the replacement of the 3 Cl by 3 CH₃ groups in the ring increases its mean distance. Changes in the mean ring distances are smaller as the effect has to be transmitted through the peptide linkage. The molecules in compounds **TCPDCA** and **TCPTCA** are linked in chains by -NH—O— hydrogen bonds (about 3 Å) between the amide groups of the molecules.

Key words: Crystal Structures; Trimethylphenyl-, Trichlorophenyl-chloroacetamides.

Introduction

Amides are of fundamental chemical interest because conjugation between nitrogen lone pair electrons and the carbonyl π-bond results in distinct physical and chemical properties [1]. The amide moiety is an important constituent of many biologically significant compounds [2]. Therefore an understanding

of the formation, properties and reactions of amides is central to future developments in areas such as polypeptide and protein chemistry. Many amides exhibit pharmacological activity, which has further stimulated interest in their chemistry. Many acetanilides also exhibit fungicidal and herbicidal activities [3 - 5].

Nuclear quadrupole resonance and crystal structural studies give valuable information on bond

Table 1. Experimental conditions for the crystal structure determination and crystallographic data of 2,4,6-(CH₃)₃C₆H₂-NHCO-CH₂Cl (**TMPMCA**); 2,4,6-Cl₃C₆H₂-NHCO-CH₂Cl (**TCPMCA**); 2,4,6-Cl₃C₆H₂-NHCO-CHCl₂ (**TCPDCA**), and 2,4,6-Cl₃C₆H₂-NHCO-CCl₃ (**TCPTCA**). Diffractometer: Nonius CAD 4; wavelength: 71.093 pm (MoK_α); Monochromator: Graphite (002); scan $\omega/2\theta = 1/1$; Refinement method: Full-matrix least-squares on F^2 .

Compound	TMPMCA	TCPMCA	TCPDCA	TCPTCA
Chemical formula	C ₁₁ H ₁₄ ClNO	C ₈ H ₅ Cl ₄ NO	C ₈ H ₄ Cl ₅ NO	C ₈ H ₃ Cl ₆ NO
Formula weight	211.71	272.94	307.38	341.82
Temperature, K	300(2)	300(2)	298(2)	298(2)
Crystal system	monoclinic	orthorhombic	orthorhombic	triclinic
Space group	P2 ₁ /n	Pna2 ₁	P2 ₁ 2 ₁ 2	P $\bar{1}$
Z	8	8	4	4
a, Å	9.029(5)	30.708(4)	16.459(3)	9.828(2)
b, Å	15.688(2)	4.685(1)	15.240(3)	11.953(2)
c, Å	16.150(3)	14.506(2)	4.640(1)	12.278(2)
α , deg.	90	90	90	71.05(1)
β , deg.	96.77 (2)	90	90	75.26(2)
γ , deg.	90	90	90	83.29(1)
Volume/Å ³	2271.6(14)	2086.9(6)	1163.9(4)	1318.4(4)
D_x /(g·cm ⁻³)	1.238	1.737	1.754	1.722
Abs. coeff., cm ⁻¹	3.03	11.03	12.20	12.90
$F(000)$	896	1088	608	672
Crystal size/(mm) ³	0.75×0.10×0.08	0.63×0.15×0.03	0.75×0.05×0.0188	0.875×0.10×0.075
θ range, deg.	1.82 to 24.98	1.33 to 25.98	1.82 to 22.98	1.80 to 24.97
Index ranges	-10 ≤ h ≤ 2, -18 ≤ k ≤ 0, -19 ≤ l ≤ 19	-37 ≤ h ≤ 37, -5 ≤ k ≤ 0, -17 ≤ l ≤ 0	-18 ≤ h ≤ 1, -16 ≤ k ≤ 1, -2 ≤ l ≤ 5	-11 ≤ h ≤ 1, -14 ≤ k ≤ 14, -14 ≤ l ≤ 14
Refl. coll.	4847	4206	1617	5350
Independ. refl.	3991	2126	1384	4643
[R_{int}]	0.0337	0.0506	0.0756	0.0246
Data	3991	2126	1384	4642
Restraints/parameters	0 / 255	1 / 259	0 / 136	120/328
Goodness-of-fit on F^2	1.022	1.027	1.068	1.048
Final R [$I > 2\sigma(I)$]	$R1 = 0.0632$, $wR2 = 0.1684$	0.0388 0.0900	0.0421 0.1027	0.0977 0.3042
R (all data)	$R1 = 0.1498$, $wR2 = 0.2149$	0.0644 0.1019	0.0644 0.1129	0.1317 0.3557
Absol. str. parameter	0.0004(14)	0.00(11)	-0.1(2)	—
Largest diff: (peak, hole), e·Å ⁻³	0.430, -0.304	0.369, -0.451	0.325, -0.432	1.018, -0.899

properties [6–9]. Thus we became interested in NQR and X-ray diffraction studies of amides in their crystalline state [10–18]. Pies, Rager, and Weiss [19] have studied ³⁵Cl NQR spectra of chloroacetanilides Cl_xC₆H_{5-x}NHCOCH_{3-y}Cl_y and observed interesting trends in their spectra. For example, 2,4,6-Cl₃C₆H₂-NHCO-CH₃ (**TCPA**) and 2,4,6-Cl₃C₆H₂-NHCO-CHCl₂ (**TCPDCA**) showed 3 and 5 ³⁵Cl NQR frequencies, respectively, as expected for 3 and 5 Cl atoms present in these compounds, but 2,4,6-Cl₃C₆H₂-NHCO-CH₂Cl (**TCPMCA**) did not show ³⁵Cl NQR resonance, while 2,4,6-Cl₃C₆H₂-NHCO-CCl₃ (**TCPTCA**) showed multiple ³⁵Cl NQR frequencies for the 6 Cl atoms

present. We have observed a single frequency for 2,4,6-(CH₃)₃C₆H₂-NHCO-CH₂Cl (**TCPMCA**), which fades out above 200 K [14]. There are no reports on structural studies of these compounds except for **TCPA** [20], to see how the -NHCO- bond parameters vary with substitution both in the benzene ring and in the side chain. Hence we thought it interesting to study the crystal structures of compounds of the type, 2,4,6-X₃C₆H₂-NHCO-CH_{3-y}Cl_y (X = CH₃ or Cl and 1 ≤ y ≤ 3) and analyse the data alongwith the crystal structures of the related compounds. We report herein our results on the compounds **TMPMCA**, **TCPMCA**, **TCPDCA**, and **TCPTCA**.

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$). $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U_{ij} tensor.

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> (eq)	Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> (eq)
2,4,6-(CH ₃) ₃ C ₆ H ₂ -NHCO-CH ₂ Cl (TMPMCA)									
Cl(1)	10528(2)	7941(1)	402(1)	98(1)	C(2)	11264(4)	7219(3)	−229(3)	65(1)
C(3)	10105(4)	6748(3)	−809(3)	48(1)	O(4)	8766(3)	6853(2)	−775(2)	61(1)
N(5)	10651(3)	6231(2)	−1345(2)	48(1)	C(6)	9738(4)	5729(3)	−1937(3)	45(1)
C(7)	9745(4)	4844(3)	−1839(3)	51(1)	C(8)	8817(4)	4371(3)	−2419(3)	57(1)
C(9)	7928(5)	4737(3)	−3070(3)	62(1)	C(10)	8000(5)	5610(3)	−3157(3)	66(1)
C(11)	8885(4)	6125(3)	−2603(3)	55(1)	C(12)	8917(6)	7073(3)	−2721(3)	77(2)
C(13)	6924(6)	4208(4)	−3675(4)	92(2)	C(14)	10691(5)	4418(3)	−1137(3)	66(1)
Cl(15)	10466(2)	794(1)	3772(1)	93(1)	C(16)	11187(4)	−204(3)	4050(3)	67(1)
C(17)	10056(4)	−913(3)	3931(3)	49(1)	O(18)	8710(3)	−770(2)	3916(2)	68(1)
N(19)	10631(3)	−1677(2)	3865(2)	50(1)	C(20)	9790(4)	−2452(3)	3814(3)	46(1)
C(21)	9972(4)	−3019(3)	4476(3)	52(1)	C(22)	9176(5)	−3785(3)	4405(3)	64(1)
C(23)	8220(6)	−3983(3)	3705(3)	65(1)	C(24)	8051(5)	−3405(3)	3073(3)	67(1)
C(25)	8825(4)	−2630(3)	3096(3)	53(1)	C(26)	8660(6)	−2039(4)	2361(3)	75(2)
C(27)	7443(7)	−4827(4)	3615(5)	111(2)	C(28)	11015(5)	−2826(4)	5248(3)	70(1)
2,4,6-Cl ₃ C ₆ H ₂ -NHCO-CHCl ₂ (TCPDCA)									
C(3)	10439(3)	2304(4)	5068(13)	39(2)	C(4)	9755(3)	1798(4)	6469(12)	32(1)
C(7)	8439(3)	1139(4)	5647(11)	29(1)	C(8)	8243(3)	311(4)	4682(12)	33(1)
C(9)	7551(3)	−125(4)	5586(14)	38(1)	C(10)	7055(3)	288(4)	7579(13)	40(2)
C(11)	7222(3)	1109(4)	8536(14)	37(1)	C(12)	7908(3)	1537(4)	7573(12)	32(1)
N(6)	9161(3)	1563(3)	4735(9)	31(1)	O(5)	9766(2)	1657(3)	9089(8)	48(1)
Cl(1)	11360(1)	2125(1)	6873(4)	50(1)	Cl(2)	10191(1)	3416(1)	5111(7)	88(1)
Cl(13)	8880(1)	−206(1)	2247(3)	50(1)	Cl(14)	6211(1)	−267(1)	8853(5)	65(1)
Cl(15)	8083(1)	2601(1)	8740(4)	48(1)					
2,4,6-Cl ₃ C ₆ H ₂ -NHCO-CH ₂ Cl (TCPMCA)									
Cl(1)	621(1)	10144(3)	−1(1)	56(1)	C(2)	234(2)	7746(10)	430(4)	45(1)
C(3)	−163(2)	9274(9)	803(4)	37(1)	O(4)	−177(2)	11853(7)	908(3)	52(1)
N(5)	−492(2)	7531(9)	1035(4)	40(1)	C(6)	−868(2)	8491(10)	1495(4)	36(1)
C(7)	−966(2)	7604(9)	2383(4)	39(1)	C(8)	−1320(2)	8564(10)	2868(4)	40(1)
C(9)	−1588(2)	10546(11)	2438(4)	43(1)	C(10)	−1516(2)	11460(12)	1555(4)	42(1)
C(11)	−1161(2)	10392(10)	1089(4)	40(1)	Cl(12)	−627(1)	5113(2)	2911(1)	51(1)
Cl(13)	−2032(1)	11893(3)	3049(1)	60(1)	Cl(14)	−1091(1)	11432(4)	−48(1)	61(1)
Cl(15)	234(1)	90(3)	3214(1)	57(1)	C(16)	639(2)	2408(11)	2832(5)	59(2)
C(17)	1016(2)	877(10)	2380(4)	40(1)	O(18)	1020(1)	−1692(6)	2269(3)	49(1)
N(19)	1341(2)	2627(8)	2119(3)	38(1)	C(20)	1707(2)	1685(9)	1621(4)	35(1)
C(21)	1800(2)	2702(9)	745(3)	35(1)	C(22)	2150(2)	1785(10)	242(4)	40(1)
C(23)	2418(2)	−232(11)	607(4)	42(1)	C(24)	2353(2)	−1263(11)	1500(4)	44(1)
C(25)	2003(2)	−268(10)	1996(4)	38(1)	Cl(26)	1451(1)	5209(3)	261(1)	47(1)
Cl(27)	2850(1)	−1563(4)	−23(1)	58(1)	Cl(28)	1947(1)	−1459(3)	3117(1)	54(1)

Experimental


Preparation and Characterisation of the Compounds

TMPMCA, **TCPMCA**, **TCPDCA** and **TCPTCA** were prepared from 2,4,6-trimethylaniline / 2,4,6-trichloroaniline and the corresponding chloroacetylchloride or chloroacetic acid and phosphoryl chloride (Aldrich, Germany) [10, 12, 19, 21]. The commercial trichloroaniline was purified by zone refining. The liquids, trimethylaniline, 2-chloro and 2,2-dichloro acetic acids were purified by double dis-

tillation. **TMPMCA** was prepared from 2,4,6-trimethylaniline and 2-chloroacetyl chloride in benzene (chemical analysis: % (found/calculated); C (62.25/62.41), H (6.52/6.67) and N (6.50/6.62)), while **TCPMCA** was prepared from 2,4,6-trichloroaniline and 2-chloroacetyl chloride in acetone. The compounds, **TCPDCA** and **TCPTCA** were prepared by treating 2,4,6-trichloroaniline with a clear mixture of 2,2-dichloroacetic acid / 2,2,2-trichloroacetic acid with phosphoryl chloride under constant stirring. The mixture was slowly warmed to expel HCl. Excess phosphoryl chloride was hydrolysed by adding cold

Table 2 (continued). 2,4,6-Cl₃C₆H₂-NHCO-CCl₃ (TCPTCA).

Atom	x	y	z	U(eq)	Atom	x	y	z	U(eq)
Cl(1A)	3058(35)	6660(57)	986(74)	143(9)	Cl(1B)	3167(32)	6311(46)	1144(66)	143(9)
Cl(1C)	3299(47)	5899(47)	1554(42)	143(9)	Cl(1D)	3872(34)	5440(26)	1658(30)	143(9)
Cl(1E)	5730(53)	5767(71)	758(98)	143(9)	Cl(1F)	4933(30)	5398(23)	1276(32)	143(9)
Cl(2A)	6208(32)	6093(38)	682(21)	79(2)	Cl(2B)	6122(32)	6151(38)	500(23)	79(2)
Cl(2C)	3012(16)	7016(19)	848(13)	79(2)	Cl(2D)	5862(50)	5758(38)	975(31)	79(2)
Cl(2E)	3510(24)	7805(27)	479(18)	79(2)	Cl(3A)	5074(43)	8467(31)	28(42)	115(3)
Cl(3B)	4350(21)	8266(16)	231(14)	115(3)	Cl(3C)	5299(43)	8355(35)	-58(42)	115(3)
Cl(3D)	6185(29)	6685(36)	162(24)	115(3)	Cl(3E)	5900(32)	7757(38)	-88(22)	115(3)
C(4)	4685(8)	6920(8)	1129(9)	85(3)	C(5)	4754(9)	7058(9)	2307(8)	62(2)
O(6)	5841(6)	6861(8)	2613(6)	78(2)	N(7)	3565(7)	7418(6)	2926(6)	57(2)
C(8)	3506(7)	7509(7)	4063(7)	51(2)	C(9)	3368(8)	6517(8)	5037(8)	55(2)
C(10)	3390(10)	6528(9)	6144(9)	67(2)	C(11)	3525(10)	7595(10)	6288(9)	69(2)
C(12)	3568(10)	8633(10)	5364(10)	73(3)	C(13)	3601(9)	8578(8)	4240(9)	65(2)
Cl(14)	3227(3)	5157(2)	4854(2)	75(1)	Cl(15)	3524(4)	7655(4)	7679(3)	114(1)
Cl(16)	3744(4)	9860(3)	3086(3)	105(1)	Cl(17)	398(4)	994(3)	5474(3)	92(1)
Cl(18)	2898(2)	2214(2)	5098(2)	71(1)	Cl(19)	308(3)	3535(2)	4779(3)	83(1)
C(20)	1035(9)	2234(7)	5627(7)	54(2)	C(21)	527(8)	2191(7)	6931(7)	51(2)
O(22)	-664(7)	1956(8)	7422(6)	79(2)	N(23)	1494(7)	2408(6)	7419(6)	54(2)
C(24)	1129(8)	2329(8)	8655(7)	54(2)	C(25)	1417(11)	1277(9)	9464(9)	69(2)
C(26)	1134(11)	1168(10)	10671(9)	76(3)	C(27)	563(10)	2119(9)	11014(8)	68(2)
C(28)	242(10)	3161(8)	10237(8)	64(2)	C(29)	539(9)	3256(7)	9047(7)	54(2)
Cl(31)	257(4)	2028(3)	12498(2)	104(1)	Cl(30)	2149(5)	77(3)	8994(4)	116(1)
Cl(32)	199(3)	4589(2)	8043(2)	79(1)					

Fig. 1. Molecular geometry of N-(2,4,6-trimethylphenyl)-2-chloroacetamide, 2,4,6-(CH₃)₃C₆H₂-NHCO-CH₂Cl (TCPTCA), with the numbering of atoms.

water drop wise under ice cold conditions. The solid was filtered under suction, washed thoroughly with water and dried. The compounds were recrystallised from ethanol several times. The purity of the com-

pounds **TCPTCA**, **TCPDCA**, and **TCPTCA** was checked by determining their melting points. The melting points in °C are (the values in parentheses are literature values [19, 21]): (**TCPTCA**) 180 (180);

Table 3. Bond lengths (Å) (standard deviation).

Connection	Bond length	Connection	Bond length
TMPMCA			
Cl(1)-C(2)	1.709(5)	C(2)-C(3)	1.513(6)
C(3)-O(4)	1.228(4)	C(3)-N(5)	1.324(5)
N(5)-C(6)	1.423(5)	C(6)-C(11)	1.394(6)
C(6)-C(7)	1.397(6)	C(7)-C(8)	1.395(6)
C(7)-C(14)	1.495(6)	C(8)-C(9)	1.372(6)
C(9)-C(10)	1.379(7)	C(9)-C(13)	1.502(7)
C(10)-C(11)	1.387(6)	C(11)-C(12)	1.501(7)
Cl(15)-C(16)	1.734(5)	C(16)-C(17)	1.507(6)
C(17)-O(18)	1.233(4)	C(17)-N(19)	1.315(5)
N(19)-C(20)	1.431(5)	C(20)-C(21)	1.386(6)
C(20)-C(25)	1.394(6)	C(21)-C(22)	1.399(7)
C(21)-C(28)	1.502(6)	C(22)-C(23)	1.374(7)
C(23)-C(24)	1.359(7)	C(23)-C(27)	1.497(8)
C(24)-C(25)	1.401(7)	C(25)-C(26)	1.500(7)
TCPMCA			
Cl(1)-C(2)	1.750(5)	C(2)-C(3)	1.514(8)
C(3)-O(4)	1.218(6)	C(3)-N(5)	1.342(7)
N(5)-C(6)	1.408(7)	C(6)-C(7)	1.386(8)
C(6)-C(11)	1.395(7)	C(7)-C(8)	1.372(8)
C(7)-Cl(12)	1.740(5)	C(8)-C(9)	1.389(8)
C(9)-C(10)	1.368(8)	C(9)-Cl(13)	1.744(6)
C(10)-C(11)	1.377(8)	C(11)-Cl(14)	1.733(6)
Cl(15)-C(16)	1.740(6)	C(16)-C(17)	1.511(8)
C(17)-O(18)	1.214(6)	C(17)-N(19)	1.347(7)
N(19)-C(20)	1.407(7)	C(20)-C(21)	1.387(7)
C(20)-C(25)	1.399(7)	C(21)-C(22)	1.368(8)
C(21)-Cl(26)	1.738(5)	C(22)-C(23)	1.361(8)
C(23)-C(24)	1.398(9)	C(23)-Cl(27)	1.726(6)
C(24)-C(25)	1.375(8)	C(25)-Cl(28)	1.728(6)
TCPA			
C(1)-C(2)	1.499(3)	C(2)-O(3)	1.221(3)
C(2)-N(4)	1.351(3)	N(4)-C(5)	1.413(3)
C(5)-C(10)	1.387(3)	C(5)-C(6)	1.398(3)
C(6)-C(7)	1.379(3)	C(6)-Cl(11)	1.722(2)
C(7)-C(8)	1.374(3)	C(8)-C(9)	1.384(3)
C(8)-Cl(12)	1.735(2)	C(9)-C(10)	1.382(3)
C(10)-Cl(13)	1.727(2)		
TCPDCA			
C(3)-Cl(1)	1.754(6)	C(3)-Cl(2)	1.744(6)
C(3)-C(4)	1.511(7)	C(4)-O(5)	1.235(7)
C(4)-N(6)	1.316(7)	O(5)-H(6N)	3.000(6)
C(7)-N(6)	1.417(7)	C(7)-C(8)	1.377(8)
C(7)-C(12)	1.390(8)	C(8)-C(9)	1.385(8)
C(8)-Cl(13)	1.731(6)	C(9)-C(10)	1.384(9)
C(10)-C(11)	1.355(8)	C(10)-Cl(14)	1.732(6)
C(11)-C(12)	1.378(8)	C(12)-Cl(15)	1.734(5)

Table 3 (continued).

Connection	Bond length	Connection	Bond length
TCPTCA			
Cl(1A)-C(4)	1.73(2)	Cl(1B)-C(4)	1.73(2)
Cl(1C)-C(4)	1.80(2)	Cl(1D)-C(4)	1.87(2)
Cl(1E)-C(4)	1.73(2)	Cl(1F)-C(4)	1.76(2)
Cl(2A)-C(4)	1.76(2)	Cl(2B)-C(4)	1.74(2)
Cl(2C)-C(4)	1.75(2)	Cl(2D)-C(4)	1.73(2)
Cl(2E)-C(4)	1.65(2)	Cl(3A)-C(4)	1.92(2)
Cl(3B)-C(4)	1.68(2)	Cl(3C)-C(4)	1.90(2)
Cl(3D)-C(4)	1.70(2)	Cl(3E)-C(4)	1.75(2)
C(4)-C(5)	1.526(13)	C(5)-O(6)	1.194(10)
C(5)-N(7)	1.331(11)	O(6)-H(7N)	2.982(9)
N(7)-C(8)	1.422(11)	C(8)-C(9)	1.374(12)
C(8)-C(13)	1.381(12)	C(9)-C(10)	1.369(13)
C(9)-Cl(14)	1.738(8)	C(10)-C(11)	1.368(14)
C(11)-C(12)	1.380(20)	C(11)-Cl(15)	1.733(10)
C(12)-C(13)	1.394(14)	C(13)-Cl(16)	1.707(10)
Cl(17)-C(20)	1.755(8)	Cl(18)-C(20)	1.781(9)
Cl(19)-C(20)	1.755(9)	C(20)-C(21)	1.536(11)
C(21)-O(22)	1.192(10)	C(21)-N(23)	1.333(9)
O(22)-H(23N)	2.998(9)	N(23)-C(24)	1.441(10)
C(24)-C(29)	1.360(12)	C(24)-C(25)	1.377(13)
C(25)-C(26)	1.401(14)	C(25)-Cl(30)	1.731(10)
C(26)-C(27)	1.350(20)	C(27)-C(28)	1.360(14)
C(27)-Cl(31)	1.739(9)	C(28)-C(29)	1.384(12)
C(29)-Cl(32)	1.727(8)		

Table 4. Bond angles (degree) (standard deviation).

Connection	Bond angle	Connection	Bond angle
TMPMCA			
C(3)-C(2)-Cl(1)	113.8(3)	O(4)-C(3)-N(5)	123.8(4)
O(4)-C(3)-C(2)	121.3(4)	N(5)-C(3)-C(2)	114.9(3)
C(3)-N(5)-C(6)	123.2(3)	C(7)-C(6)-N(5)	118.6(4)
C(11)-C(6)-N(5)	119.7(4)	C(11)-C(6)-C(7)	121.7(4)
C(8)-C(7)-C(6)	117.3(4)	C(8)-C(7)-C(14)	121.0(4)
C(6)-C(7)-C(14)	121.7(4)	C(9)-C(8)-C(7)	122.8(4)
C(8)-C(9)-C(10)	117.6(4)	C(8)-C(9)-C(13)	121.4(5)
C(10)-C(9)-C(13)	121.0(5)	C(9)-C(10)-C(11)	122.9(4)
C(10)-C(11)-C(6)	117.5(4)	C(10)-C(11)-C(12)	120.9(4)
C(6)-C(11)-C(12)	121.5(4)		
C(17)-C(16)-Cl(15)	113.9(3)	O(18)-C(17)-N(19)	124.2(4)
O(18)-C(17)-C(16)	121.3(4)	N(19)-C(17)-C(16)	114.5(3)
C(17)-N(19)-C(20)	124.6(3)	C(21)-C(20)-C(25)	121.3(4)
C(21)-C(20)-N(19)	119.1(4)	C(25)-C(20)-N(19)	119.6(4)
C(20)-C(21)-C(22)	118.3(4)	C(20)-C(21)-C(28)	121.1(4)
C(22)-C(21)-C(28)	120.5(4)	C(23)-C(22)-C(21)	121.9(5)
C(24)-C(23)-C(22)	118.1(5)	C(24)-C(23)-C(27)	120.5(5)
C(22)-C(23)-C(27)	121.3(5)	C(23)-C(24)-C(25)	123.2(4)
C(20)-C(25)-C(24)	117.2(4)	C(20)-C(25)-C(26)	121.9(4)
C(24)-C(25)-C(26)	120.8(4)		

Crystal Structure Studies

Small crystals of **TMPMCA**, **TCPMCA**, **TCPDCA** and **TCPTCA** were selected for X-ray

(**TCPDCA**) 186 (186 - 187); (**TCPTCA**) 119 (118 - 120). The compounds were further characterised by recording their infrared and/or ^{35}Cl NQR spectra and comparing the frequencies with the literature values. All other reagents employed in the preparations and purification of compounds were of analytical grade.

Table 4 (continued).

Connection	Bond angle	Connection	Bond angle
TCPMCA			
C(3)-C(2)-Cl(1)	111.8(3)	O(4)-C(3)-N(5)	123.1(5)
O(4)-C(3)-C(2)	122.8(5)	N(5)-C(3)-C(2)	114.1(4)
C(3)-N(5)-C(6)	122.8(4)	C(7)-C(6)-N(5)	121.5(5)
C(11)-C(6)-N(5)	122.2(5)	C(7)-C(6)-C(11)	116.4(5)
C(8)-C(7)-C(6)	123.4(5)	C(8)-C(7)-Cl(12)	117.9(4)
C(6)-C(7)-Cl(12)	118.7(4)	C(7)-C(8)-C(9)	117.3(5)
C(10)-C(9)-C(8)	122.3(5)	C(10)-C(9)-Cl(13)	119.2(5)
C(8)-C(9)-Cl(13)	118.5(4)	C(9)-C(10)-C(11)	118.2(5)
C(10)-C(11)-C(6)	122.4(5)	C(10)-C(11)-Cl(14)	117.5(4)
C(6)-C(11)-Cl(14)	120.1(4)		
C(17)-C(16)-Cl(15)	112.8(4)	O(18)-C(17)-N(19)	123.9(5)
O(18)-C(17)-C(16)	122.5(5)	N(19)-C(17)-C(16)	113.6(4)
C(17)-N(19)-C(20)	123.1(4)	C(21)-C(20)-C(25)	116.5(5)
C(21)-C(20)-N(19)	121.8(5)	C(25)-C(20)-N(19)	121.6(5)
C(22)-C(21)-C(20)	122.9(5)	C(22)-C(21)-Cl(26)	118.7(4)
C(20)-C(21)-Cl(26)	118.4(4)	C(23)-C(22)-C(21)	119.1(5)
C(22)-C(23)-C(24)	120.9(5)	C(22)-C(23)-Cl(27)	120.7(5)
C(24)-C(23)-Cl(27)	118.4(4)	C(25)-C(24)-C(23)	118.6(5)
C(24)-C(25)-C(20)	121.8(5)	C(24)-C(25)-Cl(28)	117.4(4)
C(20)-C(25)-Cl(28)	120.8(4)		
TCPDCA			
C(4)-C(3)-Cl(2)	108.5(4)	C(4)-C(3)-Cl(1)	111.1(4)
Cl(2)-C(3)-Cl(1)	110.4(3)	O(5)-C(4)-N(6)	124.4(5)
O(5)-C(4)-C(3)	120.1(5)	N(6)-C(4)-C(3)	115.4(5)
C(8)-C(7)-C(12)	117.4(5)	C(8)-C(7)-N(6)	121.1(5)
C(12)-C(7)-N(6)	121.4(5)	C(7)-C(8)-C(9)	122.3(5)
C(7)-C(8)-Cl(13)	119.2(4)	C(9)-C(8)-Cl(13)	118.5(5)
C(10)-C(9)-C(8)	117.9(5)	C(11)-C(10)-C(9)	121.3(5)
C(11)-C(10)-Cl(14)	120.1(5)	C(9)-C(10)-Cl(14)	118.6(4)
C(10)-C(11)-C(12)	119.8(5)	C(11)-C(12)-C(7)	121.2(5)
C(11)-C(12)-Cl(15)	118.6(4)	C(7)-C(12)-Cl(15)	120.3(4)
C(4)-N(6)-C(7)	124.4(4)		
TCPA			
O(3)-C(2)-N(4)	123.2(2)	O(3)-C(2)-C(1)	122.0(2)
N(4)-C(2)-C(1)	114.8(2)	C(2)-N(4)-C(5)	123.2(2)
C(10)-C(5)-C(6)	117.3(2)	C(10)-C(5)-N(4)	122.5(2)
C(6)-C(5)-N(4)	120.1(2)	C(7)-C(6)-C(5)	121.9(2)
C(7)-C(6)-Cl(11)	119.1(2)	C(5)-C(6)-Cl(11)	119.0(2)
C(8)-C(7)-C(6)	118.5(2)	C(7)-C(8)-C(9)	121.9(2)
C(7)-C(8)-Cl(12)	119.1(2)	C(9)-C(8)-Cl(12)	119.0(2)
C(10)-C(9)-C(8)	118.2(2)	C(9)-C(10)-C(5)	122.1(2)
C(9)-C(10)-Cl(13)	118.1(2)	C(5)-C(10)-Cl(13)	119.8(2)

diffraction and studied at room temperature. The collected intensity data were corrected for Lorentz polarisation and absorption. The positional parameters were determined by direct methods and least squares refinement (SHELXL-93) [22 - 27]. For locating the hydrogen atom positions, the C-H distances were fixed to 0.93 Å for the ring hydrogen atoms, while the side chain C-H distances were fixed to 0.96 Å for the CH₃ group, to 0.97 Å for the CH₂Cl group

Table 4 (continued). TCPTCA.

Connection	Bond angle	Connection	Bond angle
C(5)-C(4)-Cl(1A)	117.0(3)	C(5)-C(4)-Cl(1B)	115.0(2)
C(5)-C(4)-Cl(1E)	115.0(3)	C(5)-C(4)-Cl(2A)	107.0(9)
C(5)-C(4)-Cl(2B)	114.9(11)	C(5)-C(4)-Cl(2C)	116.7(8)
C(5)-C(4)-Cl(2D)	104.3(12)	C(5)-C(4)-Cl(2E)	114.8(9)
C(5)-C(4)-Cl(3B)	107.9(9)	C(5)-C(4)-Cl(3D)	120.4(14)
C(5)-C(4)-Cl(3E)	113.8(13)	Cl(1A)-C(4)-Cl(2A)	122.0(2)
Cl(1B)-C(4)-Cl(2B)	108.0(2)	Cl(2E)-C(4)-Cl(1E)	130.0(3)
Cl(3D)-C(4)-Cl(2C)	123.0(2)	O(6)-C(5)-N(7)	124.0(8)
O(6)-C(5)-C(4)	119.8(8)	N(7)-C(5)-C(4)	116.2(7)
C(5)-N(7)-C(8)	120.3(7)	C(9)-C(8)-Cl(13)	117.2(8)
C(9)-C(8)-N(7)	120.6(7)	C(13)-C(8)-N(7)	122.2(8)
C(10)-C(9)-C(8)	123.7(8)	C(10)-C(9)-Cl(14)	117.6(7)
C(8)-C(9)-Cl(14)	118.6(7)	C(11)-C(10)-C(9)	117.8(9)
C(10)-C(11)-C(12)	121.2(9)	C(10)-C(11)-Cl(15)	119.4(8)
C(12)-C(11)-Cl(15)	119.3(8)	C(11)-C(12)-C(13)	119.0(8)
C(8)-C(13)-C(12)	120.7(9)	C(8)-C(13)-Cl(16)	120.4(8)
C(12)-C(13)-Cl(16)	118.8(7)	C(21)-C(20)-Cl(17)	107.8(6)
C(21)-C(20)-Cl(18)	114.5(5)	C(21)-C(20)-Cl(19)	107.5(6)
Cl(17)-C(20)-Cl(18)	108.2(5)	Cl(17)-C(20)-Cl(19)	109.8(4)
Cl(19)-C(20)-Cl(18)	109.0(5)	O(22)-C(21)-N(23)	125.7(8)
O(22)-C(21)-C(20)	118.7(7)	N(23)-C(21)-C(20)	115.7(7)
C(21)-N(23)-C(24)	119.4(7)	C(29)-C(24)-C(25)	119.0(8)
C(29)-C(24)-N(23)	122.6(7)	C(25)-C(24)-N(23)	118.4(8)
C(24)-C(25)-C(26)	120.6(9)	C(24)-C(25)-Cl(30)	120.1(8)
C(26)-C(25)-Cl(30)	119.3(8)	C(27)-C(26)-C(25)	118.1(9)
C(26)-C(27)-C(28)	122.5(9)	C(26)-C(27)-Cl(31)	119.2(8)
C(28)-C(27)-Cl(31)	118.3(8)	C(27)-C(28)-C(29)	118.6(9)
C(28)-C(29)-C(28)	121.1(8)	C(24)-C(29)-Cl(32)	119.6(6)
C(28)-C(29)-Cl(32)	119.2(7)		

and to 0.98 Å for the CHCl₂ group. Further experimental conditions for structure determinations and refinements are given in Table 1. For comparison purposes, the compound N-(2,4,6-trichlorophenyl)-acetamide (**TCPA**) was also prepared and the crystal structures of this compound and 2,4,6-trichloroaniline (**TCA**) have been determined under identical condition.

Results and Discussion

The crystallographic data for **TMPMCA**, **TCPMCA**, **TCPDCA**, and **TCPTCA** are given in Table 1. The atomic coordinates and the mean displacement parameters are listed in Table 2. The intramolecular bond distances and bond angles are given in Tables 3 and 4, respectively. Table 5 lists the dihedral angles of **TCPDCA** and **TCPTCA**. The hydrogen coordinates, anisotropic displacement parameters and further information on the crystal structure determinations have been deposited at the Cambridge Crystallographic Data Centre, the CCDC numbers are 147743 - 147746. In Figs. 1, 3, 5 and 7 we

Table 5. Dihedral angles (degree) (standard deviations).

Connection	Angle	Connection	Angle	Connection	Angle
2,4,6-Cl ₃ C ₆ H ₂ .NHCO.CHCl ₂ (TCPDCA)					
Cl(2)-C(3)-C(4)-O(5)	-90.2(6)	Cl(1)-C(3)-C(4)-O(5)	31.3(7)	Cl(2)-C(3)-C(4)-N(6)	87.5(5)
Cl(1)-C(3)-C(4)-N(6)	-151.1(4)	C(12)-C(7)-C(8)-C(9)	-0.6(8)	N(6)-C(7)-C(8)-C(9)	178.6(5)
C(12)-C(7)-C(8)-Cl(13)	179.2(4)	N(6)-C(7)-C(8)-Cl(13)	-1.6(7)	C(7)-C(8)-C(9)-C(10)	-1.5(8)
Cl(13)-C(8)-C(9)-C(10)	178.7(4)	C(8)-C(9)-C(10)-C(11)	2.6(8)	C(8)-C(9)-C(10)-Cl(14)	-177.2(4)
C(9)-C(10)-C(11)-C(12)	-1.6(9)	Cl(14)-C(10)-C(11)-C(12)	178.2(5)	C(10)-C(11)-C(12)-C(7)	-0.6(8)
C(10)-C(11)-C(12)-Cl(15)	177.7(5)	C(8)-C(7)-C(12)-C(11)	1.7(7)	N(6)-C(7)-C(12)-C(11)	-177.5(5)
C(8)-C(7)-C(12)-Cl(15)	-176.6(4)	N(6)-C(7)-C(12)-Cl(15)	4.2(7)	O(5)-C(4)-N(6)-C(7)	2.1(9)
C(3)-C(4)-N(6)-C(7)	-175.4(5)	C(8)-C(7)-N(6)-C(4)	-117.9(6)	C(12)-C(7)-N(6)-C(4)	61.2(7)
2,4,6-Cl ₃ C ₆ H ₂ .NHCO.CCl ₃ (TCPTCA)					
Cl(2E)-C(4)-C(5)-O(6)	-150.0(2)	Cl(3B)-C(4)-C(5)-O(6)	-114.2(12)	Cl(3D)-C(4)-C(5)-O(6)	-9(2)
Cl(3E)-C(4)-C(5)-O(6)	-56(2)	Cl(1A)-C(4)-C(5)-O(6)	156(3)	Cl(1B)-C(4)-C(5)-O(6)	141(2)
Cl(2A)-C(4)-C(5)-O(6)	16(2)	Cl(2B)-C(4)-C(5)-O(6)	14(2)	Cl(1E)-C(4)-C(5)-O(6)	36(4)
Cl(2C)-C(4)-C(5)-O(6)	171(11)	Cl(2D)-C(4)-C(5)-O(6)	34(2)	Cl(1E)-C(4)-C(5)-N(7)	-145(4)
Cl(2E)-C(4)-C(5)-N(7)	29(2)	Cl(3E)-C(4)-C(5)-N(7)	123(2)	Cl(3B)-C(4)-C(5)-N(7)	64.8(11)
Cl(3D)-C(4)-C(5)-N(7)	170(2)	Cl(1A)-C(4)-C(5)-N(7)	-25(3)	Cl(1B)-C(4)-C(5)-N(7)	-40(2)
Cl(2C)-C(4)-C(5)-N(7)	-10.0(14)	Cl(2D)-C(4)-C(5)-N(7)	-147(2)	Cl(2A)-C(4)-C(5)-N(7)	-165(2)
Cl(2B)-C(4)-C(5)-N(7)	-167(2)	O(6)-C(5)-N(7)-C(8)	-4.6(14)	C(4)-C(5)-N(7)-C(8)	176.5(7)
C(5)-N(7)-C(8)-C(9)	-78.8(10)	C(5)-N(7)-C(8)-C(13)	100.3(10)	C(13)-C(8)-C(9)-C(10)	-3.3(12)
N(7)-C(8)-C(9)-C(10)	175.9(8)	C(13)-C(8)-C(9)-Cl(14)	179.4(6)	N(7)-C(8)-C(9)-Cl(14)	-1.5(10)
C(8)-C(9)-C(10)-C(11)	1.5(13)	Cl(14)-C(9)-C(10)-C(11)	179.0(7)	C(9)-C(10)-C(11)-C(12)	3.1(14)
C(9)-C(10)-C(11)-Cl(15)	178.8(7)	C(10)-C(11)-C(12)-C(13)	-5.8(14)	Cl(15)-C(11)-C(12)-C(13)	178.5(7)
C(9)-C(8)-C(13)-C(12)	0.4(12)	N(7)-C(8)-C(13)-C(12)	-178.7(8)	C(9)-C(8)-C(13)-Cl(16)	-178.9(6)
N(7)-C(8)-C(13)-Cl(16)	2.0(11)	C(11)-C(12)-C(13)-C(8)	4.0(14)	C(11)-C(12)-C(13)-Cl(16)	-176.7(8)
Cl(17)-C(20)-C(21)-O(22)	-44.3(10)	Cl(19)-C(20)-C(21)-O(22)	74.0(9)	Cl(18)-C(20)-C(21)-O(22)	-164.7(7)
Cl(17)-C(20)-C(21)-N(23)	134.6(7)	Cl(19)-C(20)-C(21)-N(23)	-107.1(7)	Cl(18)-C(20)-C(21)-N(23)	14.2(10)
O(22)-C(21)-N(23)-C(24)	1.9(13)	C(20)-C(21)-N(23)-C(24)	-176.9(7)	C(21)-N(23)-C(24)-C(29)	-87.1(10)
C(21)-N(23)-C(24)-C(25)	94.1(10)	C(29)-C(24)-C(25)-C(26)	-1.2(14)	N(23)-C(24)-C(25)-C(26)	177.5(8)
C(29)-C(24)-C(25)-Cl(30)	179.2(7)	N(23)-C(24)-C(25)-Cl(30)	-2.0(11)	C(24)-C(25)-C(26)-C(27)	1(2)
Cl(30)-C(25)-C(26)-C(27)	180.0(8)	C(25)-C(26)-C(27)-C(28)	1(2)	C(25)-C(26)-C(27)-Cl(31)	-177.7(8)
C(26)-C(27)-C(28)-C(29)	-1(2)	Cl(31)-C(27)-C(28)-C(29)	177.3(7)	C(25)-C(26)-C(29)-C(28)	0.8(13)
N(23)-C(24)-C(29)-C(28)	-177.9(7)	C(25)-C(24)-C(29)-Cl(32)	178.8(7)	N(23)-C(24)-C(29)-Cl(32)	0.1(11)
C(27)-C(28)-C(29)-C(24)	0.5(13)	C(27)-C(28)-C(29)-Cl(32)	-177.6(7)		

show the molecules of the title compounds as they appear in suitable projection with the numbering of the atoms used throughout the paper. The projections of the unit cells of **TMPMCA**, **TCPMCA**, **TCPDCA**, and **TCPTCA** are shown in Figures 2, 4, 6 and 8, respectively.

TMPMCA, **TCPMCA**, and **TCPTCA** show two molecules each in their asymmetric units. This is in agreement with the multiple lines observed in the ³⁵Cl NQR spectrum of **TCPTCA** [19]. Further, this compound shows disorder as indicated by the multiplicity of the C(side chain) - Cl bonds. **TCPTCA** does not show ³⁵Cl NQR spectra, while **TMPMCA** shows single frequency which fades out above 200 K, indicating that it may undergo phase transition well below room temperature. First we shall discuss the intramolecular geometry of the four title compounds. The individual data are given in Tables 3 and 4. Com-

paring the C(i)-C(j) bond distances within the benzene rings of the four compounds we find the following mean distances in angstrom units. The observed minimum and maximum bond lengths in Å are given in parentheses.: 1.386 (1.359, 1.401), 1.381 (1.361, 1.399), 1.378 (1.355, 1.390) and 1.375 (1.350, 1.401) Å for the compounds **TMPMCA**, **TCPMCA**, **TCPDCA**, and **TCPTCA**, respectively. We have also determined the crystal structures of **TCA** and **TCPA** under the present experimental conditions and find the mean ring distance in them as 1.386 (1.367, 1.399) Å and 1.384 (1.374, 1.398) Å, respectively. As may be seen, conversion of **TCA** into **TCPA** and gradual replacement of H-atoms in the side chain of the latter to give the compounds **TCPMCA**, **TCPDCA**, and **TCPTCA** with the side chains -CH₂Cl, -CHCl₂ and -CCl₃, respectively, slightly decreases the mean ring distances of these compounds. Changes in the




Fig. 2. Projection of the unit cell of **TPMCA** along [100] on to the *bc* plane.




Fig. 3. Molecular geometry of N-(2,4,6-trichlorophenyl)-2-chloroacetamide, $2,4,6\text{-Cl}_3\text{C}_6\text{H}_2\text{-NHCO-CH}_2\text{Cl}$ (**TCPMCA**), with the numbering of atoms.




Fig. 4. Projection of the unit cell of **TCPMCA** along [101] on to the *ac* plane.




Fig. 5. Molecular geometry of N-(2,4,6-trichlorophenyl)-2,2-dichloroacetamide, 2,4,6-Cl₃C₆H₂-NHCO-CHCl₂ (TCPDCA), with the numbering of atoms.




Fig. 6. Projection of the unit cell of TCPDCA along [110] on to the *ab* plane.

mean ring distances are smaller as the effect has to be transmitted through the peptide linkage. Further, the mean ring distances of **TMPMCA** and **TCPMCA** are 1.386 and 1.381 Å, respectively, indicating that the replacement of 3 Cl (electron withdrawing) atoms by 3 CH₃ (electron donating) groups in the ring increases the mean ring distance. The C(ring)-Cl and the C(ring) - N distances for **TCA**, **TCPA**, **TMPMCA**, **TCPMCA**, **TCPDCA**, and **TCPTCA** do not show regular trends.

As regards the bond angles, they are observed between 115.1° and 123.3° (**TCA**), 117.3° and 122.1° (**TCPA**), 117.2 and 123.2° (**TMPMCA**),

116.5° and 123.2° (**TCPMCA**), 117.4° and 122.3° (**TCPDCA**) 118.0° and 122.2° (**TCPTCA**). The C2(ring)-C1(ring)-N and C6(ring)-C1(ring)-N angles for the compounds are 122.4° and 122.0° (**TCA**), 120.1° and 122.5° (**TCPA**), 118.9 and 119.7° (**TMPMCA**), 121.7° and 121.9° (**TCPMCA**), 121.1° and 121.4° (**TCPDCA**), 119.5° and 122.4° (**TCPTCA**). Similarly the C2(ring)-C1(ring)-C6(ring) bond angles are 115.6°, 117.3°, 121.5°, 116.5°, 117.4° and 118.1° for **TCA**, **TCPA**, **TMPMCA**, **TCPMCA**, **TCPDCA** and **TCPTCA**, respectively, while the C(ring)-N-C(O) bond angles vary as 123.2° (**TCPA**), 123.9° (**TMPMCA**),




Fig. 7. Molecular geometry of N-(2,4,6-trichlorophenyl)-2,2,2-trichloroacetamide, 2,4,6-Cl₃C₆H₂-NHCO-CCl₃ (**TCPTCA**), with the numbering of atoms.




Fig. 8. Projection of the unit cell of **TCPTCA** along [101] on to the *ac* plane.

123.0° (**TCPMCA**), 124.4° (**TCPDCA**) and 119.9° (**TCPTCA**). The N-C(O)-C(side chain) angles are 114.8° (**TCPA**), 114.7° (**TMPMCA**), 113.9° (**TCPMCA**), 115.4° (**TCPDCA**) and 116.0° (**TCPTCA**), respectively.

We have recently determined the crystal structures of several N-phenyl and N-(mono-substituted phenyl)-2,2-dichloroacetamides and 2,2,2-trichloroacetamides [18]. Hence we compare here the present crystal structure data on **TMPMCA**,

TCPMCA, **TCPDCA** and **TCPTCA** with the above data. We shall first compare the results of 2,4,6-Cl₃C₆H₂-NHCO-CCl₃ (**TCPTCA**) with 2,6-Cl₂C₆H₃-NHCO-CCl₃ (**DCPTCA**) [28], 2-ClC₆H₄-NHCO-CCl₃ (**MCPTCA**) [18] and C₆H₅-NHCO-CCl₃ (**PTCA**) [11]. The average ring C-C bond distances in Å are (minimum and maximum bond distances are shown in parentheses) 1.372 (1.350, 1.401) (**TCPTCA**); 1.380 (1.372, 1.385) (**DCPTCA**), 1.383 (1.368, 1.391) (**MCPTCA**), 1.383 (1.373, 1.390)

(**PTCA**). There are no significant changes in the other bond distances on going from (**PTCA**) to (**TCPTCA**), except for C(O)-C(side chain) bond distances which decrease slightly as we go from $C_6H_5-NHCO-CCl_3$ (**PTCA**) to $2,4,6-Cl_3C_6H_2-NHCO-CCl_3$ (**TCPTCA**): 1.564 Å (**PTCA**), 1.561 Å (**MCPTCA**), 1.550 Å (**DCPTCA**) and 1.531 Å (**TCPTCA**). The C(ring)-N-C(O) and N-C(O)-C(side chain) bond angles show slight decreasing and increasing trends as we go from $C_6H_5-NHCO-CCl_3$ (**PTCA**) to $2,4,6-Cl_3C_6H_2-NHCO-CCl_3$ (**TCPTCA**): 125.4°, 114.8° (**PTCA**), 122.4°, 114.1° (**MCPTCA**), 121.4°, 114.8° (**DCPTCA**) and 119.9°, 116° (**TCPTCA**).

Comparison of the crystal structure data of $2,4,6-Cl_3C_6H_2-NHCO-CHCl_2$ (**TCPDCA**) with $2-ClC_6H_4-NHCO-CHCl_2$ (**MCPDCA**) and $C_6H_5-NHCO-CHCl_2$ (**PDCA**) revealed similar trends. The average C-C ring distances in Å are 1.384 (**PDCA**), 1.379 (**MCPDCA**) and 1.378 (**TCPDCA**). The other side chain distances generally show a slightly decreasing trend as we go from **PDCA** to **TCPDCA**. The C(ring)-N-C(O), O-C(O)-N and O-C(O)-C (side chain) bond angles show similar trends as we go from $C_6H_5-NHCO-CHCl_2$ to $2,4,6-Cl_3C_6H_2-NHCO-CHCl_2$, while the N-C(O)-C(side chain) bond angle increases by about 2°.

The following observations may be made from the careful analysis of the available crystal structure data

[11, 12, 17, 18, 20, 28, 29], but to draw general conclusions extensive data are to be collected. Our work in this direction is in progress: N-phenyl acetamide (**PA**) on conversion to either **PTCA** or **TCPA** by side chain or ring substitution changes its crystal symmetry from orthorhombic to the monoclinic. But both the side chain and ring substitutions of **PA** converts it into **TCPTCA**, and its orthorhombic symmetry is changed to the triclinic symmetry. On conversion of **PTCA** to **TCPTCA** through mono and di ring substitutions (**MCPTCA** and **DCPTCA**), the crystal symmetry changes from monoclinic to triclinic via orthorhombic and monoclinic symmetries, while conversion of **TCPA** to **TCPTCA** via mono and di- side chain substitutions by Cl (**TCPMCA** and **TCPDCA**), changes its crystal symmetry from monoclinic to triclinic through orthorhombic symmetries twice.

The molecules in $2,4,6-Cl_3C_6H_2-NHCO-CHCl_2$ (**TCPDCA**) and $2,4,6-Cl_3C_6H_2-NHCO-CCl_3$ (**TCPTCA**) are linked in chains by -NH-O- hydrogen bonds (about 3 Å) between the amide groups of the molecules.

Acknowledgements

B. T. G. thanks the Alexander von Humboldt Foundation, Bonn, Germany for a research fellowship. Support of the Deutsche Forschungsgemeinschaft is also acknowledged.

- [1] The Chemistry of Amides, ed. J. Jabicky; Interscience, London 1970.
- [2] R. Sandler and W. Karo, Organic Functional Group Preparations; Academic Press, London 1972, Vol. 3.
- [3] K. Hargreaves, J. G. Pritchard, and H. R. Dave, Chem. Rev. **70**, 439 (1970).
- [4] E. Stewart and T. H. Siddal, Chem. Rev. **70**, 517 (1970).
- [5] S. Isaacs, Chem. Soc. Rev. **5**, 181 (1976).
- [6] G. K. Semin, T. A. Babushkina, and G. G. Yakobson, Nuclear Quadrupole Resonance in Chemistry; Wiley & Sons, New York 1975.
- [7] J. A. S. Smith (Ed.), Advances in Nuclear Quadrupole Resonances; Wiley & Sons, London 1974 - 83. Vols. 1 - 5.
- [8] Al. Weiss and S. Wigand, Z. Naturforsch. **45a**, 195 (1990).
- [9] Al. Weiss, Z. Naturforsch. **48a**, 477 (1993); Acta Cryst. **B51**, 523 (1995).
- [10] B. T. Gowda and Al. Weiss, Z. Naturforsch. **49a**, 695 (1994).
- [11] S. Dou, B. T. Gowda, H. Paulus, and Al. Weiss, Z. Naturforsch. **49a**, 1136 (1994).
- [12] B. T. Gowda, S. Dou, and Al. Weiss, Z. Naturforsch. **51a**, 627 (1996).
- [13] B. T. Gowda, D. K. Bhat, H. Fuess, and Al. Weiss, Z. Naturforsch. **54a**, 261 (1999).
- [14] B. T. Gowda, D. K. Bhat, H. Fuess, and Al. Weiss, Z. Naturforsch. **54a**, 679 (1999).
- [15] B. T. Gowda, B. H. A. Kumar, and H. Fuess, Z. Naturforsch. **55a**, 721 (2000).
- [16] D. K. Bhat and B. T. Gowda, J. Indian Chem. Soc. **77**, 279 (2000).
- [17] S. Dou, H. Fuess, Al. Weiss, B. T. Gowda, and V. G. Krishnan, Z. Kristallogr. **212**, 532 (1997).
- [18] B. T. Gowda, H. Paulus, and H. Fuess, a) Z. Naturforsch. **55a**, 711 (2000) and b) Unpublished results
- [19] W. Pies, H. Rager, and Al. Weiss, Org. Mag. Reson. **3**, 147 (1971).
- [20] S. C. Nyburg, J. K. Fawcett and J. T. Szymanski, Acta Cryst. **C43**, 2452 (1987).

- [21] F. A. Bertl and L. M. Ziti, Arch. Pharm. **285**, 372 (1952).
- [22] G. M. Sheldrick: SHELXS-86. Program for the Solution of Crystal Structures, University of Göttingen, Germany 1986
- [23] G. M. Sheldrick: SHELXL-93. Program for Crystal Structure Determination, University of Göttingen, Germany 1993
- [24] Nonius Diffractometer Control Software (1993), NONIUS GmbH, Solingen, Germany 1993.
- [25] L. Spek: PLUTON-93. Program for the Display and Analysis of Crystal and Molecular Structures, University of Utrecht, The Netherlands 1993.
- [26] C. T. North, D. C. Philips, and F. S. Mathews, Acta Cryst. **A24**, 351 (1968).
- [27] D. Flack, Acta Cryst. **A39**, 876 (1983).
- [28] D. Grocke, S. Dou, and Al. Weiss, Z. Naturforsch. **47a**, 160 (1990).
- [29] C. J. Brown and D. E. C. Corbridge, Acta Cryst. **7**, 711 (1954); C. J. Brown, Acta Cryst. **21**, 442 (1966).