

Supporting Information

Alexey Slobodinyuk, Vladimir Strelnikov, Dmitriy Kiselkov, and Daria Slobodinyuk*

Synthesis of oligotetramethylene oxides with terminal amino groups as curing agents for an epoxyurethane oligomer

<https://doi.org/10.1515/znb-2021-0085>. Received June 15, 2021; accepted July 12, 2021

***Corresponding author: Daria Slobodinyuk**, Institute of Technical Chemistry, Ural Branch of the Russian Academy of Sciences, Ac. Korolev Str., 3, 614130 Perm, Russia; e-mail:

selivanovadg@gmail.com; <https://orcid.org/0000-0002-6640-1566>

Alexey Slobodinyuk, Dmitriy Kiselkov: Institute of Technical Chemistry, Ural Branch of the Russian Academy of Sciences, Ac. Korolev Str., 3, 614130 Perm, Russia; e-mail:

slobodinyuk.aleksey.ktn@mail.ru (A. Slobodinyuk), dkiselkov@yandex.ru (D. Kiselkov),

<https://orcid.org/0000-0003-0411-2125> (A. Slobodinyuk), <https://orcid.org/0000-0002-0237-2743> (D. Kiselkov)

Vladimir Strelnikov: Corresponding Member, Russian Academy of Sciences, Director of Institute of Technical Chemistry, Ural Branch of the Russian Academy of Sciences, Ac. Korolev Str., 3, 614130 Perm, Russia; e-mail: svn@itcras.ru, <https://orcid.org/0000-0003-2538-535X>

Table of Contents

¹ H NMR spectrum of PTHF.....	S2
¹ H NMR spectrum of PTMO-dibromide	S3
¹³ C NMR spectrum of PTMO-dibromide.....	S4
¹ H NMR spectrum of PTMO-diamine.....	S5
¹³ C NMR spectrum of PTMO-diamine.....	S6




Figure S1: ^1H NMR spectrum of PTHF.




Figure S2: ^1H NMR spectrum of PTMO-dibromide.




Figure S3: ^{13}C NMR spectrum of PTMO-dibromide.




Figure S4: ^1H NMR spectrum of PTMO-diamine.




Figure S5: ^{13}C NMR spectrum of PTMO-diamine.