

Isolation of the New (-)-(3*R*,4*S*)-4-Hydroxymellein from the Fungus *Septoria nodorum* Berk

Michel Devys, Michel Barbier

Institut de Chimie des Substances Naturelles, CNRS,
Avenue de la Terrasse, 91198 Gif-sur-Yvette Cedex,
France

Jean-François Bousquet, Albert Kollmann

Station de Pathologie Végétale, CNRA,
78026 Versailles Cedex, France

Z. Naturforsch. **47c**, 779–781 (1992);
received April 6/July 7, 1992

Septoria nodorum (*Phaeosphaeria nodorum*),
Fungus, Phytotoxins, Isocoumarins,
(-)-(3*R*,4*S*)-4-Hydroxymellein

The new (-)-(3*R*,4*S*)-4-hydroxymellein (**6**) is isolated from the fungus *Septoria* (*Phaeosphaeria*) *nodorum* Berk, together with the previously reported isomer **5**. The results so far obtained in this series of metabolites are discussed in relationship with biosynthetic considerations.

Introduction

The fungus *Septoria* (*Phaeosphaeria*) *nodorum* Berk is a common parasite of wheat. It is responsible for a fungal disease which seriously affects crops (septoriosis). This fungus produces in cultures a series of highly potent phytotoxins which were previously investigated. Thus, we reported on the isolation of (-)-(3*R*)-mellein (ochracin) (**4**) [1], (3*R*)-O-methylmellein, (-)-(3*R*,4*R*)-4-hydroxymellein (**5**), (-)-(3*R*)-7-hydroxymellein, mycophenolic acid [2], septorine [3], N-methoxyseptorine [4], and N-methoxyseptorinol [5]. Undoubtedly, the observed phytotoxic activity of this fungus is due to the summation of these various metabolites, some of them being endowed of a particularly strong biological activity. Mellein was shown to produce an important reduction in CO₂ net assimilation and an increase in stomatal resistance in seedling leaves of wheat [6–10].

In the present publication, the isolation of the new isomer (-)-(3*R*,4*S*) (**6**) from the culture medium of this fungus is reported. The results so far observed are discussed in relationship with the fact

that all the isocoumarins up to now isolated from *Septoria nodorum* are derived from (-)-mellein (**4**).

Results and Discussion

The culture medium of the fungus *Septoria nodorum* Berk was extracted by ethyl acetate according to a previously [11] reported method. The concentrated extract was submitted to HPLC fractionation (UV detection), leading to the known (-)-(3*R*,4*R*)-4-hydroxymellein (**5**) [2], plus the new compound (-)-(3*S*,4*R*)-4-hydroxymellein (**6**).

The structures of these substances **5** and **6** were determined on the basis of physicochemical data, by comparing with previously reported results (m.p. and $[\alpha]_D$ as represented in Fig. 1). In this series, the coupling constants of the protons at C-3 and C-4 are determinant for the attribution of the stereochemistry. As the absolute configurations were established for (-)-mellein (**1**) and (+)-(3*S*,4*S*)-4-hydroxymellein (**2**) [12, 14], it was possible to fix the configurations of the products isolated from *Septoria nodorum*. The stereochemistry of the new product **6** was confirmed by irradiation at the signal of the methyl group at 1.55 ppm, resulting in two coalescent signals (2H) for the two vicinal protons at C-3 and C-4 (one peak). Hence, it is of course deduced that the coupling constant is not noticeable between these two protons. The dihedral angle determined from a molecular model gives a value of 70° which corresponds to a coupling constant of ca. 0.3 Hz [16]. By now, three of the four possible isomers of 4-hydroxymellein are known [2, 12, 14, 15]. To our knowledge, the (+)-isomer (3*S*,4*R*) **3** still remains to be found and could exist in strains producing the corresponding (+)-mellein [14]. A diacetate of **6** was prepared which exhibited MS and ¹H NMR spectra in agreement with the proposed stereochemistry. In particular, the vicinal ³J coupling constant observed in the ¹H NMR of this diacetate is of approximately 1 Hz. It appears from the molecular models that two conformers are possible for substance **6**, each requiring some energy for transformation. The dihedral angle 3-C–H 4-C–H is of ca. 70° when the methyl group is β-axial which leads to the observed coupling constant according to the Karplus equation [16]. But in the conformer having this methyl group β-equatorial, this angle is

Reprint requests to Dr. M. Barbier.

Verlag der Zeitschrift für Naturforschung,
D-W-7400 Tübingen
0939–5075/92/0900–0779 \$ 01.30/0




Fig. 1. Absolute configurations of the two melleins (ochracins) and of the four possible 4-hydroxymelleins.

of 180°, a value which must give a higher J (between 6 and 15 Hz). Hence, two conclusions may be drawn: 1) the compound **6** isolated from *Septoria nodorum* has the 3- β -axial conformation, 2) the possibility that two conformers exist for each represented stereochemistry of **2**, **3**, **5**, **6**, can not be excluded. The observed differences in the $J_{3,4}$ -values between conformers of the *cis*-series **2** and **5** must be small (or negligible), whereas such differences should be much bigger in the *trans*-series **3** and **6**.

The presence of the two isomers **5** and **6** in the culture medium of *Septoria nodorum* is in agreement with the demonstration that the biosynthesis of (+)-(3*S*,4*S*)-4-hydroxymellein (**2**) proceeds through oxidation of (+)-mellein (**1**) by molecular oxygen [14]. The previously found (-)-O-methylmellein and (-)-7-hydroxymellein, together present in *Septoria nodorum* with (-)-mellein [2] and the (-)-4-hydroxymelleins, demonstrate of course the same biogenetic family.

Experimental

Melting points have been determined with a Kofler apparatus under the microscope and are corrected. MS were carried out on an AEI MS 50 spectrometer, ^1H NMR on a Bruker 200 MHz apparatus (CDCl_3 , δ ppm). Rotatory power was de-

termined on an electronic Perkin-Elmer polarimeter 241. TLC's on Schleicher-Schüll SiO_2 fluorescent films, 1 mm thickness for preparative purposes, UV observation with a Desaga lamp at 254 nm, or FeCl_3 spray for visualization.

The 4-hydroxymelleins **5** and **6** were extracted from the culture medium of *Septoria nodorum* according to the reported [11] method. The EtOAc concentration was submitted to HPLC (elution by ethanol-water, UV detection) and the two substances **5** and **6** recovered according to the absorption curve. SiO_2 TLC (CH_3Cl -EtOAc 5:1), **5** R_f 0.60, colour reaction with FeCl_3 : blue, **6** R_f 0.65, FeCl_3 pink, relative proportions 5:1. The physicochemical data of substance **5** are identical to the reported values [2] except for small differences in the m.p. and $(\alpha)_D$ due to more accurate technical facilities (such as electronic polarimeter). These values are reported on the Figure. **6**: MS m/z (%), 194 M^+ (100), 150 (70), 121 (80), 122 (80); ^1H NMR: 1.55 (d, 3H, CH_3), 4.50 (enlarged signal, 2H, H-3 and H-4), giving a singlet by irradiation of the methyl signal at 1.55; aromatic protons: 7.04 (d, 1H, H-5, $J = 8$ Hz), 7.55 (dd, 1H, H-6, $J = 8$ Hz), 6.95 (d, 1H, H-7, $J = 8$ Hz), 11 (s, 1H, 8-OH). The diacetate of **6** was prepared by treatment with acetic anhydride in the presence of anhydrous pyridine (10 mg **6**, 6 drops of each reagent, 1 h at 37 °C, drying under *in vacuo* overnight at room tempera-

ture). SiO₂ TLC in pentane–ethyl acetate 1:1, *R_f* 0.70, MS *m/z* (%) 278 M⁺ (2), 236 M–42⁺ (100), 194 M–42–42⁺ (45), 176 M–42–60 (100), 150 (100), 149 (75), 43 CH₃CO⁺ (95). ¹H NMR (CDCl₃): 1.35 (d, 3H, CH₃), 2.13 and 2.40 (s, 3H each, CH₃COO), 4.80 (m, 1H, H-3), 5.83 (d, 1H, H-4, *J* = 1 Hz), 7.18 (d, 1H, H-5, *J* = 8 Hz), 7.36

(d, 1H, H-7, *J* = 8 Hz), 7.66 (dd, 1H, H-6, *J* = 8 Hz).

Acknowledgements

Thanks are due to Drs B. C. Das, C. Girard and J. P. Dupuis for the EI-MS determinations carried out at Gif-sur-Yvette.

- [1] M. Devys, J. F. Bousquet, M. Skajennikoff, and M. Barbier, *Phytopathol. Z.* **81**, 92 (1974).
- [2] M. Devys, J. F. Bousquet, A. Kollmann, and M. Barbier, *Phytochemistry* **19**, 2221 (1980).
- [3] M. Devys, J. F. Bousquet, A. Kollmann, and M. Barbier, *C.R. Acad. Sci. Paris* **286**, Ser. C, 457 (1978).
- [4] M. Devys, M. Barbier, A. Kollmann, and J. F. Bousquet, *Tetrahedron Lett.* **23**, 4409 (1982).
- [5] M. Devys, M. Barbier, A. Kollmann, and J. F. Bousquet, *Phytochemistry*, in press.
- [6] J. F. Bousquet, H. Belhomme de Franqueville, A. Kollmann, and R. Fritz, *Can. J. Bot.* **58**, 2575 (1980).
- [7] J. F. Bousquet, M. Skajennikoff, O. Bethenod, and P. Chartier, *Ann. Phytopathol.* **93**, 503 (1977).
- [8] D. Laffray, J. F. Bousquet, O. Bethenod, and P. Louguet, *Agronomie* **2**, 25 (1982).
- [9] O. Bethenod, J. F. Bousquet, D. Laffray, and P. Louguet, *Agronomie* **2**, 99 (1982).
- [10] T. Touraud and J. F. Bousquet, *Can. J. Bot.* **57**, 561 (1979).
- [11] J. F. Bousquet and M. Skajennikoff, *Phytopathol. Z.* **80**, 355 (1974).
- [12] L. Camarda, L. Merlini, and G. Nasini, *Phytochemistry* **15**, 537 (1976).
- [13] M. Sasaki, Y. Kaneko, K. Oshita, H. Takamatsu, Y. Asao, and T. Yokotsuka, *Agr. Biol. Chem.* **34**, 1296 (1970).
- [14] C. Abell, A. C. Sutkowski, and J. Staunton, *J. Chem. Soc. Chem. Commun.* **1987**, 586.
- [15] D. C. Aldridge, S. Galt, D. Giles, and W. B. Turner, *J. Chem. Soc. (C)* **1971**, 1623.
- [16] L. M. Jackman, and S. Sterhell, *Applications of NMR Spectroscopy in Organic Chemistry*, p. 282, Pergamon Press, Oxford 1972.