

4-Pentadecylpyridine: A Competitive Polyphenoloxidase Inhibitor

Claus T. Fuchs and Gerhard Spiteller

Institut für Organische Chemie I, Universität Bayreuth, Universitätsstr. 30,
95440 Bayreuth, Germany

Z. Naturforsch. **52c**, 761–765 (1997); received July 30/August 19, 1997

Polyphenoloxidase, *o*-Diphenols, 4-Pentadecylpyridine, Competitive Inhibitor,
Taphrina pruni

4-Pentadecylpyridine was isolated from plums infected by *Taphrina pruni*. This fungal metabolite has been tested for its effect on polyphenoloxidase, using the polyphenoloxidase-catalysed addition of L-proline to 4-methylcatechol in a chromogenic assay. The resulting data prove that 4-pentadecylpyridine is a competitive inhibitor to polyphenoloxidase with an inhibitor constant of $3.70 \pm 0.03 \cdot 10^{-4}$ M. Further comparative measurements have demonstrated that the effect is due to the pyridine moiety. Probably reduced polyphenoloxidase-activity decreases the toxicity of *o*-diphenols to the fungus and thus increases susceptibility of plums to the fungus *Taphrina pruni*.

Introduction


Our previous investigations of the constituents of healthy plums and those infected by the fungus *Taphrina pruni* revealed that in affected fruits the content of *o*-diphenols (Fuchs, 1996), especially (+)-catechine, catechine conjugates, proanthocyanidines and caffeoyl-D-quinic acids is increased compared to unaffected fruits for a factor of about 20. Since all of these compounds are phenols they may serve as polyphenoloxidase (PPO)-substrates (Lerch, 1987; Mayer and Harel, 1979; Mayer, 1987).

Polyphenoloxidase (PPO) (tyrosinase, monophenol monooxygenase, catecholoxidase EC 1.14.18.1) is a copper containing enzyme catalysing the *o*-hydroxylation of monophenols to *o*-diphenolic compounds and their further oxidation to the toxic *o*-quinones (Lerch (1987); Scheme 2).

Furthermore, in the course of our investigations, 4-pentadecylpyridine (**2**) has been isolated from the fungus *Taphrina pruni* (Fuchs and Spiteller, 1995). As a weak base bearing long alkyl chains 4-pentadecylpyridine (**2**) is a lysosomotropic de-

tergent (Firestone and Pisano, 1979): It accumulates within lysosomes, ruptures their membranes and releases the contents of the cells into the cytoplasm.

Due to the structural analogy (Scheme 1) of this new natural compound to fusaric acid (**1**), a well-known fungal metabolite from *Fusarium* species, we suspected that **2** may develop similar biological activity as **1**. Since several communications (Becker and Pushkareva, 1972; Kern, 1972) demonstrate that **1** inhibits polyphenoloxidase and thus quinone formation, we investigated the effects of 4-pentadecylpyridine (**2**) on ppo-inhibition, assuming further consequences for the toxicity of the above identified phenols and plum susceptibility (Mason and Wassermann, 1987; Goldstein and Swain, 1963; Hayes, 1989).



Scheme 1. Fusaric acid (**1**) (5-n-butyl-2-picolinic acid) and 4-pentadecylpyridine (**2**).

Abbreviations: PPO: Polyphenoloxidase; K_I : inhibitor constant; PMMA: polymethylmethacrylate; LB: Lineweaver-Burk; K_M : Michaelis constant; v_{max} : maximal velocity.

Reprint requests to Prof. G. Spiteller.
Fax: 09 21/55 26 71.

Results

In order to study the effect of 4-pentadecylpyridine (**2**) on ppo-activity a chromogenic assay was

modified in analogy to Rzepecki *et al.* (Rzepecki and Waite, 1989). This assay is based on the ppo-catalysed addition of L-proline to 4-methylcatechol (simplest monosubstituted *o*-diphenol) (analogous Scheme 2) forming 4-*N*-prolyl-adducts.

The reaction has been used to determine N-terminal proline residues in peptide polymers (Mason and Peterson, 1955) and to detect catecholoxidase activity in nondenaturing thin layer or gel permeation chromatography of polyphenoloxidase enzymes (Jolley and Mason, 1964; Thomas *et al.*, 1978).

Figure 1 shows the Lineweaver-Burk (LB)-plots for the generation of 4-*N*-prolyl adducts after addition of different concentrations of 4-pentadecylpyridine (**2**). Changes of the slope and intercept of the $1/V$ versus $1/[S]$ plot indicate competitive respectively non-competitive inhibition. In addition the curve allows conclusions on the binding strength of an inhibitor (Engel, 1977).




Fig. 1. Lineweaver-Burk-Plot (LB) in presence of different concentrations of 4-pentadecylpyridine (**2**).

The LB-plots intersect the ordinate in one point indicating a competitive enzyme inhibition.

According to the formula

$$(1 + [I]/K_I) K_M/v_{\max} = \text{gradient of the LB-plots}$$

the inhibitor constant was calculated (Henderson, 1992; Lehninger, 1977) as mean value of four measurements with different 4-pentadecylpyridine concentrations:

$$K_I = 0.370 \cdot 10^{-4} \pm 0.003 \cdot 10^{-4} \text{ M (Table I).}$$

Table I. PPO-inhibition with 4-pentadecylpyridine **2**.

Final inhibitor concentration $[\mu\text{M}]$	Gradient of the LB-plot	$1/K_I$	K_I
0	1.125		
80	1.364	2655	0.000376
160	1.583	2544	0.000393
280	2.079	3029	0.000330
600	1.626	2710	0.000369
1400	3.975	1810	0.000552

We assume that associative effects at higher 4-pentadecylpyridine (**2**) concentrations cause a significant increase of the inhibitor constant (Engel, 1977). For this reason the highest concentration (inhibitor concentration: 1400 μM ; K_I : 0.000552 M) was neglected for mean value calculation.

As expected, further comparative measurements with equal concentrations of pyridine (80 μM and 280 μM) proved that the inhibition is exclusively due to the pyridine residue.

Experimental

PPO assay

The ppo-assay was modified in analogy to Rzepecki *et al.* (Rzepecki and Waite, 1989).

Briefly, a PMMA-cuvette was charged with a mixture of 1000 μl phosphate buffer (pH 7.5; 0.1 M), 200 μl L-proline (0.2 M in p-buffer pH 7.5) and 200 μl 4 M NaCl. Then 100 μl of a 1 μM 4-methylcatechol solution was added. Autoxidation of 4-methylcatechol was inhibited by dissolving the substrate in 0.2 M acetic acid. Finally, 700 μl water (reference) respectively X μl 4-pentadecylpyridine (**2**) ($X = 50, 100, 175, 300$ and 700 ; 10^{-2} M aqueous solution) and $(700-X)$ μl water were added. The enzymatic reaction was started by the addition of 200 μl ppo-solution (0.614 U; in p-buffer pH 7.5). Kinetics of the 4-*N*-prolyl-adduct formation was determined at 312 nm in a PMMA-cuvette.

The validity of the test system was established with cyanide, a well-known ppo-inhibitor (Wittenberg and Triplett, 1985).

UV spectroscopy

UV measurements were performed with a Shimadzu UV-160 A photometer. The absorbance at 312 nm was measured in intervals of one minute. The molar extinction coefficient (ϵ) of the 4-*N*-prolyl adduct of 4-methylcatechol is $9630 \text{ M}^{-1}\text{cm}^{-1}$ (Rzepecki and Waite, 1989).

Chemicals

L-Proline, 4-methylcatechol and tyrosinase (EC 1.14.18.1) were obtained from Fluka Chemie AG, Neu-Ulm.

Synthesis

4-Pentadecylpyridine was synthesised as previously described (Fuchs and Spiteller, 1995).


Discussion

PPO-activity increases after infection or mechanical wounding (Matta and Abbattista, 1970; Pitt, 1975; Cheung and Henderson, 1972). A similar increase of activity was observed in galls formed in a number of plants after pathogen infection (Joshi and Tandon, 1984; Tandon and Arya, 1982; Ramawat and Purohit, 1980). Furthermore, an influence on plant hormone regulation, especially indolyl acetic acid, has been implied for PPO by various experiments (Gordon and Paleg, 1961; Tomaszewski and Thimann, 1966; Sondheimer and Griffin, 1960).

Release of free fatty acids generated by membrane wounding (Galliard, 1970) induces the activation of latent PPO (Hutcheson *et al.*, 1980; Golbeck and Cammarata, 1981). Thus enhanced PPO-activity is directly correlated to host resistance respectively pathogen infectibility (Bashan *et al.*, 1987; Leon, 1971).

The assumption that PPO-oxidation plays an important role in plant defence is corroborated by the finding of increased susceptibility of plants to fungi after quinone reduction (Moustafa and Whittenbury, 1970). Furthermore, host resistance in *Ribes* to the pathogen *Spaeotheca* was found to depend on the levels and interaction of polyphenoloxidase, chlorogenic acid and a reducing agent such as ascorbic acid (Trajkowski, 1976). Mukherjee and Gosh demonstrated that ppo-inhibition is directly connected to a decrease in resistance (Mukherjee and Gosh, 1975). This resistance is directly correlated to quinone toxicity and the ability

of these quinones to add to substances with active hydrogen atoms, e.g. lysine-, cysteine- or tyrosine residues (Pierpoint *et al.*, 1977; Pierpoint, 1966; Auf' Mkolok, 1985) (Scheme 2).



Scheme 2. Toxicity of *o*-diphenols: Addition of *o*-diquinones to S-, N- and O-containing nucleophils. PPO: Polyphenoloxidase (EC 1.14.18.1).

Considering that fungal enzymes are S- and N-containing compounds, they are inactivated by the corresponding quinones (Mason and Wassermann, 1987; Goldstein and Swain, 1963; Hayes, 1989). Thus the accumulation of a high concentration of *o*-diphenols in infected plums is a defence response directed to the enzymes of the pathogen. Nevertheless, the effectiveness of this defence response is highly dependent on quinone formation respectively PPO-activity.

While competitive inhibitors influence the binding position of PPO, reagents binding to the copper atom of the enzyme are non-competitive inhibitors (e.g. cyanide and thiourea) (Wittenberg and Triplett, 1985). Our data indicate that 4-pentadecylpyridine (**2**) is a competitive inhibitor, whereas fusaric acid (**1**) (Walter, 1969) shows interaction to the copper atom of the PPO.

In conclusion, 4-pentadecylpyridine (**2**) is a strong competitive inhibitor of PPO. Its binding strength is beyond the magnitude of many natural PPO-substrates, e.g. ferulic acid (inhibitor constant: $K_I = 1.3 \text{ mM}$) (Wittenberg and Triplett, 1985; Anosike and Ayaebene, 1982; Walter, 1969; Loeffler and Zenk, 1990).

At least at the site of penetration (and high 4-pentadecylpyridine (**2**) concentration) the fungal

pyridine derivative may significantly reduce the formation of toxic *o*-quinones. Since these *o*-quinones are directly correlated with defence reactions of affected plants (Mayer, 1987; Bashan *et al.*, 1987; Leon, 1971) they may significantly increase plum susceptibility to the fungus respectively decrease plum resistance to fungal infection.

Acknowledgements

We thank Deutsche Forschungsgemeinschaft for financial support and Mr. W. Kern for purification of the solvents. Furthermore, we are obliged to Mr. R. A. Firestone for information on lysosomotropic activity of 4-pentadecylpyridine.

- Anosike, E. O. and Ayaebene, A. O. (1982), Properties of polyphenol oxidase from tubers of the yam *Dioscorea Bulbifera*. *Phytochemistry* **21**, 1889–1893.
- Auf'Molk, M., Amir, S. M., Kubota, K. and Ingbar, S. H. (1985), The active principles of plant extracts with antithyrotropic activity: Oxidation products of derivatives of 3,4-dihydroxycinnamic acid. *Endocrinology* **116**, 1677–1686.
- Bashan, J., Okon, Y. and Henis, Y. (1987), Peroxidase, polyphenoloxidase, and phenols in relation to resistance against *Pseudomonas syringae* pv. *tomato* in tomato plants. *Can. J. Bot.* **65**, 366–372.
- Becker, S. E. and Pushkareva, J. D. (1972), Nature and biosynthesis of *Fusarium* wilt toxin, its mechanism of action, and its transformation in the cotton plant. *Bodenkultur* **23**(3), 256–271.
- Cheung, K. W. K. and Henderson, H. M. (1972), Effect of physiological stress on potato polyphenol oxidase. *Phytochemistry* **11**, 1255–1260.
- Engel, P. C. (1977), *Enzyme Kinetics*. London: Chapman and Hall.
- Firestone, R. A. and Pisano, J. M. (1979), Lysosomotropic agents. 1. Synthesis and cytotoxic action of lysosomotropic detergents. *J. Med. Chem.* **22**, 1130–1133.
- Fuchs, C. and Spiteller, G. (1995), 4-Pentadecylpyridine: A metabolite from *Taphrina pruni*. *Z. Naturforsch.* **45c**, 891–893.
- Fuchs, C. (1996), Ph. D. Thesis, university of Bayreuth.
- Galliard, T. (1970), The enzymic breakdown of lipids. *Phytochemistry* **9**, 1725–1741.
- Golbeck, J. H. and Cammarata, K. V. (1981), Spinach thylakoid polyphenol oxidase. *Plant Physiol.* **67**, 977–984.
- Goldstein, J. L. and Swain, T. (1963), Changes in tannins in ripening fruits. *Phytochemistry* **2**, 371–383.
- Gordon, S. A. and Paleg, L. G. (1961), Formation of auxin from tryptophan through action of polyphenols. *Plant Physiol.* **36**, 838–845.
- Hayes, M. H. B. (ed.). (1989), *Humic substances II, in search of structure*. John Wiley & Sons.
- Henderson, P. J. F. (1992), Statistical analysis of enzyme kinetic data. In: *Enzyme Assays* (R. Eisenthal and M. J. Danson, ed.). Oxford: Oxford University Press.
- Hutcheson, S. W., Buchanan, B. B. and Montalbini, P. (1980), Polyphenol oxidation by *Vicia faba* chloroplast membranes. *Plant Physiol.* **66**, 1150–1154.
- Jolley, R. L. and Mason, H. S. (1964), The multiple forms of mushroom tyrosinase. *J. Biol. Chem.* **240**, 1489–1491.
- Joshi, S. C. and Tandon, P. (1984), Studies on certain oxidases, their isoenzymes and phenols as related to gall formation in *Leea indica*. *Biochem. Physiol. Pflanz.* **179**, 711–715.
- Kern, H. (1972), In: *Phytotoxins in Plant Disease* (R. K. S. Wood, A. Ballio and A. Graniti, ed.). London: Academic Press.
- Lehninger, A. L. (1977), *Biochemie*. Weinheim: Verlag Chemie.
- Leon, T. J. (1971), Phenoloxidase activity and fruiting body formation in *Schizophyllum commune*. *J. Bacteriol.* **106**, 162–167.
- Lerch, K. (1987), Monophenol monooxygenase from *Neurospora crassa*. *Methods Enzymol.* **142**, 165–169.
- Loeffler, S. and Zenk, M. H. (1990), The hydroxylation step in the biosynthetic pathway leading from norcoclaurine to reticuline. *Phytochemistry* **29**, 3499–3503.
- Mason, T. L. and Wassermann, B. P. (1987), Inactivation of red beet β -glucan synthase by native and oxidized phenolic compounds. *Phytochemistry* **26**, 2197–2202.
- Matta, A. and Abbattista, G. I. (1970), Differential inhibition and activation of polyphenoloxidase activity in healthy and *Fusarium*-infected tomato plants. *Phytopathol. Mediterr.* **9**, 168–175.
- Mayer, A. M. and Harel, E. (1979), Polyphenol oxidases in plants. *Phytochemistry* **18**, 193–215.
- Mayer, A. M. (1987), Polyphenol oxidases in plants – recent progress. *Phytochemistry* **26**, 11–20.
- Moustafa, F. A. and Whittenbury, R. (1970), Properties which appear to allow phytopathogenic pseudomonads to counteract plant defence. *Phytopathol. Z.* **67**, 214–217.
- Mukherjee, P. K. and Gosh, J. J. (1975), Quantitative changes in pectic substances in rice leaves infected with *Helminthosporium oryzae*. *Sci. Cult.* **41**, 433–439.

- Pierpoint, S., Ireland, R. J. and Carpenter, J. M. (1977), Modification of proteins during the oxidation of leaf phenols: reaction of potato virus X with chlorogenoquinone. *Phytochemistry* **16**, 29–34.
- Pierpoint, W. S. (1966), The enzymic oxidation of chlorogenic acid and some reactions of the quinone produced. *Biochem. J.* **98**, 567–580.
- Pitt, D. (1975), Changes in the subcellular location of catalase and *o*-diphenol oxidase during infection of potato tubers by *Phytophthora erythroseptica*. *Trans. Br. Mycol. Soc.* **65**, 91–100.
- Ramawat, K. G., Purohit, S. D. and Arya, H. C. (1980), Phenolics and oxidative enzymes in normal and gall tissues of *Gisekia*. *Sci. Cult.* **46**, 111–112.
- Rzepecki, L. and Waite, J. H. (1989), A chromogenic assay for catecholoxidases based on the addition of L-proline to quinones. *Anal. Biochem.* **179**, 375–381.
- Tandon, P. and Arya, H. C. (1982), Association of auxin protectors, peroxidase, indoleacetic acid oxidase and polyphenol oxidase in *Zizyphus* gall and normal stem tissues grown in culture. *Biochem. Physiol. Pflanz.* **177**, 114–124.
- Thomas, P., Delincee, H., and Diehl, J. F. (1978), Thin-layer isoelectric focusing of polyphenoloxidase on sephadex and its detection by the print technique. *Anal. Biochem.* **88**, 138–148.
- Tomaszewski, M. and Thimann, K. V. (1966), Interactions of phenolic acids, metallic ions and chelating agents on auxin-induced growth. *Plant Physiol.* **41**, 1443–1446.
- Trajkovski, V. (1976), The mechanism of resistance of *Ribes nigrum* to *sphaerotheca mors-uvae* (Schw.) Berk. *Swed. J. Agric. Res.* **6**, 215–223.
- Walker, J. R. L. (1969), Inhibition of the apple phenolase system through infection by *Penicillium Expansum*. *Phytochemistry* **8**, 561–566.
- Wittenberg, C. and Triplett, E. L. (1985), A detergent-activated tyrosinase from *Xenopus laevis*. *J. Biol. Chem.* **260**, 12535–12541.