

The Comparative Molecular Study between Bombycidae and Saturniidae Based on mtDNA RFLP and Cytochrome Oxidase I Gene Sequences: Implication for Molecular Evolution

Jae-Sam Hwang^a, Jin-Sung Lee^b, Tae-Won Goo^a, Hyun-Ah Kang^a,
Hae-Ryong Sohn^c, Ho-Rak Kim^a and O-Yu Kwon^{d,*}

^a National Sericulture and Entomology Research, R. D. A., Suwon 441–100, Korea

^b Genome Center, Korea Research Institute of Bioscience and Biotechnology, KIST,
P. O. Box 115 Yusong, Taejeon 305–600, Korea

^c College of Agriculture, Kyungpook National University, Taegu 702–701, Korea

^d Department of Anatomy, College of Medicine, Chungnam National University,
Taejeon 301–131, Korea. Fax: 82–42–580–8206, E-mail: oykwon@hanbat.chungnam.ac.kr

* Author for correspondence and reprint requests

Z. Naturforsch. **54c**, 587–594 (1999); received December 30, 1998/April 8, 1999

Bombycidae, Saturniidae, mtDNA RFLP, Cytochrome Oxidase I Gene

The phylogenetic relationships between *Bombyx mori* and *Bombyx mandarina* species of Bombycidae, and *Antheraea yamamai* and *Antheraea pernyi* species of Saturniidae were investigated based on mtDNA RFLP and cytochrome oxidase I gene. The sizes of the mtDNA of all the species were estimated at approximately 16 kbp $\pm$ 500 bp by total length of all the restricted fragments and no variation in size was recognized. Of the fourteen different restriction endonucleases used, *Bam*HI, *Hind*III, *Pst*I, *Eco*RI and *Xba*I showed RFLP. Among these, only *Hind*III showed RFLP between *B. mori* and *B. mandarina*. A comparative analysis of sequences was also conducted with the mitochondrial cytochrome oxidase I genes of each species. The results indicated that *B. mori* shared a 97%, 85% and 87% sequence identity with *B. mandarina*, *A. yamamai* and *A. pernyi*, respectively. *B. mandarina* shared a 87% and 88% sequence identity with *A. yamamai* and *A. pernyi*, respectively. *A. yamamai* shared 92% sequence identity with *A. pernyi*. The results of the phylogenetic analysis exhibited monophyly and confidence limits of more than 99% in all trees for both Bombycidae and Saturniidae.

Introduction

The analysis of mitochondrial DNA (mtDNA) sequence is a powerful tool for understanding insect population genetics and phylogenetics, since the small size of the mtDNA, the relatively rapid rate of its evolutionary change, and maternal inheritance all make it suitable for examining population background and evolution among closely related taxa (Moritz *et al.*, 1987; Gray, 1989; Lansman *et al.*, 1981). Recently, RFLP (Restriction Fragment Length Polymorphism), RAPD (Random Amplified Polymorphic DNA), mtDNA sequence and the sequence variation of target genes have been used in the molecular phylogenetics of insects.

Among insects, complete mtDNA sequences only exist for four insects: *Drosophila yakuba* (Clary and Wolstenholme, 1985), *Apis mellifera* (Crozier and Crozier, 1993), *Anopheles quadrimaculatus* (Mitchell *et al.*, 1993), *Anopheles gambiae*

(Beard *et al.*, 1993). It is generally recognized known that insect mtDNA consists of two ribosomal RNA genes, twenty-two transfer RNA genes and thirteen protein coding genes of the mitochondrial inner membrane respiratory complexes (Clary and Wolstenholme, 1985; Crozier and Crozier, 1993; Beard *et al.*, 1993). Further studies on mtDNA variation have also enhanced our understanding of population and evolutionary biology at both the intraspecific and interspecific levels. The PCR revolution and discovery of universal primers for mtDNA that work in a number of different species (Simon *et al.*, 1994; Lunt *et al.*, 1996; Kambhampati and Smith, 1995; Roehrdanz, 1995) have greatly increased the speed at which data can be obtained.

Silkmoths are holometabolous insects belonging to either the family Bombycidae or Saturniidae, both of which belong to the superfamily Bombycoidea. Bombycidae, *Bombyx mori* and *Bombyx mandarina*, live on mulberry leaves and make

higher quality silk fiber than Saturniidae, *Antheraea yamamai* and *Antheraea pernyi*, which live on leaves of oak tree. *B. mori* and *B. mandarina* can be crossed, and the resulting hybrid progeny shows normal fertility (Kawaguchi, 1928; Aratake *et al.*, 1973). The chromosome numbers are quite different between *A. yamamai* (31) and *A. pernyi* (49), but they can be closed to produce F1 progeny (Kawaguchi, 1934; Shimada *et al.*, 1988) which, however, are sterile due to incomplete oogenesis and other abnormalities (Shimada and Kobayashi, 1992; Kirimura, 1962). It is prerequisite to clarifying the phylogenetic relationships between Bombycidae or Saturniidae, with respect to understand physiological differences under molecular level and to create genetically transformed silkworms which may produce high quality amount of silk fiber using molecular breeding techniques. With in this mind, here, we have investigated the construction of mtDNA RFLP and sequenced the mitochondrial cytochrome oxidase I gene in *B. mori*, *B. mandarina*, *A. yamamai* and *A. pernyi* to understand the phylogenetic relationship between these species of Bombycidae and Saturniidae.

Materials and Methods

The preparation of mtDNA from silkgland was purified as described by Tamura and Aotsuka (1988). One gram of silkgland was homogenized in a 10 ml chilled homogenizing buffer [0.25 M sucrose, 10 mM EDTA, 30 mM Tris-HCl, (pH 7.5)], and centrifuged at 1,000×g for 1 min at 4°C. The supernatant was recentrifuged at 12,000×g for 10 min at 4°C. The resulting pellet of mitochondria was suspended in a 500 µl of 10 mM Tris-EDTA buffer (pH 8.0) containing 0.15 M NaCl, 0.18 N NaOH and 1% SDS. After phenol-chloroform (1:1) and ethanol extraction, the mtDNA was collected at 12,000×g for 15 min at 4°C. The resulting mtDNA was rinsed twice with 70% cold ethanol and digested with 20 µg/ml of DNase-free RNase in a 20 µl of 10 mM Tris-EDTA buffer (pH 8.0) at 37°C for 30 min. The mtDNAs were digested with 14 kind of restriction enzymes at 37°C for 2 h; *Bam*HI, *Hind*III, *Xho*I, *Sal*I, *Sma*I, *Pst*I, *Eco*RI, *Xba*I, *Eco*RV, *Dra*I, *Kpn*I, *Sca*I, *Bgl*II. Digested DNA fragments were separated on the 1% agarose gels in TBE buffer, the gels

were stained with ethidium bromide solution and were photographed using Polaroid type 667 films on a UV transilluminator.

A PCR was set up in a 50 µl volume as described by Kambhampati *et al.* (1992) in a GeneAmp PCR system 9700 (Perkin-Elmer cetus Co). Ten ng of each mtDNA, *B. mori*, *B. mandarina*, *A. yamamai* and *A. pernyi*, was used as a PCR template DNA. The primers for cytochrome oxidase I gene (Cy1 – 1,632; 5'-TGATCAAATTTATAAT-3' and Cy2 – 2,191; 5'-GGTAAAATTAAATATAAACTTC-3') were derived from the already known mtDNA sequences of *D. yakuba* (Clary and Wolstenholme, 1985), *A. mellifera* (Crozier and Crozier, 1993), and *A. quadrimacutatus* (Mitchell *et al.*, 1993) and used 5 pmol in this experiments. The temperature profile for the amplification was 95°C for 30 sec, 50°C for 1 min, then 72°C for 1 min for 35 cycles, and a final extension step of 72°C for 5 min was added. The PCR products were separated on a 1% agarose gel, and the bands of the expected size were recovered using a Gene clean II Kit (Bio 101, USA). The eluted DNA fragments were directly ligated with a pGemT Easy vector (Promega, USA) and transformed into *E. coli* JM 109. The recombinant plasmids were isolated according to procedures prescribed by the Wizard SV plasmid purification Kit (Promega, USA).

Fluorescent cycle sequencing was performed with 300–500 ng of this double stranded DNA in a 20 µl sequencing reaction mixture and prepared using an ABI PRISM Dye terminator Cycle Sequencing Ready Reaction Kit. Three clones, both orientations, were examined for each sequence.

The genetic similarity coefficient was calculated from $F = 2n_{xy}/(n_x + n_y)$ in which n_x and n_y are the number of fragments in species X and Y, respectively, where n_{xy} is the number of fragments shared by the two species (Nei and Li, 1979). Computer analyses of mtDNA sequences were carried out using the CLUSTAL W program (Thompson *et al.*, 1994). Phylogenetic trees for the data set were inferred by using the neighbor-joining program of MEGA (Saitou and Nei, 1987; Kumar *et al.*, 1993). The stability of the relationships was assessed by performing bootstrap analyses of the neighbor-joining data based on 1,000 resampling.

Results and Discussion

Silkworms, in addition to being important targets for improvement by genetic engineering, provide chances to research unique biological aspects, including developmental pattern formation and tissue specific expression. To investigate these phenomena and to genetically engineer silkworms showing better characteristics, comparison of silkworms both dominant, Bombycidae (*Bombix mori* and *Bombix mandarina*) and wild type, Saturniidae (*Antheraea yamamai* and *Antheraea pernyi*) at a molecular level is prerequisites, since little information is still available on the base of molecular level.

As described above, it is important to establish the phylogentic relationships between Bombycidae or Saturniidae on the molecular level to understand a silkworm physiology and to create a transgenic silkworm may produce a lot of useable biomatters including silk and drugs. With in this mind, here, we have established the molecular relationship of *B. mori*, *B. mandarina* and *A. yamamai*, *A. pernyi* based on those mtDNA RFLP data and mitochondrial cytochrome oxidase I gene sequences partially amplified by PCR.

Phylogenetic relationships among the *B. mori* and *B. mandarina* species of Bombycidae and *A. yamamai* and the *A. pernyi* species of Saturniidae were investigated based on mtDNA RFLP and the




Fig. 1. Determination of mitochondrial genome sizes of silkworm.

(A) Uncut form of mtDNAs purified from silkworms, separated on a 1% agarose gel. M: molecular marker of λ -Hind III, Lane 1: *Bombyx mori*, Lane 2: *Bombyx mandarina*, Lane 3: *Antheraea yamamai*, Lane 4: *Antheraea pernyi*. Arrows a: mtDNA nicked circular form, b: super coiled form, c: linear form.

(B) Electrophoretic patterns of *B. mori* mtDNA digested with five restriction endonucleases, separated on a 1% agarose gel. M: molecular marker of λ -Hind III. Lane 1: *Hind* III, 2: *Eco* R I, 3: *Bam* H I, 4: *Xba* I, 5: *Pst* I.

cytochrome oxidase I gene. The mtDNA sizes of all the species tested were estimated at approximately 16 kb $\pm$ 500 bp in length, in which the non-variation of sizes and restriction enzyme sites were observed as shown in Figs 1A and 1B. The resulting genomic sizes are similar to those previously

Table I. Numbers of restriction sites on the mtDNAs of the major silkworms.

	BamHI*	HindIII*	XhoI	SalI	SmaI	PstI*	EcoRI*	XbaI*	EcoRV	DraI	KpnI	SacI	BglI	BglII
<i>B. mori</i> (Jam305)	2	4	-	1	1	2	4	3	1	7	-	-	-	2
<i>B. mori</i> (Jam306)	2	4	-	1	1	2	4	3	1	7	-	-	-	2
<i>B. mandarina</i>	2	2	-	1	1	2	4	3	1	7	-	-	-	2
<i>A. yamamai</i>	1	4	-	1	1	2	3	1	1	7	-	-	-	2
<i>A. pernyi</i>	1	4	-	1	1	3	3	3	1	7	-	-	-	2

-: No restriction sites. *: Restriction endonuclease showing RFLPs among the five silkworm. Jam305: Japonic type. Jam306: Chinese type.

Table II. Genetic similarity values among silkworms, based on the RFLP markers of mtDNA.

	<i>B. mori</i> (Jam305)	<i>B. mori</i> (Jam306)	<i>B. mandarina</i>	<i>A. yamamai</i>	<i>A. pernyi</i>
<i>B. mori</i> (Jam305)	1.000				
<i>B. mori</i> (Jam306)	1.000	1.000			
<i>B. mandarina</i>	0.848	0.848	1.000		
<i>A. yamamai</i>	0.303	0.302	0.394	1.000	
<i>A. pernyi</i>	0.227	0.228	0.348	0.500	1.000

Jam305: Japonic type; Jam306: Chinese type.

reported for *A. gambiae* (15,363 bp), *A. quadrimaculatus* (15,455 bp), and *D. yakuba* (16,019 bp), however, they are slightly smaller than that for *D. melanogaster* (19,517 bp). Of the fourteen different restriction endonucleases used, *Bam*HI, *Hind*III, *Pst*I, *Eco*RI and *Xba*I showed RFLP (Ta-

ble I and Fig. 1B). A variation in the enzyme-digested fragments between *B. mori* strains, Japanese (Jam305) and Chinese (Jam306), was not detected (Table I). A matrix for genetic similarity was constructed based on sixty eight mtDNA RFLP markers obtained from the two families di-

Fig. 2. Alignment of PCR amplified nucleotide sequences of a part (ca. 600 bp) of the cytochrome oxidase I gene. The origins of the amplified insects are indicated to the left. Asterisks (*) indicate nucleotides common to all species. Bmocy: *Bombyx mori*, Bmacy: *Bombyx mandarina*, Aycy: *Antheraea yamamai*, Apcy: *Antheraea pernyi*.

Bmocy	TGATCAAATTTATAATACTATTGTAACAGCACATGCTTTTATTATAATTTTTTTTATAGT 60
Bmacy	TGATCAAATTTATAATACTATTGTAACAGCACATGCTTTTATTATAATTTTTTTTATAGT 60
Aycy	TGATCAAATTTATAATACTATTGTAACAGCTCATGCTTTTATTATAATTTTTTTTATAGT 60
Apcy	TGATCAAATTTATAATACTATTGTAACAGCTCATGCTTTTATTATAATTTTTTTCATAGT 60
	***** 60
Bmocy	TATACCTATTATAATTGGAGGATTGGAAATTGATTAGTTCCTCTTATACTAGGAGCACC 120
Bmacy	TATACCTATTATAATTGGAGGATTGGAAATTGATTAGTTCCTCTTATACTAGGAGCACC 120
Aycy	TATACCTATTATAATTGGAGGATTGGAAATTGACTAATTCGTTAATATTAGGAGCTCC 120
Apcy	TATACCTATCATAATTGGAGGATTGGAAATTGATTAATTCATTAATATTAGGAGCCCC 120
	***** ** 120
Bmocy	AGATATAGCATTCCCACGAATAAATAATATAAGATTTTGACTCCTACCCCCCTCCCTTAT 180
Bmacy	AGATATAGCATTCCCACGAATAAGTAATATAAGATTTTGACTCCTACCCCCCTCCCTTAT 180
Aycy	TGATATAGCTTTCCCACGAATAAATAATATAANTTTTGGATTGCTCCCCCTCTTTAAAC 180
Apcy	TGATATAGCTTTCCCACGAATAAATAATATAAGTTTTGGACTATTACCCCCCTCTTTAAAC 180
	***** ** 180
Bmocy	ATTATTAATTTCAAGAAGAATTGTAGAAAATGGTGCAGGAACAGGATGAACAGTTTACCC 240
Bmacy	ATTATTAATTTCAAGAAGAATTGTAGAAAATGGTGCAGGAACAGGATGAACAGTTTACCC 240
Aycy	TTTATTAATTTCTAGAAGAATTGTAGAAAATGGAGCTGGAACGGATGAACAGTTTATCC 240
Apcy	CCTATTAATCTCCAGAAGAATTGTAGAAAATGGAGCTGGAACGGATGAACAGTTTACCC 240
	***** ** 240
Bmocy	CCCACTTTCATCTAATATCGCACATAGAGGAAGATCCGTAGATCTTGCTATTTTTTCACT 300
Bmacy	CCCACTTTCATCTAATATTGCACATAGAGGAAGATCCGTAGATCTTGCTATTTTTTCACT 300
Aycy	CCCCCTTCTTCAAATATTGCTCATGGAGGCTCTTCAGTAAATCTTGCTATTTTTTCATT 300
Apcy	CCCTCTCTTCAAATATTGCTCATGGAGGATCTTCAGTAGATCTTGCTATTTTTTCCCT 300
	*** ** 300

Bmocy	ACATTTAGCAGGTATTTTCATCAATTATAGGAGCAATTAATTTTATTACAACAATAATTAA 360
Bmacy	ACATTTAGCAGGTATTTTCATCAATTATAGGAGCAATTAATTTTATTACAACAATAATTAA 360
Ayey	ACATCTGCAGGTATTTCTTCAATTTTAGGAGCAATTAATTTTATTACTACAATTATTAA 360
Apycy	TCATCTGCAGGTATTTTCGTCATTTTAGGAGCAATTAATTTTATTACTACAATTATTAA 360 *** * ***** ***** ***** * * * * * 360
Bmocy	TATACGATTAATAATATATCATTTGATCAATTACCCTTATTTGTATGAGCTGTATCGGA 420
Bmacy	TATGCGATTAATAATATATCATTTGATCAATTACCCTTATTTGTATGAGCTGTAG-GAA 420
Ayey	TATACAAATAATAATTTATCATTTGATCAAATACCTTTATTTGTTGAGCTGTGCG-GAA 420
Apycy	TATACGAATAATAATTTATCATTTGATCAAATACCTTTATTTGTCTGAGCTGTTG-GAA 420 *** * * ***** ***** ***** ***** * * 420
Bmocy	TTACAGCATTTT-ATTATTATTATCACTACCCGTTTTAGCTGGAGCTATTACAATATTAT 480
Bmacy	TTACAGCATTTTTATTATTATTATCACTACCTGTTT-AGCTGGAGCTATTACAATATTAT 480
Ayey	TTACAGCCTTTTTACTTCTTTTATCATTACCTGCTTTAGCTGGAGCTATTACAATATTAC 480
Apycy	TTACAGCTTCTTACTTCTTTTATCATTACCTGTTTATAGCTGGAGCTATTACTATACTTT 480 ***** ** * * * * ***** ***** * * ***** ***** * * 480
Bmocy	TAACAGATCGAGACTTAAATACATCATTTTTTGATCCTGCTGGAGGAGGAGACCCAATT 540
Bmacy	TAACAGATCGAAACTTAAATACATCATTTTTTGATCCTGCTGGAGGAGGAGACCCAATT 540
Ayey	TAACAGATCGAAATTTAAATACTTCTTTCTTTGATCCTGCTGGAGGGGAGATCCTATTC 540
Apycy	TAACAGATCGAAACTTAAATACTTCTTTTTTTGATCCTGCTGGTGGAGGAGATCCAAATT 540 ***** * ***** ** * * ***** ***** * * ***** * * 540
Bmocy	TATATCAACATTTATTTTGAATTTTGGACATCCTGAAGTTTATATTTTAATTTTACC 598
Bmacy	TATATCAAC-TTTATTTTGATTTTTTGGGCATCCTGAAGTTTATATTTTAATTTTACC 598
Ayey	TTTATCAACATTTATTTTGAATTTTTTGGTCACCCTGAAGTTTATATTTTAATTTTACC 598
Apycy	TATATCAACATTTATTTTGAATTTTTTGGTCATCCAGAAGTTTATATTTTAATTTTACC 598 * ***** ***** ***** * * * * ***** ***** ***** 598

gested with the fourteen restriction endonucleases by using the equation of Nei and Li (1979). The genetic similarity coefficients varied from 0.227 to 1.000 with an average of 0.498. *B. mori* (Jam305) shared a 0.848, 0.303 and 0.227 genetic similarity coefficient with *B. mandarina*, *A. yamamai*, *A. pernyi*, respectively. *B. mandarina* shared a 0.394 and 0.348 genetic similarity coefficient with *A. yama-*

mai and *A. pernyi*, respectively, and the genetic similarity coefficient between *A. yamamai* and *A. pernyi* was 0.500 (Table II). As a result, the genetic similarity coefficient between *B. mori* and *B. mandarina* from the four silkmoths was determined at 0.848 which indicates a highly homogeneous genetic background.

The nucleotide sequence homology of a mitochondrial cytochrome oxidase I gene from the four silkmoths was investigated using PCR. The resulting PCR products of the cytochrome oxidase I gene were about 600 bp corresponding to the position of 1,617~2,213 in the cytochrome oxidase I gene of the *D. yakuba* (Clary and Wolstenholme, 1985). The sequences of the four amplified fragments were aligned using the CLUSTAL W program as shown in Fig. 2. As a result, *B. mori* shared a 97%, 85% and 87% identity with *B. mandarina*, *A. yamamai* and *A. pernyi*, respectively. *B. mandarina* shared a 87% and 88% identity with both *A. yamamai* and *A. pernyi*, and *A. yamamai* shared a 92% identity with *A. pernyi*. These sequences were compared with four insect species previously reported on (Table III). These sequences were also used for a phylogenetic analysis.

All the trees showed monophyly for both Bombycidae and Saturniidae (Fig. 3). The confidence limits for the monophyly of these trees were estimated using bootstrapping tests. The confidence limits for both Bombycidae and Saturniidae were more than 99% for all trees.

Until now, many studies have examined the relationships between Bombycidae and Saturniidae. *B. mori* and *B. mandarina* can be crossed, and the resulting hybrid progeny shows normal fertility (Kawaguchi, 1928; Aratake *et al.*, 1973). Studies of the fibroin gene (Kusuda *et al.*, 1986), ribosome gene (Makawa *et al.*, 1988), the immunological properties of arylphorin (Shimada *et al.*, 1992) and the arylphorin gene (Shimada *et al.*, 1995), all suggest close genetic relationships between *B. mori* and *B. mandarina*. Although their chromosome numbers are quite different (*yamamai*=31, *pernyi*=

Table III. Levels of similarity based on cytochrome oxidase I gene sequences in silkmoths and other insects.

Organisms	% Cytochrome oxidase I gene sequence similarity							
	1	2	3	4	5	6	7	8
<i>Bombyx mori</i>	100							
<i>Bombyx mandarina</i>	97	100						
<i>Antheraea yamamai</i>	85	87	100					
<i>Antheraea pernyi</i>	87	88	92	100				
<i>Drosophila yakuba</i>	80	81	82	82	100			
<i>Drosophila melanogaster</i>	82	83	82	82	91	100		
<i>Anopheles gambiae</i>	82	82	82	82	86	87	100	
<i>Apis mellifera</i>	76	78	75	75	75	75	76	100

1. *Bombyx mori*, 2. *Bombyx mandarina*, 3. *Antheraea yamamai*, 4. *Antheraea pernyi*, 5. *Drosophila yakuba*, 6. *Drosophila melanogaster*, 7. *Anopheles gambiae*, 8. *Apis mellifera*.




Fig. 3. Dendrogram of the phylogenetic relationships in silkmoths and other insects based on mtDNA cytochrome oxidase I sequence data.

49), *A. yamamai* and *A. pernyi* can be closed to produce F1 progeny (Kawaguchi, 1934; Shimada *et al.*, 1988). The progeny, however, are sterile due to incomplete oogenesis and other abnormalities (Shimada and Kobayashi, 1992). Kirimura (1962) showed that the fibroin of *A. yamamai* and *A. pernyi* are clearly similar. It is generally known that a mitochondria genome in a higher eukaryote is higher than a somatic genome on the evolutionary change-speed. In addition, Kimura (1968) reported in his paper on 'Evolutionary rate at the molecular level' that evolution on the molecular level does not depend on Darwinian selection but rather on the result of a spontaneous change of genetic matters in the unimportant genetic region. In other words, natural selection on the molecular level also chooses optimal best adaptability in response to ever changing environments in the evolutionary process. Therefore, genetic variation requires a minimum of change in the less important

regions of genes. Here, we investigated the phylogenetic relationships between the *B. mori* and *B. mandarina* species of the Bombycidae and the *A. yamamai* and *A. pernyi* species of the Saturniidae based on mtDNA RFLP and the cytochrome oxidase I gene. From the results of the mtDNA RFLP and the cytochrome oxidase I sequence homology, we can assume a closer genetic relationship between the *B. mori* and *B. mandarina* species than between *A. yamamai* and *A. pernyi*. Although this report is not exhaustive to prove that wild silkworm, *B. mandarina* is a possible ancestor of the domesticated *B. mori*, it does provide data to understand the mtDNA relationship of *B. mori* and *B. mandarina*. In addition, this study indicates that the mtDNA gene can serve as a good basis for the phylogenetic analysis of silkmths. At least mtDNA RFLP and cytochrome oxidase I gene sequences can provide evidence on the phylogenetic relationships among genus and families.

- Aratake Y. and Kayamura T. (1973), "Scattered" and "small sized" silkworm strains selected from the progeny of hybrid between *Bombyx mori* and *Theophila mandarina*. J. Seric. Sci. Jpn. **42**, 331–339.
- Beard C. B., Hamm D. M. and Collins F. H. (1993), The mitochondrial genome of the mosquito *Anopheles gambiae*: DNA sequence, genome organization, and comparisons with mitochondrial sequences of other insects. Insect Mol. Biol. **2**, 103–114.
- Clary D. O. and Wolstenholme D. R. (1985), The mitochondrial molecule of *Drosophila yakuba*: nucleotide sequence, gene organization and genetic code. J. Mol. Evol. **22**, 252–271.
- Crozier R. H. and Crozier Y. C. (1993), The mitochondrial genome of the honeybee, *Apis mellifera*: Complete sequence and genome organization, Genetics **133**, 97–117.
- Gray M. W. (1989), Origin and evolution of mitochondrial DNA. Annu. Rev. Cell. Biol. **5**, 25–50.
- Kambhampati S., Black W. C. and Rai K. S. (1992), Random amplified polymorphic DNA of mosquitoes: techniques, applications and analysis. J. Med. Entomol. **29**, 939–945.
- Kambhampati S. and Smith P. T. (1995), PCR primers for the amplification of four insect mitochondrial gene fragments. Insect Mol. Biol. **4**, 233–236.
- Kawaguchi E. (1928), Zytologische Untersuchungen am Seidenspinner und seinen Verwandten. I. Gametogenese von *Bombyx mori* L. und *B. mandarina* M. und ihrer Bastarde. Zeitschrift für Zellforschung und Mikroskopische Anatomie **7**, 519–552.
- Kawaguchi E. (1934), Zytologische Untersuchungen am Seidenspinner und seinen Verwandten. II. Spermatogenese bei *Antheraea yamamai* Guerin, *Antheraea pernyi* Guerin und ihren Bastarden. Jpn. J. Genet. **10**, 135–151.
- Kimura M. (1968), Evolutionary rate at the molecular level. Nature **56**, 217.
- Kirimura J. (1962), Studies on amino acid composition and chemical structure of silk protein by microbiological determination. Bull. Sericul. Exp. Stat. **17**, 447–552.
- Kumar S., Tamura K. and Nei M. (1993), MEGA: Molecular Evolutionary Genetics Analysis. Version 1.0. The Pennsylvania State University, University Park, PA.
- Kusuda J., Tazim Y., Onumaru K., Ninaki O. and Suzuki Y. (1986), The sequence around the 5' end of the fibroin gene from the wild silkworm, *Bombyx mandarina*, and comparison with that of the domesticated species, *B. mori*. Mol. Gen. Genet. **203**, 359–364.
- Lansman R. A., Shade R. O., Shapira J. F. and Avise J. C. (1981), The use of restriction endonucleases to measure mitochondrial sequence relatedness in natural populations. III. Techniques and potential application. J. Mol. Evol. **17**, 214–226.
- Lunt D., Zhang D. X., Szymura J. M. and Hewitt G. M. (1996), The insect cytochrome oxidase I gene: evolutionary patterns and conserved primers for phylogenetic studies, Insect Mol. Biol. **5**, 153–165.

- Makawa H., Takada N., Mikitani K., Ogura I., Miyajima N., Fujiwara H., Kobayashi M. and Ninaki O. (1988), Nucleolus organizers in the wild silkworm *Bombyx mandarina* and the domesticated silkworm *B. mori*. *Chromosoma* **96**, 263–269.
- Mitchell S. E., Cockburn A. F. and Seawright J. A. (1993), The mitochondrial genome of *Anopheles quadrimaculatus* species A: complete nucleotide sequence and genome organization. *Genome* **36**, 1058–1073.
- Moritz C., Dowling T. E. and Brown W. M. (1987), Evolution of animal mitochondrial DNA: Relevance for population biology and systematics. *Annu. Rev. Ecol. Syst.* **18**, 269–292.
- Nei M. and Li W. (1979), Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proc. Natl. Acad. Sci. USA*, **76**, 5219–5273.
- Roehrdanz R. L. (1995), Amplification of complete insect mitochondrial genome in two easy pieces. *Insect Mol. Biol.* **4**, 169–172.
- Saitou N. and Nei M. (1987), The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**, 406–425.
- Shimada T., Kobayashi J., Miyata T., Nagata M. and Kobayashi M. (1992), Comparison of arylphorins among species belonging to Saturniidae and Bombycidae (Lepidoptera) by immunodiffusion analyses. *Jpn. J. Appl. Entomol. Zool.* **36**, 119–126.
- Shimada T. and Kobayashi M. (1992), Fertility of F1 hybrids between *Antheraea yamamai* (Guerin-Meneville) and *Antheraea pernyi* (G.-M.). *Wild Silkmooths '91*, Akai H., Kato Y., Kiuchi M., Kobayashi J. eds., International Society for Wild Silkmooths, p185.
- Shimada T., Kurimoto Y. and Kobayashi M. (1995), Phylogenetic relationships of silkmooths inferred from sequence data of the arylphorin gene. *Mol. Phylogen. Evol.* **4**, 223–234.
- Shimada T., Yamauchi H. and Kobayashi M. (1988), Diapause of the inter-specific hybrids between *Antheraea yamamai* (Guerin-Meneville) and *A. pernyi* (G.-M.) (Lepidoptera: Saturniidae). *Jpn. J. Appl. Entomol. Zool.* **32**, 120–125.
- Simon C., Frati F., Beckenbach A., Crespi B., Liu H. and Flook P. (1994), Evolution, weighting and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain reaction primers. *Annu. Entomol. Soc. Amer.* **87**, 654–701.
- Tamura K. and Aotsuka T. (1988), Rapid isolation method of animal mitochondrial DNA by the alkaline lysis procedure. *Biochem. Genetics* **26**, 815–819.
- Thompson J. D., Higgins G. and Gibson T. J. (1994), CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, positions-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**, 4673–4680.