

## ***In vitro* Antiplasmodial Investigation of Medicinal Plants from El Salvador<sup>§</sup>**

Inga Köhler<sup>a</sup>, Kristina Jenett-Siems<sup>a,\*</sup>, Karsten Siems<sup>b</sup>, Marco Antonio Hernández<sup>c</sup>, Ricardo A. Ibarra<sup>c</sup>, Walter G. Berendsohn<sup>d</sup>, Ulrich Bienzle<sup>e</sup>, and Eckart Eich<sup>a</sup>

<sup>a</sup> Institut für Pharmazie (Pharmazeutische Biologie), Freie Universität Berlin, Königin-Luise-Strasse 2–4, D-14195 Berlin, Germany. E-mail: kjsiems@zedat.fu-berlin.de

<sup>b</sup> Analyticon Discovery GmbH, Hermannswerder Haus 17, D-14473 Potsdam, Germany

<sup>c</sup> ProBioTec S. A. de C. V., Apdo. 2827 C.G., San Salvador, El Salvador

<sup>d</sup> Botanischer Garten und Botanisches Museum Berlin-Dahlem, Königin-Luise-Strasse 6–8, D-14195 Berlin, Germany

<sup>e</sup> Institut für Tropenmedizin, Medizinische Fakultät Charité, Humboldt Universität zu Berlin, Spandauer Damm 30, D-14050 Berlin, Germany

\* Author for correspondence and reprint requests

Z. Naturforsch. **57c**, 277–281 (2002); received December 12, 2001/January 9, 2002

*Plasmodium falciparum*, *Calea tenuifolia*, Flavones

*In vitro* antiplasmodial activities of extracts from *Albizia saman*, Fabaceae, *Calea tenuifolia* (*C. zacatechichi*), Asteraceae, *Hymenaea courbaril*, Fabaceae, *Jatropha curcas*, Euphorbiaceae, *Momordica charantia*, Cucurbitaceae, and *Moringa oleifera*, Moringaceae were evaluated. From the lipophilic extract of *C. tenuifolia* five active flavones were obtained. 4',5-Dihydroxy-7-methoxyflavone [genkwanin] and 5-hydroxy-4',7-dimethoxyflavone [apigenin 4',7-dimethylether] exhibited the strongest antiplasmodial activity against a chloroquine-sensitive strain (poW) and a chloroquine-resistant strain (Dd2) of *Plasmodium falciparum* (IC<sub>50</sub> values: 17.1–28.5 µM). Furthermore octadeca-9,12-dienoic acid [linoleic acid] [IC<sub>50</sub> values of 21.8 µM (poW) and 31.1 µM (Dd2)] and octadeca-9,12,15-trienoic acid (α-linolenic acid) were isolated.

### **Introduction**

Malaria is still the most dangerous parasitic infectious disease which causes millions of deaths every year. In many countries where it is endemic the traditional medical methods hold a strong part in the public health care system. For safety reasons phytochemical investigations on medicinal plants traditionally used as antimalarials are urgently needed. In this context we are evaluating several species from El Salvador. Results of a bioassay-guided fractionation of *Exostema mexicanum* (Rubiaceae) were already described previously (Köhler *et al.*, 2000). In the present study we investigated another six traditional medicinal plants used as antimalarial or antipyretic remedies (Morton *et al.*, 1981): *Albizia [Samanea] saman* (Fabaceae), *Calea tenuifolia* (Asteraceae), *Hymenaea courbaril* (Fabaceae), *Jatropha curcas* (Euphorbiaceae), *Momordica charantia* (Cucurbitaceae), and *Moringa oleifera* (Moringaceae). Uses in traditional medicine and previously isolated

classes of constituents from these species are given in Table I. In our screening program a crude extract from the leaves of *Calea tenuifolia* showed the most promising antiplasmodial activity. Thus, a bioassay-guided fractionation was carried out in order to isolate and characterize the major anti-protozoan principles. Phytochemical and pharmacological investigations of the other active extracts will be part of further studies.

*Calea tenuifolia* Kunth is the correct name for the species commonly designated as *C. zacatechichi* Schlecht. It is a plant species of extensive popular medicinal use in Mexico (Díaz *et al.*, 1976). “Zacatechichi” (Nahuatl language) means “bitter grass”. It is also known as “dream herb”, “zacate de perro” (Spanish for dog’s grass), “hoja de dios” (God’s leaf), and thle-pela-kano (Chontal) (Rätsch, 1998). The shrub, 1–1.5 m in height, is native to dry forests from central Mexico to Costa Rica at 1500–1800 m (Morton, 1981). The leaves of *C. tenuifolia* are famed as a febrifuge, e.g. aqueous decoctions are given to patients in hospitals (Martinez, 1959). Mixe Indians are using such preparations against haemorrhage and malaria (Heinrich, 1989); it is also a popular remedy

<sup>§</sup> Part 3 in the series ‘Herbal remedies traditionally used against malaria’, for Part 2 see Kraft *et al.*, 2000.

against bilharziose and diarrhoea (Baytelman, 1979). Furthermore, this species is used by the Chontal Indians to produce or to enhance dreams of a divinatory nature (Mayagoitia *et al.*, 1986).

## Experimental

### General experimental procedures

For fractionation, a column containing reversed phase material (LiChroprep® RP-18, 40-63 µm) was used. Preparative high performance liquid chromatography (HPLC) was performed on a Knauer Eurochrom 2000 equipped with an Eurosphere 100 C-18 (10 µm, 22 × 250 mm) column. For preparative thin layer chromatography (TLC) aluminum sheets (20 × 20 cm) coated with silica gel 60 F<sub>254</sub> were used. Mass spectra were determined with a Finnigan MAT CH7A (220 °C, ionisation 70 eV) and <sup>1</sup>H NMR spectra were obtained using acetone-*d*<sub>6</sub> and MeOD as solvents with a Bruker AVANCE DPX 400 (400 MHz, TMS as internal standard). To evaluate the bioassays we used an Inotech cell harvester and for determination of IC<sub>50</sub> values a liquid scintillation counter Wallac 1450 MicroBeta plus.


### Plant material

*Albizia saman*, leaves collected May 7, 1995, at the roadside in San Pedro Mashuat, La Paz, El Salvador; voucher specimens deposited in the herbaria B, LAGU, MEXU, and MO; duplicate at MEXU authenticated by M. Sousa. *Calea tenuifolia*, leaves collected October 27, 1996, La Palma, Chalatenango, El Salvador, authenticated by González. *Hymenaea courbaril*, bark and leaves collected June 24, 1995, cantón Calderitas, Apastepaque, San Vicente, El Salvador, authenticated by González. *Jatropha curcas*, leaves collected in April, 1996, roadside at Rosario de Mora, San Salvador, El Salvador; voucher specimens authenticated by J.C. González were deposited in the herbaria B, ITIC, LAGU, and MO. *Momordica charantia*, stems with leaves collected June 8, 1996, at Laguna de Chanmico, San Juan Opico, La Libertad, El Salvador, voucher specimens authenticated by González were deposited in the herbaria B, ITIC, LAGU, and MO. *Moringa oleifera*, leaves, collected February 1, 1998, at Comalapa, road to International Airport, La Paz, El Salvador,

voucher specimens authenticated by Hernández were deposited at the herbarium LAGU.

### Extraction and isolation

In a screening program, the air dried plant material (20 g) was crushed and extracted three times for 2 h with 150 ml petrol-EtOAc (1:1, V/V) at room temperature to gain the lipophilic extracts. Afterwards the plant material was again air dried and treated three times with 150 ml MeOH-H<sub>2</sub>O (8:2) to afford the hydrophilic extracts. Additionally, air dried plant material of *Calea tenuifolia* was extracted for 2 h with 150 ml H<sub>2</sub>O under reflux to gain the aqueous extract. For further investigation of *C. tenuifolia*, air dried leaves (300 g) were extracted with petrol-EtOAc (1:1, V/V) and MeOH. The oily residue from the lipophilic extraction was subjected to column chromatography on RP 18 material and sequentially eluted with MeOH-H<sub>2</sub>O mixtures of decreasing polarity (up to 90% MeOH), MeOH, and CHCl<sub>3</sub>. Fractions, eluting with MeOH-H<sub>2</sub>O 8:2 and 9:1, proved to be most active in the antiplasmodial assay and were further purified by preparative HPLC with MeOH-H<sub>2</sub>O mixtures. Comparison of its spectroscopic data with literature values led to the identification of **1** as 5,7-dihydroxy-3',4'-dimethoxyflavone (luteolin 3',4'-dimethyl ether) (Nakanishi *et al.*, 1985). Compound **2** was identified as 4',5,7-trihydroxyflavone (apigenin) by <sup>1</sup>H-NMR and MS spectra and comparison with an authentic natural sample. Spectroscopic data from **3** were identical with literature data for 4',5-dihydroxy-7'-methoxyflavone



Flavone	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>
5,7-Dihydroxy-3',4'-dimethoxyflavone ( <b>1</b> )	OCH <sub>3</sub>	OCH <sub>3</sub>	OH
4',5,7-Trihydroxyflavone ( <b>2</b> )	H	OH	OH
4',5-Dihydroxy-7-methoxyflavone ( <b>3</b> )	H	OH	OCH <sub>3</sub>
5-Hydroxy-3',4',7-trimethoxyflavone ( <b>4</b> )	OCH <sub>3</sub>	OCH <sub>3</sub>	OCH <sub>3</sub>
5-Hydroxy-4',7-dimethoxyflavone ( <b>5</b> )	H	OCH <sub>3</sub>	OCH <sub>3</sub>

(genkwanin) (Brieskorn *et al.*, 1968). Compound **4** was identified as 5-hydroxy-3',4',7-trimethoxyflavone (luteolin 3',4',7-trimethyl ether) (Nakanishi *et al.*, 1985), **5** as 5-hydroxy-4',7-dimethoxyflavone (apigenin 4',7-dimethylether) (Silva *et al.*, 1971), **6** as octadeca-9,12,15-trienoic acid ( $\alpha$ -linolenic acid) (Bhacca *et al.*, 1963), and **7** as octadeca-9,12-dienoic acid (linoleic acid) (Gunstone, 1995). Fractionation of the inactive aqueous extract by column chromatography on RP 18 material with MeOH-H<sub>2</sub>O mixtures of decreasing polarity (up to 80% MeOH) led to active fractions, which contained compounds **1** and **3**.

#### Antiplasmodial activity

The antiplasmodial assay was performed by means of the microculture radioisotope technique as described previously (Jenett-Siems *et al.*, 2000). The concentration at which growth was inhibited by 50% (IC<sub>50</sub>) was estimated by interpolation. IC<sub>50</sub> values > 50 µg/ml for extracts and IC<sub>50</sub> values > 25 µg/ml for fractions, respectively, were considered inactive (O'Neill *et al.*, 1985).

#### Results and Discussion

Of the six plant species tested, lipophilic crude extracts of *C. tenuifolia*, *H. coubaril*, *M. oleifera*, and *M. charantia* showed significant antiplasmodial activity *in vitro* with IC<sub>50</sub> values between 6

Table I. *In vitro* antiplasmodial activity of plant extracts against *Plasmodium falciparum*.

Plant family/Species	Local uses	Previously isolated compounds	Part used	Extract	Mean IC <sub>50</sub> values [µg/ml] <sup>a</sup>	
					poW	Dd2
Asteraceae						
<i>Calea tenuifolia</i> Kunth (syn.: <i>C. zacatechichi</i> Schlecht.)	febrifuge (Martinez, 1959) haemorrhage, malaria (Heinrich, 1989) bilharziose, diarrhoea (Baytelman, 1979) enhancing dreams (Mayagoita <i>et al.</i> , 1986)	sesquiterpene lactones (Ortega <i>et al.</i> , 1970; Bohlmann <i>et al.</i> , 1981; Quijano <i>et al.</i> , 1977 and 1978; Herz <i>et al.</i> , 1980) flavonoids (Herz <i>et al.</i> , 1980)	leaves	lipophilic	10.4	24.3
				methanolic	19.7	19.5
				aqueous	> 50	> 50
Cucurbitaceae						
<i>Momordica charantia</i> L.	antimalarial (Muñoz <i>et al.</i> , 2000) hypoglycemic (Matsuda <i>et al.</i> , 1998)	oleanolic acid glycosides (Matsuda <i>et al.</i> , 1998)	whole plant	lipophilic	10.3	9.4
				methanolic	> 50	> 50
Euphorbiaceae						
<i>Jatropha curcas</i> L.	molluscicidal activity (Liu <i>et al.</i> , 1997) febrifuge (Morton, 1981)	phorbol esters (Liu <i>et al.</i> , 1997)	aerial parts	lipophilic	> 50	> 50
				methanolic	> 50	> 50
Fabaceae						
<i>Samanea saman</i> (Jacq.) Merr. (syn.: <i>Albizia saman</i> (Jacq.) F.v. Muell.)	antimalarial use of the related species <i>Albizia amara</i> Boiv. (Watt <i>et al.</i> , 1962)	terpenoids (Varshney <i>et al.</i> , 1985)	leaves	lipophilic	> 50	> 50
				methanolic	46.4	10.0
			bark	lipophilic	> 50	> 50
				methanolic	35.5	7.3
Fabaceae						
<i>Hymenaea courbaril</i> L.	substitute for quinine, rheumatism (Morton, 1981)	terpenoids (Khoo <i>et al.</i> , 1973)	stems	lipophilic	11.8	11.8
				methanolic	> 50	> 50
Moringaceae						
<i>Moringa oleifera</i> Lam.	antibiotic (Faizi <i>et al.</i> , 1995) antitumor (Murakami <i>et al.</i> , 1998)	isothiocyanates (Faizi <i>et al.</i> , 1995) thiocarbamate (Murakami <i>et al.</i> , 1998)	flowers	lipophilic	6.0	16.3
				methanolic	> 50	> 50
			leaves	lipophilic	7.8	15.4
				methanolic	> 50	> 50
			stems	lipophilic	> 50	> 50
				methanolic	> 50	> 50

<sup>a</sup> Tested in triplicates, internal standard: see Table II.

and 25 µg/ml (Table I). Methanolic crude extracts of *C. tenuifolia* and *S. saman* displayed an activity with IC<sub>50</sub> values ranging from 7 to 36 µg/ml. Bioactivity-guided fractionation of the lipophilic extract of *C. tenuifolia* led to the isolation of five flavones (**1-5**). To the best of our knowledge, these flavones were isolated from *C. tenuifolia* for the first time. All compounds showed activities against *P. falciparum*, with IC<sub>50</sub> values in a range between 4 and 40 µM (Table II). From all active fractions we isolated flavones. Certain active fractions additionally contained fatty acids as by-products. Two of them could be isolated and characterized as **6** and **7**. The antimalarial properties of unsaturated fatty acids were described previously: linoleic acid and α-linolenic acid seem to inhibit parasite growth in culture and *in vivo* (Krugliak *et al.*, 1995). Results of another study demonstrated that the antiplasmodial activity of the fatty acids is dependent in part

on the degree of unsaturation (Kumaratilake *et al.*, 1992). Since flavones were isolated from all active fractions of *C. tenuifolia*, we assume, that this class of compounds represents the major antiprozoan principle. Thus, our results may represent a rational explanation for a potential antimalarial effect of the leaves of *C. tenuifolia*. Furthermore, fractionation of the aqueous extract led to the detection of the flavones **1** and **3** in active fractions. This result can account for the ethnomedicinal use, because an aqueous leaf decoction is used in traditional Central American medicine.

#### Acknowledgements

This study was supported by a grant from Freie Universität Berlin (KFN-AX VIII-1) to Inga Köhler, and from Deutsche Pharmazeutische Gesellschaft to Kristina Jenett-Siems.

Table II. *In vitro* antiplasmodial activity of compounds isolated from the leaves of *Calea tenuifolia* against *Plasmodium falciparum*.

Compound	Mean IC <sub>50</sub> values <sup>a</sup>				
	poW		Dd2		
	[µg/ml]	[µM]	[µg/ml]	[µM]	
5,7-Dihydroxy-3',4'-dimethoxyflavone ( <b>1</b> ) [luteolin 3',4'-dimethyl ether]	13.6	43.3	10.5	33.4	
4',5,7-Trihydroxyflavone ( <b>2</b> ) [apigenin]	14.6	54.1	25.0	92.6	
4',5-Dihydroxy-7-methoxyflavone ( <b>3</b> ) [genkwanin]	5.4	19.0	8.1	28.5	
5-Hydroxy-3',4',7-trimethoxyflavone ( <b>4</b> ) [luteolin 3',4',7-trimethyl ether]	5.9	18.0	n.d.	n.d.	
5-Hydroxy-4',7-dimethoxyflavone ( <b>5</b> ) [apigenin 4',7-dimethylether]	6.0	20.1	5.1	17.1	
Octadeca-9,12,15-trienoic acid ( <b>6</b> ) [α-linolenic acid]	13.8	49.6	39.5	142.0	
Octadeca-9,12-dienoic acid ( <b>7</b> ) [linoleic acid]	6.1	21.8	8.7	31.1	
Artemisinin	0.0008	0.003	0.004	0.015	
Choroquine × 2 H <sub>3</sub> PO <sub>4</sub>	0.008	0.015	0.073	0.14	

<sup>a</sup> Tested in triplicate; n.d.: not determined.

- Baytelman B. (1979), Etnobotanica en el estado de Morelos. Instituto Nacional de Antropología Historia. Centro Regional de Morelos, Cuernavaca (México).
- Bhacca, NS. (1963), High resolution NMR-spectra catalog. Varian Associates. Palo Alto, 267.
- Bohlmann F., Fritz U., King R. M. and Robinson H. (1981), Fourteen heliangolides from *Calea* species. *Phytochemistry* **20**, 743–749.
- Brieskorn C. H. and Michel H. (1968), Flavone aus dem Blatt von *Rosmarinus officinalis* L. *Tetrahedron Lett.* **30**, 3447–3448.
- Díaz J. L. (1976), Uso de las plantas medicinales de México. Monografías Científicas II. Ed. Instituto Mexicano para el Estudio de las Plantas Medicinales. Mexico City (México).
- Faizi S., Siddiqui B. S., Saleem R., Siddiqui S., Aftab K. and Gilani A. (1995), Fully acetylated carbamate and hypotensive thiocarbamate glycosides from *Moringa oleifera*. *Phytochemistry* **38**, 957–963.
- Gunstone F. D., Harwood J. L. and Padley F. P. (1995), *The Lipid Handbook*. Chapman & Hall, London (UK).
- Heinrich M. (1989), Ethnobotanik der Tieflandmixe (Oaxaca, Mexico) und phytochemische Untersuchung von *Capraria biflora* L. (Scrophulariaceae). Dissertationes Botanicae No. 144. J. Cramer in Gebr. Borntraeger Verlagsbuchhdlg., Berlin und Stuttgart (Germany).
- Herz W. and Narendra K. (1980), Sesquiterpene lactones of *Calea zacatechichi* and *C. urticiflora*. *Phytochemistry* **19**, 593–597.
- Jenett-Siems K., Siems K., Jakupovic J., Solis P. N., Gupta M. P., Mockenhaupt F. P., Bienzle U. and Eich E. (2000), Sipandinolide: A butenolide including a novel type of carbon skeleton from *Siparuna andina*. *Planta Med.* **66**, 384.
- Khoo S. F. (1973), Structure and stereochemistry of the diterpenes of *Hymenaea courbaril* (Caesalpinoideae) seed pod resin. *Tetrahedron* **29**, 3379–3388.
- Köhler I., Jenett-Siems K., Mockenhaupt F. P., Siems K., Jakupovic J., González J. C., Hernández M. A., Ibarra R. A., Berendsohn W. G., Bienzle U. and Eich E. (2000), *In vitro* antiplasmodial activity of 4-phenylcoumarins from *Exostema mexicanum*. *Planta Med.* **66**, 89–91.
- Kraft C., Jenett-Siems K., Siems K., Gupta M. P., Bienzle U. and Eich E. (2000), Antiplasmodial activity of isoflavones from *Andira inermis*. *J. Ethnopharmacol.* **73**, 131–135.
- Krugliak M., Deharo E., Shalmiev G., Sauvain M., Moretti C. and Ginsburg H. (1995), Antimalarial effects of C18 fatty acids on *Plasmodium falciparum* in culture and on *Plasmodium vinckei petteri* and *Plasmodium yoelii nigeriensis* *in vivo*. *Exp. Parasitol.* **81**, 97–105.
- Kumaratilake L. M., Robinson B. S., Ferrante A. and Poulos A. (1992), Antimalarial properties of n-3 and n-6 polyunsaturated fatty acids: *in vitro* effects on *Plasmodium falciparum* and *in vivo* effects on *P. berghei*. *J. Clin. Invest.* **89**, 961–967.
- Liu S. Y., Sporer F., Wink M., Jourdane J., Hennig R., Li Y. L. and Ruppel A. (1997), Anthraquinones in *Rheum palmatum* and *Rumex dentatus* (Polygonaceae), and phorbol esters in *Jatropha curcas* (Euphorbiaceae) with molluscicidal activity against the schistosome vector snails *Oncomelania*, *Biomphalaria* and *Bulinus*. *Trop. Med. Int. Health* **2**, 179–188.
- Matsuda H., Li Y., Murakami T., Matsumura N., Yamahara J. and Yoshikawa M. (1998), Antidiabetic principles of natural medicines. III. Structure-related inhibitory activity and action mode of oleanolic acid glycosides on hypoglycemic activity. *Chem. Pharm. Bull.* **46**, 1399–1403.
- Martínez M. (1959), Las Plantas Medicinales de México. 4<sup>th</sup> ed. Ediciones Botas. Mexico City, 347–349.
- Mayagoitia L., Díaz J. L. and Contreras C. M. (1986), Psychopharmacologic analysis of an alleged oneirogenic plant: *Calea zacatechichi*. *J. Ethnopharmacol.* **18**, 229–243.
- Morton J. F. (1981), Atlas of Medicinal Plants of Middle America. Bahamas to Yucatan, Springfield (IL), C. Thomas (USA).
- Muñoz V., Sauvain M., Bourdy G., Callapa J., Rojas I., Vargas L., Tae A. and Deharo E. (2000), The search for natural bioactive compounds through a multidisciplinary approach in Bolivia. Part II. Antimalarial activity of some plants used by Mosetene Indians. *J. Ethnopharmacol.* **69**, 139–155.
- Murakami H., Kitazono Y., Jiwajinda S., Koshimizu K. and Ohigashiz H. (1998), Niaziminin, a thiocarbamate from the leaves of *Moringa oleifera*, holds a strict structural requirement for inhibition of tumor-promoter-induced Epstein-Barr virus activation. *Planta Med.* **64**, 319–323.
- Nakanishi T., Ogaki I. A., Murata H. and Nishi M. (1985), Flavonoids of *Striga asiatica*. *J. Nat. Prod.* **48**, 491–493.
- O'Neill M. J., Bray D. H., Boardmann P., Phillipson J. D. and Warhurst D. C. (1985), Plants as sources of antimalarial drugs part 1. *In vitro* method for the evaluation of crude extracts from plants. *Planta Med.* **51**, 394–398.
- Ortega A., Vivar R., Díaz E. and Romo J. (1970), Determinación de las estructuras de la calaxina y de la ciliarina, nuevos germacranólidos furanónicos. *Rev. Latinoam. Quím.* **1**, 81–85.
- Quijano L., Calderón J. S. and Ríos T. (1977), Los componentes de la *Calea zacatechichi* estructura de los caleocromenos A, Y, B. *Rev. Latinoam. Quím.* **8**, 90–93.
- Quijano L., Vivar R. and Ríos T. (1978), Los componentes de la *Calea zacatechichi* estructura de las caleinas A, Y, B. *Rev. Latinoam. Quím.* **9**, 86–89.
- Rätsch C. (1998), Enzyklopädie der psychoaktiven Pflanzen. AT-Verlag. Aarau, pp. 116–118.
- Silva M., Mundaca J. M. and Sammes P. G. (1971), Flavonoid and triterpene constituents of *Baccharis rhomboidalis*. *Phytochemistry* **10**, 1942–1943.
- Varshney I. P. and Vyas P. (1985), Samanin-C and samanin-E new saponins from *Pithecolobium saman*. *Fitterapia* **56**, 281–283.
- Watt J. M. and Breyer-Brandwijk M. G. (1962), The Medicinal and Poisonous Plants of Southern and Eastern Africa, Edinburgh London, E.&S. Livingstone LTD.