

Effect of the Number of *rol* Genes Integrations on Phenotypic Variation in Hairy Root-Derived *Hypericum perforatum* L. Plants

Hedviga Komarovská^a, Ján Košuth^a, Annalisa Giovannini^b, Andrija Smelcerovic^{c,d}, Sebastian Zuehlke^c, and Eva Čellárová^{a,*}

^a P. J. Šafárik University, Faculty of Science, Institute of Biology and Ecology, Mánesova 23, 041 54 Košice, Slovakia. Fax: +421 55 6337353. E-mail: eva.cellarova@upjs.sk

^b CRA-FSO, Experimental Unit for Floriculture and Ornamental Species, corso Inglese 508, 18038, Sanremo (Imperia), Italy

^c Institute of Environmental Research, Technical University of Dortmund, Otto-Hahn-Str. 6, D-44221 Dortmund, Germany

^d Present address: Department of Pharmacy, Faculty of Medicine, University of Nis, Bulevar Dr Zorana Djindjica 81, 18000 Nis, Serbia

* Author for correspondence and reprint requests

Z. Naturforsch. **65c**, 701–712 (2010); received April 19/July 23, 2010

The extent of phenotypic variation of St. John's wort (*Hypericum perforatum* L.) plants transformed with wild agropine ATCC 15834 *Agrobacterium rhizogenes* plasmid was evaluated with respect to the number of *rol* genes integrations. The transfer of T_L-DNA to plant explants during each transformation event was incomplete with different *rolA*, *rolB*, and *rolC* copy numbers. Along with typical features representing the hairy root syndrome, an altered size, number and density of dark and translucent glands, changes in ability to synthesize secondary metabolites, and reduced fertility were observed. The highest copy number of transferred *rol* genes resulted in weak expression of transgenic character and comparable quantitative parameters with the controls. Only 1 out of 11 transgenic clones was able to produce seed progeny and not more than 4 out of its 35 offsprings were positive for *rolC* gene integration. Sterility of the clones was due to retarded development of both gametophytes.

Key words: St. John's Wort, Transgenes Copy Number, Secondary Metabolites

Introduction

Hypericum perforatum L. is a medicinal plant used in folk medicine since ancient times. The extract is employed for the treatment of mild to moderate depression. *H. perforatum* possesses specialized structures, translucent and dark glands which are the sites for storage of phloroglucinols – hyperforin and its derivatives and naphthodianthrones, especially hypericin and pseudohypericin, respectively (Briskin and Gawienowski, 2001; Soelberg *et al.*, 2007). These two groups of biologically active compounds are of great interest due to their anticancer, antidepressive, and antiviral effects (Kubin *et al.*, 2005; Beerhues, 2006).

Agrobacterium rhizogenes rol genes located in T-DNA are responsible for the altered phenotype of transgenic plants (Nilsson and Olsson, 1997) including reduced apical dominance, dwarfing, shorter internodes, increased branching, higher rooting potential, smaller wrinkled leaves, reduced fertility, altered onset of flowering, and

flower morphology (Casanova *et al.*, 2005; Christey, 2001). Integration of these genes can also affect the production of secondary metabolites as they may function as potential activators of the secondary metabolism in transformed cells (Bulgakov, 2008). The extent of the hairy root syndrome depends not only on the species and individual genotype but also on unique transformation events. The expression of transgene/s can be influenced by the site of integration, also called position effect (Kooter *et al.*, 1999). Several T-DNA copies can be integrated into the plant genome, even from different bacterial cells (De Block and Debrouwer, 1991). However, a higher copy number of transgenes increases the risk of transgene silencing (Kooter *et al.*, 1999). The hairy root phenotype may thus not be inherited; the progeny of regenerants often does not express the T-DNA genes or does not display the parent phenotype even though the T-DNA is present in their genome (reviewed in Yin *et al.*, 2004).

Although genetic transformation of several plant species became a routine, the possibilities of genetic modification via *Agrobacterium*-mediated transformation in *H. perforatum* are still limited. According to Franklin *et al.* (2008) the main problem hindering transformation of this species is its strong recalcitrance to *Agrobacterium* infection. In spite of this there are few reports on successful *A. rhizogenes*-mediated transformation of *H. perforatum* (Di Guardo *et al.*, 2003; Vinterhalter *et al.*, 2006) including our recent paper on agrobacterial transformation of two other *Hypericum* species (Komarovská *et al.*, 2009). However, the effect of transgenes on the phenotype of the transgenic *Hypericum* plants was not assessed in detail so far.

The aim of the present work was to evaluate the impact of the number of *rol* genes integrations on the phenotype of transgenic *H. perforatum*. For such purpose (i) integration of *A. rhizogenes* T-DNA, its copy number and expression, (ii) the extent of the hairy root syndrome at phenotype level including secondary metabolite production in relationship to the *rol* gene copy number, and (iii) transgenic plant seed formation and transmission of the T-DNA into the offspring were studied.

Material and Methods

Plant material

Shoots of eleven transgenic clones and two non-transformed controls of *Hypericum perforatum* L. grown *in vitro* were used for the analyses. The transgenic root-derived clones R₁ and R₂ and leaf-derived L₁–L₉ were originated from hairy root cultures obtained from one mother *H. perforatum* genotype (C_M) by transformation with *Agrobacterium rhizogenes* strain ATCC 15834 (American Type Cultures Collection, Manassas, USA). Control clone C_L was untransformed, *i.e.* regenerated from C_M clone leaf. The clones R₁ and R₂ and L₁–L₉ were derived from individual hairy roots developed on root and leaf cuttings, respectively. Transgenic plants spontaneously regenerated on hormone-free basal medium. All clones were derived at CRA-FSO, Experimental Unit for Floriculture and Ornamental Species, Sanremo (Imperia), Italy as published earlier by Di Guardo *et al.* (2003). The mother clone originated from a wild habitat in Sanremo.

Culture conditions

The plant material was cultured on basal medium containing mineral salts according to Linsmaier and Skoog (1965) supplemented with B5 vitamins (Gamborg *et al.*, 1968), 100 mg l⁻¹ myo-inositol, 2 mg l⁻¹ glycine, 30 g l⁻¹ sucrose, and 6 g l⁻¹ agar; pH value was adjusted to 5.6 before autoclaving. The plantlets were kept in a culture room at (23 ± 1) °C, a 16/8-h photoperiod, fluorescent lightening of 33 μmol m⁻² s⁻¹, and 70% relative humidity. They were subcultured in a six-week interval.

Adaptation to ex vitro conditions

Transgenic and control *in vitro* cultivated plantlets with differentiated roots were transferred to aerated hydroponic vessels placed in a plastic container. Containers were gradually opened during 2–3 weeks to lower the humidity until the plants were able to grow under laboratory conditions. Four weeks after hydroponic cultivation the plants were transferred to autoclaved soil.

Detection of transgenes integrations and their expression

Integration and expression of *A. rhizogenes* T-DNA genes (*rolA*, *B*, *C*, *D* and *aux1*, 2) at the transcript level in the studied transgenic clones were detected by PCR and RT-PCR with gene-specific primers (Table I). Total genomic DNA and RNA were isolated from 100 mg of fresh plant material using the DNeasy Plant Mini Kit (Qiagen, Valencia, USA) and according to Jaakola *et al.* (2001), respectively. The isolated RNA was treated with DNase I (Invitrogen, Carlsbad, USA) to digest contaminating DNA. Reverse transcription (RT) was performed using 200 U M-MLV reverse transcriptase (Promega, Madison, USA) and 10 mM anchored oligoT primer at 42 °C according to the manufacturer's instructions. The number of integrated *rol* genes was measured by quantitative real time PCR (qPCR) and calculated according to Mason *et al.* (2002). Briefly, the relative amount of the *rolA*, *rolB*, and *rolC* genes was quantified by qPCR and normalized to an endogenous gene standard. Since all the transgenic clones were derived from one maternal genotype, the transformed clones shared, except the integrated T-DNA, the same genes and their copy number; any single-copy gene of *Hypericum* could be

Table I. Nucleotide sequence of primers used for detection of *rolA*, *B*, *C*, *D* and *aux1*, 2 genes in transgenic clones.

Primer	Nucleotide sequence [nt]	Product length [bp]	Tm [°C]	Reference
<i>rolA</i> for	5'-GTTAGGCGTGCAAAGGCCAAG-3'	239	60	Zdravković-Korać <i>et al.</i> (2003)
<i>rolA</i> rev	5'-TGCGTATTAATCCCGTAGGTC-3'			
<i>rolB</i> F1	5'-AAAGTCTGCTATCATCTCTCTATG-3'	348	60	Zdravković-Korać <i>et al.</i> (2003)
<i>rolB</i> R1	5'-AAAGAAGGTGCAAGCTACCTCTCT-3'			
<i>rolC</i> F1	5'-AAATGCGAAGTAGGCGCTCCG-3'	190	60	Di Guardo <i>et al.</i> (2003)
<i>rolC</i> R1	5'-TACGTGCGACTGCCCCGACGATGATG-3'			
<i>rolD</i> F3	5'-CTGAGCGTGTGGCTCATG-3'	101	60	Designed according to published <i>rolD</i> sequence
<i>rolD</i> R3	5'-GGAGGTAAAGACGAAGGACAGAG-3'			
<i>rolD</i> F1	5'-AGCTCTCAACGGCTTCATGTCGAT-3'	575	60	Zdravković-Korać <i>et al.</i> (2003)
<i>rolD</i> R1	5'-CTATTCCAACAGGACCTTGCCAAT-3'			
<i>aux1</i> for	5'-CATAGGATCGCCTCACAGGT-3'	198	61	Designed according to <i>aux1</i> (GenBank acc. no. DQ782955)
<i>aux1</i> rev	5'-CGTTGCTTGATGTCAGGAGA-3'			
<i>aux2</i> for	5'-AACGATAATAGCCCGCTGTG-3'	217	61	Designed according to <i>aux2</i> (GenBank acc. no. DQ782955)
<i>aux2</i> rev	5'-CGTCTTGGGTTTGTGGTTCT-3'			

used as endogenous standard. In this experiment a genomic fragment of the *hyp1* gene (the candidate gene of hypericin biosynthesis) was used for template normalization to count the *rol/hyp1* ratio. From the normalized relative amount of *rol* genes in all transgenic clones the theoretical value for one integration of each *rol* gene (the virtual calibrator, $r_{i\text{rolA}} - r_{i\text{rolC}}$) was calculated, and on the base of this value the final number of integrations was estimated. The values of virtual calibrator were 0.55 (*rolA*), 0.67 (*rolB*), and 0.69 (*rolC*). The PCR, RT-PCR, and qPCR reaction/amplification conditions were as described previously in Koperdákóvá *et al.* (2009a).

Evaluation of some quantitative parameters of plant organs

Plant height, length and number of internodes, axillary shoot number, and length of the longest root were scored on each transgenic and non-transgenic clone. The plants with at least one root longer than 1 cm were scored as rooting and those with at least one flower as flowering. Leaf area and number of dark and translucent glands were determined on two uppermost fully developed leaf pairs. The leaves were photographed under a stereomicroscope with a digital camera (ARTRAY ARTCAM), and the leaf area and length (*l*) and width (*w*) of black nodules were measured by UTHSCSA Image Tool 3.00 and Quick PHOTO Camera 2.1 software. The area of each dark gland was calculated as $\frac{1}{4}\pi lw$. The average area of one dark gland was calculated as

quotient of area of all glands on leaf and number of glands on leaf. The overall dark gland area on leaf is a sum of area of all glands on the leaf. Gland density (number *N* of glands per mm² of leaf blade) was calculated as quotient of number of glands on leaf and leaf area. From each clone 20 *in vitro* micropropagated plants were evaluated 6 weeks after subculture.

Preparation of the extracts and determination of secondary metabolite contents

Extraction of hypericins and hyperforin from *H. perforatum* has been optimized by Smelcerovic *et al.* (2006). The extractions were done using dry plant material (mass 0.39–0.61 g) and 10 ml of methanol by direct sonication. Sonications were performed for 20 min, using a Branson Sonifier B-12 apparatus (Branson, Heusenstamm, Germany) operating at 20 kHz and 60 W. Extracts were finally separated from the residual plant material by filtration (folded filters, grade 597 ½, Schweitzer & Schuell, Dassel, Germany). The residues were washed with 10-ml aliquots of methanol. Filtrates were then combined and methanol was added up to 25 ml. The extracts were stored in the dark at –20 °C until analysis was performed. Extractions were done in duplicate.

The composition of the extracts was analysed using an LC-ESI-MS-MS method slightly modified by Kusari *et al.* (2009) and Bonkanka *et al.* (2008). Chromatographic conditions were optimized for separation of hypericin, pseudohypericin, and hyperforin. External calibration was

Table II. Calculated copy number of *rolA*, *B*, *C* genes in transgenic *H. perforatum* clones. Normalized quantity (Norm) represents ratio of relative quantity of *rolX* to endogenous reference. The calculated copy number (*rolX*/*r_i*) is obtained as the ratio between the normalized quantity of *rolX* gene and the virtual calibrator *r_i*. The value of virtual calibrator for each gene was calculated according to Mason *et al.* (2002). The values \pm 95% confidence interval are shown. Integer numbers in the interval represent the expected number of integrations (int). The clones/genes with no integer in such interval are marked (*); in such case the copy number is estimated as the nearest integer to the calculated copy number.

Clone	<i>rolA</i> ($r_{i\text{rolA}} = 0.55$)			<i>rolB</i> ($r_{i\text{rolB}} = 0.67$)			<i>rolC</i> ($r_{i\text{rolC}} = 0.69$)		
	Norm \pm SD	<i>rolA</i> / <i>r_i</i>	int	Norm \pm SD	<i>rolB</i> / <i>r_i</i>	int	Norm \pm SD	<i>rolC</i> / <i>r_i</i>	int
R ₁	1.85 \pm 0.20	3.37 \pm 0.71	3–4	2.21 \pm 0.43	3.30 \pm 1.27	3–4	1.81 \pm 0.27	2.62 \pm 0.77	2–3
R ₂	0.58 \pm 0.23	1.05 \pm 0.83	1	0.71 \pm 0.04	1.06 \pm 0.11	1	0.70 \pm 0.05	1.02 \pm 0.13	1
L ₁	0.47 \pm 0.08	0.85 \pm 0.29	1	0.56 \pm 0.03	0.84 \pm 0.08	1*	0.46 \pm 0.11	0.66 \pm 0.32	1*
L ₂	0.62 \pm 0.05	1.12 \pm 0.17	1	0.74 \pm 0.04	1.11 \pm 0.11	1	0.65 \pm 0.06	0.95 \pm 0.17	1
L ₃	1.10 \pm 0.25	2.00 \pm 0.89	2	0.95 \pm 0.17	1.41 \pm 0.49	1	0.99 \pm 0.16	1.43 \pm 0.45	1
L ₄	1.49 \pm 0.17	2.70 \pm 0.61	3	1.56 \pm 0.21	2.33 \pm 0.60	2	1.48 \pm 0.19	2.15 \pm 0.55	2
L ₅	1.55 \pm 0.22	2.82 \pm 0.78	3	1.35 \pm 0.18	2.02 \pm 0.52	2	1.35 \pm 0.19	1.96 \pm 0.54	2
L ₆	2.64 \pm 0.33	4.80 \pm 1.17	4–5	1.74 \pm 0.31	2.59 \pm 0.91	2–3	1.25 \pm 0.25	1.81 \pm 0.72	2
L ₇	0.72 \pm 0.05	1.30 \pm 0.17	1*	0.95 \pm 0.19	1.42 \pm 0.55	1	0.89 \pm 0.05	1.29 \pm 0.14	1*
L ₈	0.51 \pm 0.05	0.93 \pm 0.18	1	0.66 \pm 0.02	0.98 \pm 0.04	1	0.69 \pm 0.01	1.00 \pm 0.01	1
L ₉	1.64 \pm 0.13	2.98 \pm 0.48	3	1.97 \pm 0.31	2.93 \pm 0.90	3	1.80 \pm 0.16	2.61 \pm 0.46	3

performed in the range 0.1–50 $\mu\text{g/ml}$ for pseudohypericin (Alexis Biochemicals, Lausen, Switzerland), hypericin (Sigma, Steinheim, Germany), and hyperforin (Sigma). Standard solutions were stored in the dark at -20°C . The correlation coefficient for the calibration curves was > 0.989 . All procedures were carried out in the dark. The relative standard deviation (RSD) of the analytical method was determined by eighth injections of an extract and was below 6% for all compounds.

Histological observation of reproductive tissues

Flower buds and flowers of the transgenic clone R₁ and control clone C_M were collected at different developmental stages: small buds without visible petals, buds with yellow petals, buds just before anthesis, fully opened flowers and overblown flowers. The plant material was fixed in Carnoy's solution for 24 h. After dehydration in butanol and ethanol series and embedding in paraffin, flower sections (7 μm) were cut with a microtome, dried on slides, deparaffinized in toluene, and then stained with alcian blue and safranin. Slides were observed under a light microscope (Olympus C011). Photographs were taken with a C-3020ZOOM digital camera (Olympus).

Statistical analyses

The differences between the clones were evaluated by ANOVA or Kruskal-Wallis tests, depend-

ing on the result of the Levene test. This was followed by multiple comparisons of mean ranks. The results were evaluated at the 0.05 significance level using Statistica 7 software.

Results

In this work plants of eleven transgenic *H. perforatum* clones, L₁–L₉ and R₁ and R₂, initiated from hairy roots developed on leaf (L) and root (R) cuttings were characterized. The phenotypic changes were correlated with the *rol* genes copy number. Mother genotype (C_M) and non-transgenic clone C_L regenerated from mother clone leaf were used as controls.

Detection and quantification of T-DNA integrations

Integration of T-DNA genes was determined by PCR with specific primers. While the *rolA*, *B*, *C*, *D* genes localized on T_L-DNA were found in all hairy root-derived clones, transfer of *aux1* from T_R-DNA was proved only in L₉ and *aux2* was missing in all samples. Expression of the integrated genes with an exception of the *rolD* gene was confirmed by RT-PCR. Re-analysis of *rolD* with another set of PCR primers (*rolD*-F3/R3), delimiting longer, 575 bp long *rolD* fragment, did not reveal its transfer. The presence of the shorter 101 bp long *rolD* PCR product and lack of the




Fig. 1. Phenotypic variation of transgenic clones in comparison with the controls: C_M, mother clone; C_L, non-transformed clone regenerated from mother clone leaf; R₁, R₂, transgenic clones derived from root cuttings; L₁–L₉, transgenic clones derived from leaf cuttings.

longer one indicated incomplete transfer of the respective T-DNA region.

The transgenic clones differed in the number of *rolA*, *B*, *C* genes integrations (Table II). The clones R₂, L₁, L₂, L₃, L₇, and L₈ possessed only one or two integrations; in the remaining clones more than three integrations occurred. The highest number of all *rol* genes was detected in R₁ and L₉ clones. The clones shared either the same number of integrated *rolA*, *B*, *C* genes (R₂, L₁, L₂, L₇, L₈, and L₉), *i.e.* the part of the T_L-DNA covering *rolA*, *B*, *C* genes was completely transferred, or the number of *rolA*, *B*, *C* genes was descending from *rolA* to *rolC* indicating an incomplete transfer of this region.

Phenotype of in vitro cultivated transgenic plantlets

The extent of hairy root syndrome expression observed among the transgenic plants varied (Fig. 1). The least pronounced appeared in the clones R₁ and L₉ with the highest number of *rol* genes integrations.

As seen from Figs. 2a–f the number of *rol* genes integrations influenced almost all quantitative characteristics studied with an exception of

number of internodes (Fig. 2b). The plants with a higher copy number resembled the controls while those with only one or two integrations expressed the features typical for hairy root syndrome more intensively, including reduced flowering (Fig. 2g). Also formation of roots was reduced in transgenic clones (Fig. 2h).

Variation of dark and translucent glands and secondary metabolite content

Evaluation of size, number, and density of dark glands as a site for hypericin accumulation, and number and density of translucent glands on leaf serving for storage of hyperforin revealed that most of these characters are likely affected by transformation as well as transgenes copy number (Figs. 3a–e). The significantly smaller leaf blade of transgenic clones compared to controls was characterized by a higher dark gland density (Fig. 3d). It was reflected, with an exception of clone L₃, in the higher total content of hypericins (hypericin and pseudohypericin) (Fig. 3f, Table III), although a clear correlation between hypericin content and number or size of dark glands was not proved. The majority of the transgenic clones accumulated less hyperforin than controls




Fig. 2. Some quantitative parameters of transgenic and non-transgenic *H. perforatum* clones: (a) plant height; (b) number of internodes; (c) internode length; (d) leaf area; (e) number of axillary shoots; (f) length of the longest root in plant; (g) percentage of flowering plants; (h) percentage of rooting plants. Small letters indicate homogenous groups.

but without unambiguous relationship between the amount of hyperforin and translucent glands number and density.

Histology of gametophytes in transgenic clones and transfer of T-DNA into the seed progeny

Only three transgenic clones, R₁, R₂ and L₂, were successfully adapted to *ex vitro* conditions. The problems with *ex vitro* adaptation were mainly due to poor rooting of the plants and a fragile root system. Although all the *ex vitro* adapted clones flowered, only the clone R₂ produced seed progeny. To elucidate the reasons of sterility of transgenic plants, a histological study of pistils and anthers in different stages of development of the clone R₁ was performed. The results revealed several anatomical differences in the development of male and female gametophytes compared to the mother non-transgenic clone C_M.

The development of both gametophytes in transgenic plants lagged behind the control in all evaluated stages. Finally, mature pollen grains were not observed in transgenic flowers, even after anthesis. The zygotes and early stages of embryo development were visible only in a few ovules of overblown flowers, and endosperm development was not observed in any of the transgenic ovules.

To assess the inheritance and the ability of transgenic *Hypericum* plants to transmit the foreign T-DNA into the offspring we studied the seed progeny of the only fertile transgenic clone (R₂). The screening for the presence of the *rolC* gene revealed its integration only in 4 out of 35 tested offsprings.

Discussion

Despite several reports on the efficient *A. rhizogenes*-mediated transformation of *H. perforatum* (Di Guardo *et al.*, 2003; Vinterhalter *et al.*, 2006), only limited information on the effect of transgenes and their copy number on the phenotype of transgenic *H. perforatum* plants is avail-

Table III. Secondary metabolite content (mg/g dry weight) in control and transgenic clones.

Clone	Pseudohypericin	Hypericin	Hyperforin
R ₁	2.63	0.84	5.68
R ₂	2.59	0.47	3.99
L ₁	3.82	0.63	5.49
L ₂	2.70	0.53	3.88
L ₃	1.83	0.34	3.51
L ₄	3.12	0.50	4.50
L ₅	2.17	0.53	1.94
L ₆	3.18	0.56	3.44
L ₇	2.73	0.44	4.86
L ₈	2.82	0.47	3.94
L ₉	3.17	0.61	5.72
C _L	1.89	0.39	8.07
C _M	2.15	0.37	5.24

able. In our study transgenic clones regenerated from hairy roots derived by *A. rhizogenes* ATCC 15834 were characterized with respect to the phenotypic alterations related to the transgene copy number. The hairy root syndrome with different extent was observed in all transgenic clones. Contrary to our results all transgenic clones of *H. perforatum* transformed by *A. rhizogenes* plasmid A4M70GUS were similar in appearance to the controls (Vinterhalter *et al.*, 2006). However, in the present work the copy number of the *rol* genes was not studied and so their silencing effect could not be excluded.

In the genome of all transgenic clones analysed in our study the insertion and expression of the *rolA*, *B*, *C* genes from T_L-DNA was proved. The left part of agropine Ri plasmid T-DNA (T_L-DNA) is composed of 18 ORFs (open reading frames), where ORFs 10, 11, 12 and 15 correspond to *rolA*, *B*, *C*, *D* genes (White *et al.*, 1985; Slighton *et al.*, 1986). The physical distance between *rolA*, *B*, *C* and *rolD* in T_L-DNA might explain the problems with the transfer of the whole *rolD* gene in all samples. The *rolA*, *B*, *C* genes copy number was descending in the direction from *rolA* to *rolC* in several clones. Similar results with incomplete T_L-DNA transfer were reported for some of the *Coffea arabica* hairy roots lines derived by Alpizar *et al.* (2008). The integration of the T_R-DNA was confirmed only in clone L₉ by the detection of *aux1* gene. Incorporation of only T_L-DNA, essential for hairy root induction, into the plant genome is a well-known feature of transformation with agropine strains of *A. rhizogenes* (Jouanin *et al.*, 1987), as recently proved *e.g.* by studies of




Fig. 3. Some quantitative parameters of dark and translucent glands on leaves: (a) dark gland area; (b) dark gland number on leaves; (c) overall dark gland area on leaf; (d) dark gland density on leaf; (e) number of translucent glands on leaf; (f) relationship between hypericin content and dark gland density. Small letters indicate homogenous groups.

Alpizar *et al.* (2008) and Christensen *et al.* (2008). Also in our recent experiments with *H. tomentosum* and *H. tetrapterum* transformation T_R-DNA was integrated only into the genome of several hairy root lines (Komarovská *et al.*, 2009).

Most of the clones possessed one or two copies of the *rol* genes. The transgenic clones with the highest *rol* genes copy number, R₁ and L₉, resembled the phenotypic quantitative parameters of the control. This suggests that in the transgenic

clones with higher transgene copy number the silencing of the introduced genes might giving and gave rise to a suppressed hairy root phenotype. Post-transcriptional silencing has been registered frequently in the transgenic lines of eastern white pine with three or more copies but never with only one T-DNA copy (Tang *et al.*, 2007). However in our experiment, a complete switch off of the *rol* gene expression in the clones R₁ and L₉ was not observed since the *rolA*, *B*, and *C* gene transcripts were detected and several symptoms of the hairy root regenerated plants noticed as well. The plants with the selfsame number of integrations did not always resemble the same phenotype. It was markedly expressed in the clones L₄ and L₅ which showed an identical number of *rol* genes but a different phenotype. One of the reasons is the position effect. Tang *et al.* (2007) demonstrated by *in situ* hybridization that foreign genes became silenced if two and more copies of the transgene are introduced in the same chromosome. This may be also the case for the L₄–L₆ *H. perforatum* transgenic clones which possessed several T-DNA integrations and had also the intensive hairy root phenotype. However, in these clones incomplete *rolA*, *B*, *C* transfer occurred.

Transgenic plants derived from two different organs of the same mother plant were compared. According to our findings, it seems that the origin of the explants does not influence the habitus of transgenic plants, but may affect rooting of *H. perforatum*. While both of the root segment-derived clones (R₁ and R₂) were able to root and grow under *ex vitro* conditions, only one out of nine leaf explant-derived clones (L₁–L₉) was capable of rooting and acclimation. Unlike our results plants with introduced *rol* genes are usually characterized by increased rooting ability which was used for improvement of rooting of woody plants such as apple or cherry tree (Sedira *et al.*, 2001; Gutiérrez-Pesce *et al.*, 1998).

H. perforatum plants are an important source of hypericins which are accumulated in dark glands (Briskin and Gawienowski, 2001). The transformation events in our experiment influenced mainly the size of the dark glands and the total leaf area covered by them. The dark gland density was changed mainly due to the altered leaf size of the transgenic clones. The literature data on correlation between hypericin content and dark gland density are inconsistent. While Southwell and Campbell (1991) and Cirak *et al.* (2006) re-

ported a close relationship, Čellárová *et al.* (1994) and Walker *et al.* (2001) did not find any association. Kirakosyan *et al.* (2003) observed a markedly different content of hypericin in several lines possessing a comparable number of dark glands on leaf. Neither our results confirmed a close correlation between dark gland number or size and hypericin content, although a certain relationship between the dark gland density on leaf blade and hypericin content was found. The accumulation of hypericin may be affected also by the structure of the dark glands as it was found that the bigger and better developed cells of the peripheral layer of dark glands accumulated higher amounts of hypericin (Kornfeld *et al.*, 2007).

Although the dark glands are well known as hypericin accumulation sites, the function of translucent glands as sites of hyperforin accumulation was published only recently (Soelberg *et al.*, 2007). As the translucent gland density on leaves seems to be genetically conserved, the reduced leaf area of transgenic clones leads to a decreased number of these glands. The results of our experiments did not confirm any relationship between hyperforin content and density or number of translucent glands. The capacity of the glands to store hyperforin might be therefore affected or altered by transformation. According to Karppinen and Hohtola (2008) and our preliminary findings (data not shown) the roots are also capable of hyperforin production/accumulation as well as indicating that the translucent glands on the aerial parts might not be the only structures where hyperforin is accumulated.

Genetic transformation by wild-type *A. rhizogenes* strains may affect the biosynthetic potential as the *rol* genes act as potential activators of the secondary metabolism (Bulgakov, 2008). Although the hairy root cultures represent a more promising system, there are also several data showing an increased secondary metabolites production in intact transgenic plants as well. For example, Phongprueksapattana *et al.* (2008) recently reported a significantly higher concentration of mitragynin in transgenic *Mitragyna speciosa*. However, transgenic plants of *H. perforatum* analysed in this study did not appear profitable for production of any tested compound even though hypericin and pseudohypericin contents were slightly increased. A more perspective application of transformation within the genus *Hypericum* would be the use of gene transfer as a means to study the gene func-

tion *in situ* via introduction of new genes or silencing/inactivation of the candidate genes. Such an approach necessitates a transformation system with high efficacy and reproducibility which is not available for *H. perforatum* yet. Unlike *H. perforatum*, we have recently reported higher efficiency of *Agrobacterium*-mediated transformation in two other *Hypericum* species (Komarovská *et al.*, 2009), which implies the successful use of such approach.

The hairy root-regenerated transgenic plants are usually partially or completely sterile (Casanova *et al.*, 2005; Christey, 2001). Transformation negatively affected the flowering and seed production also in transgenic *H. perforatum*. As found in our previous experiment the floral parts of the transgenic *ex vitro* cultivated *H. perforatum* clone were significantly smaller than the non-transgenic ones (Koperdákóvá *et al.*, 2009b). Histological observations showed that in transgenic plants the development of both gametophytes was retarded. Moreover, no signs of endosperm presence were encountered. Reduced number of flowers and fertility of the transgenic *H. perforatum* clones along with the problems with acclimation did not enable a deeper study of the T-DNA transmission into the seed progeny. A reduced number of inflorescences per plant was observed, for example, in two hairy root-derived *Helichrysum* clones; this symptom was transmitted into the *rolC* positive seed progeny (Giovannini *et al.*, 2008). For transgenic *rolD*-positive plants a higher number of flowers and fruits and earlier onset of flowering is typical (Mauro *et al.*, 1996; Bettini *et al.*, 2003). Successful integration of the whole *rolD* would therefore positively influence the flowering in *H. perforatum*.

Conclusion

This study provides an insight into phenotypic alterations of *H. perforatum* transgenic plants.

The level of expression of the hairy root phenotype in the transgenic plants was markedly affected by the copy number of integrated *rol* genes. The phenotype of transgenic clones with the highest number of *rol* genes integrations resembled the controls; but the hairy root syndrome was not completely suppressed in these plants. The origin of the explants used for transformation influenced the ability of transgenic clones to adapt to *ex vitro* conditions since better acclimation was observed in clones originating from root segments. T-DNA transfer into the seed progeny as well as seed production ability of transgenic plants was considerably reduced. The future of transgenesis in this medicinal plant or genus can be seen mainly in studying the function of the candidate genes of secondary metabolite pathways. An efficient transformation system should be used as a tool for introducing and/or switching off the genes of interest, which can enable the determination of their function.

Acknowledgement

This work was supported by Slovak Research and Development Agency under contract Nos. VVCE-0001-07, 0321-07, LPP 0015-07, and LPP-0021-09, by Scientific Grant Agency of the Slovak Republic (1/0049/08), and by Structural Funds of EU under contract No. 007/20092.1/OPVaV.

The authors appreciate the contribution of Dr. Anna Macková to interpretation of histological data, Jana Koperdákóvá, PhD, for part of experimental design and statistical evaluation, and diploma students Ivana Masárová Kubalcová and Zuzana Oravcová for collecting the data. We would also like to thank Bc. Ľudmila Majšniarová, Mrs. Zdenka Lacková, and Mrs. Alžbeta Maková for technical assistance.

Alpizar E., Dechamp E., Lapeyre-Montes F., Guilhaumon C., Bertrand B., Jourdan C., Lashermes P., and Etienne H. (2008). *Agrobacterium rhizogenes*-transformed roots of coffee (*Coffea arabica*) – conditions for long-term proliferation, and morphological and molecular characterization. *Ann. Bot. London* **101**, 921–940.

Beerhues L. (2006). Hyperforin. *Phytochemistry* **67**, 2201–2207.

Bettini P., Michelotti S., Bindi D., Giannini R., Capuana M., and Buiatti M. (2003). Pleiotropic effect of the insertion of the *Agrobacterium rhizogenes rolD* gene in tomato (*Lycopersicon esculentum* Mill.). *Theor. Appl. Genet.* **107**, 831–836.

Bonkanka C. X., Smelcerovic A., Zuehlke S., Rabanal R. M., Spittler M., and Sánchez-Mateo C. C. (2008). HPLC-MS analysis and anti-oedematogenic activity

- of *Hypericum grandifolium* Choisy (Hypericaceae). *Planta Med.* **74**, 719–725.
- Briskin D. P. and Gawienowski M. C. (2001), Differential effects of light and nitrogen on production of hypericins and leaf glands in *Hypericum perforatum*. *Plant. Physiol. Biochem.* **39**, 1075–1081.
- Bulgakov V. P. (2008), Functions of *rol* genes in plant secondary metabolism. *Biotechnol. Adv.* **26**, 318–324.
- Casanova E., Trillas M. I., Moysset L., and Vainstein A. (2005), Influence of *rol* genes in floriculture. *Biotechnol. Adv.* **23**, 3–39.
- Čellárová E., Daxnerová Z., Kimáková K., and Halušková J. (1994), The variability of the hypericin content in the regenerants of *Hypericum perforatum*. *Acta Biotechnol.* **14**, 267–274.
- Christensen B., Sriskandarajah S., Serek M., and Müller R. (2008), Transformation of *Kalanchoe blossfeldiana* with *rol*-genes is useful in molecular breeding towards compact growth. *Plant Cell. Rep.* **27**, 1485–1495.
- Christey M. C. (2001), Use of Ri-mediated transformation for production of transgenic plants. *In Vitro Cell. Dev. Biol. Plant* **37**, 687–700.
- Cirak C., Sağlam B., Ayan A. K., and Kevseroğlu K. (2006), Morphogenetic and diurnal variation of hypericin in some *Hypericum* species from Turkey during the course of ontogenesis. *Biochem. Syst. Ecol.* **34**, 1–13.
- De Block M. and Debrouwer D. (1991), Two T-DNA's co-transformed into *Brassica napus* by a double *Agrobacterium tumefaciens* infection are mainly integrated at the same locus. *Theor. Appl. Genet.* **82**, 257–263.
- Di Guardo A., Čellárová E., Koperdákóvá J., Pistelli L., Ruffoni B., Allavena A., and Giovannini A. (2003), Hairy root induction and plant regeneration in *Hypericum perforatum* L. *J. Genet. Breed.* **57**, 269–278.
- Franklin G., Conceição L. F. R., Kombrink E., and Dias A. C. P. (2008), *Hypericum perforatum* plant cells reduce *Agrobacterium* viability during co-cultivation. *Planta* **227**, 1401–1408.
- Gamborg O. L., Miller R. A., and Ojima K. (1968), Nutrient requirements of suspension cultures of soybean root cultures. *Exp. Cell. Res.* **50**, 151–158.
- Giovannini A., Mascarello C., Pipino L., and Nostro A. (2008), *Agrobacterium rhizogenes*-mediated transformation in Mediterranean *Helichrysum*. *Transgenic Plant J.* **2**, 54–61.
- Gutiérrez-Pesce P., Taylor K., Muleo R., and Rugini E. (1998), Somatic embryogenesis and shoot regeneration from transgenic roots of the cherry rootstock Colt (*Prunus avium* × *P. pseudocerasus*) mediated by pRi 1855 T-DNA of *Agrobacterium rhizogenes*. *Plant. Cell. Rep.* **17**, 574–580.
- Jaakola L., Pirttilä A. M., Halonen M., and Hohtola A. (2001), Isolation of high quality RNA from bilberry (*Vaccinium myrtillus* L.) fruit. *Mol. Biotechnol.* **19**, 201–204.
- Jouanin L., Guerche D., Pamboukdjian N., Tourneur C., Casse-Delbart F., and Tourneur J. (1987), Structure of T-DNA in plants regenerated from roots transformed by *Agrobacterium rhizogenes* strain A4. *Mol. Gen. Genet.* **206**, 387–392.
- Karppinen K. and Hohtola A. (2008), Molecular cloning and tissue-specific expression of two cDNAs encoding polyketide synthases from *Hypericum perforatum*. *J. Plant Physiol.* **165**, 1079–1086.
- Kirakosyan A., Gibson D. M., and Sirvent T. (2003), A comparative study of *Hypericum perforatum* plants as sources of hypericins and hyperforins. *J. Herbs Spices Med.* **10**, 73–88.
- Komarovská H., Giovannini A., Košuth J., and Čellárová E. (2009), *Agrobacterium rhizogenes*-mediated transformation of *Hypericum tomentosum* L. and *Hypericum tetrapterum* Fries. *Z. Naturforsch.* **64c**, 864–868.
- Kooter J. M., Matzke M. A., and Meyer P. (1999), Listening to the silent genes: transgene silencing, gene regulation and pathogen control. *Trends Plant Sci.* **4**, 340–347.
- Koperdákóvá J., Katkovčínová Z., Košuth J., Giovannini A., and Čellárová E. (2009a), Differences in morphogenetic response to PGRs between transformed and non-transformed *Hypericum perforatum* L. clones. *Acta Biol. Cracov. Bot.* **51**, 61–70.
- Koperdákóvá J., Komarovská H., Košuth J., Giovannini A., and Čellárová E. (2009b), Characterization of hairy root-phenotype in transgenic *Hypericum perforatum* L. clones. *Acta Physiol. Plant.* **31**, 351–358.
- Kornfeld A., Kaufman P. B., Lu C. R., Gibson D. M., Bolling S. F., Warber S. L., Chang S. C., and Kirakosyan A. (2007), The production of hypericins in two selected *Hypericum perforatum* shoot cultures is related to differences in black gland structure. *Plant. Physiol. Biochem.* **45**, 24–32.
- Kubin A., Wierrani F., Burner U., Alth G., and Grunberger W. (2005), Hypericin – The facts about a controversial agent. *Curr. Pharm. Design* **11**, 233–253.
- Kusari S., Zühlke S., Borsch T., and Spiteller M. (2009), Positive correlations between hypericin and putative precursors detected in the quantitative secondary metabolite spectrum of *Hypericum*. *Phytochemistry* **70**, 1222–1232.
- Linsmaier E. M. and Skoog F. (1965), Organic growth factor requirements of tobacco tissue cultures. *Physiol. Plant.* **18**, 100–127.
- Mason G., Provero P., Vaira A. M., and Accotto G. P. (2002), Estimating the number of integrations in transformed plants by quantitative real-time PCR. *BMC Biotech.* **2**, 20.
- Mauro M. L., Trovato M., De Paolis A., Gallelli A., Costantino P., and Altamura M. M. (1996), The plant oncogene *rolD* stimulates flowering in transgenic tobacco plants. *Dev. Biol.* **180**, 693–700.
- Nilsson O. and Olsson O. (1997), Getting to the root: The role of the *Agrobacterium rhizogenes rol* genes in formation of hairy roots. *Physiol. Plant.* **100**, 463–473.
- Phongprueksapattana S., Putalun W., Keawpradub N., and Wungsintaweekul J. (2008), *Mitragyna speciosa*: Hairy root culture for triterpenoid production and high yield of mitragynine by regenerated plants. *Z. Naturforsch.* **63c**, 691–698.
- Sedira M., Holfors A., and Welander M. (2001), Protocol for transformation of the apple rootstock Jork 9 with the *rol B* gene and its influence on rooting. *Plant Cell. Rep.* **20**, 517–524.
- Slightom J. L., Durand-Tardif M., Jouanin L., and Tepfer D. (1986), Nucleotide sequence analysis of T_L-DNA of *Agrobacterium rhizogenes* agropine type plasmid.

- Identification of open reading frames. *J. Biol. Chem.* **261**, 108–121.
- Smelcerovic A., Spiteller M., and Zuehlke S. (2006), Comparison of methods for the exhaustive extraction of hypericins, flavonoids, and hyperforin from *Hypericum perforatum* L. *J. Agric. Food Chem.* **54**, 2750–2753.
- Soelberg J., Jørgensen L. B., and Jäger A. K. (2007), Hyperforin accumulates in the translucent glands of *Hypericum perforatum*. *Ann. Bot. London* **99**, 1097–1100.
- Southwell I. A. and Campbell M. H. (1991), Hypericin content variation in *Hypericum perforatum* L., Australia. *Phytochemistry* **30**, 475–478.
- Tang W., Newton R. J., and Weidner D. A. (2007), Genetic transformation and gene silencing mediated by multiple copies of a transgene in eastern white pine. *J. Exp. Bot.* **58**, 545–554.
- Vinterhalter B., Ninkovic S., Cingel A., and Vinterhalter D. (2006), Shoot and root culture of *Hypericum perforatum* L. transformed with *Agrobacterium rhizogenes* A4M70GUS. *Biol. Plant.* **50**, 767–770.
- Walker L., Sirvent T., Gibson D., and Vance N. (2001), Regional differences in hypericin and pseudo-hypericin concentrations and five morphological traits among *Hypericum perforatum* plants in the North-western United States. *Can. J. Bot.* **79**, 1248–1255.
- White F. F., Taylor B. H., Huffman G. A., Gordon M. P., and Nester E. W. (1985), Molecular and genetic analysis of the transferred DNA regions of the root-inducing plasmid of *Agrobacterium rhizogenes*. *J. Bacteriol.* **164**, 33–44.
- Yin Z., Plader W., and Malepszy S. (2004), Transgene inheritance in plants. *J. Appl. Genet.* **45**, 127–144.
- Zdravković-Korać S., Čalić D., Druart P. H., and Radojevic L. (2003), The horse chestnut lines harboring the *rol* genes. *Biol. Plant.* **47**, 487–491.