

Crystal structure of 5-methoxy-6-hydroxy-2,3-dihydro-7*H*-dibenzo[*de,h*]-quinolin-7-one, C₁₇H₁₃NO₃


E. Sobarzo-Sánchez^{*I}, B. K. Cassels^I, L. Castedo^{II} and L. Valencia-Matarranz^{III}

^I Universidad de Chile, Facultad de Ciencias, Departamento de Química, Casilla 653 Santiago, Chile

^{II} Universidad de Santiago de Compostela, Facultad de Química, Departamento de Química Orgánica y Unidad Asociada al C.S.I.C., ES-15706 Santiago de Compostela, Spain

^{III} Universidad de Santiago de Compostela, Facultad de Química, Departamento de Química Inorgánica, ES-15782 Santiago de Compostela, Spain

Received November 7, 2002, accepted and available on-line January 2, 2003; CCDC-No. 1267/967



Abstract

C₁₇H₁₃NO₃, orthorhombic, *P*2₁*nb* (No. 33), *a* = 8.7927(8) Å, *b* = 30.165(2) Å, *c* = 4.9136(6) Å, *V* = 1303.2 Å³, *Z* = 4, *R*_{gt}(*F*) = 0.053, *wR*_{ref}(*F*²) = 0.161, *T* = 293 K.

Source of material

To a solution of phthalaldehydic acid (23 g, 153 mmol) in toluene (150 mL) was added homoveratrylamine (25 mL, 153 mmol) and the mixture was refluxed with stirring under a Dean-Stark trap for 2 hours at 413 K. The reaction mixture was then treated with polyphosphoric acid (80 g) and kept at 373 K for 10 minutes. The red mixture was dissolved in 100 mL water, neutralized with NH₃ and extracted with chloroform (700 mL). The chloroform extract was dried over anhydrous sodium sulphate, concentrated to dryness, and the residue subjected to flash chromatography on silica gel, eluting with hexane-ethyl acetate 95:5 (v/v) to give 5-methoxy-6-hydroxy-2,3-dihydro-7*H*-dibenzo[*de,h*]quinolin-7-one (3.83 g, yield 9%) crystallized in MeOH as red needles.

Experimental details

The low ratio of *N*(*hkl*)_{gt}/*N*(*param*) of about 5 is due to the low quality of the crystal and to its partial descomposition.

Discussion

The 5-methoxy-6-hydroxy-2,3-dihydro-oxoisoporphine molecule is largely planar with two methylene carbons, C2 and C3, forming a torsion angle C6–N1–C2–C3 of 25.9(10)°. The methoxyl group carbon is coplanar with the aromatic ring with a C16–O18–C19 angle of 117.5(4)°. An intramolecular hydrogen bond is also observed *d*(O21...H20) = 1.824 Å since the bond *d*(H20–O20) is considerably shorter (0.820 Å).

Table 1. Data collection and handling.

Crystal:	red prism, size 0.12 × 0.16 × 0.56 mm
Wavelength:	Cu K α radiation (1.54184 Å)
μ :	8.05 cm ⁻¹
Diffractometer, scan mode:	Enraf Nonius CAD4, $\omega/2\theta$
2 θ _{max} :	146.8°
<i>N</i> (<i>hkl</i>) _{measured} , <i>N</i> (<i>hkl</i>) _{unique} :	1493, 1472
Criterion for <i>I</i> _{obs} , <i>N</i> (<i>hkl</i>) _{gt} :	<i>I</i> _{obs} > 2 σ (<i>I</i> _{obs}), 1093
<i>N</i> (<i>param</i>) _{refined} :	193
Programs:	SIR-97 [1], SHELXL-97 [2], ORTEP-3 [3], WinGX [4]

Table 2. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
H(2A)	4a	0.2483	0.5856	0.4957	0.120
H(2B)	4a	0.3582	0.5696	0.7246	0.120
H(3A)	4a	0.3147	0.6348	0.8385	0.103
H(3B)	4a	0.3096	0.6521	0.5385	0.103
H(11)	4a	1.0544	0.5850	−0.0316	0.066
H(12)	4a	0.9909	0.5268	−0.3196	0.074
H(13)	4a	0.7483	0.4970	−0.3005	0.071
H(14)	4a	0.5739	0.5240	0.0006	0.066
H(15)	4a	0.4926	0.6976	0.9669	0.067
H(19A)	4a	0.6042	0.7304	1.3022	0.109
H(19B)	4a	0.5892	0.7710	1.1048	0.109
H(19C)	4a	0.7131	0.7712	1.3343	0.109
H(20)	4a	1.0205	0.6820	0.6045	0.122

* Correspondence author (e-mail: esobarzo@icaro.dic.uchile.cl)

Table 3. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> ₁₁	<i>U</i> ₂₂	<i>U</i> ₃₃	<i>U</i> ₁₂	<i>U</i> ₁₃	<i>U</i> ₂₃
N(1)	4a	0.4574(4)	0.5726(1)	0.360(1)	0.045(2)	0.063(2)	0.101(4)	−0.010(2)	0.009(3)	−0.014(3)
C(2)	4a	0.3503(5)	0.5891(2)	0.568(2)	0.058(3)	0.092(4)	0.150(8)	−0.020(3)	0.028(4)	−0.031(5)
C(3)	4a	0.3639(5)	0.6327(2)	0.662(2)	0.045(2)	0.111(4)	0.100(5)	−0.010(3)	0.012(3)	−0.036(4)
C(4)	4a	0.5239(4)	0.6493(1)	0.690(1)	0.039(2)	0.063(2)	0.052(3)	0.001(2)	0.001(2)	−0.007(2)
C(5)	4a	0.6324(4)	0.6292(1)	0.521(1)	0.039(2)	0.042(2)	0.043(2)	−0.001(1)	−0.003(2)	0.008(2)
C(6)	4a	0.5878(4)	0.5914(1)	0.350(1)	0.039(2)	0.049(2)	0.048(2)	−0.001(2)	−0.006(2)	0.002(2)
C(7)	4a	0.7818(4)	0.6456(1)	0.5189(9)	0.039(2)	0.038(2)	0.041(2)	0.002(1)	−0.001(2)	0.001(2)
C(8)	4a	0.8949(4)	0.6268(1)	0.336(1)	0.044(2)	0.048(2)	0.051(3)	−0.002(2)	0.000(2)	0.006(2)
C(9)	4a	0.8504(4)	0.5902(1)	0.1522(9)	0.049(2)	0.040(2)	0.043(2)	0.006(2)	0.001(2)	0.004(2)
C(10)	4a	0.7034(4)	0.5726(1)	0.1618(9)	0.049(2)	0.040(2)	0.041(2)	0.003(2)	−0.004(2)	0.004(2)
C(11)	4a	0.9569(5)	0.5730(1)	−0.027(1)	0.056(2)	0.055(2)	0.054(3)	0.007(2)	0.014(2)	0.007(2)
C(12)	4a	0.9195(6)	0.5381(1)	−0.198(1)	0.080(3)	0.054(2)	0.052(3)	0.017(2)	0.014(3)	0.000(2)
C(13)	4a	0.7746(5)	0.5203(1)	−0.186(1)	0.079(3)	0.047(2)	0.051(3)	0.006(2)	−0.006(3)	−0.004(2)
C(14)	4a	0.6699(5)	0.5369(1)	−0.008(1)	0.062(2)	0.048(2)	0.054(3)	−0.001(2)	−0.008(3)	−0.001(2)
C(15)	4a	0.5656(5)	0.6842(1)	0.857(1)	0.056(2)	0.061(3)	0.052(3)	0.010(2)	0.004(3)	−0.004(2)
C(16)	4a	0.7129(5)	0.6994(1)	0.864(1)	0.055(2)	0.049(2)	0.047(2)	0.001(2)	−0.007(2)	−0.005(2)
C(17)	4a	0.8219(4)	0.6805(1)	0.692(1)	0.043(2)	0.050(2)	0.055(3)	−0.003(2)	0.000(2)	0.000(2)
O(18)	4a	0.7648(3)	0.7329(1)	1.0245(9)	0.073(2)	0.062(2)	0.066(2)	0.000(2)	0.001(2)	−0.019(2)
C(19)	4a	0.6592(6)	0.7530(2)	1.206(1)	0.092(4)	0.065(3)	0.060(3)	0.022(3)	−0.005(3)	−0.021(3)
O(20)	4a	0.9639(4)	0.6977(1)	0.696(1)	0.067(2)	0.087(2)	0.091(3)	−0.022(2)	0.006(2)	−0.027(2)
O(21)	4a	1.0283(3)	0.6412(1)	0.331(1)	0.045(2)	0.075(2)	0.091(3)	−0.010(1)	0.014(2)	−0.018(2)

Acknowledgments. E. S.-S. thanks Fundación Andes for a doctoral fellowship. This work was funded in part by FONDECYT Grant No 2010056. Intensity measurements were performed at the Unidade de Raios X, RIAIDT, University of Santiago de Compostela, Spain.

References

- Altomare, A.; Burla, M. C.; Camalli, M.; Carrozzini, B.; Cascarano, G. L.; Giacovazzo, C.; Guagliardi, A.; Moliterni, A. G. G.; Polidori, G.; Rizzi, R.: Sir97: A new tool for crystal structure determination and refinement. *J. Appl. Crystallogr.* **32** (1999) 115-119.
- Sheldrick, G. M.: SHELXL-97. Program for the Refinement of Crystal Structures. University of Göttingen, Germany 1997.
- Farrugia, L. J.: ORTEP-3 for Windows - a version of ORTEP-III with Graphical User Interface (GUI). *J. Appl. Crystallogr.* **30** (1997) 565.
- Farrugia, L. J.: WinGX - Version 1.63.02. An integrated System of Windows Programs for the Solution, Refinement and Analysis of Single Crystals X-Ray Diffraction Data. *J. Appl. Crystallogr.* **32** (1999) 837-838.