


Crystal structure of 4,5-(ethoxyethylenedithio)-1,3-dithiole-2-thione, $C_7H_8OS_5$

G. Xue, W.-T. Yu and Q. Fang*

Shandong University, State Key Laboratory of Crystal Materials, Jinan, Shandong Province, 250100 P. R. China

Received May 2, 2003, accepted and available on-line July 29, 2003; CCDC-No. 1267/1082



Abstract

$C_7H_8OS_5$, monoclinic, $P12_1/c1$ (No. 14), $a = 8.3699(7)$ Å, $b = 17.573(2)$ Å, $c = 7.5305(7)$ Å, $\beta = 103.584(7)^\circ$, $V = 1076.6$ Å³, $Z = 4$, $R_{\text{gt}}(F) = 0.033$, $wR_{\text{ref}}(F^2) = 0.082$, $T = 293$ K.

Source of material

$(\text{Bu}_4\text{N})_2[\text{Zn}(\text{dmit})_2]$ was stirred with 1,2-dibromoethyl ethyl ether in excess acetone at room temperature. The orange solid was obtained, and it was extracted with benzene, washed with aqueous sodium hydrogencarbonate, and then with water. After drying over Na_2SO_4 , the benzene was evaporated, and the residue recrystallized from a mixed solvent of petroleum ether and chloroform (3:2) to give the title compound [1].

Discussion

Electronic donor BEDT-TTF [bis(ethylenedithio)-tetrathiafulvalene] and its variety of charge-transfer salts have attracted great interest for their electronic conductivity or superconductivity [2]. A replacement of the two ethylene units of BEDT-TTF with carbon-carbon double bonds can give rise to another attractive donor named as BVDT-TTF [bis(vinylenedithio)-tetrathiafulvalene] [3]. In the course of researching new BVDT-TTF based molecular electronic conductors, we obtained the crystal structure of title compound which serves as the precursor of BVDT-TTF.

In title compound, the terminal $\text{S1}-\text{C1}$ bond length of $1.647(3)$ Å is slightly longer than a typical $\text{S}=\text{C}$ double bond. The $\text{C2}-\text{C3}$ bond is a double bond with the length of $1.349(3)$ Å. The bond lengths of $\text{S4}-\text{C4}$ and $\text{S5}-\text{C5}$ to the methylene carbon atoms are $1.801(3)$ Å and $1.835(2)$ Å, respectively, showing a character-

istic of single $\text{S}-\text{C}$ bond. The remaining six $\text{S}-\text{C}$ bonds are in the range of $1.720(3)$ Å – $1.754(2)$ Å which are between typical single $\text{S}-\text{C}$ and double $\text{S}=\text{C}$ bonds. Thus, the 4,5-dimercapto-1,3-dithiole-2-thione (dmit) moiety in title compound is highly conjugated, which is similar to the case of other dmit derivatives [4–6]. Least-squares planes calculations show that the planarity of dmit moiety in the title compound is quite good. There are many $\text{S}\cdots\text{O}$, $\text{S}\cdots\text{C}$ interactions between neighboring molecules. The distances of $\text{S3}\cdots\text{O1}$, $\text{S4}\cdots\text{O1}$ and $\text{S3}\cdots\text{C2}$ are $3.320(2)$ Å, $3.182(2)$ Å and $3.464(2)$ Å, respectively. Above intermolecular distances between the molecules are shorter than the sum of the van der Waals radii, so the crystal packing is considerably stabilized.

Table 1. Data collection and handling.

Crystal:	pale-yellow plate, size $0.08 \times 0.24 \times 0.4$ mm
Wavelength:	Mo $K\alpha$ radiation (0.71073 Å)
μ :	10.32 cm^{-1}
Diffractionmeter, scan mode:	Bruker P4, ω
$2\theta_{\text{max}}$:	54.98°
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$:	3202, 2465
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2\sigma(I_{\text{obs}})$, 1834
$N(\text{param})_{\text{refined}}$:	119
Program:	SHELXTL [7]

Table 2. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	x	y	z	U_{iso}
H(4A)	4e	0.4932	0.5686	0.3484	0.052
H(4B)	4e	0.5241	0.5390	0.2173	0.052
H(5A)	4e	0.7623	0.5700	0.2951	0.048
H(6A)	4e	0.9271	0.5615	0.0918	0.064
H(6B)	4e	0.8931	0.6490	0.0606	0.064
H(7A)	4e	0.9730	0.5976	−0.1882	0.082
H(7B)	4e	0.7916	0.6270	−0.2500	0.082
H(7C)	4e	0.8265	0.5397	−0.2185	0.082

* Correspondence author (e-mail: fangqi@icm.sdu.edu.cn)

Table 3. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	x	y	z	U ₁₁	U ₂₂	U ₃₃	U ₁₂	U ₁₃	U ₂₃
S(1)	4e	0.1517(1)	0.90920(4)	−0.0029(1)	0.0611(5)	0.0498(4)	0.0610(5)	0.0121(3)	0.0122(4)	0.0051(4)
S(2)	4e	0.17686(8)	0.73991(4)	−0.01138(9)	0.0376(3)	0.0460(4)	0.0449(4)	−0.0038(3)	0.0089(3)	−0.0032(3)
S(3)	4e	0.45681(8)	0.82697(3)	0.17200(9)	0.0437(3)	0.0368(3)	0.0441(4)	−0.0042(3)	0.0112(3)	−0.0061(3)
S(4)	4e	0.33451(8)	0.58917(4)	0.0582(1)	0.0405(3)	0.0368(3)	0.0551(4)	−0.0086(3)	0.0154(3)	−0.0047(3)
S(5)	4e	0.68031(8)	0.69669(4)	0.2883(1)	0.0398(3)	0.0438(4)	0.0508(4)	−0.0028(3)	0.0047(3)	−0.0075(3)
O(1)	4e	0.6899(2)	0.58769(9)	0.0349(2)	0.0324(8)	0.047(1)	0.043(1)	−0.0027(7)	0.0120(7)	−0.0025(8)
C(1)	4e	0.2560(3)	0.8295(1)	0.0485(3)	0.044(1)	0.043(1)	0.036(1)	−0.001(1)	0.014(1)	0.000(1)
C(2)	4e	0.3518(3)	0.6873(1)	0.0891(3)	0.039(1)	0.037(1)	0.039(1)	−0.004(1)	0.014(1)	−0.002(1)
C(3)	4e	0.4842(3)	0.7282(1)	0.1743(3)	0.038(1)	0.038(1)	0.037(1)	−0.001(1)	0.013(1)	−0.002(1)
C(4)	4e	0.5119(3)	0.5585(1)	0.2283(3)	0.050(1)	0.038(1)	0.046(2)	−0.003(1)	0.018(1)	0.003(1)
C(5)	4e	0.6716(3)	0.5971(1)	0.2136(3)	0.042(1)	0.036(1)	0.041(1)	0.000(1)	0.009(1)	0.003(1)
C(6)	4e	0.8554(3)	0.5986(2)	0.0175(4)	0.033(1)	0.067(2)	0.063(2)	−0.006(1)	0.015(1)	−0.007(2)
C(7)	4e	0.8622(3)	0.5900(2)	−0.1770(4)	0.043(1)	0.062(2)	0.066(2)	0.000(1)	0.024(1)	−0.002(2)

Acknowledgments. This work was supported by the National Natural Science Foundation of China (Grant No. 20172034) and by a grant for the State Key Program of China.

References

1. Nakamura, T.; Nogami, T.; Shirota, Y.: Syntheses of new multi-sulfur 1,2-dithiolene complexes. Ni- and Pd-complexes of 1,4-dithiin-2,3-dithiolate. *Bull. Chem. Soc. Jpn.* **60** (1987) 3447-3449.
2. Williams, J. M.; Ferraro, J. R.; Thorn, R.J.; Carlson, K. D.; Geiser, U.; Wang, H. H.; Kini, A. M.; Whangbo, M.-H.: *Organic superconductors (including Fullerenes)*. Prentice Hall, New Jersey (1992).
3. Nakamura, T.; Iwasaka, S.-I.; Nakano, H.; Inoue, K.; Nogami, T.; Mikawa, H.: Syntheses of bis(vinylenedithio)tetrathiafulvalene and bis(methylvinylenedithio)tetrathiafulvalene. *Bull. Chem. Soc. Jpn.* **60** (1987) 365-368.
4. Fang, Q.; Yu, W.-T.; Xu, J.-H.; Lei, H.; Jiang, M.-H.: 5,6-Dihydro-5,6-diphenyl-1,3-dithiol[4,5-*b*]-[1,4]dithiine-2-thione (DHPT-DTT), C₁₇H₁₂S₅. *Acta Crystallogr. C* **50** (1994) 1519-1522.
5. Drew, M. G. B.; Kisenyi, J. M.; Parish, R. V.: Complexes of 2,5,7,9-tetra-thiabicyclo[4.3.0]non-1(6)-ene-8-thione(L¹). Part 1. Synthesis and antimony-121 mossbauer spectroscopic characterisation of SbCl₃(L¹)_{1.5} crystal and molecular structures of SbCl₃(L¹)_{1.5} and uncomplexed L¹. *J. Chem. Soc., Dalton Trans.* (1987) 1605-1609.
6. Fang, Q.; Jiang, M.-H.; Qu, Z.; Cai, J.-H.; Lei, H.; Yu, W.-T.; Zhuo, Z.: Non-linear optical properties of DMIT derivatives. *J. Mater. Chem.* **4** (1994) 1041-1045.
7. Bruker: SHELXTL. Structure Determination Programs. Version 5.10, Bruker AXS, Inc., Madison, Wisconsin, USA 1997.