

Crystal structure of bis(2,6-diaminopyridinium) diaqua-bis-(2,6-pyridinedicarboxylato)-bis(2,6-pyridinedicarboxylato)-dibismuthate(III) tetrahydrate, (C₂₈H₁₆O₁₈N₄Bi₂)(C₅H₈N₃)₂ · 4H₂O


M. Ranjbar^I, H. Aghabozorg^{*,II} and A. Moghimi^{III}

¹ Iranian Research Organization for Science and Technology (IROST), Department of Chemical Industries, Tehran, Iran

II Teacher Training University, Department of Chemistry, 49 Mofateh Avenue, 15614 Tehran, Iran

III University of Imam Hossein, Department of Chemistry, Tehran, Iran

Received June 7, 2003, accepted in revised form and available on-line October 7, 2003; CCDC-No. 1267/1099



Abstract

$C_{38}H_{40}Bi_2N_{10}O_{22}$, triclinic, $P\bar{1}$ (No. 2), $a = 9.699(1)$ Å, $b = 10.500(1)$ Å, $c = 10.890(1)$ Å, $\alpha = 85.587(3)^\circ$, $\beta = 88.128(3)^\circ$, $\gamma = 84.257(3)^\circ$, $V = 1099.9$ Å³, $Z = 1$, $R_{\text{gt}}(F) = 0.028$, $wR_{\text{ref}}(F^2) = 0.061$, $T = 295$ K.

Source of material

The title compound was synthesised by the reaction between ligand LH_2 , $[\text{H}_2\text{pyda}]^{2+}[\text{pydc}]^{2-}$, (pyda = 2,6-pyridinediamine and H_2pydc = 2,6-pyridinedicarboxylic acid) [1], and bismuth subnitrate $[\text{4BiNO}_3(\text{OH})_2 \cdot \text{BiOOH}]$ in aqueous solution. After two weeks, colorless crystals were isolated.

Experimental details

There is a positive residual density of $1.31 \text{ e} \cdot \text{\AA}^{-3}$ near the N5 center (distance $0.36 \text{ \AA}$), which can be the result of several factors that contribute to the bias of the data. Also, a refinement of a model considering a possible disordering of N5 failed. This together with the relative small displacement factor of N5 do not support the hypothesis that the residual electron density near N5 is the consequence of disordering.

Discussion

Bismuth complexes are of interest in the treatment of gastric ulcer [2], and may also be useful for the treatment of other diseases. Bi(III) exhibits a highly variable coordination number (three-ten)

and often an irregular coordination geometry [3]. The coordination chemistry of Bi(III) with carboxylates and aminocarboxylates is dominated by intermolecular interactions which leads to polymeric structures [3]. Continuing the synthesis of a novel self-assembling pyridine containing system ligand LH₂ [1], we become interested in investigating the influence of the counter ion [H₂pyda]²⁺ in the complexation process and succeeded in reporting some of them [4-6]. Here we report the crystal structure of a novel Bi(III) complex. The molecular structure consists of two metal fragments linked via the central four-membered Bi₂O₂ ring. The five Bi—O bonds are in the range 2.327(3) Å to 2.589(3) Å and the two Bi—N1 and Bi—N2 distances are 2.435(3) Å and 2.555(3) Å, respectively. It is interesting to note that, the existence of a stereochemically active lone pair on the Bi(III) atoms. These distances are in agreement with a previous report [7]. There is a significant difference between the N—C bonds in the 2,6-diaminopyridinium cation. However, such a difference is not uncommon in published structures containing this cationic fragment in the presence of heavy ions [8]. Discussing the self-assembled binuclear complex of Bi(III), the extensive intermolecular hydrogen bonding between coordinated and uncoordinated water molecules, two [Hpyda]⁺ and one [Bi₂(pydc)₄(H₂O)₂]²⁻ species has to be mentioned. The intermolecular forces existing between anionic and cationic units in the Bi(III) complex consist of both H-bonding and ion-pairing.

* Correspondence author (e-mail: haghbozorg@yahoo.com)

Table 1. Data collection and handling.

Crystal:	colorless prism, size 0.17 × 0.24 × 0.37 mm
Wavelength:	Mo <i>K</i> _α radiation (0.71073 Å)
μ :	80.89 cm ⁻¹
Diffractometer, scan mode:	Bruker SMART 1000 CCD, ϕ/ω
$2\theta_{\max}$:	60.16°
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$:	24752, 6425
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 5386
$N(\text{param})_{\text{refined}}$:	331
Programs:	SHELXTL [9], SADABS [10]

Table 2. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
H(9A)	2i	0.6268	0.6808	0.3959	0.06
H(9B)	2i	0.5878	0.6135	0.3000	0.06
H(10A)	2i	0.7895	0.6069	0.6247	0.06
H(10B)	2i	0.6677	0.6197	0.5772	0.06
H(11A)	2i	0.3748	0.5470	0.2514	0.06
H(11B)	2i	0.3786	0.5969	0.1471	0.06
H(3N)	2i	0.3574	0.1206	0.2271	0.06
H(4A)	2i	0.2908	-0.0731	0.2692	0.06
H(4B)	2i	0.3798	-0.1912	0.2392	0.06
H(5A)	2i	0.4040	0.3391	0.1853	0.06
H(5B)	2i	0.5377	0.3670	0.1227	0.06
H(2)	2i	0.6153	-0.0015	0.4529	0.06
H(3)	2i	0.8126	-0.1337	0.3971	0.06
H(4)	2i	1.0134	-0.0430	0.3356	0.06
H(9)	2i	1.1049	0.8211	0.1260	0.06
H(10)	2i	1.0054	0.8080	-0.0612	0.06
H(11)	2i	0.8508	0.6540	-0.0817	0.06
H(16)	2i	0.6106	-0.1709	0.1347	0.06
H(17)	2i	0.7639	-0.0292	0.0593	0.06
H(18)	2i	0.7086	0.1900	0.0643	0.06

Table 3. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> ₁₁	<i>U</i> ₂₂	<i>U</i> ₃₃	<i>U</i> ₁₂	<i>U</i> ₁₃	<i>U</i> ₂₃
Bi(1)	2i	0.83147(1)	0.44620(1)	0.38360(1)	0.02483(7)	0.02614(7)	0.02813(7)	-0.00382(5)	-0.00108(5)	-0.00110(5)
O(1)	2i	1.0432(2)	0.3312(2)	0.3095(2)	0.027(1)	0.033(1)	0.037(1)	-0.008(1)	0.006(1)	-0.002(1)
O(2)	2i	1.1662(3)	0.1444(3)	0.2828(3)	0.025(1)	0.043(2)	0.067(2)	0.006(1)	0.016(1)	-0.001(1)
O(3)	2i	0.6059(2)	0.3772(2)	0.4415(2)	0.026(1)	0.028(1)	0.055(2)	-0.002(1)	0.006(1)	-0.004(1)
O(4)	2i	0.4694(3)	0.2198(3)	0.4753(3)	0.026(1)	0.040(2)	0.067(2)	-0.007(1)	0.013(1)	-0.007(1)
O(5)	2i	1.0135(3)	0.6120(2)	0.4273(2)	0.038(1)	0.032(1)	0.026(1)	-0.011(1)	-0.004(1)	0.006(1)
O(6)	2i	1.1546(3)	0.7558(3)	0.3537(2)	0.043(2)	0.044(2)	0.038(2)	-0.021(1)	-0.002(1)	-0.001(1)
O(7)	2i	0.7465(3)	0.4116(2)	0.1925(2)	0.045(2)	0.035(1)	0.030(1)	-0.013(1)	-0.008(1)	-0.002(1)
O(8)	2i	0.7090(3)	0.4657(3)	-0.0043(3)	0.071(2)	0.060(2)	0.041(2)	-0.016(2)	-0.029(2)	-0.004(1)
O(9)	2i	0.6557(3)	0.6433(2)	0.3322(2)	0.032(1)	0.039(1)	0.041(2)	0.004(1)	-0.001(1)	0.000(1)
O(10)	2i	0.7217(3)	0.5621(3)	0.6150(3)	0.039(2)	0.050(2)	0.088(3)	-0.004(1)	-0.018(2)	0.002(2)
O(11)	2i	0.4146(3)	0.6084(3)	0.2151(3)	0.052(2)	0.058(2)	0.050(2)	-0.010(2)	-0.013(1)	-0.010(2)
N(1)	2i	0.8210(3)	0.2155(2)	0.3806(2)	0.022(1)	0.025(1)	0.023(1)	0.002(1)	0.000(1)	-0.002(1)
N(2)	2i	0.9202(3)	0.5891(3)	0.2037(2)	0.027(2)	0.027(1)	0.024(1)	0.001(1)	-0.002(1)	-0.003(1)
N(3)	2i	0.4349(3)	0.0924(3)	0.1941(3)	0.023(2)	0.043(2)	0.041(2)	-0.003(1)	0.005(1)	-0.011(1)
N(4)	2i	0.3666(4)	-0.1087(4)	0.2386(4)	0.049(2)	0.050(2)	0.068(3)	-0.020(2)	0.014(2)	-0.003(2)
N(5)	2i	0.4828(3)	0.3141(3)	0.1522(3)	0.020(2)	0.049(2)	0.036(2)	0.022(1)	0.003(1)	-0.008(1)
C(1)	2i	0.7042(3)	0.1646(3)	0.4159(3)	0.026(2)	0.028(2)	0.025(2)	-0.004(1)	-0.000(1)	-0.001(1)
C(2)	2i	0.6965(4)	0.0331(3)	0.4259(3)	0.026(2)	0.031(2)	0.040(2)	-0.005(2)	0.002(2)	-0.001(2)
C(3)	2i	0.8147(4)	-0.0450(3)	0.3939(4)	0.040(2)	0.024(2)	0.056(3)	-0.005(2)	0.001(2)	-0.002(2)
C(4)	2i	0.9340(4)	0.0088(3)	0.3563(4)	0.029(2)	0.024(2)	0.047(2)	0.004(1)	0.003(2)	-0.003(2)
C(5)	2i	0.9340(3)	0.1403(3)	0.3507(3)	0.024(2)	0.026(2)	0.029(2)	-0.001(1)	-0.001(1)	0.000(1)
C(6)	2i	1.0587(4)	0.2096(3)	0.3116(3)	0.028(2)	0.033(2)	0.028(2)	-0.000(2)	0.003(1)	-0.004(1)
C(7)	2i	0.5841(3)	0.2598(3)	0.4472(3)	0.022(2)	0.033(2)	0.031(2)	-0.002(1)	0.001(1)	-0.000(2)
C(8)	2i	1.0075(3)	0.6778(3)	0.2158(3)	0.021(2)	0.026(2)	0.028(2)	0.002(1)	0.002(1)	0.002(1)
C(9)	2i	1.0426(4)	0.7612(4)	0.1163(3)	0.043(2)	0.034(2)	0.035(2)	-0.009(2)	0.003(2)	0.005(2)
C(10)	2i	0.9840(4)	0.7527(4)	0.0054(3)	0.056(3)	0.048(2)	0.026(2)	-0.010(2)	0.005(2)	0.009(2)
C(11)	2i	0.8920(4)	0.6618(4)	-0.0074(3)	0.046(2)	0.047(2)	0.025(2)	-0.005(2)	-0.004(2)	0.003(2)
C(12)	2i	0.8633(4)	0.5814(3)	0.0959(3)	0.032(2)	0.035(2)	0.026(2)	0.001(2)	-0.003(2)	-0.003(2)
C(13)	2i	0.7649(4)	0.4790(3)	0.0924(3)	0.038(2)	0.031(2)	0.035(2)	-0.000(2)	-0.008(2)	-0.007(2)
C(14)	2i	1.0650(3)	0.6830(3)	0.3414(3)	0.026(2)	0.025(2)	0.027(2)	-0.001(1)	0.001(1)	-0.000(1)
C(15)	2i	0.4641(4)	-0.0367(3)	0.1908(3)	0.036(2)	0.035(2)	0.032(2)	-0.010(2)	-0.002(2)	-0.002(2)
C(16)	2i	0.5884(4)	-0.0830(4)	0.1384(4)	0.036(2)	0.032(2)	0.051(2)	0.006(2)	-0.004(2)	-0.000(2)
C(17)	2i	0.6788(4)	0.0013(4)	0.0930(4)	0.030(2)	0.053(2)	0.050(2)	-0.002(2)	0.011(2)	-0.009(2)
C(18)	2i	0.6466(4)	0.1333(4)	0.0964(4)	0.027(2)	0.046(2)	0.045(2)	-0.012(2)	0.004(2)	-0.004(2)
C(19)	2i	0.5209(4)	0.1792(4)	0.1473(3)	0.032(2)	0.036(2)	0.040(2)	-0.008(2)	0.000(2)	-0.005(2)

Acknowledgments. This work was supported by the Teacher Training University. We are particularly grateful to G. V. Grintselev-Knyazev from General and Technical Chemistry Division Academy of Sciences of Russia for X-ray crystallographic data.

References

1. Moghimi, A.; Ranjbar, M.; Aghabozorg, H.; Jalali, F.; Shamsipur, M.; Yap, G. P. A.; Rahbarnoohi, H.: A novel pyridine containing self-assembling system: Synthesis, Characterization, X-ray Crystal Structure, ¹³C Solid Phase NMR and Solution Studies. *J. Mol. Struct.* **605** (2002) 133-149.
2. Sadler, P. J.; Li, H.; Sun, H.: Coordination Chemistry of metals in medicine. *Coord. Chem. Rev.* **185-186** (1999) 689-709.
3. Mahjoub, A. R.; Morsali, A.: Refinement of the crystal structure of *trans*-bis(1,10-phenanthroline)bismuth(III) tribromide, C₂₄H₁₆BiBr₃N₄. *Z. Kristallogr. NCS* **217** (2002) 513-514.
4. Ranjbar, M.; Aghabozorg, H.; Moghimi, A.; Yanovsky, A.: Crystal Structure of 2,6-diaminopyridinium bis(2,6-pyridinedicarboxylato) chromium(III) 2,6-pyridinedicarboxylic acid hemihydrate, C₂₆H₁₈CrN₆O₁₂ · 0.5H₂O. *Z. Kristallogr. NCS* **216** (2001) 626-628.
5. Ranjbar, M.; Aghabozorg, H.; Moghimi, A.: Crystal structure of bis(2,6-diaminopyridinium) tetrachloropalladate(II), (C₅N₃H₈)₂PdCl₄. *Z. Kristallogr. NCS* **218** (2003) 75-76.
6. Ranjbar, M.; Taghavipur, M.; Aghabozorg, H.; Moghimi, A.; Jalali, F.; Shamsipur, M.: Synthesis, X-ray Crystal Structure and Solution Studies of Cu(II) Complexes of a Pyridine Containing Self-assembling System. *Pol. J. Chem.* **76** (2002) 785-794.
7. Ranjbar, M.; Aghabozorg, H.; Moghimi, A.; Yanovsky, A.: Crystal Structure of Bi(III) Complex of a Pyridine Containing Self-assembling System. *Anal. Sci.* **17** (2001) 4164-4165.
8. Liu, S. H.; Chen, J. D.; Liou, L. S.; Wang, J. C.: Synthesis and crystal structures of (C₅H₇N₃Br)₃CuBr₄ and (C₅H₈N₃)CuBr₂: An isolated tetrahedral CuBr₄³⁻ anion. *Inorg. Chem.* **40** (2001) 6499-6501.
9. Sheldrick, G. M.: SHELXTL v. 5.10, Structure Determination Software Suite, Bruker AXS, Madison, Wisconsin, USA 1998.
10. Sheldrick, G. M.: SADABS v. 2.01, Bruker/Siemens Area Detector Absorption Correction Program, Bruker AXS, Madison, Wisconsin, USA 1998.