


Crystal structure of *trans*-diaqua-di(bis(pyrazolylmethane)copper(II) dinitrate, $\text{Cu}(\text{C}_7\text{H}_{10}\text{N}_4)_2(\text{OH}_2)_2(\text{NO}_3)_2$

K. A. Van Langenberg^I, B. Moubaraki^I, K. S. Murray^I and E. R. T. Tiekink^{*,II,I}

^I Monash University, School of Chemistry, PO Box 23, Clayton, Victoria 3800, Australia

^{II} The University of Adelaide, Department of Chemistry, Australia 5005

Received May 14, 2003, accepted and available on-line July 29, 2003; CCDC-No. 1267/1088



295 K), which were compatible with a mononuclear Cu(II) structure; and by IR spectroscopy which showed the characteristic bands of the ligand and anion.

Discussion

The Cu atom in the cation lies on a crystallographic centre of inversion and exists within a N_4O_2 donor set defined by two chelating bpm ligands and two oxygen atoms derived from two water molecules. The coordination is axially-distorted octahedral with the O atoms in axial positions. Key geometric parameters: $d(\text{Cu}—\text{N}11) = 1.994(2) \text{ \AA}$, $d(\text{Cu}—\text{N}21) = 2.028(2) \text{ \AA}$, $d(\text{Cu}—\text{O}1) = 2.471(2) \text{ \AA}$ and a chelate angle of $88.90(9)^\circ$. Each water molecule forms two hydrogen bonds, one each to two nitrate anions that, in turn are linked to a centrosymmetrically related water molecule so that twelve-membered $[\text{H}—\text{O}—\text{H} \cdots \text{O}—\text{N}(\text{O})—\text{O} \cdots]_2$ rings are formed as highlighted in the lower view of the Figure. Thus, the complex cations and anions are linked into chains via hydrogen bonding contacts. The parameters associated with the hydrogen bonding are $d(\text{O}1—\text{H} \cdots \text{O}2) = 2.02 \text{ \AA}$, $d(\text{O}1 \cdots \text{O}2) = 2.871(5) \text{ \AA}$ and the angle at H is 168° , and $d(\text{O}1—\text{H} \cdots \text{O}4^i) = 2.06 \text{ \AA}$, $d(\text{O}1 \cdots \text{O}4^i) = 2.839(5) \text{ \AA}$ and angle of 149° ; $i: -x, -y, -1-z$.

Table 1. Data collection and handling.

Crystal:	blue block, size $0.16 \times 0.19 \times 0.40 \text{ mm}$
Wavelength:	Mo K_α radiation ($0.71073 \text{ \AA}$)
μ :	11.09 cm^{-1}
Diffractometer, scan mode:	Rigaku AFC6R, $\omega/2\theta$
$2\theta_{\text{max}}$:	55.2°
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$:	2567, 2406
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 2088
$N(\text{param})_{\text{refined}}$:	152
Programs:	teXsan [2], SAPI91 [3], SHELXL-97 [4], DIFABS [5], PLATON [6], ORTEP [7]

Abstract

$\text{C}_{14}\text{H}_{20}\text{CuN}_{10}\text{O}_8$, triclinic, $P\bar{1}$ (No. 2), $a = 8.498(1) \text{ \AA}$, $b = 9.167(2) \text{ \AA}$, $c = 8.323(3) \text{ \AA}$, $\alpha = 103.41(3)^\circ$, $\beta = 101.87(3)^\circ$, $\gamma = 117.10(1)^\circ$, $V = 523.9 \text{ \AA}^3$, $Z = 1$, $R_{\text{gt}}(F) = 0.041$, $wR_{\text{ref}}(F^2) = 0.122$, $T = 293 \text{ K}$.

Source of material

The complex was used as a precursor to prepare mixed $(\text{bpm})_2\text{Cu}(\text{II})$ -hexacyanoiron(III)-bridged molecule-based magnetic materials for comparison to other $\text{Ni}(\text{II})(\text{bpm})_2\text{-Fe}(\text{CN})_6$ systems [1]. The complex was characterized by magnetic moment data ($1.88 \mu\text{B}$ at

Table 2. Atomic coordinates and displacement parameters (in $\text{\AA}^2$).

Atom	Site	x	y	z	U_{iso}
H(1o)	2i	0.0741	0.1283	-0.2893	0.070
H(2o)	2i	0.0732	-0.0288	-0.3518	0.070
H(1a)	2i	-0.4898	-0.1680	-0.4138	0.070
H(1b)	2i	-0.2731	-0.0925	-0.3755	0.070
H(11)	2i	-0.0652	0.3112	0.1879	0.056
H(12)	2i	-0.3484	0.3217	0.0563	0.056
H(13)	2i	-0.5360	0.0709	-0.2328	0.056
H(21)	2i	-0.2155	-0.4159	-0.0420	0.056
H(22)	2i	-0.5539	-0.6329	-0.2310	0.056
H(23)	2i	-0.6575	-0.4801	-0.3966	0.056

* Correspondence author (e-mail: chmert@nus.edu.sg)

^I Current address: The National University of Singapore, Department of Chemistry, Singapore 117543

Table 3. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	x	y	z	U ₁₁	U ₂₂	U ₃₃	U ₁₂	U ₁₃	U ₂₃
Cu	1a	0	0	0	0.0333(2)	0.0310(2)	0.0363(3)	0.0166(2)	0.0113(2)	0.0109(2)
O(1)	2i	0.0813(4)	0.0468(3)	-0.2597(3)	0.095(2)	0.081(2)	0.051(1)	0.062(2)	0.046(1)	0.039(1)
O(2)	2i	0.0551(4)	0.3358(4)	-0.3092(5)	0.053(1)	0.075(2)	0.110(3)	0.020(1)	-0.007(2)	0.038(2)
O(3)	2i	-0.1683(5)	0.3787(5)	-0.4004(5)	0.102(3)	0.091(2)	0.124(3)	0.069(2)	-0.003(2)	0.003(2)
O(4)	2i	-0.2040(4)	0.1316(4)	-0.5204(4)	0.072(2)	0.066(2)	0.064(2)	0.027(1)	0.015(1)	0.007(1)
N(1)	2i	-0.1018(4)	0.2847(3)	-0.4109(3)	0.048(1)	0.047(1)	0.047(1)	0.021(1)	0.012(1)	0.017(1)
N(11)	2i	-0.1627(3)	0.1010(3)	-0.0337(3)	0.039(1)	0.038(1)	0.032(1)	0.0214(9)	0.0128(8)	0.0110(8)
N(12)	2i	-0.3243(3)	0.0213(3)	-0.1751(3)	0.038(1)	0.038(1)	0.035(1)	0.0212(9)	0.0143(8)	0.0168(9)
N(21)	2i	-0.2322(3)	-0.2442(3)	-0.1469(3)	0.036(1)	0.035(1)	0.035(1)	0.0167(8)	0.0120(8)	0.0144(8)
N(22)	2i	-0.3855(3)	-0.2699(3)	-0.2677(3)	0.036(1)	0.036(1)	0.029(1)	0.0172(8)	0.0132(8)	0.0104(8)
C(1)	2i	-0.3706(4)	-0.1278(3)	-0.3240(3)	0.043(1)	0.040(1)	0.031(1)	0.020(1)	0.013(1)	0.016(1)
C(11)	2i	-0.1599(4)	0.2339(3)	0.0778(4)	0.052(2)	0.039(1)	0.040(1)	0.025(1)	0.022(1)	0.014(1)
C(12)	2i	-0.3185(4)	0.2405(4)	0.0061(4)	0.059(2)	0.047(1)	0.056(2)	0.035(1)	0.034(1)	0.025(1)
C(13)	2i	-0.4207(4)	0.1028(4)	-0.1534(4)	0.046(1)	0.051(2)	0.056(2)	0.032(1)	0.023(1)	0.030(1)
C(21)	2i	-0.2909(4)	-0.3952(3)	-0.1199(4)	0.047(1)	0.037(1)	0.041(1)	0.021(1)	0.017(1)	0.018(1)
C(22)	2i	-0.4808(4)	-0.5178(4)	-0.2247(4)	0.045(1)	0.036(1)	0.048(2)	0.016(1)	0.023(1)	0.017(1)
C(23)	2i	-0.5371(4)	-0.4335(3)	-0.3160(3)	0.037(1)	0.039(1)	0.035(1)	0.015(1)	0.016(1)	0.009(1)

Acknowledgment. The Australian Research Council is thanked for support.

References

1. Van Langenberg, K.; Batten, S. R.; Berry, K. J.; Hockless, D. C. R.; Moubaraki, B.; Murray, K. S.: Structure and magnetism of a bimetallic pentanuclear cluster [(Ni(bpm)₂)₃(Fe(CN)₆)₂]·7H₂O (bpm = Bis(1-pyrazolyl)methane). The role of the hydrogen-bonded 7H₂O "cluster" in long-range magnetic ordering. *Inorg. Chem.* **36** (1997) 5006-5015.
2. teXsan: Single Crystal Structure Analysis Software. Version 1.04. Molecular Structure Corporation. The Woodlands, TX, USA, 1997.
3. Fan H.-F.: SAPI91. Structure Analysis Programs with Intelligent Control, Rigaku Corporation, Tokyo, Japan 1991.
4. Sheldrick, G. M.: SHELXL-97. Program for crystal structure refinement. University of Göttingen, Germany 1997.
5. Walker, N.; Stuart, D.: An empirical method for correcting diffractometer data for absorption effects. *Acta Crystallogr.* **A39** (1983) 158-166.
6. Spek, T.: PLATON. A Multipurpose Crystallographic Tool, Utrecht University, Utrecht, The Netherlands 2000.
7. Johnson, C.K.: ORTEP. Report ORNL-5138, Oak Ridge National Laboratory, Tennessee, USA 1976.